

Chapter 7: One-factor Multi-sample Experiments

Introductory Statistics for Engineering Experimentation

Peter R. Nelson, Marie Coffin and Karen A.F. Copeland

Slides by Douglas Bates

Outline

7.1 Basic Inference

7.5 Analysis of variance

Multiple comparisons

Outline

7.1 Basic Inference

7.5 Analysis of variance

Multiple comparisons

Outline

7.1 Basic Inference

7.5 Analysis of variance

Multiple comparisons

Factors with more than two levels

- In the previous chapter we discussed inference for comparative experiments on two groups.
- In chapter 3 we showed how to model a continuous response as a function of one or more factors that could have multiple levels. We used the *R* function `aov` to estimate the cell means or the “effects” of the levels of each factor.
- The statistical assessment of whether or not the effects are significant is usually based on an *analysis of variance*; hence the name `aov` for the model-fitting function and the name `anova` for the extractor function that produces the analysis of variance table.
- The text book emphasizes a technique called the analysis of means (ANOM). This is not a widely-used technique.
- We will focus on the analysis of variance and another approach called `multiple comparisons` for follow-up analysis.

Overview of techniques

- In chapter 2 we used graphical methods, such as comparative dotplots and comparative density plots, to display a continuous response as it depends on levels of a factor. In section 4.4 we also discussed normal probability plots, which can be used as comparative plots for such data.
- In chapter 3 we discussed fitting models of the form

$$Y_{ij} = \mu_i + \epsilon_{ij}, \quad i = 1, \dots, I; \quad j = 1, \dots, n_i$$

or

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad i = 1, \dots, I; \quad j = 1, \dots, n_i$$

- These are the same model; just two different ways of writing it. The first is called the *cell means* form and the second is the *effects* form.
- We check for differences in the mean response in two stages: first we check if all the means could be equal and, if we reject this hypothesis, we check for which levels of the factor produce significantly different means.

R functions used in this chapter

- Preliminary plots are obtained with `dotplot`, `bwplot`, `densityplot` and `qqmath`, all in the comparative form. Model fits assume that the variances in the groups are more-or-less equal. Hence we check the plots for equal variances as well as equal means.
- We use `aov` to fit the model, `summary` or `anova` to obtain the analysis of variance table, and `model.tables` to obtain estimates of the cell means or the effects.
- If we reject the hypothesis $H_0 : \mu_1 = \mu_2 = \dots = \mu_I$ (or, equivalently, $H_0 : \alpha_1 = \alpha_2 = \dots = \alpha_I = 0$) then we use `TukeyHSD` to perform multiple comparisons using Tukey's Honest Significant Difference method.
- We assess residual plots obtained with `plot(fm, which = 1)` and `plot(fm, which = 2)`



Section 7.1, Basic Inference

- We begin by plotting the data, preferably with comparative dotplots or comparative normal probability plots. (The text uses an error-bar chart in figure 7.1 but these are less informative than those mentioned above.)
- Group means and s^2 , the mean square error, are evaluated by fitting an `aov` model and using `summary` or `anova`. The degrees of freedom for s^2 , $n_1 + n_2 + \cdots + n_I - I$, is given in the table.
- You could use critical values from a T_ν distribution to calculate confidence intervals on the individual means (p. 248) but the practice is discouraged.

Examples 7.1.1 and 7.1.2

```
> reac <-
+   data.frame(yield = c(3.3, 4.0, 4.7,
+                       4.0, 5.0, 4.3, 5.5,
+                       5.3, 6.5, 6.4, 7.0, 7.7),
+             cat = factor(rep(LETTERS[1:3], c(3,4,5))))
> summary(fm1 <- aov(yield ~ cat, reac))
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
cat	2	14.7012	7.3506	12.099	0.002813
Residuals	9	5.4680	0.6076		



Section 7.5 Analysis of Variance

- The F ratio quoted in the analysis of variance table is the ratio of $MS_{\text{treatment}}$ to the mean square for error, MS_e . It is a “signal-to-noise” ratio based on the differences between groups versus the differences within groups.
- Both the numerator and the denominator have degrees of freedom associated with them. In this case they are $I - 1$ (numerator) and $N - I$ (denominator) where N is the total number of observations ($N = n_1 + n_2 + \cdots + n_I$)
- The p-value is calculated from a theoretical distribution for this quantity, written F_{ν_1, ν_2} . The **R** functions for this distribution are **df**, **pf**, **qf** and **rf**.
- The hypothesis being tested is “are all the means the same?” versus “are there any differences?”. Symbolically $H_0 : \mu_1 = \mu_2 = \cdots = \mu_I$, in the cell means form, or $H_0 : \alpha_1 = \alpha_2 = \cdots = \alpha_I = 0$ in the effects form.

Multiple comparisons

- If we reject H_0 in the analysis of variance the natural follow-up question is “so which group means are significantly different”.
- It is tempting to use a series of t-tests to compare each pair of groups but doing so will inflate the probability of a false positive.
- There are several ways of controlling for this inflated false positive probability. We will use Tukey's Honest Significant Differences, **TukeyHSD**, which is preferred when the groups are of equal importance. (Other methods are used when we have, say, a control group that we wish to compare with each of several treatments.)

Example 7.1.1 cont'd

```
> TukeyHSD(fm1)
```

```
  Tukey multiple comparisons of means
```

```
    95% family-wise confidence level
```

```
Fit: aov(formula = yield ~ cat, data = reac)
```

```
$cat
```

	diff	lwr	upr	p adj
B-A	0.70	-0.9621403	2.362140	0.4954811
C-A	2.58	0.9906899	4.169310	0.0036432
C-B	1.88	0.4201254	3.339875	0.0144223