

# **lmer for SAS PROC MIXED Users**

Douglas Bates  
Department of Statistics  
University of Wisconsin – Madison  
Bates@wisc.edu

## **1 Introduction**

The `lmer` function from the `lme4` package for R is used to fit linear mixed-effects models. It is similar in scope to the SAS procedure PROC MIXED described in Littell et al. (1996).

A file on the SAS Institute web site (<http://www.sas.com>) contains all the data sets in the book and all the SAS programs used in Littell et al. (1996). We have converted the data sets from the tabular representation used for SAS PROC MIXED to the `data.frame` objects used by `lmer`. To help users familiar with SAS PROC MIXED get up to speed with `lmer` more quickly, we provide transcripts of some `lmer` analyses paralleling the SAS PROC MIXED analyses in Littell et al. (1996).

In this paper we highlight some of the similarities and differences of `lmer` analysis and SAS PROC MIXED analysis.

## **2 Similarities between lmer and SAS PROC MIXED**

Both SAS PROC MIXED and `lmer` can fit linear mixed-effects models expressed in the Laird-Ware formulation. For a single level of grouping Laird and Ware (1982) write the  $n_i$ -dimensional response vector  $\mathbf{y}_i$  for the  $i$ th experimental

unit as

$$\begin{aligned} \mathbf{y}_i &= \mathbf{X}_i\boldsymbol{\beta} + \mathbf{Z}_i\mathbf{b}_i + \boldsymbol{\epsilon}_i, \quad i = 1, \dots, M \\ \mathbf{b}_i &\sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}), \quad \boldsymbol{\epsilon}_i \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I}) \end{aligned} \tag{1}$$

where  $\boldsymbol{\beta}$  is the  $p$ -dimensional vector of *fixed effects*,  $\mathbf{b}_i$  is the  $q$ -dimensional vector of *random effects*,  $\mathbf{X}_i$  (of size  $n_i \times p$ ) and  $\mathbf{Z}_i$  (of size  $n_i \times q$ ) are known fixed-effects and random-effects regressor matrices, and  $\boldsymbol{\epsilon}_i$  is the  $n_i$ -dimensional *within-group error* vector with a spherical Gaussian distribution. The assumption  $\text{Var}(\boldsymbol{\epsilon}_i) = \sigma^2 \mathbf{I}$  can be relaxed using additional arguments in the model fitting.

The basic specification of the model requires a linear model expression for the fixed effects and a linear model expression for the random effects. In **SAS PROC MIXED** the fixed-effects part is specified in the `model` statement and the random-effects part in the `random` statement. In `lmer` the fixed effects and the random effects are both specified as terms in the `formula` argument to `lmer`.

Both **SAS PROC MIXED** and `lmer` allow a mixed-effects model to be fit by maximum likelihood (`method = ml` in SAS) or by maximum residual likelihood, sometimes also called restricted maximum likelihood or REML. This is the default criterion in **SAS PROC MIXED** and in `lmer`. To get ML estimates use the optional argument `REML=FALSE` in the call to `lmer`.

### 3 Important differences

The output from **PROC MIXED** typically includes values of the Akaike Information Criterion (AIC) and Schwartz’s Bayesian Criterion (SBC). These are used to compare different models fit to the same data. The output of the `summary` function applied to the object created by `lmer` also produces values of AIC and BIC but the definitions used in older versions of **PROC MIXED** are different from those used in more recent versions of **PROC MIXED** and in `lmer`. In `lmer` the definitions are such that “smaller is better”. In some older versions of **PROC MIXED** the definitions are such that “bigger is better”.

When models are fit by REML, the values of AIC, SBC (or BIC) and the log-likelihood can only be compared between models with exactly the same fixed-effects structure. When models are fit by maximum likelihood these criteria can be compared between any models fit to the same data. That is,

these quality-of-fit criteria can be used to evaluate different fixed-effects specifications or different random-effects specifications or different specifications of both fixed effects and random effects.

We encourage developing and testing the model using likelihood ratio tests or the AIC and BIC criteria. Once a form for both the random effects and the fixed effects has been determined, the model can be refit with `REML = TRUE` if the restricted estimates of the variance components are desired. Note that the `update` function provides a convenient way of refitting a model with changes to one or more arguments.

## 4 Data manipulation

Both PROC MIXED and `lmer` work with data in a tabular form with one row per observation. There are, however, important differences in the internal representations of variables in the data.

In SAS a qualitative factor can be stored either as numerical values or alphanumeric labels. When a factor stored as numerical values is used in PROC MIXED it is listed in the `class` statement to indicate that it is a factor. In S this information is stored with the data itself by converting the variable to a factor when it is first stored. If the factor represents an ordered set of levels, it should be converted to an `ordered` factor.

For example the SAS code

```
data animal;
  input trait animal y;
  datalines;
1 1 6
1 2 8
1 3 7
2 1 9
2 2 5
2 3 .
;
```

would require that the `trait` and `animal` variables be specified in a `class` statement in any model that is fit.

In R these data could be read from a file, say `animal.dat`, and converted to factors by

```
animal <- within(read.table("animal.dat", header = TRUE),
{
```

```

        trait <- factor(trait)
        animal <- factor(animal)
    })

```

In general it is a good idea to check the types of variables in a data frame before working with it. One way of doing this is to apply the function `data.class` to each variable in turn using the `sapply` function.

```

> sapply(Animal, data.class)
      Sire      Dam AvgDailyGain
"factor"  "factor"  "numeric"
> str(Animal)
'data.frame':      20 obs. of  3 variables:
 $ Sire      : Factor w/ 5 levels "1","2","3","4",...: 1 1 1 1 2 2 2 2 3 3 ...
 $ Dam       : Factor w/ 2 levels "1","2": 1 1 2 2 1 1 2 2 1 1 ...
 $ AvgDailyGain: num  2.24 1.85 2.05 2.41 1.99 1.93 2.72 2.32 2.33 2.68 ...
- attr(*, "ginfo")=List of 7
 ..$ formula      :Class 'formula' length 3 AvgDailyGain ~ 1 | Sire/Dam
 .. .. ..- attr(*, ".Environment")=<environment: R_GlobalEnv>
 ..$ order.groups:List of 2
 .. ..$ Sire: logi TRUE
 .. ..$ Dam : logi TRUE
 ..$ FUN          :function (x)
 ..$ outer        : NULL
 ..$ inner        : NULL
 ..$ labels       :List of 1
 .. ..$ AvgDailyGain: chr "Average Daily Weight Gain"
 ..$ units        : list()

```

## 4.1 Unique levels of factors

Designs with nested grouping factors are indicated differently in the two languages. An example of such an experimental design is the semiconductor experiment described in section 2.2 of Littell et al. (1996) where twelve wafers are assigned to four experimental treatments with three wafers per treatment. The levels for the wafer factor are 1, 2, and 3 but the wafer factor is only meaningful within the same level of the treatment factor, *et*. There is nothing associating wafer 1 of the third treatment group with wafer 1 of the first treatment group.

In SAS this nesting of factors is denoted by `wafer(et)`. In S the nesting is written with `ET/Wafer` and read “wafer within ET”. If both levels of

nested factors are to be associated with random effects then this is all you need to know. You would use an expression with a "/" in the grouping factor part of the formula in the call to `lmer` object. The random effects term would be either

```
(1 | ET:Wafer)
```

or, equivalently

```
(1 | ET:Wafer) + (1 | ET)
```

In this case, however, there would not usually be any random effects associated with the “experimental treatment” or `ET` factor. The only random effects are at the `Wafer` level. It is necessary to create a factor that will have unique levels for each `Wafer` within each level of `ET`. One way to do this is to assign

```
> Semiconductor <- within(Semiconductor, Grp <- factor(ET:Wafer))
```

after which we could specify a random effects term of `(1 | Grp)`. Alternatively, we can use the explicit term

```
(1 | ET:Wafer)
```

## 4.2 General approach

As a general approach to importing data into R for mixed-effects analysis you should:

- Create a `data.frame` with one row per observation and one column per variable.
- Use `factor` or `as.factor` to explicitly convert any ordered factors to class `ordered`.
- Use `ordered` or `as.ordered` to explicitly convert any ordered factors to class `ordered`.
- If necessary, use interaction terms to create a factor with unique levels from inner nested factors.
- Plot the data. Plot it several ways. The use of lattice graphics is closely integrated with the `lme4` library. Lattice plots can provide invaluable insight into the structure of the data. Use them.

## 5 Contrasts

When comparing estimates produced by SAS PROC MIXED and by `lmer` one must be careful to consider the contrasts that are used to define the effects of factors. In SAS a model with an intercept and a qualitative factor is defined in terms of the intercept and the indicator variables for all but the last level of the factor. The default behaviour in S is to use the Helmert contrasts for the factor. On a balanced factor these provide a set of orthogonal contrasts. In R the default is the “treatment” contrasts which are almost the same as the SAS parameterization except that they drop the indicator of the first level, not the last level.

When in doubt, check which contrasts are being used with the `contrasts` function.

To make comparisons easier, you may find it worthwhile to declare

```
> options(contrasts = c(factor = "contr.SAS", ordered = "contr.poly"))
```

at the beginning of your session.

## References

- Nan M. Laird and James H. Ware. Random-effects models for longitudinal data. *Biometrics*, 38:963–974, 1982.
- Ramon C. Littell, George A. Milliken, Walter W. Stroup, and Russell D. Wolfinger. *SAS System for Mixed Models*. SAS Institute, Inc., 1996.

## A AvgDailyGain

```
> print(xyplot(adg ~ Treatment | Block, AvgDailyGain, type = c("g", "p", "r")
+         xlab = "Treatment (amount of feed additive)",
+         ylab = "Average daily weight gain (lb.)", aspect = "xy",
+         index.cond = function(x, y) coef(lm(y ~ x))[1]))

> ## compare with output 5.1, p. 178
> (fmlAdg <- lmer(adg ~ (Treatment - 1)*InitWt + (1 | Block), AvgDailyGain))
Linear mixed model fit by REML ['lmerMod']
Formula: adg ~ (Treatment - 1) * InitWt + (1 | Block)
Data: AvgDailyGain
```




Figure 1: Average daily weight gain

REML criterion at convergence: 65.3268

Random effects:

Groups	Name	Std.Dev.
Block	(Intercept)	0.5092
Residual		0.2223

Number of obs: 32, groups: Block, 8

Fixed Effects:

Treatment0	Treatment10	Treatment20
0.439137	1.426118	0.479628
Treatment30	InitWt	Treatment0:InitWt
0.200107	0.004448	-0.002154
Treatment10:InitWt	Treatment20:InitWt	
-0.003365	-0.001082	

> anova(fm1Adg) # checking significance of terms

Analysis of Variance Table

	Df	Sum Sq	Mean Sq	F value
Treatment	4	5.7248	1.43119	28.9543
InitWt	1	0.5495	0.54953	11.1175
Treatment:InitWt	3	0.1381	0.04603	0.9312

> ## common slope model

> (fm2Adg <- lmer(adg ~ InitWt + Treatment + (1 | Block), AvgDailyGain))

Linear mixed model fit by REML ['lmerMod']

Formula: adg ~ InitWt + Treatment + (1 | Block)

Data: AvgDailyGain

REML criterion at convergence: 36.3373

Random effects:

Groups	Name	Std.Dev.
Block	(Intercept)	0.4908
Residual		0.2238

Number of obs: 32, groups: Block, 8

Fixed Effects:

(Intercept)	InitWt	Treatment0	Treatment10	Treatment20
0.80111	0.00278	-0.55207	-0.06857	-0.08813

> anova(fm2Adg)

Analysis of Variance Table

	Df	Sum Sq	Mean Sq	F value
InitWt	1	0.51455	0.51455	10.275
Treatment	3	1.52670	0.50890	10.162

> (fm3Adg <- lmer(adg ~ InitWt + Treatment - 1 + (1 | Block), AvgDailyGain))



```

Linear mixed model fit by REML ['lmerMod']
Formula: adg ~ InitWt + Treatment - 1 + (1 | Block)
Data: AvgDailyGain
REML criterion at convergence: 36.3373
Random effects:
Groups      Name      Std.Dev.
Block      (Intercept) 0.4908
Residual                    0.2238
Number of obs: 32, groups: Block, 8
Fixed Effects:
      InitWt      Treatment0      Treatment10      Treatment20      Treatment30
      0.00278      0.24903      0.73254      0.71298      0.80111

```

## B BIB

```

> print(xyplot(y ~ x | Block, BIB, groups = Treatment, type = c("g", "p"),
+           aspect = "xy", auto.key = list(points = TRUE, space = "right",
+           lines = FALSE)))

```

```

> ## compare with Output 5.7, p. 188

```

```

> (fm1BIB <- lmer(y ~ Treatment * x + (1|Block), BIB))

```

```

Linear mixed model fit by REML ['lmerMod']

```

```

Formula: y ~ Treatment * x + (1 | Block)

```

```

Data: BIB

```

```

REML criterion at convergence: 104.8945

```

```

Random effects:

```

```

Groups      Name      Std.Dev.
Block      (Intercept) 4.272
Residual                    1.096

```

```

Number of obs: 24, groups: Block, 8

```

```

Fixed Effects:

```

```

      (Intercept)      Treatment1      Treatment2      Treatment3      x
      22.36784      4.42949      -0.43737      6.27864      0.44255
Treatment1:x      Treatment2:x      Treatment3:x
      -0.22377      0.05338      -0.17918

```

```

> anova(fm1BIB)      # strong evidence of different slopes

```

```

Analysis of Variance Table

```

```

      Df      Sum Sq      Mean Sq      F value
Treatment      3      23.447      7.816      6.5110
x              1     136.809     136.809     113.9695
Treatment:x     3      18.427      6.142      5.1169

```




Figure 2: Balanced incomplete block design

```

> ## compare with Output 5.9, p. 193
> (fm2BIB <- lmer(y ~ Treatment + x:Grp + (1|Block), BIB))
Linear mixed model fit by REML ['lmerMod']
Formula: y ~ Treatment + x:Grp + (1 | Block)
Data: BIB
REML criterion at convergence: 99.177
Random effects:
Groups      Name          Std.Dev.
Block      (Intercept)  4.304
Residual                    1.019
Number of obs: 24, groups: Block, 8
Fixed Effects:
(Intercept)      Treatment1      Treatment2      Treatment3      x:Grp13
      20.9452          5.3414          1.1356          8.1810          0.2395
      x:Grp24
      0.4892
> anova(fm2BIB)
Analysis of Variance Table
              Df  Sum Sq Mean Sq F value
Treatment    3   23.424    7.808  7.5236
x:Grp         2  154.733   77.367  74.5471

```

## C Bond

```

> ## compare with output 1.1 on p. 6
> (fm1Bond <- lmer(pressure ~ Metal + (1|Ingot), Bond))
Linear mixed model fit by REML ['lmerMod']
Formula: pressure ~ Metal + (1 | Ingot)
Data: Bond
REML criterion at convergence: 107.7902
Random effects:
Groups      Name          Std.Dev.
Ingot      (Intercept)  3.383
Residual                    3.220
Number of obs: 21, groups: Ingot, 7
Fixed Effects:
(Intercept)      Metalc      Metali
      71.1000      -0.9143      4.8000
> anova(fm1Bond)
Analysis of Variance Table
              Df Sum Sq Mean Sq F value
Metal         2   131.9    65.95  6.3587

```

## D Cultivation

```
> str(Cultivation)
'data.frame':      24 obs. of  4 variables:
 $ Block: Factor w/ 4 levels "1","2","3","4": 1 1 1 1 1 1 2 2 2 2 ...
 $ Cult : Factor w/ 2 levels "a","b": 1 1 1 2 2 2 1 1 1 2 ...
 $ Inoc : Factor w/ 3 levels "con","dea","liv": 1 2 3 1 2 3 1 2 3 1 ...
 $ drywt: num  27.4 29.7 34.5 29.4 32.5 34.4 28.9 28.7 33.4 28.7 ...
- attr(*, "ginfo")=List of 7
 ..$ formula      :Class 'formula' length 3 drywt ~ 1 | Block/Cult
 .. ..- attr(*, ".Environment")=<environment: R_GlobalEnv>
 ..$ order.groups:List of 2
 .. ..$ Block: logi TRUE
 .. ..$ Cult : logi TRUE
 ..$ FUN          :function (x)
 ..$ outer        : NULL
 ..$ inner        :List of 1
 .. ..$ Cult:Class 'formula' length 2 ~Inoc
 .. ..- attr(*, ".Environment")=<environment: R_GlobalEnv>
 ..$ labels       :List of 1
 .. ..$ drywt: chr "Yield"
 ..$ units        : list()
> xtabs(~Block+Cult, Cultivation)
      Cult
Block a b
  1 3 3
  2 3 3
  3 3 3
  4 3 3
> (fmlCult <- lmer(drywt ~ Inoc * Cult + (1|Block) + (1|Cult), Cultivation))
Linear mixed model fit by REML ['lmerMod']
Formula: drywt ~ Inoc * Cult + (1 | Block) + (1 | Cult)
Data: Cultivation
REML criterion at convergence: 68.4874
Random effects:
 Groups   Name                Std.Dev.
Block    (Intercept)  1.099
Cult     (Intercept)  1.105
Residual                    1.094
Number of obs: 24, groups: Block, 4; Cult, 2
Fixed Effects:
```

(Intercept)	Inoccon	Inocdea	Culta	Inoccon:Culta
33.525	-5.500	-2.875	-0.375	0.250

Inocdea:Culta  
-1.025

```
> anova(fm1Cult)
```

Analysis of Variance Table

	Df	Sum Sq	Mean Sq	F value
Inoc	2	118.176	59.088	49.3908
Cult	1	0.182	0.182	0.1517
Inoc:Cult	2	1.826	0.913	0.7631

```
> (fm2Cult <- lmer(drywt ~ Inoc + Cult + (1|Block) + (1|Cult), Cultivation))
```

Linear mixed model fit by REML ['lmerMod']

Formula: drywt ~ Inoc + Cult + (1 | Block) + (1 | Cult)

Data: Cultivation

REML criterion at convergence: 73.7535

Random effects:

Groups	Name	Std.Dev.
Block	(Intercept)	1.101
Cult	(Intercept)	1.070
Residual		1.078

Number of obs: 24, groups: Block, 4; Cult, 2

Fixed Effects:

(Intercept)	Inoccon	Inocdea	Culta
33.6542	-5.3750	-3.3875	-0.6333

```
> anova(fm2Cult)
```

Analysis of Variance Table

	Df	Sum Sq	Mean Sq	F value
Inoc	2	118.176	59.088	50.8069
Cult	1	0.188	0.188	0.1616

```
> (fm3Cult <- lmer(drywt ~ Inoc + (1|Block) + (1|Cult), Cultivation))
```

Linear mixed model fit by REML ['lmerMod']

Formula: drywt ~ Inoc + (1 | Block) + (1 | Cult)

Data: Cultivation

REML criterion at convergence: 75.6778

Random effects:

Groups	Name	Std.Dev.
Block	(Intercept)	1.1013
Cult	(Intercept)	0.3219
Residual		1.0784

Number of obs: 24, groups: Block, 4; Cult, 2

Fixed Effects:

(Intercept)	Inoccon	Inocdea
33.338	-5.375	-3.388

```
> anova(fm3Cult)
Analysis of Variance Table
      Df Sum Sq Mean Sq F value
Inoc   2 118.18   59.088   50.807
```

## E Demand

```
> ## compare to output 3.13, p. 132
> (fm1Demand <-
+  lmer(log(d) ~ log(y) + log(rd) + log(rt) + log(rs) + (1|State) + (1|Year),
+      Demand))
Linear mixed model fit by REML ['lmerMod']
Formula: log(d) ~ log(y) + log(rd) + log(rt) + log(rs) + (1 | State) +
      Data: Demand
REML criterion at convergence: -240.1653
Random effects:
      Groups      Name      Std.Dev.
      Year      (Intercept) 0.01627
      State      (Intercept) 0.17177
      Residual              0.03342
Number of obs: 77, groups: Year, 11; State, 7
Fixed Effects:
      (Intercept)      log(y)      log(rd)      log(rt)      log(rs)
      -1.28382      1.06978      -0.29533      0.03988      -0.32673
```

## F HR

```
> ## linear trend in time
> (fm1HR <- lmer(HR ~ Time * Drug + baseHR + (Time|Patient), HR))
Linear mixed model fit by REML ['lmerMod']
Formula: HR ~ Time * Drug + baseHR + (Time | Patient)
      Data: HR
REML criterion at convergence: 767.607
Random effects:
      Groups      Name      Std.Dev. Corr
      Patient      (Intercept) 7.787
      Time          6.147      -0.56
      Residual          4.936
Number of obs: 120, groups: Patient, 24
Fixed Effects:
```

```

(Intercept)      Time      Druga      Drugb      baseHR
    33.9776    -3.1970    3.5992    7.0912    0.5434
Time:Druga  Time:Drugb
    -7.5013    -3.9894
> anova(fm1HR)
Analysis of Variance Table

      Df Sum Sq Mean Sq F value
Time    1  379.23   379.23  15.5671
Drug     2   92.88    46.44   1.9064
baseHR   1  533.27   533.27  21.8905
Time:Drug 2   72.12    36.06   1.4802
> ## remove interaction
> (fm3HR <- lmer(HR ~ Time + Drug + baseHR + (Time|Patient), HR))
Linear mixed model fit by REML ['lmerMod']
Formula: HR ~ Time + Drug + baseHR + (Time | Patient)
Data: HR
REML criterion at convergence: 779.8283
Random effects:
Groups   Name      Std.Dev. Corr
Patient (Intercept) 7.846
        Time       6.400  -0.57
Residual              4.936
Number of obs: 120, groups: Patient, 24
Fixed Effects:
(Intercept)      Time      Druga      Drugb      baseHR
    36.0463    -7.0273    -0.4524    4.9365    0.5434
> anova(fm3HR)
Analysis of Variance Table

      Df Sum Sq Mean Sq F value
Time    1  364.02   364.02  14.9431
Drug     2   92.88    46.44   1.9064
baseHR   1  533.27   533.27  21.8905
> ## remove Drug term
> (fm4HR <- lmer(HR ~ Time + baseHR + (Time|Patient), HR))
Linear mixed model fit by REML ['lmerMod']
Formula: HR ~ Time + baseHR + (Time | Patient)
Data: HR
REML criterion at convergence: 791.1481
Random effects:
Groups   Name      Std.Dev. Corr

```

```

Patient (Intercept) 7.939
          Time      6.400    -0.55
Residual              4.936
Number of obs: 120, groups: Patient, 24
Fixed Effects:
(Intercept)          Time          baseHR
      36.9313      -7.0273      0.5508
> anova(fm4HR)
Analysis of Variance Table
      Df Sum Sq Mean Sq F value
Time    1 364.03  364.03  14.943
baseHR   1 534.87  534.87  21.956

```

## G Mississippi

```

> ## compare with output 4.1, p. 142
> (fm1Miss <- lmer(y ~ 1 + (1 | influent), Mississippi))
Linear mixed model fit by REML ['lmerMod']
Formula: y ~ 1 + (1 | influent)
Data: Mississippi
REML criterion at convergence: 252.3511
Random effects:
Groups   Name      Std.Dev.
influent (Intercept) 7.958
Residual              6.531
Number of obs: 37, groups: influent, 6
Fixed Effects:
(Intercept)
      21.22
> ## compare with output 4.2, p. 143
> (fm1MLMiss <- lmer(y ~ 1 + (1 | influent), Mississippi, REML=FALSE))
Linear mixed model fit by maximum likelihood ['lmerMod']
Formula: y ~ 1 + (1 | influent)
Data: Mississippi
      AIC      BIC    logLik  deviance
262.5570 267.3898 -128.2785 256.5570
Random effects:
Groups   Name      Std.Dev.
influent (Intercept) 7.159
Residual              6.534

```



```

Number of obs: 37, groups: influent, 6
Fixed Effects:
(Intercept)
      21.22
> ranef(fm1MLMiss)           # BLUP's of random effects on p. 144
$influent
(Intercept)
1      0.3097833
2     -6.5772239
3     -3.7862717
4      2.8826693
5     -5.8435163
6     13.0145592

attr(,"class")
[1] "ranef.mer"
> ranef(fm1Miss)           # BLUP's of random effects on p. 142
$influent
(Intercept)
1      0.309286
2     -6.719325
3     -3.897940
4      2.946101
5     -6.012976
6     13.374854

attr(,"class")
[1] "ranef.mer"
> VarCorr(fm1Miss)         # compare to output 4.7, p. 148
Groups   Name             Std.Dev.
influent (Intercept)  7.9576
Residual                6.5313
> ## compare to output 4.8 and 4.9, pp. 150-152
> (fm2Miss <- lmer(y ~ Type + (1 | influent), Mississippi))
Linear mixed model fit by REML ['lmerMod']
Formula: y ~ Type + (1 | influent)
Data: Mississippi
REML criterion at convergence: 234.5246
Random effects:
Groups   Name             Std.Dev.

```

```

influent (Intercept) 3.869
Residual              6.520
Number of obs: 37, groups: influent, 6
Fixed Effects:
(Intercept)          Type1          Type2
           36.40         -20.80         -16.46
> anova(fm2Miss)
Analysis of Variance Table
      Df Sum Sq Mean Sq F value
Type  2 541.75  270.88   6.3715

```

## H Multilocation

```

> str(Multilocation)
'data.frame':      108 obs. of  7 variables:
 $ obs      : num  3 4 6 7 9 10 12 16 19 20 ...
 $ Location: Factor w/ 9 levels "A","B","C","D",...: 1 1 1 1 1 1 1 1 1 1 ...
 $ Block    : Factor w/ 3 levels "1","2","3": 1 1 1 1 2 2 2 2 3 3 ...
 $ Trt      : Factor w/ 4 levels "1","2","3","4": 3 4 2 1 2 1 3 4 1 2 ...
 $ Adj      : num  3.16 3.12 3.16 3.25 2.71 ...
 $ Fe       : num  7.1 6.68 6.83 6.53 8.25 ...
 $ Grp      : Factor w/ 27 levels "1:A","1:B","1:C",...: 1 1 1 1 10 10 10 10 19
- attr(*, "ginfo")=List of 7
 ..$ formula      :Class 'formula' length 3 Adj ~ 1 | Location/Block
 .. ..- attr(*, ".Environment")=<environment: R_GlobalEnv>
 ..$ order.groups:List of 2
 .. ..$ Location: logi TRUE
 .. ..$ Block    : logi TRUE
 ..$ FUN          :function (x)
 ..$ outer        : NULL
 ..$ inner        :List of 1
 .. ..$ Block:Class 'formula' length 2 ~Trt
 .. ..- attr(*, ".Environment")=<environment: R_GlobalEnv>
 ..$ labels       :List of 1
 .. ..$ Adj: chr "Adjusted yield"
 ..$ units        : list()
> ### Create a Block %in% Location factor
> Multilocation$Grp <- with(Multilocation, Block:Location)
> (fm1Mult <- lmer(Adj ~ Location * Trt + (1|Grp), Multilocation))

```

```

Linear mixed model fit by REML ['lmerMod']
Formula: Adj ~ Location * Trt + (1 | Grp)
Data: Multilocation
REML criterion at convergence: 10.6462
Random effects:
Groups      Name      Std.Dev.
Grp      (Intercept) 0.07496
Residual              0.18595
Number of obs: 108, groups: Grp, 27
Fixed Effects:
(Intercept)      LocationA      LocationB      LocationC
2.35923      0.64930      0.06643      0.54533
LocationD      LocationE      LocationF      LocationG
0.37413      0.55000      0.99810      0.36057
LocationH      Trt1      Trt2      Trt3
1.01403      0.22720      -0.00140      0.42323
LocationA:Trt1 LocationB:Trt1 LocationC:Trt1 LocationD:Trt1
-0.18853      -0.27523      -0.04000      -0.53513
LocationE:Trt1 LocationF:Trt1 LocationG:Trt1 LocationH:Trt1
-0.26297      -0.27153      0.20323      -0.14953
LocationA:Trt2 LocationB:Trt2 LocationC:Trt2 LocationD:Trt2
-0.09347      -0.32273      0.08960      -0.29693
LocationE:Trt2 LocationF:Trt2 LocationG:Trt2 LocationH:Trt2
-0.30693      -0.30993      -0.10860      -0.33060
LocationA:Trt3 LocationB:Trt3 LocationC:Trt3 LocationD:Trt3
-0.40247      -0.56550      -0.12247      -0.54840
LocationE:Trt3 LocationF:Trt3 LocationG:Trt3 LocationH:Trt3
-0.32863      -0.46257      -0.25297      -0.37203
> anova(fm1Mult)
Analysis of Variance Table
              Df Sum Sq Mean Sq F value
Location      8 6.9476 0.86845 25.1150
Trt           3 1.2217 0.40725 11.7774
Location:Trt 24 0.9966 0.04152 1.2008
> (fm2Mult <- lmer(Adj ~ Location + Trt + (1|Grp), Multilocation))
Linear mixed model fit by REML ['lmerMod']
Formula: Adj ~ Location + Trt + (1 | Grp)
Data: Multilocation
REML criterion at convergence: -6.0011
Random effects:

```

```

Groups   Name          Std.Dev.
Grp      (Intercept) 0.07131
Residual                0.19161
Number of obs: 108, groups: Grp, 27
Fixed Effects:
(Intercept)   LocationA   LocationB   LocationC   LocationD
      2.53296      0.47818      -0.22443      0.52712      0.02902
LocationE     LocationF     LocationG     LocationH      Trt1
      0.32537      0.73709      0.32098      0.80099      0.05834
      Trt2          Trt3
      -0.18802      0.08379
> (fm3Mult <- lmer(Adj ~ Location + (1|Grp), Multilocation))
Linear mixed model fit by REML ['lmerMod']
Formula: Adj ~ Location + (1 | Grp)
Data: Multilocation
REML criterion at convergence: 9.8205
Random effects:
Groups   Name          Std.Dev.
Grp      (Intercept) 0.04067
Residual                0.22459
Number of obs: 108, groups: Grp, 27
Fixed Effects:
(Intercept)   LocationA   LocationB   LocationC   LocationD
      2.52149      0.47818      -0.22443      0.52712      0.02902
LocationE     LocationF     LocationG     LocationH
      0.32537      0.73709      0.32098      0.80099
> (fm4Mult <- lmer(Adj ~ Trt + (1|Grp), Multilocation))
Linear mixed model fit by REML ['lmerMod']
Formula: Adj ~ Trt + (1 | Grp)
Data: Multilocation
REML criterion at convergence: 31.5057
Random effects:
Groups   Name          Std.Dev.
Grp      (Intercept) 0.3330
Residual                0.1916
Number of obs: 108, groups: Grp, 27
Fixed Effects:
(Intercept)      Trt1          Trt2          Trt3
      2.86567      0.05834      -0.18802      0.08379
> (fm5Mult <- lmer(Adj ~ 1 + (1|Grp), Multilocation))

```

Linear mixed model fit by REML ['lmerMod']

Formula: Adj ~ 1 + (1 | Grp)

Data: Multilocation

REML criterion at convergence: 47.3273

Random effects:

Groups	Name	Std.Dev.
Grp	(Intercept)	0.3279
Residual		0.2246

Number of obs: 108, groups: Grp, 27

Fixed Effects:

(Intercept)	2.854
-------------	-------

> anova(fm2Mult)

Analysis of Variance Table

	Df	Sum Sq	Mean Sq	F value
Location	8	7.3768	0.92209	25.115
Trt	3	1.2217	0.40725	11.092

> (fm2MultR <- lmer(Adj ~ Trt + (Trt - 1|Location) + (1|Block), Multilocation +  
+ verbose = TRUE))

(NM)	20:	f = 61.9198 at	1.00545	0.00545455	0.00545455	0.00545455	1.00545
(NM)	40:	f = 59.561 at	1.00545	0.00261822	0.0403054	0.0344762	1.00004
(NM)	60:	f = 54.0031 at	1.02396	0.00929341	0.138615	0.0780559	0.98756
(NM)	80:	f = 39.4344 at	1.0713	0.0373298	0.307821	0.20942	1.00342
(NM)	100:	f = 32.0819 at	1.19531	0.0751749	0.728702	0.453329	0.994316
(NM)	120:	f = 26.5195 at	1.25611	0.366584	0.603141	0.407144	1.05837
(NM)	140:	f = 24.4381 at	1.3621	0.665294	0.738454	0.547998	1.1212
(NM)	160:	f = 24.4381 at	1.3621	0.665294	0.738454	0.547998	1.1212
(NM)	180:	f = 24.2193 at	1.3499	0.65325	0.7649	0.517308	1.10627
(NM)	200:	f = 24.0942 at	1.40234	0.763815	0.855302	0.571687	1.11842
(NM)	220:	f = 22.6873 at	1.45124	0.675951	1.03427	0.50153	1.03942
(NM)	240:	f = 20.7867 at	1.66872	0.913098	1.33985	0.45005	1.02703
(NM)	260:	f = 20.689 at	1.82264	1.01623	1.60798	0.378413	0.979706
(NM)	280:	f = 20.1255 at	1.80393	1.03898	1.49164	0.454599	1.02789
(NM)	300:	f = 18.352 at	1.83323	1.03575	1.48799	0.419642	1.02885
(NM)	320:	f = 16.2411 at	1.79771	0.847523	1.50297	0.306929	0.979897
(NM)	340:	f = 14.2286 at	1.94231	1.04428	1.73304	0.366229	0.995187
(NM)	360:	f = 14.0233 at	2.0624	1.07514	1.86884	0.342522	0.955776
(NM)	380:	f = 13.9003 at	2.0849	1.18424	1.98312	0.34784	0.95409
(NM)	400:	f = 13.676 at	2.18803	1.23441	2.11555	0.297887	0.930266
(NM)	420:	f = 13.6232 at	2.04961	1.10473	1.89215	0.305718	0.955703

(NM) 440:	f = 13.5372 at	2.12781	1.19533	1.99379	0.296788	0.951442	0.68556	
(NM) 460:	f = 13.4904 at	2.20434	1.27342	2.05497	0.275888	0.950807	0.950807	0
(NM) 480:	f = 13.4707 at	2.2032	1.26208	2.03241	0.262177	0.953144	0.953144	0
(NM) 500:	f = 13.4025 at	2.26975	1.27669	2.10891	0.239464	0.935694	0.935694	0
(NM) 520:	f = 13.2212 at	2.36945	1.25332	2.13275	0.207875	0.916464	0.916464	0
(NM) 540:	f = 13.1398 at	2.43055	1.23094	2.08102	0.200132	0.915996	0.915996	0
(NM) 560:	f = 13.1364 at	2.43347	1.21358	2.06449	0.194451	0.91487	0.91487	0
(NM) 580:	f = 13.1154 at	2.39312	1.19974	2.0126	0.186435	0.926064	0.926064	0
(NM) 600:	f = 13.1054 at	2.42565	1.22996	2.06537	0.190615	0.920009	0.920009	0
(NM) 620:	f = 13.0991 at	2.41147	1.20897	2.01764	0.189463	0.926611	0.926611	0
(NM) 640:	f = 13.0922 at	2.42964	1.22439	2.02352	0.179988	0.928005	0.928005	0
(NM) 660:	f = 13.0639 at	2.43315	1.23191	2.01856	0.173076	0.93278	0.93278	0
(NM) 680:	f = 13.016 at	2.3713	1.22969	2.00382	0.188877	0.946905	0.946905	0
(NM) 700:	f = 12.9083 at	2.29075	1.13328	1.86767	0.177206	0.978849	0.978849	0
(NM) 720:	f = 12.8075 at	2.21898	1.09171	1.89915	0.169841	0.984197	0.984197	0
(NM) 740:	f = 12.7992 at	2.19757	1.10341	1.86916	0.178646	0.995988	0.995988	0
(NM) 760:	f = 12.7897 at	2.17796	1.13273	1.90752	0.179228	1.0011	1.0011	0
(NM) 780:	f = 12.7809 at	2.15056	1.1046	1.84015	0.174026	1.0153	1.0153	0
(NM) 800:	f = 12.7804 at	2.12732	1.09436	1.82856	0.183473	1.01688	1.01688	0
(NM) 820:	f = 12.7787 at	2.13662	1.08847	1.83066	0.177947	1.01481	1.01481	0
(NM) 840:	f = 12.7773 at	2.15028	1.10128	1.84278	0.176254	1.01445	1.01445	0
(NM) 860:	f = 12.7725 at	2.15922	1.10712	1.85667	0.177219	1.01065	1.01065	0
(NM) 880:	f = 12.7655 at	2.14918	1.11286	1.86305	0.180553	1.01155	1.01155	0
(NM) 900:	f = 12.7537 at	2.18932	1.12635	1.88674	0.172552	1.00228	1.00228	0.54552
(NM) 920:	f = 12.7291 at	2.17076	1.16096	1.86147	0.185085	1.00762	1.00762	0.59752
(NM) 940:	f = 12.7026 at	2.16097	1.13991	1.83463	0.175591	1.01189	1.01189	0
(NM) 960:	f = 12.6284 at	2.10113	1.17069	1.80221	0.198297	1.01406	1.01406	0
(NM) 980:	f = 12.5335 at	2.03733	1.16124	1.7379	0.207672	1.02195	1.02195	0
(NM) 1000:	f = 12.384 at	2.11367	1.13749	1.78552	0.197318	0.966393	0.966393	0
(NM) 1020:	f = 12.233 at	2.08328	1.2079	1.73936	0.240806	0.936401	0.936401	0
(NM) 1040:	f = 12.173 at	2.11078	1.22627	1.74331	0.261277	0.893935	0.893935	0
(NM) 1060:	f = 12.1041 at	2.07633	1.32742	1.77528	0.279233	0.897747	0.897747	0
(NM) 1080:	f = 12.035 at	2.1325	1.3692	1.80872	0.291651	0.833455	0.833455	0
(NM) 1100:	f = 12.0201 at	2.15982	1.41312	1.79734	0.287904	0.829344	0.829344	0
(NM) 1120:	f = 12.0057 at	2.14337	1.40507	1.78759	0.27548	0.839779	0.839779	0
(NM) 1140:	f = 12.0009 at	2.13059	1.39954	1.76789	0.291957	0.824309	0.824309	0
(NM) 1160:	f = 11.9937 at	2.12354	1.41228	1.75392	0.287247	0.82142	0.82142	0
(NM) 1180:	f = 11.9881 at	2.08833	1.39056	1.74483	0.293739	0.8131	0.8131	0.5975
(NM) 1200:	f = 11.9855 at	2.08809	1.38455	1.74466	0.288707	0.820793	0.820793	0
(NM) 1220:	f = 11.985 at	2.08004	1.37932	1.7397	0.287017	0.821874	0.821874	0

(NM) 1240:	f = 11.9837 at	2.08386	1.40108	1.74754	0.289483	0.816459	0.58
(NM) 1260:	f = 11.9837 at	2.08386	1.40108	1.74754	0.289483	0.816459	0.58
(NM) 1280:	f = 11.9837 at	2.08996	1.39612	1.74955	0.288525	0.818697	
(NM) 1300:	f = 11.9836 at	2.08841	1.39984	1.75106	0.289973	0.817218	
(NM) 1320:	f = 11.9835 at	2.08973	1.40261	1.75006	0.289534	0.815948	
(NM) 1340:	f = 11.9835 at	2.08891	1.40083	1.74946	0.289551	0.816277	
(NM) 1360:	f = 11.9834 at	2.09111	1.40111	1.74937	0.289615	0.815641	
(NM) 1380:	f = 11.9832 at	2.08996	1.4	1.74752	0.288629	0.815924	
(NM) 1400:	f = 11.9822 at	2.08829	1.39455	1.74649	0.287041	0.816721	
(NM) 1420:	f = 11.9812 at	2.09074	1.3991	1.75231	0.28809	0.814558	
(NM) 1440:	f = 11.9789 at	2.10291	1.39203	1.76035	0.284577	0.813817	
(NM) 1460:	f = 11.9784 at	2.10452	1.38288	1.76606	0.28424	0.816039	
(NM) 1480:	f = 11.9734 at	2.13221	1.41877	1.77934	0.285026	0.804117	
(NM) 1500:	f = 11.9698 at	2.13289	1.42208	1.79342	0.285478	0.802824	
(NM) 1520:	f = 11.9693 at	2.13183	1.42029	1.78343	0.284999	0.801275	
(NM) 1540:	f = 11.9664 at	2.14521	1.42269	1.79235	0.284746	0.796568	0.5550
(NM) 1560:	f = 11.963 at	2.1601	1.41335	1.79853	0.28385	0.796978	0
(NM) 1580:	f = 11.9542 at	2.17395	1.4072	1.79487	0.283013	0.801184	
(NM) 1600:	f = 11.9287 at	2.18982	1.40132	1.80705	0.289973	0.807197	
(NM) 1620:	f = 11.9026 at	2.22527	1.42112	1.83807	0.306969	0.808488	
(NM) 1640:	f = 11.8661 at	2.23857	1.41239	1.84436	0.319065	0.812582	
(NM) 1660:	f = 11.7914 at	2.18134	1.37563	1.78018	0.322485	0.827421	
(NM) 1680:	f = 11.6448 at	2.19173	1.30767	1.71513	0.380578	0.846455	
(NM) 1700:	f = 11.4468 at	2.10631	1.33488	1.64624	0.491925	0.885251	
(NM) 1720:	f = 11.0259 at	2.06191	1.39526	1.62123	0.543078	0.872248	0.8321
(NM) 1740:	f = 10.0475 at	2.10941	1.35954	1.55582	0.778509	0.938	
(NM) 1760:	f = 9.27268 at	1.95803	1.27274	1.31405	1.07539	1.04039	1.294
(NM) 1780:	f = 8.91509 at	2.03355	1.24606	1.42214	1.17822	1.08941	
(NM) 1800:	f = 8.29709 at	2.03352	1.15665	1.38263	1.12801	1.04244	1.110
(NM) 1820:	f = 8.11779 at	1.95655	1.18711	1.38354	1.08507	1.01596	0.984
(NM) 1840:	f = 8.0726 at	1.96253	1.17907	1.41558	1.07713	1.02363	0.95632
(NM) 1860:	f = 7.96664 at	1.99183	1.22938	1.52611	1.02639	0.996848	0.9385
(NM) 1880:	f = 7.87922 at	1.92371	1.18389	1.46657	1.0442	1.01993	0.9363
(NM) 1900:	f = 7.85737 at	1.91944	1.19355	1.48516	1.0842	1.03747	0.9910
(NM) 1920:	f = 7.83673 at	1.89171	1.18225	1.47285	1.07146	1.04115	0.9860
(NM) 1940:	f = 7.79845 at	1.8534	1.13028	1.43044	1.0534	1.05866	0.9755
(NM) 1960:	f = 7.78101 at	1.82894	1.09835	1.38622	1.02775	1.06669	0.961
(NM) 1980:	f = 7.75116 at	1.83584	1.1105	1.41578	1.02365	1.06507	0.9481
(NM) 2000:	f = 7.69582 at	1.8218	1.06492	1.37267	0.982198	1.0768	0.9252
(NM) 2020:	f = 7.66571 at	1.82851	1.04047	1.35675	1.01455	1.09577	0.9551

(NM) 2040: f = 7.54979 at	1.89459	1.0323	1.36912	1.01424	1.1004	0.9405
(NM) 2060: f = 7.4556 at	1.88425	1.03975	1.37355	0.99562	1.1156	0.9405
(NM) 2080: f = 7.27001 at	1.90729	1.08678	1.45478	1.0164	1.14602	0.9405
(NM) 2100: f = 7.14887 at	1.87107	1.12161	1.46779	1.00976	1.14649	0.9405
(NM) 2120: f = 7.06088 at	1.82261	1.1938	1.44906	1.09486	1.17023	0.9714
(NM) 2140: f = 6.8876 at	1.85403	1.22	1.46988	1.10555	1.18647	0.9405
(NM) 2160: f = 6.8087 at	1.9333	1.26647	1.53804	1.17783	1.17679	0.9405
(NM) 2180: f = 6.64797 at	1.94072	1.32191	1.53544	1.20588	1.20712	0.9405
(NM) 2200: f = 6.38235 at	1.89789	1.2275	1.50345	1.12161	1.17124	0.9405
(NM) 2220: f = 5.28253 at	1.93002	1.26748	1.46829	1.25486	0.822962	0.6494
(NM) 2240: f = 3.91833 at	1.786	1.39324	1.45804	1.30288	0.496145	0.4527
(NM) 2260: f = 3.67151 at	1.69175	1.47508	1.47088	1.41085	0.264411	0.4120
(NM) 2280: f = 3.47942 at	1.78106	1.39349	1.46496	1.20416	0.55617	0.4868
(NM) 2300: f = 3.07981 at	1.76468	1.45347	1.53864	1.24433	0.386478	0.2988
(NM) 2320: f = 2.90588 at	1.80163	1.60534	1.58663	1.45143	0.316265	0.2988
(NM) 2340: f = 2.46793 at	1.76705	1.56042	1.6135	1.35563	0.278209	0.2988
(NM) 2360: f = 2.44314 at	1.72804	1.54535	1.63088	1.33512	0.212492	0.2729
(NM) 2380: f = 2.44314 at	1.72804	1.54535	1.63088	1.33512	0.212492	0.2729
(NM) 2400: f = 2.39768 at	1.75619	1.51737	1.6167	1.28883	0.308789	0.2988
(NM) 2420: f = 2.38472 at	1.76387	1.49057	1.60428	1.30066	0.27666	0.2988
(NM) 2440: f = 2.36176 at	1.75308	1.50101	1.61453	1.30375	0.288	0.2988
(NM) 2460: f = 2.34672 at	1.74961	1.51342	1.60297	1.31362	0.268366	0.2988
(NM) 2480: f = 2.29346 at	1.75058	1.50786	1.60221	1.3135	0.31407	0.2988
(NM) 2500: f = 2.28385 at	1.74011	1.48772	1.58804	1.31791		0.2988
(NM) 2520: f = 2.25515 at	1.733	1.49054	1.60464	1.33439	0.318	0.2988
(NM) 2540: f = 2.21317 at	1.72278	1.472	1.60045	1.36301	0.359	0.2988
(NM) 2560: f = 2.16328 at	1.71638	1.49166	1.63418	1.35117	0.367	0.2988
(NM) 2580: f = 2.08334 at	1.68688	1.44607	1.62126	1.35717	0.298	0.2988
(NM) 2600: f = 2.01409 at	1.65951	1.43489	1.62315	1.35938	0.305	0.2988
(NM) 2620: f = 1.85329 at	1.67381	1.48828	1.62979	1.36232	0.25	0.2988
(NM) 2640: f = 1.68403 at	1.67739	1.5014	1.6188	1.42776	0.26925	0.2988
(NM) 2660: f = 1.57893 at	1.72673	1.54207	1.65795	1.47801	0.239536	0.2988
(NM) 2680: f = 1.45909 at	1.80751	1.59036	1.68377	1.5488	0.223	0.2988
(NM) 2700: f = 1.45909 at	1.80751	1.59036	1.68377	1.5488	0.223	0.2988
(NM) 2720: f = 1.45524 at	1.84096	1.60635	1.692	1.56536	0.232	0.2988
(NM) 2740: f = 1.43092 at	1.84902	1.60903	1.71071	1.55575	0.242	0.2988
(NM) 2760: f = 1.42293 at	1.8751	1.62999	1.73329	1.58505	0.243	0.2988
(NM) 2780: f = 1.41627 at	1.89083	1.64553	1.76646	1.59434	0.254	0.2988
(NM) 2800: f = 1.41378 at	1.8863	1.65314	1.7692	1.61093	0.254	0.2988
(NM) 2820: f = 1.41179 at	1.88883	1.65235	1.7613	1.60597	0.253	0.2988



(NM) 2840: f = 1.41021 at	1.89963	1.65661	1.76696	1.60809	0.251
(NM) 2860: f = 1.409 at	1.90136	1.66553	1.77076	1.61045	0.24458
(NM) 2880: f = 1.40878 at	1.9069	1.67131	1.77717	1.61419	0.248
(NM) 2900: f = 1.40842 at	1.902	1.67066	1.77808	1.61408	0.243593
(NM) 2920: f = 1.4083 at	1.90099	1.66836	1.77466	1.61526	0.2445
(NM) 2940: f = 1.40828 at	1.90092	1.66878	1.77711	1.61481	0.243
(NM) 2960: f = 1.40825 at	1.90121	1.66846	1.77586	1.61502	0.244
(NM) 2980: f = 1.40824 at	1.90081	1.66759	1.77577	1.61388	0.244
(NM) 3000: f = 1.40823 at	1.90101	1.66889	1.77659	1.61519	0.244
(NM) 3020: f = 1.40823 at	1.90119	1.66822	1.77666	1.61482	0.244798
(NM) 3040: f = 1.40822 at	1.90094	1.66844	1.7763	1.61458	0.244
(NM) 3060: f = 1.40822 at	1.90115	1.66893	1.7766	1.61523	0.244
(NM) 3080: f = 1.40821 at	1.9008	1.66867	1.77643	1.61494	0.244
(NM) 3100: f = 1.4082 at	1.90176	1.66958	1.7773	1.61504	0.2443
(NM) 3120: f = 1.4082 at	1.9021	1.66982	1.77743	1.61538	0.2444
(NM) 3140: f = 1.4082 at	1.9018	1.66955	1.77703	1.61523	0.2444
(NM) 3160: f = 1.40819 at	1.90181	1.66938	1.77671	1.61458	0.244
(NM) 3180: f = 1.40817 at	1.90154	1.66914	1.77648	1.61453	0.244
(NM) 3200: f = 1.40814 at	1.90032	1.66857	1.7752	1.61483	0.244
(NM) 3220: f = 1.40809 at	1.90062	1.67033	1.77586	1.61587	0.243
(NM) 3240: f = 1.40807 at	1.89957	1.66932	1.77425	1.61517	0.243
(NM) 3260: f = 1.40805 at	1.90052	1.66912	1.77525	1.61539	0.243
(NM) 3280: f = 1.40801 at	1.90091	1.66965	1.77572	1.61659	0.242
(NM) 3300: f = 1.40796 at	1.89989	1.66908	1.77465	1.61649	0.242
(NM) 3320: f = 1.40792 at	1.89971	1.66804	1.77371	1.61497	0.243
(NM) 3340: f = 1.40784 at	1.90049	1.66807	1.77337	1.61482	0.242
(NM) 3360: f = 1.40778 at	1.8972	1.66547	1.77056	1.61254	0.241
(NM) 3380: f = 1.40767 at	1.89954	1.66661	1.77186	1.61357	0.239924
(NM) 3400: f = 1.40753 at	1.90223	1.66933	1.77562	1.61564	0.240
(NM) 3420: f = 1.40746 at	1.90183	1.66957	1.77647	1.61605	0.239
(NM) 3440: f = 1.40741 at	1.89991	1.66778	1.77392	1.61371	0.240
(NM) 3460: f = 1.40738 at	1.8993	1.66861	1.77343	1.61425	0.242165
(NM) 3480: f = 1.40736 at	1.89931	1.66829	1.77409	1.61438	0.241
(NM) 3500: f = 1.40733 at	1.89951	1.66868	1.77397	1.61521	0.242
(NM) 3520: f = 1.40731 at	1.89942	1.66843	1.77309	1.61477	0.243
(NM) 3540: f = 1.4073 at	1.89932	1.66881	1.77392	1.61556	0.2430
(NM) 3560: f = 1.40729 at	1.89888	1.66851	1.77348	1.61514	0.243
(NM) 3580: f = 1.40726 at	1.89832	1.66749	1.77253	1.61423	0.243
(NM) 3600: f = 1.40724 at	1.89834	1.66711	1.77324	1.61424	0.243
(NM) 3620: f = 1.40722 at	1.89699	1.66568	1.77168	1.61269	0.244

(NM) 3640:	f = 1.4072 at	1.89649	1.66565	1.77197	1.61277	0.2451
(NM) 3660:	f = 1.4072 at	1.89656	1.6649	1.77179	1.6122	0.2450
(NM) 3680:	f = 1.40719 at	1.89713	1.66557	1.7727	1.6127	0.245
(NM) 3700:	f = 1.40719 at	1.89733	1.6659	1.77268	1.61292	0.245
(NM) 3720:	f = 1.40718 at	1.89751	1.66595	1.77286	1.61253	
(NM) 3740:	f = 1.40718 at	1.89749	1.66565	1.77277	1.61208	0.245
(NM) 3760:	f = 1.40718 at	1.89772	1.66592	1.77289	1.61235	0.245
(NM) 3780:	f = 1.40718 at	1.89765	1.66583	1.77309	1.61239	0.245
(NM) 3800:	f = 1.40717 at	1.89761	1.66569	1.77298	1.61252	0.24
(NM) 3820:	f = 1.40717 at	1.89769	1.66567	1.77308	1.61245	0.245
(NM) 3840:	f = 1.40717 at	1.89766	1.6657	1.77305	1.61247	0.245
(NM) 3860:	f = 1.40717 at	1.8978	1.66587	1.77317	1.61248	0.245
(NM) 3880:	f = 1.40717 at	1.89776	1.66588	1.77319	1.6126	0.245
(NM) 3900:	f = 1.40717 at	1.8977	1.66589	1.77311	1.61264	0.245
(NM) 3920:	f = 1.40717 at	1.89786	1.66594	1.77321	1.61262	0.245
(NM) 3940:	f = 1.40717 at	1.89777	1.66586	1.77315	1.61256	0.245
(NM) 3960:	f = 1.40717 at	1.89757	1.66571	1.77304	1.6125	
(NM) 3980:	f = 1.40717 at	1.89738	1.66553	1.77297	1.61223	
(NM) 4000:	f = 1.40717 at	1.89739	1.66557	1.77291	1.61225	0.245
(NM) 4020:	f = 1.40717 at	1.89723	1.66537	1.77279	1.61205	0.24
(NM) 4040:	f = 1.40717 at	1.89723	1.66544	1.77279	1.61206	
(NM) 4060:	f = 1.40717 at	1.89714	1.66541	1.77272	1.61191	
(NM) 4080:	f = 1.40717 at	1.89699	1.66536	1.77256	1.61183	0.245
(NM) 4100:	f = 1.40717 at	1.89685	1.66529	1.77237	1.61172	
(NM) 4120:	f = 1.40717 at	1.89688	1.66525	1.77242	1.61175	
(NM) 4140:	f = 1.40717 at	1.89693	1.66522	1.77247	1.61167	
(NM) 4160:	f = 1.40717 at	1.89704	1.66532	1.77252	1.61176	
(NM) 4180:	f = 1.40717 at	1.89698	1.6653	1.7725	1.61172	
(NM) 4200:	f = 1.40717 at	1.89701	1.66532	1.77251	1.61177	
(NM) 4220:	f = 1.40717 at	1.89698	1.66529	1.7725	1.61175	
(NM) 4240:	f = 1.40717 at	1.897	1.66532	1.77252	1.61178	0.245
(NM) 4260:	f = 1.40717 at	1.897	1.66531	1.77252	1.61179	
(NM) 4280:	f = 1.40717 at	1.89699	1.66531	1.77252	1.61179	
(NM) 4300:	f = 1.40717 at	1.897	1.66531	1.77251	1.61177	
(NM) 4320:	f = 1.40717 at	1.897	1.66531	1.77251	1.61177	
(NM) 4340:	f = 1.40717 at	1.897	1.66532	1.77252	1.61179	
(NM) 4360:	f = 1.40717 at	1.89701	1.66532	1.77252	1.61178	
(NM) 4380:	f = 1.40717 at	1.89701	1.66532	1.77252	1.61178	
(NM) 4400:	f = 1.40717 at	1.89702	1.66532	1.77252	1.61178	
(NM) 4420:	f = 1.40717 at	1.89702	1.66532	1.77252	1.61179	

(NM) 4440: f = 1.40717 at	1.89703	1.66532	1.77252	1.61179
(NM) 4460: f = 1.40717 at	1.89705	1.66534	1.77254	1.6118
(NM) 4480: f = 1.40717 at	1.89705	1.66534	1.77254	1.61181
(NM) 4500: f = 1.40717 at	1.89704	1.66534	1.77254	1.6118
(NM) 4520: f = 1.40717 at	1.89703	1.66533	1.77253	1.61179
(NM) 4540: f = 1.40717 at	1.89704	1.66534	1.77254	1.6118
(NM) 4560: f = 1.40717 at	1.89702	1.66532	1.77252	1.6118
(NM) 4580: f = 1.40717 at	1.89703	1.66532	1.77252	1.6118
(NM) 4600: f = 1.40717 at	1.89701	1.66529	1.7725	1.61177 0.245
(NM) 4620: f = 1.40717 at	1.897	1.66529	1.7725	1.61177
(NM) 4640: f = 1.40717 at	1.897	1.66529	1.77249	1.61177
(NM) 4660: f = 1.40717 at	1.89697	1.66527	1.77245	1.61175
(NM) 4680: f = 1.40717 at	1.89697	1.66527	1.77245	1.61175
(NM) 4700: f = 1.40717 at	1.89697	1.66527	1.77245	1.61174
(NM) 4720: f = 1.40717 at	1.89697	1.66527	1.77245	1.61174
(NM) 4740: f = 1.40717 at	1.89697	1.66526	1.77245	1.61173
(NM) 4760: f = 1.40717 at	1.89698	1.66527	1.77245	1.61174
(NM) 4780: f = 1.40717 at	1.89698	1.66526	1.77245	1.61174
(NM) 4800: f = 1.40717 at	1.89699	1.66527	1.77246	1.61174
(NM) 4820: f = 1.40717 at	1.89698	1.66527	1.77246	1.61175
(NM) 4840: f = 1.40717 at	1.89699	1.66527	1.77246	1.61175
(NM) 4860: f = 1.40717 at	1.89698	1.66527	1.77245	1.61175
(NM) 4880: f = 1.40717 at	1.89698	1.66527	1.77245	1.61175
(NM) 4900: f = 1.40717 at	1.89698	1.66527	1.77246	1.61175
(NM) 4920: f = 1.40717 at	1.89698	1.66527	1.77245	1.61175
(NM) 4940: f = 1.40717 at	1.89698	1.66527	1.77246	1.61175
(NM) 4960: f = 1.40717 at	1.89698	1.66527	1.77245	1.61175
(NM) 4980: f = 1.40717 at	1.89698	1.66527	1.77245	1.61175
(NM) 5000: f = 1.40717 at	1.89698	1.66528	1.77245	1.61175
(NM) 5020: f = 1.40717 at	1.89698	1.66527	1.77245	1.61176
(NM) 5040: f = 1.40717 at	1.89698	1.66527	1.77245	1.61175
(NM) 5060: f = 1.40717 at	1.89698	1.66527	1.77245	1.61175
(NM) 5080: f = 1.40717 at	1.89698	1.66526	1.77244	1.61174
(NM) 5100: f = 1.40717 at	1.89698	1.66527	1.77244	1.61175
(NM) 5120: f = 1.40717 at	1.89699	1.66527	1.77244	1.61175
(NM) 5140: f = 1.40717 at	1.897	1.66528	1.77245	1.61175
(NM) 5160: f = 1.40717 at	1.897	1.66529	1.77245	1.61177
(NM) 5180: f = 1.40717 at	1.89701	1.66529	1.77245	1.61177
(NM) 5200: f = 1.40717 at	1.89701	1.66529	1.77245	1.61177
(NM) 5220: f = 1.40717 at	1.897	1.66529	1.77245	1.61176

(NM) 5240: f = 1.40717 at	1.89701	1.66529	1.77246	1.61176
(NM) 5260: f = 1.40717 at	1.89701	1.66529	1.77246	1.61177
(NM) 5280: f = 1.40717 at	1.89701	1.66529	1.77246	1.61177
(NM) 5300: f = 1.40717 at	1.89703	1.6653	1.77247	1.61177
(NM) 5320: f = 1.40717 at	1.89703	1.66531	1.77248	1.61177
(NM) 5340: f = 1.40717 at	1.89703	1.66531	1.77247	1.61177
(NM) 5360: f = 1.40717 at	1.89704	1.66531	1.77249	1.61179
(NM) 5380: f = 1.40717 at	1.89703	1.66531	1.77248	1.61178
(NM) 5400: f = 1.40717 at	1.89703	1.6653	1.77247	1.61177 0.245
(NM) 5420: f = 1.40717 at	1.89703	1.66531	1.77248	1.61178
(NM) 5440: f = 1.40717 at	1.89701	1.6653	1.77247	1.61176
(NM) 5460: f = 1.40717 at	1.89702	1.66532	1.77247	1.61177
(NM) 5480: f = 1.40717 at	1.89702	1.66531	1.77246	1.61177
(NM) 5500: f = 1.40717 at	1.897	1.66531	1.77246	1.61176
(NM) 5520: f = 1.40717 at	1.89701	1.66532	1.77246	1.61178
(NM) 5540: f = 1.40717 at	1.89701	1.66531	1.77245	1.61178
(NM) 5560: f = 1.40717 at	1.897	1.66531	1.77246	1.61178
(NM) 5580: f = 1.40717 at	1.89702	1.66533	1.77251	1.61181
(NM) 5600: f = 1.40717 at	1.89704	1.66533	1.77251	1.61181
(NM) 5620: f = 1.40717 at	1.89705	1.66534	1.77252	1.61182 0.245
(NM) 5640: f = 1.40717 at	1.89705	1.66534	1.77251	1.61184
(NM) 5660: f = 1.40717 at	1.89706	1.66534	1.77251	1.61183
(NM) 5680: f = 1.40717 at	1.89706	1.66534	1.77252	1.61185
(NM) 5700: f = 1.40717 at	1.89706	1.66534	1.77252	1.61184
(NM) 5720: f = 1.40717 at	1.89705	1.66534	1.77252	1.61184
(NM) 5740: f = 1.40717 at	1.89705	1.66533	1.77251	1.61183
(NM) 5760: f = 1.40717 at	1.89705	1.66534	1.77252	1.61184
(NM) 5780: f = 1.40717 at	1.89705	1.66534	1.77252	1.61184
(NM) 5800: f = 1.40717 at	1.89705	1.66534	1.77252	1.61184
(NM) 5820: f = 1.40717 at	1.89705	1.66534	1.77252	1.61184
(NM) 5840: f = 1.40717 at	1.89705	1.66534	1.77252	1.61184
(NM) 5860: f = 1.40717 at	1.89705	1.66534	1.77252	1.61183
(NM) 5880: f = 1.40717 at	1.89705	1.66534	1.77252	1.61184 0.245
(NM) 5900: f = 1.40717 at	1.89701	1.66531	1.77249	1.61181
(NM) 5920: f = 1.40717 at	1.89704	1.66534	1.77251	1.61184
(NM) 5940: f = 1.40717 at	1.89704	1.66534	1.77254	1.61186
(NM) 5960: f = 1.40717 at	1.89706	1.66536	1.77257	1.61189
(NM) 5980: f = 1.40717 at	1.89702	1.66532	1.77257	1.61188
(NM) 6000: f = 1.40717 at	1.897	1.66529	1.77255	1.61183
(NM) 6020: f = 1.40717 at	1.89705	1.66535	1.7726	1.6118

(NM) 6040: f = 1.40717 at	1.89709	1.66539	1.77266	1.61181	
(NM) 6060: f = 1.40717 at	1.8971	1.66539	1.77266	1.61179	
(NM) 6080: f = 1.40717 at	1.89706	1.66536	1.77261	1.61173	
(NM) 6100: f = 1.40717 at	1.89712	1.66539	1.77263	1.61177	
(NM) 6120: f = 1.40717 at	1.89707	1.66535	1.77261	1.61178	
(NM) 6140: f = 1.40717 at	1.89713	1.66538	1.77263	1.6118	
(NM) 6160: f = 1.40717 at	1.89722	1.66545	1.77273	1.61184	
(NM) 6180: f = 1.40717 at	1.89718	1.66538	1.77269	1.61177	
(NM) 6200: f = 1.40717 at	1.89717	1.66536	1.77267	1.61175	
(NM) 6220: f = 1.40717 at	1.89715	1.66536	1.77267	1.61174	
(NM) 6240: f = 1.40717 at	1.89711	1.66531	1.77265	1.61169	0.245
(NM) 6260: f = 1.40717 at	1.89713	1.66532	1.77265	1.61169	
(NM) 6280: f = 1.40717 at	1.89712	1.66529	1.77266	1.6117	
(NM) 6300: f = 1.40717 at	1.89715	1.66532	1.77268	1.61172	
(NM) 6320: f = 1.40717 at	1.8972	1.66534	1.77269	1.61174	
(NM) 6340: f = 1.40717 at	1.89727	1.66536	1.7727	1.61175	
(NM) 6360: f = 1.40717 at	1.89725	1.66532	1.77269	1.61173	
(NM) 6380: f = 1.40717 at	1.89724	1.66535	1.77271	1.61175	
(NM) 6400: f = 1.40717 at	1.89728	1.66537	1.77274	1.61178	
(NM) 6420: f = 1.40717 at	1.89727	1.66537	1.77273	1.61177	
(NM) 6440: f = 1.40717 at	1.89727	1.66537	1.77275	1.61176	
(NM) 6460: f = 1.40717 at	1.89725	1.66533	1.77273	1.61173	
(NM) 6480: f = 1.40717 at	1.89726	1.66534	1.77275	1.61174	
(NM) 6500: f = 1.40717 at	1.89724	1.6653	1.77272	1.6117	
(NM) 6520: f = 1.40717 at	1.89722	1.66527	1.77271	1.6117	0.245
(NM) 6540: f = 1.40717 at	1.89716	1.66522	1.77267	1.61167	0.245
(NM) 6560: f = 1.40717 at	1.89716	1.66523	1.77264	1.61168	0.245
(NM) 6580: f = 1.40717 at	1.89714	1.66519	1.77262	1.6117	0.245
(NM) 6600: f = 1.40717 at	1.89716	1.66521	1.77263	1.61171	0.245
(NM) 6620: f = 1.40717 at	1.89714	1.6652	1.77261	1.6117	0.245
(NM) 6640: f = 1.40717 at	1.8972	1.66525	1.77265	1.61174	0.245
(NM) 6660: f = 1.40717 at	1.89719	1.66524	1.77264	1.61174	0.245
(NM) 6680: f = 1.40717 at	1.89722	1.66527	1.77267	1.61177	0.245
(NM) 6700: f = 1.40717 at	1.8972	1.66524	1.77266	1.61176	0.245
(NM) 6720: f = 1.40717 at	1.89722	1.66524	1.77269	1.61176	0.245
(NM) 6740: f = 1.40717 at	1.8973	1.6653	1.77273	1.61182	0.245
(NM) 6760: f = 1.40717 at	1.89731	1.66529	1.77278	1.61178	0.245
(NM) 6780: f = 1.40717 at	1.89732	1.66532	1.77275	1.61181	0.245
(NM) 6800: f = 1.40717 at	1.89724	1.66527	1.77268	1.61173	0.245
(NM) 6820: f = 1.40717 at	1.89728	1.66528	1.77269	1.61174	0.245

(NM) 6840: f = 1.40717 at	1.89731	1.66522	1.77269	1.61171	0.245
(NM) 6860: f = 1.40717 at	1.89726	1.66519	1.77263	1.61168	0.245
(NM) 6880: f = 1.40717 at	1.89733	1.66523	1.77265	1.61165	0.245
(NM) 6900: f = 1.40717 at	1.89739	1.66521	1.77272	1.61162	0.245
(NM) 6920: f = 1.40717 at	1.89734	1.66511	1.77263	1.61151	0.245
(NM) 6940: f = 1.40717 at	1.89722	1.66508	1.77252	1.61147	0.245
(NM) 6960: f = 1.40717 at	1.89718	1.66503	1.77246	1.61143	
(NM) 6980: f = 1.40717 at	1.89724	1.66501	1.7725	1.61145	0.245
(NM) 7000: f = 1.40717 at	1.89728	1.66506	1.77254	1.61151	0.245
(NM) 7020: f = 1.40717 at	1.89724	1.66504	1.77252	1.61149	0.245
(NM) 7040: f = 1.40717 at	1.89715	1.66503	1.77247	1.61147	
(NM) 7060: f = 1.40716 at	1.89717	1.66508	1.77257	1.61153	
(NM) 7080: f = 1.40716 at	1.8972	1.66511	1.77257	1.61156	0.245
(NM) 7100: f = 1.40716 at	1.89729	1.66522	1.77264	1.61163	
(NM) 7120: f = 1.40716 at	1.89735	1.66526	1.77276	1.61173	
(NM) 7140: f = 1.40716 at	1.89731	1.66522	1.77273	1.61165	
(NM) 7160: f = 1.40716 at	1.89736	1.66523	1.77276	1.61167	
(NM) 7180: f = 1.40716 at	1.8974	1.66525	1.77275	1.61164	
(NM) 7200: f = 1.40716 at	1.8974	1.66525	1.77276	1.61165	
(NM) 7220: f = 1.40716 at	1.89744	1.6653	1.77277	1.61169	
(NM) 7240: f = 1.40716 at	1.89747	1.66529	1.77278	1.61168	
(NM) 7260: f = 1.40716 at	1.89745	1.66525	1.77273	1.61162	
(NM) 7280: f = 1.40716 at	1.89749	1.66528	1.77279	1.6117	
(NM) 7300: f = 1.40716 at	1.89754	1.6653	1.77287	1.61172	
(NM) 7320: f = 1.40716 at	1.89758	1.66534	1.77292	1.61177	
(NM) 7340: f = 1.40716 at	1.89756	1.66539	1.77291	1.61177	0.245
(NM) 7360: f = 1.40716 at	1.89741	1.66532	1.77285	1.61167	0.245
(NM) 7380: f = 1.40716 at	1.8974	1.66536	1.77286	1.61182	
(NM) 7400: f = 1.40716 at	1.89723	1.66539	1.77283	1.61175	
(NM) 7420: f = 1.40716 at	1.8974	1.66546	1.77288	1.61184	0.24
(NM) 7440: f = 1.40716 at	1.89741	1.66554	1.77297	1.61192	
(NM) 7460: f = 1.40716 at	1.89749	1.66565	1.77299	1.61202	
(NM) 7480: f = 1.40716 at	1.89752	1.66569	1.77306	1.61206	0.245
(NM) 7500: f = 1.40716 at	1.89756	1.66571	1.77306	1.61204	
(NM) 7520: f = 1.40716 at	1.89747	1.66578	1.77298	1.61198	0.246
(NM) 7540: f = 1.40716 at	1.89744	1.66565	1.7729	1.61172	0.246
(NM) 7560: f = 1.40716 at	1.89734	1.66547	1.77271	1.61159	0.246
(NM) 7580: f = 1.40716 at	1.89734	1.66547	1.77271	1.61159	0.246
(NM) 7600: f = 1.40716 at	1.89744	1.66554	1.77284	1.61169	0.246
(NM) 7620: f = 1.40716 at	1.8974	1.66549	1.77274	1.61164	0.246

(NM) 7640:	f = 1.40716 at	1.89731	1.66542	1.77268	1.61155	0.246
(NM) 7660:	f = 1.40716 at	1.89732	1.66543	1.7727	1.6116	0.246
(NM) 7680:	f = 1.40716 at	1.89729	1.66546	1.7727	1.61162	0.246
(NM) 7700:	f = 1.40716 at	1.8973	1.66543	1.7727	1.6116	0.246
(NM) 7720:	f = 1.40716 at	1.89736	1.66547	1.77272	1.61164	0.246
(NM) 7740:	f = 1.40716 at	1.89736	1.66548	1.77273	1.61166	0.246
(NM) 7760:	f = 1.40716 at	1.89734	1.6655	1.77273	1.61166	0.246
(NM) 7780:	f = 1.40716 at	1.89741	1.66553	1.77275	1.61169	0.246
(NM) 7800:	f = 1.40716 at	1.8974	1.66557	1.77274	1.61173	0.246
(NM) 7820:	f = 1.40716 at	1.89738	1.66553	1.7727	1.61169	0.246
(NM) 7840:	f = 1.40716 at	1.89737	1.66545	1.77265	1.6116	0.246
(NM) 7860:	f = 1.40716 at	1.89746	1.66552	1.77269	1.61168	0.246
(NM) 7880:	f = 1.40716 at	1.89739	1.66546	1.77265	1.61166	0.246
(NM) 7900:	f = 1.40716 at	1.89757	1.66565	1.77281	1.61182	0.246
(NM) 7920:	f = 1.40716 at	1.89754	1.6656	1.77279	1.61178	0.246
(NM) 7940:	f = 1.40716 at	1.89756	1.66571	1.77285	1.61185	0.246
(NM) 7960:	f = 1.40716 at	1.89759	1.66567	1.77291	1.61178	0.246
(NM) 7980:	f = 1.40716 at	1.89774	1.66576	1.773	1.61185	0.246
(NM) 8000:	f = 1.40716 at	1.89763	1.66569	1.77298	1.6118	0.246
(NM) 8020:	f = 1.40716 at	1.89766	1.66574	1.77299	1.61185	0.246
(NM) 8040:	f = 1.40716 at	1.89767	1.66572	1.773	1.61185	0.246
(NM) 8060:	f = 1.40716 at	1.89768	1.66576	1.77304	1.6119	0.246
(NM) 8080:	f = 1.40716 at	1.89767	1.66576	1.77304	1.61191	0.246
(NM) 8100:	f = 1.40716 at	1.8977	1.66575	1.77305	1.61198	0.246
(NM) 8120:	f = 1.40716 at	1.8977	1.66575	1.77305	1.61198	0.246
(NM) 8140:	f = 1.40716 at	1.89769	1.66573	1.77303	1.61198	0.246
(NM) 8160:	f = 1.40716 at	1.8977	1.6657	1.77302	1.61195	0.246
(NM) 8180:	f = 1.40716 at	1.89765	1.66568	1.773	1.61193	0.246
(NM) 8200:	f = 1.40716 at	1.8977	1.66571	1.77303	1.61195	0.246
(NM) 8220:	f = 1.40716 at	1.89767	1.66567	1.77301	1.61193	0.246
(NM) 8240:	f = 1.40716 at	1.8977	1.6657	1.77304	1.61195	0.246
(NM) 8260:	f = 1.40716 at	1.89769	1.66569	1.77303	1.61195	0.246
(NM) 8280:	f = 1.40716 at	1.8977	1.6657	1.77303	1.61195	0.246
(NM) 8300:	f = 1.40716 at	1.8977	1.6657	1.77304	1.61196	0.246
(NM) 8320:	f = 1.40716 at	1.8977	1.66571	1.77304	1.61196	0.246
(NM) 8340:	f = 1.40716 at	1.89769	1.6657	1.77303	1.61195	0.246
(NM) 8360:	f = 1.40716 at	1.8977	1.66571	1.77305	1.61197	0.246
(NM) 8380:	f = 1.40716 at	1.8977	1.6657	1.77304	1.61196	0.246
(NM) 8400:	f = 1.40716 at	1.89769	1.66571	1.77303	1.61196	0.246
(NM) 8420:	f = 1.40716 at	1.89769	1.6657	1.77304	1.61197	0.246

(NM) 8440: f = 1.40716 at	1.89769	1.66571	1.77304	1.61197	0.246
(NM) 8460: f = 1.40716 at	1.89772	1.66572	1.77305	1.61199	0.246
(NM) 8480: f = 1.40716 at	1.89769	1.66571	1.77304	1.61198	0.246
(NM) 8500: f = 1.40716 at	1.89768	1.6657	1.77304	1.61198	0.246
(NM) 8520: f = 1.40716 at	1.89769	1.66571	1.77304	1.61199	0.246
(NM) 8540: f = 1.40716 at	1.89769	1.6657	1.77304	1.61198	0.246
(NM) 8560: f = 1.40716 at	1.89768	1.6657	1.77304	1.61198	0.246
(NM) 8580: f = 1.40716 at	1.89768	1.6657	1.77304	1.61198	0.246
(NM) 8600: f = 1.40716 at	1.89767	1.6657	1.77304	1.61197	0.246
(NM) 8620: f = 1.40716 at	1.89768	1.66569	1.77304	1.61196	0.246
(NM) 8640: f = 1.40716 at	1.89768	1.66569	1.77305	1.61196	0.246
(NM) 8660: f = 1.40716 at	1.89768	1.6657	1.77305	1.61197	0.246
(NM) 8680: f = 1.40716 at	1.89766	1.66568	1.77304	1.61196	0.246
(NM) 8700: f = 1.40716 at	1.89767	1.66567	1.77303	1.61196	0.24
(NM) 8720: f = 1.40716 at	1.89765	1.66567	1.77302	1.61197	0.246
(NM) 8740: f = 1.40716 at	1.89765	1.66566	1.77302	1.61196	0.246
(NM) 8760: f = 1.40716 at	1.89765	1.66566	1.77302	1.61196	0.246
(NM) 8780: f = 1.40716 at	1.89764	1.66564	1.77301	1.61194	0.246
(NM) 8800: f = 1.40716 at	1.89762	1.66563	1.77298	1.61193	0.246
(NM) 8820: f = 1.40716 at	1.89759	1.6656	1.77294	1.61189	0.246
(NM) 8840: f = 1.40716 at	1.89759	1.66561	1.77295	1.6119	0.246886
(NM) 8860: f = 1.40716 at	1.89758	1.66558	1.77293	1.61188	0.246
(NM) 8880: f = 1.40716 at	1.89757	1.66558	1.77291	1.61186	0.246
(NM) 8900: f = 1.40716 at	1.89759	1.66558	1.77291	1.61186	0.246
(NM) 8920: f = 1.40716 at	1.89759	1.66559	1.77293	1.61186	0.246
(NM) 8940: f = 1.40716 at	1.89759	1.6656	1.77293	1.61187	0.246
(NM) 8960: f = 1.40716 at	1.8976	1.6656	1.77292	1.61187	0.246
(NM) 8980: f = 1.40716 at	1.89759	1.66559	1.77291	1.61187	0.246
(NM) 9000: f = 1.40716 at	1.89761	1.66561	1.77293	1.61188	0.246901
(NM) 9020: f = 1.40716 at	1.89762	1.66561	1.77292	1.61188	0.246
(NM) 9040: f = 1.40716 at	1.89761	1.66559	1.77289	1.61186	0.246
(NM) 9060: f = 1.40716 at	1.89759	1.66559	1.7729	1.61187	0.246
(NM) 9080: f = 1.40716 at	1.89751	1.66553	1.77283	1.61179	0.246
(NM) 9100: f = 1.40716 at	1.89752	1.66554	1.77286	1.61183	0.246
(NM) 9120: f = 1.40716 at	1.89755	1.66557	1.77286	1.61184	0.247
(NM) 9140: f = 1.40715 at	1.89756	1.66556	1.77289	1.6119	0.246
(NM) 9160: f = 1.40715 at	1.89762	1.66556	1.77287	1.61192	0.247
(NM) 9180: f = 1.40715 at	1.8976	1.66553	1.77286	1.61193	0.247
(NM) 9200: f = 1.40715 at	1.89763	1.66551	1.77285	1.61193	0.247
(NM) 9220: f = 1.40715 at	1.89759	1.66548	1.77279	1.61188	0.246



(NM) 9240: f = 1.40715 at	1.89754	1.66537	1.77273	1.61185	0.246
(NM) 9260: f = 1.40715 at	1.89756	1.66539	1.77278	1.61188	0.246
(NM) 9280: f = 1.40715 at	1.89763	1.66542	1.77283	1.61193	0.246
(NM) 9300: f = 1.40715 at	1.89771	1.66545	1.77287	1.61195	0.247
(NM) 9320: f = 1.40715 at	1.89776	1.66548	1.77291	1.61197	0.247
(NM) 9340: f = 1.40715 at	1.89776	1.66541	1.77289	1.61193	0.247
(NM) 9360: f = 1.40715 at	1.89773	1.66537	1.77284	1.61189	0.247
(NM) 9380: f = 1.40715 at	1.89772	1.66537	1.77282	1.61187	0.247
(NM) 9400: f = 1.40715 at	1.89768	1.66534	1.7728	1.61186	0.247
(NM) 9420: f = 1.40715 at	1.89759	1.66526	1.7727	1.61179	0.247
(NM) 9440: f = 1.40715 at	1.89766	1.66527	1.77268	1.61183	0.247
(NM) 9460: f = 1.40715 at	1.89766	1.66526	1.77263	1.61187	0.247
(NM) 9480: f = 1.40715 at	1.8977	1.66529	1.77264	1.61191	0.247
(NM) 9500: f = 1.40715 at	1.89776	1.66534	1.77266	1.61199	0.247
(NM) 9520: f = 1.40715 at	1.89769	1.66543	1.77272	1.61209	0.247
(NM) 9540: f = 1.40715 at	1.89776	1.66538	1.77268	1.61206	0.247
(NM) 9560: f = 1.40715 at	1.89779	1.66544	1.77269	1.61207	0.247
(NM) 9580: f = 1.40715 at	1.89783	1.66547	1.7727	1.61206	0.247
(NM) 9600: f = 1.40715 at	1.89777	1.66547	1.77265	1.612	0.247
(NM) 9620: f = 1.40715 at	1.89782	1.66551	1.77269	1.61207	0.247
(NM) 9640: f = 1.40715 at	1.89777	1.66548	1.77262	1.61201	0.247
(NM) 9660: f = 1.40715 at	1.89774	1.66545	1.77262	1.61202	0.247
(NM) 9680: f = 1.40715 at	1.89771	1.6654	1.77259	1.61197	0.247
(NM) 9700: f = 1.40715 at	1.89769	1.66542	1.77261	1.61195	0.247
(NM) 9720: f = 1.40715 at	1.89769	1.66538	1.7726	1.61195	0.247
(NM) 9740: f = 1.40715 at	1.89761	1.66538	1.77262	1.61192	0.247
(NM) 9760: f = 1.40715 at	1.89768	1.66544	1.77265	1.61204	0.247
(NM) 9780: f = 1.40715 at	1.89759	1.66532	1.77266	1.612	0.247
(NM) 9800: f = 1.40715 at	1.89761	1.6654	1.77268	1.61206	0.247
(NM) 9820: f = 1.40715 at	1.89744	1.66538	1.77258	1.61206	
(NM) 9840: f = 1.40715 at	1.89719	1.66536	1.77257	1.61205	
(NM) 9860: f = 1.40715 at	1.89726	1.66533	1.77241	1.61197	
(NM) 9880: f = 1.40715 at	1.89734	1.66533	1.77245	1.61203	
(NM) 9900: f = 1.40715 at	1.8972	1.6653	1.77242	1.61202	
(NM) 9920: f = 1.40715 at	1.89724	1.66524	1.7724	1.61199	
(NM) 9940: f = 1.40715 at	1.89722	1.66522	1.77238	1.612	
(NM) 9960: f = 1.40715 at	1.89726	1.66522	1.77241	1.612	
(NM) 9980: f = 1.40715 at	1.89724	1.66523	1.7724	1.61198	
(NM) 10000: f = 1.40715 at	1.89725	1.66521	1.77239	1.61198	

Linear mixed model fit by REML ['lmerMod']

Formula: Adj ~ Trt + (Trt - 1 | Location) + (1 | Block)

Data: Multilocation

REML criterion at convergence: 1.4071

Random effects:

Groups	Name	Std.Dev.	Corr
Location	Trt1	0.3687	
	Trt2	0.3271	0.99
	Trt3	0.3451	1.00 1.00
	Trt4	0.3378	0.93 0.97 0.95
Block	(Intercept)	0.0000	
Residual		0.1943	

Number of obs: 108, groups: Location, 9; Block, 3

Fixed Effects:

(Intercept)	Trt1	Trt2	Trt3
2.86567	0.05834	-0.18802	0.08379

## I PBIB

```
> str(PBIB)
```

```
'data.frame':      60 obs. of  3 variables:
```

```
$ response : num  2.4 2.5 2.6 2 2.7 2.8 2.4 2.7 2.6 2.8 ...
```

```
$ Treatment: Factor w/ 15 levels "1","10","11",...: 7 15 1 5 11 13 14 1 2 1 ...
```

```
$ Block : Factor w/ 15 levels "1","10","11",...: 1 1 1 1 8 8 8 8 9 9 ...
```

```
- attr(*, "ginfo")=List of 7
```

```
..$ formula :Class 'formula' length 3 response ~ Treatment | Block
```

```
.. .. - attr(*, ".Environment")=<environment: R_GlobalEnv>
```

```
..$ order.groups: logi TRUE
```

```
..$ FUN :function (x)
```

```
..$ outer : NULL
```

```
..$ inner : NULL
```

```
..$ labels : list()
```

```
..$ units : list()
```

```
> ## compare with output 1.7 pp. 24-25
```

```
> (fm1PBIB <- lmer(response ~ Treatment + (1 | Block), PBIB))
```

Linear mixed model fit by REML ['lmerMod']

Formula: response ~ Treatment + (1 | Block)

Data: PBIB

REML criterion at convergence: 51.9849

Random effects:

Groups	Name	Std.Dev.
--------	------	----------

```

Block      (Intercept) 0.2157
Residual                    0.2925
Number of obs: 60, groups: Block, 15
Fixed Effects:
(Intercept)  Treatment1  Treatment10  Treatment11  Treatment12
  2.891311    -0.073789    -0.400250     0.007387     0.161510
Treatment13  Treatment14  Treatment15  Treatment2  Treatment3
 -0.273542    -0.400000    -0.032078    -0.485996    -0.436368
Treatment4   Treatment5   Treatment6   Treatment7   Treatment8
 -0.107482    -0.086413     0.019382    -0.102327    -0.109706

```

## J SIMS

```

> str(SIMS)
'data.frame':      3691 obs. of  3 variables:
 $ Pretot: num  29 38 31 31 29 23 23 33 30 32 ...
 $ Gain   : num  2 0 6 6 5 9 7 2 1 3 ...
 $ Class  : Factor w/ 190 levels "1","10","100",...: 1 1 1 1 1 1 1 1 1 1 ...
- attr(*, "ginfo")=List of 7
 ..$ formula      :Class 'formula' length 3 Gain ~ Pretot | Class
 .. ..- attr(*, ".Environment")=<environment: R_GlobalEnv>
 ..$ order.groups: logi TRUE
 ..$ FUN          :function (x)
 ..$ outer        : NULL
 ..$ inner        : NULL
 ..$ labels       :List of 2
 .. ..$ Pretot: chr "Sum of pre-test core item scores"
 .. ..$ Gain   : chr "Gain in mathematics achievement score"
 ..$ units       : list()
> ## compare to output 7.4, p. 262
> (fmlSIMS <- lmer(Gain ~ Pretot + (Pretot | Class), SIMS))
Linear mixed model fit by REML ['lmerMod']
Formula: Gain ~ Pretot + (Pretot | Class)
Data: SIMS
REML criterion at convergence: 22380.57
Random effects:
Groups   Name             Std.Dev. Corr
Class    (Intercept)  3.80651
         Pretot      0.09593  -0.64
Residual                    4.71548

```

Number of obs: 3691, groups: Class, 190

Fixed Effects:

(Intercept)	Pretot
7.060	-0.186