

Introduction to `rstpm2`

Mark Clements

September 20, 2012

1 Cure model

For cure, we use the melanoma dataset used by Andersson and colleagues for cure models with Stata's `stpm2` (see <http://www.pauldickman.com/survival/>).

Initially, we merge the patient data with the all cause mortality rates.

```
> require(rstpm2)
> data(popmort)
> data(colon)
> popmort2 <- transform(popmort, exitage=age, exityear=year, age=NULL, year=NULL)
> colon2 <- within(colon, {
+   status <- ifelse(surv_mm>120.5, 1, status)
+   tm <- pmin(surv_mm, 120.5)/12
+   exit <- dx+tm*365.25
+   sex <- as.numeric(sex)
+   exitage <- pmin(floor(age+tm), 99)
+   exityear <- floor(yydx+tm)
+ })
> colon2 <- merge(colon2, popmort2)
```

For comparisons, we fit the relative survival model without and with cure.

```
> summary(fit0 <- stpm2(Surv(tm, status %in% 2:3)~I(year8594=="Diagnosed 85-94"),
+                       data=colon2,
+                       bhazard=colon2$rate, df=5))

> summary(fit <- stpm2(Surv(tm, status %in% 2:3)~I(year8594=="Diagnosed 85-94"),
+                      data=colon2,
+                      bhazard=colon2$rate,
+                      df=5, cure=TRUE))
```

Maximum likelihood estimation

Call:

```
mle2(minuslogl = negll, start = init, vecpar = TRUE, control = list(
  parscale = c(0.1, 0.1, 0.1, 0.1, 0.1, 0.1, 0.1), maxit = 300),
  lower = -Inf, upper = Inf)
```

Coefficients:

Estimate Std. Error z value

```

(Intercept) -3.977326 0.054778 -72.6078
I(year8594 == "Diagnosed 85-94")TRUE -0.155613 0.025102 -6.1992
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)1 3.323188 0.053166 62.5064
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)2 3.628637 0.053160 68.2594
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)3 1.634854 0.022465 72.7748
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)4 6.592018 0.111504 59.1193
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)5 3.371806 0.042788 78.8029
Pr(z)
(Intercept) < 2.2e-16 ***
I(year8594 == "Diagnosed 85-94")TRUE 5.675e-10 ***
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)1 < 2.2e-16 ***
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)2 < 2.2e-16 ***
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)3 < 2.2e-16 ***
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)4 < 2.2e-16 ***
nsx(log(tm), df = 5, cure = TRUE, log = TRUE)5 < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

-2 log L: 42190.77

```
> predict(fit, head(colon2), se.fit=TRUE)
```

	Estimate	lower	upper
1	0.8610835	0.8542896	0.8675857
2	0.7934666	0.7850101	0.8016338
3	0.6967422	0.6863187	0.7068971
4	0.8610835	0.8542896	0.8675857
5	0.8221253	0.8143224	0.8296357
6	0.8610835	0.8542896	0.8675857

The estimate for the year parameter from the model without cure is within three significant figures with that in Stata. For the predictions, the Stata model gives:

	surv	surv_lci	surv_uci
1.	.86108264	.8542898	.8675839
2.	.79346526	.7850106	.8016309
3.	.69674037	.6863196	.7068927
4.	.86108264	.8542898	.8675839
5.	.82212425	.8143227	.8296332
6.	.86108264	.8542898	.8675839

We can estimate the proportion of failures at the end of follow-up using:

```

> newdata.eof <- data.frame(year8594 = unique(colon2$year8594),
+                             tm=max(colon2$tm))
> 1-predict(fit0, newdata.eof, type="surv", se.fit=TRUE)

```


	Estimate	lower	upper
1	0.6063672	0.6211742	0.5916005
2	0.5515138	0.5661344	0.5370100

```
> 1-predict(fit, newdata.eof, type="surv", se.fit=TRUE)
```

	Estimate	lower	upper
1	0.5913307	0.6055051	0.5772138
2	0.5350813	0.5485401	0.5217406

We can plot the predicted survival estimates:

```
> plot(fit0,newdata=data.frame(year8594 = "Diagnosed 85-94"), ylim=0:1,
+       xlab="Time since diagnosis (years)")
> plot(fit0,newdata=data.frame(year8594 = "Diagnosed 75-84"),
+       add=TRUE,line.col="red",rug=FALSE)
> plot(fit,newdata=data.frame(year8594 = "Diagnosed 85-94"),
+       add=TRUE,ci=FALSE,lty=2,rug=FALSE)
> plot(fit,newdata=data.frame(year8594="Diagnosed 75-84"),
+       add=TRUE,rug=FALSE,line.col="red",ci=FALSE,lty=2)
> legend("topright",c("85-94 without cure","75-84 without cure",
+                     "85-94 with cure","75-84 with cure"),
+       col=c(1,2,1,2), lty=c(1,1,2,2), bty="n")
```



And the hazard curves:

```
> plot(fit0,newdata=data.frame(year8594 = "Diagnosed 85-94"),
+       ylim=c(0,0.5), type="hazard",
```

```

+       xlab="Time since diagnosis (years)")
> plot(fit0,newdata=data.frame(year8594 = "Diagnosed 75-84"),
+       type="hazard",
+       add=TRUE,line.col="red",rug=FALSE)
> plot(fit,newdata=data.frame(year8594 = "Diagnosed 85-94"),
+       type="hazard",
+       add=TRUE,ci=FALSE,lty=2,rug=FALSE)
> plot(fit,newdata=data.frame(year8594="Diagnosed 75-84"),
+       type="hazard",
+       add=TRUE,rug=FALSE,line.col="red",ci=FALSE,lty=2)
> legend("topright",c("85-94 without cure","75-84 without cure",
+       "85-94 with cure","75-84 with cure"),
+       col=c(1,2,1,2), lty=c(1,1,2,2), bty="n")

```

