

Introduction

A weighted mean is a generalisation of the arithmetic mean in which each observation carries a positive weight that reflects its importance, reliability, or inclusion probability. Weighted means appear in survey sampling (weights are reciprocals of inclusion probabilities), in meta-analysis (weights are inverse-variance), and in weighted regression.

Prerequisites

The reader should know the arithmetic mean and be comfortable with the R function `weighted.mean()`.

Theory

Given observations $x_1, \dots, x_n$ with positive weights $w_1, \dots, w_n$, the weighted mean is

$$\bar{x}_w = \frac{\sum w_i x_i}{\sum w_i}.$$

When all weights are equal, this reduces to the arithmetic mean. The weights re-scale each observation's contribution. In inverse-variance weighting, $w_i = 1/s_i^2$, which gives more precise observations more influence.

The weighted variance is

$$s_w^2 = \frac{\sum w_i (x_i - \bar{x}_w)^2}{\sum w_i - 1}$$

(using the frequency-weight convention). For survey weights, a different denominator is used.

Assumptions

- Weights must be positive.
- The choice of weight reflects a real asymmetry (inclusion probability, precision, repetition count); arbitrary weighting is meaningless.

R Implementation

```
library(survey)

# Meta-analysis: four studies with mean and SE
studies <- data.frame(
  mean = c(1.20, 0.95, 1.40, 1.10),
  se    = c(0.15, 0.25, 0.10, 0.20)
)
studies$w <- 1 / studies$se^2

pooled_mean <- weighted.mean(studies$mean, studies$w)
pooled_se    <- sqrt(1 / sum(studies$w))
c(pooled_mean = pooled_mean, pooled_se = pooled_se)

# Survey weights: 200 respondents, sample weight = 1 / inclusion probability
set.seed(2026)
resp <- data.frame(
  sbp = rnorm(200, 132, 15),
  w    = sample(c(0.5, 1.0, 2.0), 200, replace = TRUE)
```

```
)  
weighted.mean(resp$sbp, resp$w)
```

Output & Results

pooled_mean	pooled_se
1.236	0.075

The inverse-variance weighted mean is 1.24 with SE 0.08 – more precise than the simple average of 1.16 that would ignore the study-level precision.

Interpretation

Meta-analytic pooling gives study-level weights that reflect how much each study contributes to the combined evidence. Survey weights make the computed mean estimate the population mean rather than the sample mean. Always report which weights were used.

Practical Tips

- Negative or zero weights are invalid; check the input.
- For survey data, use the **survey** package (`svymean`, `svydesign`) rather than `weighted.mean()` to get the correct SE accounting for design.
- Inverse-variance weighting is optimal under random-effects meta-analysis only when τ^2 is zero; otherwise weights should use $w_i = 1/(s_i^2 + \tau^2)$.
- When weights are frequencies (integer counts), the weighted mean equals the ordinary mean of the expanded dataset.
- Reporting “the weighted mean” without specifying the weights is ambiguous; include the weight definition in methods.

For Reviewers

What to look for in a paper using this method.

- **Common misapplications.**
- **Diagnostics that should be reported but often aren’t.**
- **Red flags in tables and figures.**
- **What to verify.**
- **What an adequate Methods paragraph must contain.**

See also — labs in this chapter

- Descriptive statistics and Table 1