

8. Nash Equilibrium

Definition, existence, computation, and multiplicity of Nash equilibrium — the central solution concept in non-cooperative game theory.

9. Nash Equilibrium

Learning objectives

- Define Nash equilibrium in terms of best responses and unilateral deviations.
- State Nash's existence theorem and understand why mixed strategies guarantee existence.
- Compute pure-strategy and mixed-strategy Nash equilibria for 2×2 games in R.
- Visualize best-response correspondences and identify equilibria graphically.

9.1. Motivation

Consider two competing firms choosing prices simultaneously. Each firm wants to maximize profit, but profit depends on the rival's price. Firm A cannot simply optimize in isolation — it must reason about Firm B's choice, knowing that Firm B is reasoning about Firm A in exactly the same way.

This mutual consistency requirement is the heart of Nash equilibrium: a strategy profile where no player can improve their payoff by changing their strategy alone, given what everyone else is doing. It is the most widely used solution concept in non-cooperative game theory, and the starting point for almost every applied analysis of strategic interaction.

9.2. Theory

9.2.1. Best responses

Given a finite game $G = (N, (A_i)_{i \in N}, (u_i)_{i \in N})$ with player set N , action sets A_i , and payoff functions u_i , player i 's **best response** to a strategy profile a_{-i} of the other players is any action that maximizes i 's payoff:

$$BR_i(a_{-i}) = \arg \max_{a_i \in A_i} u_i(a_i, a_{-i}) \quad (9.1)$$

9.2.2. Nash equilibrium defined

i Definition: Nash Equilibrium

A strategy profile $a^* = (a_1^*, \dots, a_n^*)$ is a **Nash equilibrium** if every player's strategy is a best response to the others:

$$u_i(a_i^*, a_{-i}^*) \geq u_i(a_i, a_{-i}^*) \quad \forall a_i \in A_i, \forall i \in N \quad (9.2)$$

Equivalently, $a_i^* \in BR_i(a_{-i}^*)$ for all i .

The definition captures a stability property: at a Nash equilibrium, no player has a profitable **unilateral** deviation. This does not mean the outcome is socially optimal (the Prisoner's Dilemma shows otherwise), nor that players coordinate easily — it means the profile is self-enforcing once reached.

9.2.3. Existence

Nash (1950) proved that every finite game — one with finitely many players and finitely many actions per player — has at least one Nash equilibrium, possibly in mixed strategies. The proof uses Kakutani's fixed-point theorem applied to the best-response correspondence.

 **Theorem: Nash's Existence Theorem**

Every finite game has at least one Nash equilibrium in mixed strategies.

The theorem does not guarantee a *pure-strategy* equilibrium. The game Matching Pennies, for instance, has no pure-strategy equilibrium but has a unique mixed equilibrium where each player randomizes 50-50 (see Chapter 11).

9.2.4. Multiplicity

Games can have zero, one, or many pure-strategy equilibria:

- **Matching Pennies** — zero pure, one mixed.
- **Battle of the Sexes** — two pure, one mixed.
- **Prisoner's Dilemma** — one pure (Defect, Defect), which is also the unique NE overall.

When multiple equilibria exist, selecting among them is a separate problem — the **equilibrium selection** problem — addressed by refinements such as subgame perfection (Chapter 13), trembling-hand perfection, and risk dominance.

9.3. Implementation in R

We implement two approaches: a pure-R solver for 2×2 games and a visualization of best-response correspondences.

9.3.1. Finding pure-strategy Nash equilibria

```
source(here::here("R", "solvers.R"))

# Prisoner's Dilemma payoff matrices
# Rows = Player 1 actions (C, D), Cols = Player 2 actions (C, D)
A <- matrix(c(3, 0, 5, 1), nrow = 2, byrow = TRUE,
            dimnames = list(c("C", "D"), c("C", "D")))
B <- matrix(c(3, 5, 0, 1), nrow = 2, byrow = TRUE,
```

```
dimnames = list(c("C", "D"), c("C", "D")))

cat("Player 1 payoffs:\n")
```

```
#> Player 1 payoffs:
```

```
A
```

```
#>   C D
#> C 3 0
#> D 5 1
```

```
cat("\nPlayer 2 payoffs:\n")
```

```
#>
```

```
#> Player 2 payoffs:
```

```
B
```

```
#>   C D
#> C 3 5
#> D 0 1
```

```
pure_ne <- solve_2x2_pure_nash(A, B)
cat("\nPure-strategy Nash equilibria:\n")
```

```
#>
```

```
#> Pure-strategy Nash equilibria:
```

```
for (eq in pure_ne) {
  cat(sprintf(" (%s, %s) with payoffs (%d, %d)\n",
              rownames(A)[eq[1]], colnames(A)[eq[2]],
              A[eq[1], eq[2]], B[eq[1], eq[2]]))
}
```

```
#> (D, D) with payoffs (1, 1)
```

9.3.2. Battle of the Sexes — multiple equilibria

```
# Battle of the Sexes
A_bos <- matrix(c(3, 0, 0, 2), nrow = 2, byrow = TRUE,
                dimnames = list(c("Opera", "Football"), c("Opera", "Football")))
B_bos <- matrix(c(2, 0, 0, 3), nrow = 2, byrow = TRUE,
                dimnames = list(c("Opera", "Football"), c("Opera", "Football")))

cat("Battle of the Sexes:\n")
```

9. Nash Equilibrium

#> Battle of the Sexes:

```
cat("Player 1 payoffs:\n")
```

#> Player 1 payoffs:

```
A_bos
```

```
#>      Opera Football
#> Opera      3      0
#> Football   0      2
```

```
cat("\nPlayer 2 payoffs:\n")
```

#>

#> Player 2 payoffs:

```
B_bos
```

```
#>      Opera Football
#> Opera      2      0
#> Football   0      3
```

```
pure_bos <- solve_2x2_pure_nash(A_bos, B_bos)
mixed_bos <- solve_2x2_mixed_nash(A_bos, B_bos)
```

```
cat("\nPure-strategy NE:\n")
```

#>

#> Pure-strategy NE:

```
for (eq in pure_bos) {
  cat(sprintf("  (%s, %s)\n", rownames(A_bos)[eq[1]], colnames(A_bos)[eq[2]]))
}
```

#> (Opera, Opera)

#> (Football, Football)

```
cat(sprintf("\nMixed-strategy NE: p = %.2f, q = %.2f\n",
  mixed_bos$p, mixed_bos$q))
```

#>

#> Mixed-strategy NE: p = 0.60, q = 0.40

9.3.3. Best-response correspondence plot

```

# For a 2x2 game, let p = P(Player 1 plays row 1), q = P(Player 2 plays col 1)
# Player 2's expected payoff from col 1: p * B[1,1] + (1-p) * B[2,1]
# Player 2's expected payoff from col 2: p * B[1,2] + (1-p) * B[2,2]
# Player 2 is indifferent when these are equal

p_seq <- seq(0, 1, length.out = 200)

# Player 1's best response: BR1(q)
# EU1(row1) = q*A[1,1] + (1-q)*A[1,2] = q*3
# EU1(row2) = q*A[2,1] + (1-q)*A[2,2] = (1-q)*2
# Indifferent when 3q = 2(1-q) => q = 2/5
br1_q <- 2 / (3 + 2) # q at which P1 is indifferent

# Player 2's best response: BR2(p)
# EU2(col1) = p*B[1,1] + (1-p)*B[2,1] = 2p
# EU2(col2) = p*B[1,2] + (1-p)*B[2,2] = 3(1-p)
# Indifferent when 2p = 3(1-p) => p = 3/5
br2_p <- 3 / (2 + 3) # p at which P2 is indifferent

# Build best-response data
br_data <- tibble(
  q = c(0, br1_q, br1_q, br1_q, 1),
  p_br1 = c(0, 0, 0.5, 1, 1),
  type = "Player 1 BR"
)

br2_data <- tibble(
  p = c(0, br2_p, br2_p, br2_p, 1),
  q_br2 = c(0, 0, 0.5, 1, 1),
  type = "Player 2 BR"
)

# Equilibrium points
eq_points <- tibble(
  p = c(0, br2_p, 1),
  q = c(0, br1_q, 1),
  label = c("(Football, Football)", "Mixed NE", "(Opera, Opera)")
)

p_fig <- ggplot() +
  geom_path(data = br_data, aes(x = q, y = p_br1),
    colour = okabe_ito[1], linewidth = 1.2) +
  geom_path(data = br2_data, aes(x = q_br2, y = p),
    colour = okabe_ito[2], linewidth = 1.2) +
  geom_point(data = eq_points, aes(x = q, y = p),
    size = 3, colour = okabe_ito[6]) +
  geom_text(data = eq_points, aes(x = q, y = p, label = label),
    vjust = -1, size = 3) +
  annotate("text", x = 0.85, y = 0.15, label = "BR (q)",
    colour = okabe_ito[1], fontface = "bold", size = 4) +

```

9. Nash Equilibrium

```
annotate("text", x = 0.15, y = 0.85, label = "BR (p)",
        colour = okabe_ito[2], fontface = "bold", size = 4) +
scale_x_continuous(name = "q (Player 2: prob. of Opera)",
                  limits = c(-0.05, 1.15), breaks = seq(0, 1, 0.2)) +
scale_y_continuous(name = "p (Player 1: prob. of Opera)",
                  limits = c(-0.05, 1.25), breaks = seq(0, 1, 0.2)) +
theme_publication() +
labs(title = "Best-Response Correspondences: Battle of the Sexes")

p_fig
save_pub_fig(p_fig, "best-response-bos")
```




Figure 9.1.: Best-response correspondences for the Battle of the Sexes. Player 1's best response (orange) and Player 2's best response (blue) intersect at three Nash equilibria: two pure-strategy and one mixed.

9.4. Worked example

Consider a coordination game where two drivers approach each other on a road and must choose to swerve Left or Right:

```
A_coord <- matrix(c(1, 0, 0, 1), nrow = 2, byrow = TRUE,
                dimnames = list(c("Left", "Right"), c("Left", "Right")))
B_coord <- A_coord # symmetric game

cat("Coordination Game (symmetric):\n")
```

```
#> Coordination Game (symmetric):
```

```
A_coord
```

```
#>      Left Right
#> Left    1    0
#> Right   0    1
```

```
eq_coord <- support_enumeration(A_coord, B_coord)
```

```
cat("\nPure-strategy NE:\n")
```

```
#>
#> Pure-strategy NE:
```

```
for (eq in eq_coord$pure) {
  cat(sprintf("  (%s, %s)\n", rownames(A_coord)[eq[1]], colnames(A_coord)[eq[2]]))
}
```

```
#> (Left, Left)
#> (Right, Right)
```

```
if (!is.null(eq_coord$mixed)) {
  cat(sprintf("\nMixed NE: p = %.2f, q = %.2f\n",
              eq_coord$mixed$p, eq_coord$mixed$q))
  cat("Expected payoff at mixed NE: ",
      eq_coord$mixed$p * eq_coord$mixed$q * 1 +
      (1 - eq_coord$mixed$p) * (1 - eq_coord$mixed$q) * 1, "\n")
}
```

```
#>
#> Mixed NE: p = 0.50, q = 0.50
#> Expected payoff at mixed NE: 0.5
```

The coordination game has two pure NE — (Left, Left) and (Right, Right) — and a mixed NE at $p = q = 0.5$ with expected payoff 0.5, which is worse than either pure equilibrium. This illustrates that mixed equilibria can be Pareto-dominated by pure ones: the randomization creates a risk of miscoordination.

9.5. Extensions

Nash equilibrium is the foundation on which most of the rest of this book builds:

- **Mixed strategies** (Chapter 11) formalize the randomization used in the mixed NE above.
- **Extensive-form games** (Chapter 13) introduce sequential moves and **subgame perfect equilibrium**, a refinement of Nash equilibrium.

9. Nash Equilibrium

- **Bayesian games** (Chapter 15) extend Nash equilibrium to settings with incomplete information.
- **Repeated games** (Chapter 17) show how the folk theorem expands the set of equilibria when games are played repeatedly.

For a comprehensive treatment of Nash equilibrium and its refinements, see Osborne (2004) (Chapters 2–4).

Exercises

1. **Stag Hunt.** Consider the Stag Hunt game with payoffs: both hunt Stag yields (4, 4), both hunt Hare yields (3, 3), one hunts Stag while the other hunts Hare yields (0, 3) for the Stag hunter and (3, 0) for the Hare hunter. Find all pure-strategy and mixed-strategy Nash equilibria. Which equilibrium is risk-dominant? Which is payoff-dominant?
2. **Three equilibria.** Construct a 2×2 game that has exactly two pure-strategy Nash equilibria and one mixed-strategy Nash equilibrium. Verify your answer using the `support_enumeration()` function from `R/solvers.R`.
3. **Best-response plot.** Modify the best-response correspondence code in Section 9.3 to plot the correspondences for the game of Chicken (payoffs: (0, 0) for mutual swerve, (−1, 1) and (1, −1) for one swerving, (−5, −5) for neither swerving). How many equilibria does the plot reveal?

Solutions appear in Appendix H.

10. Mixed Strategies and the Indifference Principle

Why rational players sometimes randomize, how the indifference principle pins down mixing probabilities, and how to compute and visualize mixed-strategy Nash equilibria in R.

11. Mixed Strategies and the Indifference Principle

Learning objectives

- Explain why pure-strategy equilibria sometimes fail to exist and why randomization resolves the problem.
- State and apply the indifference principle: at a mixed Nash equilibrium, each player's mixture makes every opponent indifferent across the strategies in their support.
- Compute mixed-strategy Nash equilibria for 2x2 games by hand and with `solve_2x2_mixed_nash()`.
- Visualize expected payoffs as a function of mixing probability and identify the indifference point graphically.

11.1. Motivation

Not every strategic situation has a stable deterministic outcome. Consider a penalty kick in football. The striker chooses left or right; the goalkeeper dives left or right. If the goalkeeper could predict the striker's choice, she would always save the shot. If the striker could predict the goalkeeper, he would always score. Any predictable pattern is exploitable, so both players must be unpredictable — they must *randomize*.

This is not a peculiarity of sports. Firms randomizing the timing of sales, tax authorities choosing which returns to audit, and poker players blending bluffs with genuine bets all face the same logic. Whenever the interests of the players are sufficiently opposed that no pure-strategy profile is self-enforcing (see Chapter 9), stability requires mixing.

The profound insight of Nash (1950) is that this resolution is always available: every finite game possesses at least one Nash equilibrium when players are allowed to randomize. This chapter develops the theory behind that guarantee, introduces the *indifference principle* that makes mixed equilibria tractable, and implements the computations in R.

11.2. Theory

11.2.1. Mixed strategies defined

In a normal-form game (Chapter 7), a **pure strategy** is a definite choice of action. A **mixed strategy** σ_i for player i is a probability distribution over i 's action set A_i . We write $\sigma_i(a_i)$ for the probability that player i plays action a_i , with $\sigma_i(a_i) \geq 0$ for all a_i and $\sum_{a_i \in A_i} \sigma_i(a_i) = 1$.

The **support** of a mixed strategy is the set of actions played with positive probability: