

# Deep Generative Models

## Lecture 7

Roman Isachenko



Ozon Masters

Spring, 2021

# Dequantization

- ▶ Images are discrete data, pixels lie in the  $\{0, 255\}$  integer domain (the model is  $P(\mathbf{x}|\theta) = \text{Categorical}(\pi(\theta))$ ).
- ▶ Flow is a continuous model (it works with continuous data  $\mathbf{x}$ ).

By fitting a continuous density model to discrete data, one can produce a degenerate solution with all probability mass on discrete values.


How to convert a discrete data distribution to a continuous one?

## Uniform dequantization

$$\mathbf{x} \sim \text{Categorical}(\pi)$$

$$\mathbf{u} \sim U[0, 1]$$

$$\mathbf{y} = \mathbf{x} + \mathbf{u} \sim \text{Continuous}$$



# Uniform dequantization

## Statement


Fitting continuous model  $p(\mathbf{y}|\boldsymbol{\theta})$  on uniformly dequantized data  $\mathbf{y} = \mathbf{x} + \mathbf{u}$ ,  $\mathbf{u} \sim U[0, 1]$  is equivalent to maximization of a lower bound on log-likelihood for a discrete model:

$$P(\mathbf{x}|\boldsymbol{\theta}) = \int_{U[0,1]} p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u}$$

## Proof

$$\begin{aligned}\mathbb{E}_{\pi} \log p(\mathbf{y}|\boldsymbol{\theta}) &= \int \pi(\mathbf{y}) \log p(\mathbf{y}|\boldsymbol{\theta}) d\mathbf{y} = \\ &= \sum \pi(\mathbf{x}) \int_{U[0,1]} \log p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u} \leq \\ &\leq \sum \pi(\mathbf{x}) \log \int_{U[0,1]} p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u} = \\ &= \sum \pi(\mathbf{x}) \log P(\mathbf{x}|\boldsymbol{\theta}) = \mathbb{E}_{\pi} \log P(\mathbf{x}|\boldsymbol{\theta}).\end{aligned}$$

# Variational dequantization



- ▶  $p(\mathbf{y}|\boldsymbol{\theta})$  assign uniform density to unit hypercubes  $\mathbf{x} + U[0, 1]$  (left fig).
- ▶ Neural network density models are smooth function approximators (right fig).
- ▶ Smooth dequantization is more natural.

How to perform the smooth dequantization?

# Flow++

## Variational dequantization

Introduce variational dequantization noise distribution  $q(\mathbf{u}|\mathbf{x})$  and treat it as an approximate posterior.

## Variational lower bound

$$\begin{aligned}\log P(\mathbf{x}|\boldsymbol{\theta}) &= \left[ \log \int q(\mathbf{u}|\mathbf{x}) \frac{p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta})}{q(\mathbf{u}|\mathbf{x})} d\mathbf{u} \right] \geq \\ &\geq \int q(\mathbf{u}|\mathbf{x}) \log \frac{p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta})}{q(\mathbf{u}|\mathbf{x})} d\mathbf{u} = \mathcal{L}(q, \boldsymbol{\theta}).\end{aligned}$$

## Uniform dequantization bound

$$\log P(\mathbf{x}|\boldsymbol{\theta}) = \log \int_{U[0,1]} p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u} \geq \int_{U[0,1]} \log p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta}) d\mathbf{u}.$$

## Variational lower bound

$$\mathcal{L}(q, \theta) = \int q(\mathbf{u}|\mathbf{x}) \log \frac{p(\mathbf{x} + \mathbf{u}|\theta)}{q(\mathbf{u}|\mathbf{x})} d\mathbf{u}.$$

Let  $\mathbf{u} = h(\epsilon, \phi)$  is a flow model with base distribution  $\epsilon \sim p(\epsilon) = \mathcal{N}(0, \mathbf{I})$ :

$$q(\mathbf{u}|\mathbf{x}) = p(h^{-1}(\mathbf{u}, \phi)) \cdot \left| \det \frac{\partial h^{-1}(\mathbf{u}, \phi)}{\partial \mathbf{u}} \right|.$$

Then

$$\log P(\mathbf{x}|\theta) \geq \mathcal{L}(\phi, \theta) = \int p(\epsilon) \log \left( \frac{p(\mathbf{x} + h(\epsilon, \phi)|\theta)}{p(\epsilon) \cdot \left| \det \frac{\partial h(\epsilon, \phi)}{\partial \epsilon} \right|^{-1}} \right) d\epsilon.$$


## Variational lower

$$\log P(\mathbf{x}|\boldsymbol{\theta}) \geq \int p(\boldsymbol{\epsilon}) \log \left( \frac{p(\mathbf{x} + h(\boldsymbol{\epsilon}, \phi))}{p(\boldsymbol{\epsilon}) \cdot \left| \det \frac{\partial h(\boldsymbol{\epsilon}, \phi)}{\partial \boldsymbol{\epsilon}} \right|^{-1}} \right) d\boldsymbol{\epsilon}.$$


- ▶ If  $p(\mathbf{x} + \mathbf{u}|\boldsymbol{\theta})$  is also a flow model, it is straightforward to calculate stochastic gradient of this ELBO.
- ▶ Uniform dequantization is a special case of variational dequantization ( $q(\mathbf{u}|\mathbf{x}) = U[0, 1]$ ). The gap between  $\log P(\mathbf{x}|\boldsymbol{\theta})$  and the derived ELBO is  $KL(q(\mathbf{u}|\mathbf{x})||p(\mathbf{u}|\mathbf{x}))$ .
- ▶ In the case of uniform dequantization the model unnaturally places uniform density over each hypercube  $\mathbf{x} + U[0, 1]$  due to inexpressive distribution  $q$ .

Table 1. Unconditional image modeling results in bits/dim

Model family	Model	CIFAR10	ImageNet 32x32	ImageNet 64x64
Non-autoregressive	RealNVP (Dinh et al., 2016)	3.49	4.28	—
	Glow (Kingma & Dhariwal, 2018)	3.35	4.09	3.81
	IAF-VAE (Kingma et al., 2016)	3.11	—	—
	<b>Flow++ (ours)</b>	<b>3.08</b>	<b>3.86</b>	<b>3.69</b>
Autoregressive	Multiscale PixelCNN (Reed et al., 2017)	—	3.95	3.70
	PixelCNN (van den Oord et al., 2016b)	3.14	—	—
	PixelRNN (van den Oord et al., 2016b)	3.00	3.86	3.63
	Gated PixelCNN (van den Oord et al., 2016c)	3.03	3.83	3.57
	PixelCNN++ (Salimans et al., 2017)	2.92	—	—
	Image Transformer (Parmar et al., 2018)	2.90	3.77	—
	PixelSNAIL (Chen et al., 2017)	2.85	3.80	3.52



(a) PixelCNN



(b) Flow++



# VAE limitations

- ▶ Poor variational posterior distribution (encoder)

$$q(\mathbf{z}|\mathbf{x}, \phi) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_{\phi}(\mathbf{x}), \boldsymbol{\sigma}_{\phi}^2(\mathbf{x})).$$

- ▶ Poor prior distribution

$$p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I}).$$

- ▶ Poor probabilistic model (decoder)

$$p(\mathbf{x}|\mathbf{z}, \theta) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_{\theta}(\mathbf{z}), \boldsymbol{\sigma}_{\theta}^2(\mathbf{z})).$$

- ▶ Loose lower bound

$$\log p(\mathbf{x}|\theta) - \mathcal{L}(q, \theta) = (?).$$

# Importance Sampling

## Generative model

$$\begin{aligned} p(\mathbf{x}|\theta) &= \int p(\mathbf{x}, \mathbf{z}|\theta) d\mathbf{z} = \int \left[ \frac{p(\mathbf{x}, \mathbf{z}|\theta)}{q(\mathbf{z}|\mathbf{x})} \right] q(\mathbf{z}|\mathbf{x}) d\mathbf{z} \\ &= \int f(\mathbf{x}, \mathbf{z}) q(\mathbf{z}|\mathbf{x}) d\mathbf{z} = \mathbb{E}_{\mathbf{z} \sim q(\mathbf{z}|\mathbf{x})} f(\mathbf{x}, \mathbf{z}) \end{aligned}$$

Here  $f(\mathbf{x}, \mathbf{z}) = \frac{p(\mathbf{x}, \mathbf{z}|\theta)}{q(\mathbf{z}|\mathbf{x})}$ .

## ELBO

$$\begin{aligned} \log p(\mathbf{x}|\theta) &= \log \mathbb{E}_{\mathbf{z} \sim q(\mathbf{z}|\mathbf{x})} f(\mathbf{x}, \mathbf{z}) \geq \mathbb{E}_{\mathbf{z} \sim q(\mathbf{z}|\mathbf{x})} \log f(\mathbf{x}, \mathbf{z}) = \\ &= \mathbb{E}_{\mathbf{z} \sim q(\mathbf{z}|\mathbf{x})} \log \frac{p(\mathbf{x}, \mathbf{z}|\theta)}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}(q, \theta). \end{aligned}$$

Could we choose better  $f(\mathbf{x}, \mathbf{z})$ ?

$$p(\mathbf{x}|\boldsymbol{\theta}) = \int p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) d\mathbf{z} = \int \left[ \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x})} \right] q(\mathbf{z}|\mathbf{x}) d\mathbf{z} = \mathbb{E}_{\mathbf{z} \sim q(\mathbf{z}|\mathbf{x})} f(\mathbf{x}, \mathbf{z})$$

Let define

$$f(\mathbf{x}, \mathbf{z}_1, \dots, \mathbf{z}_K) = \frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k|\boldsymbol{\theta})}{q(\mathbf{z}_k|\mathbf{x})}$$

$$\mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x})} f(\mathbf{x}, \mathbf{z}_1, \dots, \mathbf{z}_K) = p(\mathbf{x}|\boldsymbol{\theta})$$

## ELBO

$$\begin{aligned} \log p(\mathbf{x}|\boldsymbol{\theta}) &= \log \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x})} f(\mathbf{x}, \mathbf{z}, \dots, \mathbf{z}_K) \geq \\ &\geq \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x})} \log f(\mathbf{x}, \mathbf{z}, \dots, \mathbf{z}_K) = \\ &= \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x})} \log \left[ \frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k|\boldsymbol{\theta})}{q(\mathbf{z}_k|\mathbf{x})} \right] = \mathcal{L}_K(q, \boldsymbol{\theta}). \end{aligned}$$

# IWAE

## VAE objective

$$\log p(\mathbf{x}|\boldsymbol{\theta}) \geq \mathcal{L}(q, \boldsymbol{\theta}) = \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x})} \rightarrow \max_{q, \boldsymbol{\theta}}$$

$$\mathcal{L}(q, \boldsymbol{\theta}) = \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x})} \left( \frac{1}{K} \sum_{k=1}^K \log \frac{p(\mathbf{x}, \mathbf{z}_k|\boldsymbol{\theta})}{q(\mathbf{z}_k|\mathbf{x})} \right) \rightarrow \max_{q, \boldsymbol{\theta}}.$$

## IWAE objective

$$\mathcal{L}_K(q, \boldsymbol{\theta}) = \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x})} \log \left( \frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k|\boldsymbol{\theta})}{q(\mathbf{z}_k|\mathbf{x})} \right) \rightarrow \max_{q, \boldsymbol{\theta}}.$$

If  $K = 1$ , these objectives coincide.

## Theorem

1.  $\log p(\mathbf{x}|\boldsymbol{\theta}) \geq \mathcal{L}_K(q, \boldsymbol{\theta}) \geq \mathcal{L}_M(q, \boldsymbol{\theta})$ , for  $K \geq M$ ;
2.  $\log p(\mathbf{x}|\boldsymbol{\theta}) = \lim_{K \rightarrow \infty} \mathcal{L}_K(q, \boldsymbol{\theta})$  if  $\frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x})}$  is bounded.

## Proof of 1.

$$\begin{aligned}
 \mathcal{L}_K(q, \boldsymbol{\theta}) &= \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K} \log \left( \frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k|\boldsymbol{\theta})}{q(\mathbf{z}_k|\mathbf{x})} \right) = \\
 &= \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K} \log \mathbb{E}_{k_1, \dots, k_M} \left( \frac{1}{M} \sum_{m=1}^M \frac{p(\mathbf{x}, \mathbf{z}_{k_m}|\boldsymbol{\theta})}{q(\mathbf{z}_{k_m}|\mathbf{x})} \right) \geq \\
 &\geq \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K} \mathbb{E}_{k_1, \dots, k_M} \log \left( \frac{1}{M} \sum_{m=1}^M \frac{p(\mathbf{x}, \mathbf{z}_{k_m}|\boldsymbol{\theta})}{q(\mathbf{z}_{k_m}|\mathbf{x})} \right) = \\
 &= \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_M} \log \left( \frac{1}{M} \sum_{m=1}^M \frac{p(\mathbf{x}, \mathbf{z}_m|\boldsymbol{\theta})}{q(\mathbf{z}_m|\mathbf{x})} \right) = \mathcal{L}_M(q, \boldsymbol{\theta})
 \end{aligned}$$

$$\frac{a_1 + \dots + a_K}{K} = \mathbb{E}_{k_1, \dots, k_M} \frac{a_{k_1} + \dots + a_{k_M}}{M}, \quad k_1, \dots, k_M \sim U[1, K]$$

Burda Y., Grosse R., Salakhutdinov R. Importance Weighted Autoencoders, 2015

## Theorem

1.  $\log p(\mathbf{x}|\boldsymbol{\theta}) \geq \mathcal{L}_K(q, \boldsymbol{\theta}) \geq \mathcal{L}_M(q, \boldsymbol{\theta})$ , for  $K \geq M$ ;
2.  $\log p(\mathbf{x}|\boldsymbol{\theta}) = \lim_{K \rightarrow \infty} \mathcal{L}_K(q, \boldsymbol{\theta})$  if  $\frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x})}$  is bounded.

## Proof of 2.

Consider r.v.  $\xi_K = \frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k|\boldsymbol{\theta})}{q(\mathbf{z}_k|\mathbf{x})}$ .

If summands are bounded, then (from the strong law of large numbers)

$$\xi_K \xrightarrow[K \rightarrow \infty]{a.s.} \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x})} = p(\mathbf{x}|\boldsymbol{\theta}).$$

Hence  $\mathcal{L}_K(q, \boldsymbol{\theta}) = \mathbb{E} \log \xi_K$  converges to  $\log p(\mathbf{x}|\boldsymbol{\theta})$  as  $K \rightarrow \infty$ .

$$\log p(\mathbf{x}|\boldsymbol{\theta}) \geq \mathcal{L}_K(q, \boldsymbol{\theta}) \geq \mathcal{L}(q, \boldsymbol{\theta})$$

If  $K > 1$  the bound could be tighter.

$$\mathcal{L}(q, \boldsymbol{\theta}) = \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x})};$$

$$\mathcal{L}_K(q, \boldsymbol{\theta}) = \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x})} \log \left( \frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k|\boldsymbol{\theta})}{q(\mathbf{z}_k|\mathbf{x})} \right).$$

- ▶  $\mathcal{L}_1(q, \boldsymbol{\theta}) = \mathcal{L}(q, \boldsymbol{\theta})$ ;
- ▶  $\mathcal{L}_\infty(q, \boldsymbol{\theta}) = \log p(\mathbf{x}|\boldsymbol{\theta})$ .
- ▶ Which  $q^*(\mathbf{z}|\mathbf{x})$  gives  $\mathcal{L}(q^*, \boldsymbol{\theta}) = \log p(\mathbf{x}|\boldsymbol{\theta})$ ?
- ▶ Which  $q^*(\mathbf{z}|\mathbf{x})$  gives  $\mathcal{L}(q^*, \boldsymbol{\theta}) = \mathcal{L}_K(q, \boldsymbol{\theta})$ ?

## Theorem

The VAE objective is equal to IWAE objective

$$\mathcal{L}(q_{EW}, \theta) = \mathcal{L}_K(q, \theta)$$

for the following variational distribution

$$q_{EW}(\mathbf{z}|\mathbf{x}) = \mathbb{E}_{\mathbf{z}_2, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x})} q_{IW}(\mathbf{z}|\mathbf{x}, \mathbf{z}_{2:K}),$$


where

$$q_{IW}(\mathbf{z}|\mathbf{x}, \mathbf{z}_{2:K}) = \frac{\frac{p(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})}}{\frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k)}{q(\mathbf{z}_k|\mathbf{x})}} q(\mathbf{z}|\mathbf{x}) = \frac{p(\mathbf{x}, \mathbf{z})}{\frac{1}{K} \left( \frac{p(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} + \sum_{k=2}^K \frac{p(\mathbf{x}, \mathbf{z}_k)}{q(\mathbf{z}_k|\mathbf{x})} \right)}.$$




# IWAE

## IWAE posterior



## IWAE samples



# IWAE

## Objective

$$\mathcal{L}_K(q, \theta) = \mathbb{E}_{\mathbf{z}_1, \dots, \mathbf{z}_K \sim q(\mathbf{z}|\mathbf{x}, \phi)} \log \left( \frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k | \theta)}{q(\mathbf{z}_k | \mathbf{x}, \phi)} \right) \rightarrow \max_{\phi, \theta}.$$

## Gradient

$$\Delta_K = \nabla_{\theta, \phi} \log \left( \frac{1}{K} \sum_{k=1}^K \frac{p(\mathbf{x}, \mathbf{z}_k | \theta)}{q(\mathbf{z}_k | \mathbf{x}, \phi)} \right), \quad \mathbf{z}_k \sim q(\mathbf{z} | \mathbf{x}, \phi).$$

## Theorem


$$\text{SNR}_K = \frac{\mathbb{E}[\Delta_K]}{\sigma(\Delta_K)}; \quad \text{SNR}_K(\theta) = O(\sqrt{K}); \quad \text{SNR}_K(\phi) = O\left(\sqrt{\frac{1}{K}}\right).$$

Hence, increasing  $K$  vanishes gradient signal of inference network  $q(\mathbf{z}|\mathbf{x}, \phi)$ .

# IWAE

## Theorem

$$\text{SNR}_K = \frac{\mathbb{E}[\Delta_K]}{\sigma(\Delta_K)}; \quad \text{SNR}_K(\theta) = O(\sqrt{K}); \quad \text{SNR}_K(\phi) = O\left(\sqrt{\frac{1}{K}}\right).$$



- ▶ IWAE makes the variational bound tighter and extends the class of variational distributions.
- ▶ Gradient signal becomes really small, training is complicated.
- ▶ IWAE is very popular technique as a quality measure for VAE models.

# VAE limitations

- ▶ Poor variational posterior distribution (encoder)

$$q(\mathbf{z}|\mathbf{x}, \phi) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_{\phi}(\mathbf{x}), \boldsymbol{\sigma}_{\phi}^2(\mathbf{x})).$$

- ▶ Poor prior distribution

$$p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I}).$$

- ▶ Poor probabilistic model (decoder)

$$p(\mathbf{x}|\mathbf{z}, \theta) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_{\theta}(\mathbf{z}), \boldsymbol{\sigma}_{\theta}^2(\mathbf{z})).$$

- ▶ Loose lower bound

$$\log p(\mathbf{x}|\theta) - \mathcal{L}(q, \theta) = (?).$$

## ELBO interpretations

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(\phi, \boldsymbol{\theta}) + KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) \geq \mathcal{L}(\phi, \boldsymbol{\theta}).$$

$$\mathcal{L}(\phi, \boldsymbol{\theta}) = \int q(\mathbf{z}|\mathbf{x}, \phi) \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x}, \phi)} d\mathbf{z}.$$

- Evidence minus posterior KL

$$\mathcal{L}(q, \boldsymbol{\theta}) = \log p(\mathbf{x}|\boldsymbol{\theta}) - KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})).$$

- Average negative energy plus entropy

$$\begin{aligned}\mathcal{L}(q, \boldsymbol{\theta}) &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} [\log p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) - \log q(\mathbf{z}|\mathbf{x}, \phi)] \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \log p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) + \mathbb{H}[q(\mathbf{z}|\mathbf{x}, \phi)].\end{aligned}$$

- Average reconstruction minus KL to prior

$$\begin{aligned}\mathcal{L}(q, \boldsymbol{\theta}) &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} [\log p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) + \log p(\mathbf{z}) - \log q(\mathbf{z}|\mathbf{x}, \phi)] \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \log p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) - KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z})).\end{aligned}$$

# ELBO surgery

$$\frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(q, \theta) = \frac{1}{n} \sum_{i=1}^n [\mathbb{E}_{q(\mathbf{z}|\mathbf{x}_i)} \log p(\mathbf{x}_i|\mathbf{z}, \theta) - KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z}))].$$

## Theorem

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) = KL(q(\mathbf{z})||p(\mathbf{z})) + \mathbb{I}_q[\mathbf{x}, \mathbf{z}],$$

- ▶  $q(\mathbf{z}) = \frac{1}{n} \sum_{i=1}^n q(\mathbf{z}|\mathbf{x}_i)$  – **aggregated** posterior distribution.
- ▶  $\mathbb{I}_q[\mathbf{x}, \mathbf{z}]$  – mutual information between  $\mathbf{x}$  and  $\mathbf{z}$  under empirical data distribution and distribution  $q(\mathbf{z}|\mathbf{x})$ .
- ▶ First term pushes  $q(\mathbf{z})$  towards the prior  $p(\mathbf{z})$ .
- ▶ Second term reduces the amount of information about  $\mathbf{x}$  stored in  $\mathbf{z}$ .

# ELBO surgery

## Theorem

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) = KL(q(\mathbf{z})||p(\mathbf{z})) + \mathbb{I}_q[\mathbf{x}, \mathbf{z}].$$

## Proof

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) &= \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z}|\mathbf{x}_i)}{p(\mathbf{z})} d\mathbf{z} = \\ &= \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z})q(\mathbf{z}|\mathbf{x}_i)}{p(\mathbf{z})q(\mathbf{z})} d\mathbf{z} = \int \frac{1}{n} \sum_{i=1}^n q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} + \\ &+ \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z}|\mathbf{x}_i)}{q(\mathbf{z})} d\mathbf{z} = KL(q(\mathbf{z})||p(\mathbf{z})) + \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||q(\mathbf{z})) \end{aligned}$$

# ELBO surgery

## Theorem

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) = KL(q(\mathbf{z})||p(\mathbf{z})) + \mathbb{I}_q[\mathbf{x}, \mathbf{z}],$$

## Proof (continued)

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z}_i)) = KL(q(\mathbf{z})||p(\mathbf{z})) + \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||q(\mathbf{z}))$$

It could be shown (exercise):

$$\mathbb{I}_q[\mathbf{x}, \mathbf{z}] = \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||q(\mathbf{z})) \in [0, \log n].$$



# Summary

- ▶ Uniform dequantization is the simplest form of dequantization.
- ▶ Variational dequantization is a more natural type that was proposed in Flow++ model.
- ▶ The IWAE could get the tighter lower bound to the likelihood, but the training of such model becomes more difficult.
- ▶ The ELBO surgery reveals insights about a prior distribution in VAE. The optimal prior is the aggregated posterior.