

Deep Generative Models

Lecture 12

Roman Isachenko

Moscow Institute of Physics and Technology

2023, Autumn

Recap of previous lecture

Forward gaussian diffusion process

Let $\mathbf{x}_0 = \mathbf{x} \sim \pi(\mathbf{x})$, $\beta_t \in (0, 1)$, $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^t \alpha_s$.

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1} + \sqrt{\beta_t} \cdot \boldsymbol{\epsilon}, \quad \text{where } \boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I});$$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \cdot \boldsymbol{\epsilon}, \quad \text{where } \boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I}).$$


$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\mathbf{x}_t | \sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \cdot \mathbf{I});$$

$$q(\mathbf{x}_t | \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_t | \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_t) \cdot \mathbf{I}).$$

1. $\mathbf{x}_0 = \mathbf{x} \sim \pi(\mathbf{x})$;
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1} + \sqrt{\beta_t} \cdot \boldsymbol{\epsilon}$, where $\boldsymbol{\epsilon} \sim \mathcal{N}(0, 1)$, $t \geq 1$;
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, 1)$, where $T \gg 1$.

If we are able to invert this process, we will get the way to sample $\mathbf{x} \sim \pi(\mathbf{x})$ using noise samples $p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.

Recap of previous lecture



Reverse gaussian diffusion process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)q(\mathbf{x}_{t-1}|\mathbf{x}_0)}{q(\mathbf{x}_t|\mathbf{x}_0)} = \mathcal{N}(\tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I})$$

- ▶ $q(\mathbf{x}_{t-1})$, $q(\mathbf{x}_t)$ are intractable.
- ▶ If β_t is small enough, $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ will be Gaussian (Feller, 1949).
- ▶ $q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)$ defines how to denoise a noisy image \mathbf{x}_t with access to the completely denoised image \mathbf{x}_0 .

Recap of previous lecture

Let's define the reverse process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) \approx p(\mathbf{x}_{t-1}|\mathbf{x}_t, \theta) = \mathcal{N}(\mathbf{x}_{t-1} | \mu_\theta(\mathbf{x}_t, t), \sigma_\theta^2(\mathbf{x}_t, t))$$

Forward process

1. $\mathbf{x}_0 = \mathbf{x} \sim \pi(\mathbf{x})$;

2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1} + \sqrt{\beta_t} \cdot \boldsymbol{\epsilon}$,
where $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$, $t \geq 1$;

3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.


Reverse process

1. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$;

2. $\mathbf{x}_{t-1} = \sigma_\theta(\mathbf{x}_t, t) \cdot \boldsymbol{\epsilon} + \mu_\theta(\mathbf{x}_t, t)$;

3. $\mathbf{x}_0 = \mathbf{x} \sim \pi(\mathbf{x})$;

Gaussian diffusion model as VAE



Recap of previous lecture

- ▶ $\mathbf{z} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$ is a latent variable.
- ▶ Variational posterior distribution

$$q(\mathbf{z}|\mathbf{x}) = q(\mathbf{x}_1, \dots, \mathbf{x}_T|\mathbf{x}_0) = \prod_{t=1}^T q(\mathbf{x}_t|\mathbf{x}_{t-1}).$$

- ▶ Generative distribution and prior

$$p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) = p(\mathbf{x}_0|\mathbf{x}_1, \boldsymbol{\theta}); \quad p(\mathbf{z}|\boldsymbol{\theta}) = \prod_{t=2}^T p(\mathbf{x}_{t-1}|\mathbf{x}_t, \boldsymbol{\theta}) \cdot p(\mathbf{x}_T)$$

ELBO

$$\log p(\mathbf{x}|\boldsymbol{\theta}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}(q, \boldsymbol{\theta}) \rightarrow \max_{q, \boldsymbol{\theta}}$$

$$\begin{aligned} \mathcal{L}(q, \boldsymbol{\theta}) &= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p(\mathbf{x}_0|\mathbf{x}_1, \boldsymbol{\theta}) - \textcolor{violet}{KL}(q(\mathbf{x}_T|\mathbf{x}_0) || p(\mathbf{x}_T)) - \\ &\quad - \sum_{t=2}^T \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \textcolor{violet}{KL}(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) || p(\mathbf{x}_{t-1}|\mathbf{x}_t, \boldsymbol{\theta}))}_{\mathcal{L}_t} \end{aligned}$$

Outline

1. Denoising Diffusion Probabilistic Model (DDPM)

Reparametrization of gaussian diffusion model

Overview of DDPM

2. Langevin dynamic and SDE basics

Outline

1. Denoising Diffusion Probabilistic Model (DDPM)

Reparametrization of gaussian diffusion model

Overview of DDPM

2. Langevin dynamic and SDE basics

Outline

1. Denoising Diffusion Probabilistic Model (DDPM)

Reparametrization of gaussian diffusion model

Overview of DDPM

2. Langevin dynamic and SDE basics

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} KL(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) || p(\mathbf{x}_{t-1}|\mathbf{x}_t, \theta))$$

\mathcal{L}_t is the mean of KL between two normal distributions:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1} | \tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p(\mathbf{x}_{t-1}|\mathbf{x}_t, \theta) = \mathcal{N}(\mathbf{x}_{t-1} | \mu_\theta(\mathbf{x}_t, t), \sigma_\theta^2(\mathbf{x}_t, t) \mathbf{I})$$

Here

$$\tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1 - \bar{\alpha}_{t-1})}{1 - \bar{\alpha}_t} \mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1 - \alpha_t)}{1 - \bar{\alpha}_t} \mathbf{x}_0;$$

$$\tilde{\beta}_t = \frac{(1 - \alpha_t)(1 - \bar{\alpha}_{t-1})}{1 - \bar{\alpha}_t} = \text{const.}$$

Let assume

$$\sigma_\theta^2(\mathbf{x}_t, t) = \tilde{\beta}_t \mathbf{I} \quad \Rightarrow \quad p(\mathbf{x}_{t-1}|\mathbf{x}_t, \theta) = \mathcal{N}(\mathbf{x}_{t-1} | \mu_\theta(\mathbf{x}_t, t), \tilde{\beta}_t \mathbf{I}).$$

Reparametrization of DDPM

$$q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1} | \tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I});$$
$$p(\mathbf{x}_{t-1} | \mathbf{x}_t, \theta) = \mathcal{N}(\mathbf{x}_{t-1} | \mu_\theta(\mathbf{x}_t, t), \tilde{\beta}_t \mathbf{I}).$$

Use the formula for KL between two normal distributions:

$$\begin{aligned}\mathcal{L}_t &= \mathbb{E}_{q(\mathbf{x}_t | \mathbf{x}_0)} KL\left(\mathcal{N}(\tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}) || \mathcal{N}(\mu_\theta(\mathbf{x}_t, t), \tilde{\beta}_t \mathbf{I})\right) \\ &= \mathbb{E}_{q(\mathbf{x}_t | \mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \|\tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0) - \mu_\theta(\mathbf{x}_t, t)\|^2 \right]\end{aligned}$$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \boldsymbol{\epsilon} \quad \Rightarrow \quad \mathbf{x}_0 = \frac{\mathbf{x}_t - \sqrt{1 - \bar{\alpha}_t} \boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}$$

$$\begin{aligned}\tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0) &= \frac{\sqrt{\alpha_t}(1 - \bar{\alpha}_{t-1})}{1 - \bar{\alpha}_t} \mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1 - \alpha_t)}{1 - \bar{\alpha}_t} \mathbf{x}_0 \\ &= \frac{1}{\sqrt{\alpha_t}} \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \boldsymbol{\epsilon}\end{aligned}$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \|\tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0) - \mu_\theta(\mathbf{x}_t, t)\|^2 \right]$$

Reparametrization

$$\tilde{\mu}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{1}{\sqrt{\alpha_t}} \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \epsilon$$

$$\mu_\theta(\mathbf{x}_t, t) = \frac{1}{\sqrt{\alpha_t}} \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \epsilon_\theta(\mathbf{x}_t, t)$$

$$\mathcal{L}_t = \mathbb{E}_{\epsilon \sim \mathcal{N}(0, I)} \left[\frac{(1 - \alpha_t)^2}{2\tilde{\beta}_t \alpha_t (1 - \bar{\alpha}_t)} \|\epsilon - \epsilon_\theta(\mathbf{x}_t, t)\|^2 \right]$$

$$= \mathbb{E}_{\epsilon \sim \mathcal{N}(0, I)} \left[\frac{(1 - \alpha_t)^2}{2\tilde{\beta}_t \alpha_t (1 - \bar{\alpha}_t)} \|\epsilon - \epsilon_\theta(\sqrt{\bar{\alpha}_t} \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \epsilon, t)\|^2 \right]$$

At each step of reverse diffusion process we try to predict the noise ϵ that we used in the forward diffusion process!

Reparametrization of DDPM

$$\mathcal{L}(q, \theta) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p(\mathbf{x}_0|\mathbf{x}_1, \theta) - KL(q(\mathbf{x}_T|\mathbf{x}_0)||p(\mathbf{x}_T)) - \\ - \sum_{t=2}^T \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} KL(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)||p(\mathbf{x}_{t-1}|\mathbf{x}_t, \theta))}_{\mathcal{L}_t}$$

$$\mathcal{L}_t = \mathbb{E}_{\epsilon \sim \mathcal{N}(0, \mathbf{I})} \left[\frac{(1 - \alpha_t)^2}{2\tilde{\beta}_t \alpha_t (1 - \bar{\alpha}_t)} \left\| \epsilon - \epsilon_\theta(\sqrt{\bar{\alpha}_t} \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \epsilon, t) \right\|^2 \right]$$

Simplified objective

$$\mathcal{L}_{\text{simple}} = \mathbb{E}_{t \sim U[2, T]} \mathbb{E}_{\epsilon \sim \mathcal{N}(0, \mathbf{I})} \left\| \epsilon - \epsilon_\theta(\sqrt{\bar{\alpha}_t} \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \epsilon, t) \right\|^2$$

Outline

1. Denoising Diffusion Probabilistic Model (DDPM)

Reparametrization of gaussian diffusion model

Overview of DDPM

2. Langevin dynamic and SDE basics

Denoising diffusion probabilistic model (DDPM)

DDPM is a VAE model

- ▶ Encoder is a fixed Gaussian Markov chain $q(\mathbf{x}_1, \dots, \mathbf{x}_T | \mathbf{x}_0)$.
- ▶ Latent variable is a hierarchical (in each step the dim. of the latent equals to the dim of the input).
- ▶ Decoder is a simple Gaussian model $p(\mathbf{x}_0 | \mathbf{x}_1, \theta)$.
- ▶ Prior distribution is given by parametric Gaussian Makov chain $p(\mathbf{x}_{t-1} | \mathbf{x}_t, \theta)$.

Forward process

1. $\mathbf{x}_0 = \mathbf{x} \sim \pi(\mathbf{x})$;
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1} + \sqrt{\beta_t} \cdot \boldsymbol{\epsilon}$,
where $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$, $t \geq 1$;
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.

Reverse process

1. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$;
2. $\mathbf{x}_{t-1} = \sigma_\theta(\mathbf{x}_t, t) \cdot \boldsymbol{\epsilon} + \mu_\theta(\mathbf{x}_t, t)$;
3. $\mathbf{x}_0 = \mathbf{x} \sim \pi(\mathbf{x})$;

Denoising diffusion probabilistic model (DDPM)

Training

1. Get the sample $\mathbf{x}_0 \sim \pi(\mathbf{x})$.
2. Sample timestamp $t \sim U[1, T]$ and the noise $\epsilon \sim \mathcal{N}(0, \mathbf{I})$.
3. Get noisy image $\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \epsilon$.
4. Compute loss $\mathcal{L}_{\text{simple}} = \|\epsilon - \epsilon_{\theta}(\mathbf{x}_t, t)\|^2$.

Sampling



1. Sample $\mathbf{x}_T \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute mean of $p(\mathbf{x}_{t-1} | \mathbf{x}_t, \theta) = \mathcal{N}(\mu_{\theta}(\mathbf{x}_t, t), \tilde{\beta}_t \mathbf{I})$:

$$\mu_{\theta}(\mathbf{x}_t, t) = \frac{1}{\sqrt{\alpha_t}} \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \epsilon_{\theta}(\mathbf{x}_t, t)$$

3. Get denoised image $\mathbf{x}_{t-1} = \mu_{\theta}(\mathbf{x}_t, t) + \sqrt{\tilde{\beta}_t} \cdot \epsilon$, where $\epsilon \sim \mathcal{N}(0, \mathbf{I})$.

Denoising diffusion probabilistic model (DDPM)

Samples



Outline

1. Denoising Diffusion Probabilistic Model (DDPM)

Reparametrization of gaussian diffusion model

Overview of DDPM

2. Langevin dynamic and SDE basics

Langevin dynamic

Imagine that we have some generative model $p(\mathbf{x}|\theta)$.

Statement

Let \mathbf{x}_0 be a random vector. Then under mild regularity conditions for small enough η samples from the following dynamics

$$\mathbf{x}_{t+1} = \mathbf{x}_t + \eta \frac{1}{2} \nabla_{\mathbf{x}_t} \log p(\mathbf{x}_t | \theta) + \sqrt{\eta} \cdot \boldsymbol{\epsilon}, \quad \boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I}).$$

will comes from $p(\mathbf{x}|\theta)$.

What do we get if $\boldsymbol{\epsilon} = \mathbf{0}$?

Energy-based model

$$p(\mathbf{x}|\theta) = \frac{\hat{p}(\mathbf{x}|\theta)}{Z_\theta}, \quad \text{where } Z_\theta = \int \hat{p}(\mathbf{x}|\theta) d\mathbf{x}$$

$$\nabla_{\mathbf{x}} \log p(\mathbf{x}|\theta) = \nabla_{\mathbf{x}} \log \hat{p}(\mathbf{x}|\theta) - \nabla_{\mathbf{x}} \log Z_\theta = \nabla_{\mathbf{x}} \log \hat{p}(\mathbf{x}|\theta)$$

Gradient of normalized density equals to gradient of unnormalized density.

Stochastic differential equation (SDE)

Let define stochastic process $\mathbf{x}(t)$ with initial condition $\mathbf{x}(0) \sim p_0(\mathbf{x})$:

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

- ▶ $\mathbf{f}(\mathbf{x}, t)$ is the **drift** function of $\mathbf{x}(t)$.
- ▶ $g(t)$ is the **diffusion** coefficient of $\mathbf{x}(t)$.
- ▶ If $g(t) = 0$ we get standard ODE.
- ▶ $\mathbf{w}(t)$ is the standard Wiener process (Brownian motion):
 1. $\mathbf{w}(0) = 0$ (almost surely);
 2. $\mathbf{w}(t)$ has independent increments;
 3. $\mathbf{w}(t) - \mathbf{w}(s) \sim \mathcal{N}(0, (t-s)\mathbf{I})$.
- ▶ $d\mathbf{w} = \mathbf{w}(t+dt) - \mathbf{w}(t) = \mathcal{N}(0, \mathbf{I} \cdot dt) = \epsilon \cdot \sqrt{dt}$, where $\epsilon \sim \mathcal{N}(0, \mathbf{I})$.

Note: In contrast to ODE, initial condition $\mathbf{x}(0)$ does not uniquely determine the process trajectory.

Stochastic differential equation (SDE)

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}, \quad d\mathbf{w} = \boldsymbol{\epsilon} \cdot \sqrt{dt}, \quad \boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I}).$$

- ▶ At each moment t we have the density $p(\mathbf{x}(t), t)$.
- ▶ How to get distribution $p(\mathbf{x}, t)$ for $\mathbf{x}(t)$?

Theorem (Kolmogorov-Fokker-Planck)

Evolution of the distribution $p(\mathbf{x}, t)$ is given by the following ODE:

$$\frac{\partial p(\mathbf{x}, t)}{\partial t} = \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} [\mathbf{f}(\mathbf{x}, t)p(\mathbf{x}, t)] + \frac{1}{2}g^2(t)\frac{\partial^2 p(\mathbf{x}, t)}{\partial \mathbf{x}^2} \right)$$

Note: This is the generalization of KFP theorem that we used in continuous-in-time NF.

Langevin SDE (special case)

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

$$d\mathbf{x} = \frac{1}{2}\frac{\partial}{\partial \mathbf{x}} \log p(\mathbf{x}, t)dt + \mathbf{1} \cdot d\mathbf{w}$$

Langevin SDE (special case)

$$d\mathbf{x} = \frac{1}{2} \frac{\partial}{\partial \mathbf{x}} \log p(\mathbf{x}, t) dt + \mathbf{1} \cdot d\mathbf{w}$$

Let apply KFP theorem.

$$\begin{aligned}\frac{\partial p(\mathbf{x}, t)}{\partial t} &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[p(\mathbf{x}, t) \frac{1}{2} \frac{\partial}{\partial \mathbf{x}} \log p(\mathbf{x}, t) \right] + \frac{1}{2} \frac{\partial^2 p(\mathbf{x}, t)}{\partial \mathbf{x}^2} \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\frac{1}{2} \frac{\partial}{\partial \mathbf{x}} p(\mathbf{x}, t) \right] + \frac{1}{2} \frac{\partial^2 p(\mathbf{x}, t)}{\partial \mathbf{x}^2} \right) = 0\end{aligned}$$

The density $p(\mathbf{x}, t) = \text{const}(t)!$

Discretized Langevin SDE

$$\mathbf{x}_{t+1} - \mathbf{x}_t = \eta \frac{1}{2} \frac{\partial}{\partial \mathbf{x}} \log p(\mathbf{x}, t) + \sqrt{\eta} \cdot \boldsymbol{\epsilon}, \quad \eta \approx dt.$$

Langevin dynamic

$$\mathbf{x}_{t+1} = \mathbf{x}_t + \eta \frac{1}{2} \nabla_{\mathbf{x}} \log p(\mathbf{x} | \theta) + \sqrt{\eta} \cdot \boldsymbol{\epsilon}, \quad \eta \approx dt.$$

Stochastic differential equation (SDE)


Statement

Let \mathbf{x}_0 be a random vector. Then samples from the following dynamics

$$\mathbf{x}_{t+1} = \mathbf{x}_t + \eta \frac{1}{2} \nabla_{\mathbf{x}_t} \log p(\mathbf{x}_t | \theta) + \sqrt{\eta} \cdot \epsilon, \quad \epsilon \sim \mathcal{N}(0, \mathbf{I}).$$

will come from $p(\mathbf{x} | \theta)$ under mild regularity conditions for small enough η and large enough t .

The density $p(\mathbf{x} | \theta)$ is a **stationary** distribution for this SDE.



Summary

- ▶ DDPM is a VAE model with hierarchical latent variables.
- ▶ At each step DDPM predicts the noise that was used in the forward diffusion process.
- ▶ DDPM is really slow, because we have to apply the model T times.
- ▶ Langevin dynamics allows to sample from the energy-based model using the score function (due to the existence of stationary distribution of SDE).