

Deep Generative Models

Lecture 13

Roman Isachenko



2025, Spring

Recap of previous lecture

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

Variance Exploding SDE (NCSN)

$$d\mathbf{x} = \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}, \quad \mathbf{f}(\mathbf{x}, t) = 0, \quad g(t) = \sqrt{\frac{d[\sigma^2(t)]}{dt}}$$

Variance grows since $\sigma(t)$ is a monotonically increasing function.

Variance Preserving SDE (DDPM)

$$d\mathbf{x} = -\frac{1}{2}\beta(t)\mathbf{x}(t)dt + \sqrt{\beta(t)} \cdot d\mathbf{w}$$

$$\mathbf{f}(\mathbf{x}, t) = -\frac{1}{2}\beta(t)\mathbf{x}(t), \quad g(t) = \sqrt{\beta(t)}$$

Variance is preserved if $\mathbf{x}(0)$ has a unit variance.

Recap of previous lecture

Discrete-in-time objective

$$\mathbb{E}_{\pi(\mathbf{x}_0)} \mathbb{E}_{t \sim U\{1, T\}} \mathbb{E}_{q(\mathbf{x}_t | \mathbf{x}_0)} \| \mathbf{s}_{\theta, t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t | \mathbf{x}_0) \|_2^2$$

Continuous-in-time objective

$$\mathbb{E}_{\pi(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0, 1]} \mathbb{E}_{q(\mathbf{x}(t) | \mathbf{x}(0))} \| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t) | \mathbf{x}(0)) \|_2^2$$

NCSN

$$q(\mathbf{x}(t) | \mathbf{x}(0)) = \mathcal{N} (\mathbf{x}(0), [\sigma^2(t) - \sigma^2(0)] \cdot \mathbf{I})$$


DDPM

$$q(\mathbf{x}(t) | \mathbf{x}(0)) = \mathcal{N} \left(\mathbf{x}(0) e^{-\frac{1}{2} \int_0^t \beta(s) ds}, \left(1 - e^{-\int_0^t \beta(s) ds} \right) \cdot \mathbf{I} \right)$$

Recap of previous lecture

Sampling

Solve reverse SDE using numerical solvers (SDESolve).



- ▶ Discretization of the reverse SDE gives us the ancestral sampling.
- ▶ If we use probability flow instead of SDE than the reverse ODE gives us the DDIM sampling.

Recap of previous lecture

Let consider ODE dynamic $\mathbf{x}(t)$ in time interval $t \in [0, 1]$ with $\mathbf{x}_0 \sim p_0(\mathbf{x}) = p(\mathbf{x})$, $\mathbf{x}_1 \sim p_1(\mathbf{x}) = \pi(\mathbf{x})$.

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t), \quad \text{with initial condition } \mathbf{x}(0) = \mathbf{x}_0.$$

KFP theorem (continuity equation)

$$\frac{\partial p_t(\mathbf{x})}{\partial t} = -\operatorname{div}(\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x})) \Leftrightarrow \frac{d \log p_t(\mathbf{x}(t))}{dt} = -\operatorname{tr}\left(\frac{\partial \mathbf{f}(\mathbf{x}(t), t)}{\partial \mathbf{x}(t)}\right)$$

Solving the continuity equation using the adjoint method is complicated and unstable process.

Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{f}(\mathbf{x}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_\theta$$

Recap of previous lecture

Let's introduce the latent variable \mathbf{z} :

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{z})p(\mathbf{z})d\mathbf{z}$$
$$\frac{\partial p_t(\mathbf{x}|\mathbf{z})}{\partial t} = -\text{div}(\mathbf{f}(\mathbf{x}, \mathbf{z}, t)p_t(\mathbf{x}|\mathbf{z})).$$

- ▶ $p_t(\mathbf{x}|\mathbf{z})$ is a **conditional probability path**;
- ▶ $\mathbf{f}(\mathbf{x}, \mathbf{z}, t)$ is a **conditional vector field**.

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t) \quad \Rightarrow \quad \frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, \mathbf{z}, t)$$

Theorem

The following vector field generates the probability path $p_t(\mathbf{x})$.

$$\mathbf{f}(\mathbf{x}, t) = \mathbb{E}_{p_t(\mathbf{z}|\mathbf{x})}\mathbf{f}(\mathbf{x}, \mathbf{z}, t) = \int \mathbf{f}(\mathbf{x}, \mathbf{z}, t) \frac{p_t(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{p_t(\mathbf{x})} d\mathbf{z}$$

Recap of previous lecture

Flow Matching (FM)


$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{f}(\mathbf{x}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditional Flow Matching (CFM)

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Theorem

If $\text{supp}(p_t(\mathbf{x})) = \mathbb{R}^m$, then the optimal value of FM objective is equal to the optimal value of CFM objective.



Outline

1. Conditional Flow Matching

Conical gaussian paths

Linear interpolation

Link with score-based models

Outline

1. Conditional Flow Matching

Conical gaussian paths

Linear interpolation

Link with score-based models

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_\theta(x, t)\|^2 \rightarrow \min_{\theta}$$

What is left?

- ▶ How to choose the conditioning latent variable z ?
- ▶ How to define $p_t(x|z)$ which follows the constraints?

Gaussian conditional probability path

$$p_t(x|z) = \mathcal{N}(\mu_t(z), \sigma_t^2(z))$$

- ▶ There is an infinite number of vector fields that generate any particular probability path.
- ▶ Let consider the following dynamics:

$$x_t = \mu_t(z) + \sigma_t(z) \odot x_0, \quad x_0 \sim p_0(x) = \mathcal{N}(0, I)$$

Flow Matching

Gaussian conditional probability path

$$p_t(\mathbf{x}|\mathbf{z}) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{z}), \boldsymbol{\sigma}_t^2(\mathbf{z})) ; \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{z}) + \boldsymbol{\sigma}_t(\mathbf{z}) \odot \mathbf{x}_0$$

Theorem

$$\mathbf{f}(\mathbf{x}, \mathbf{z}, t) = \boldsymbol{\mu}'_t(\mathbf{z}) + \frac{\boldsymbol{\sigma}'_t(\mathbf{z})}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x} - \boldsymbol{\mu}_t(\mathbf{z}))$$

Proof

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, \mathbf{z}, t); \quad \mathbf{x}_0 = \frac{1}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{z}))$$

$$\frac{d\mathbf{x}}{dt} = \boldsymbol{\mu}'_t(\mathbf{z}) + \boldsymbol{\sigma}'_t(\mathbf{z}) \odot \mathbf{x}_0 = \boldsymbol{\mu}'_t(\mathbf{z}) + \frac{\boldsymbol{\sigma}'_t(\mathbf{z})}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x} - \boldsymbol{\mu}_t(\mathbf{z}))$$

Outline

1. Conditional Flow Matching

Conical gaussian paths

Linear interpolation

Link with score-based models

Endpoint conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditioning latent variable

Let choose $\mathbf{z} = \mathbf{x}_1$. Then $p(\mathbf{z}) = p_1(\mathbf{x}_1)$.

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{x}_1) p_1(\mathbf{x}_1) d\mathbf{x}_1$$

We need to ensure boundary constraints:

$$\begin{cases} p(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); (= \mathcal{N}(0, \mathbf{I})) \\ \pi(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}). \end{cases} \Rightarrow \begin{cases} p_0(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(0, \mathbf{I}); \\ p_1(\mathbf{x}|\mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1). \end{cases}$$

Conical gaussian paths


$$p_0(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(0, \mathbf{I}); \quad p_1(\mathbf{x}|\mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1).$$

Gaussian conditional probability path

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_1) \odot \mathbf{x}_0.$$

Let consider straight conditional paths

$$\begin{cases} \boldsymbol{\mu}_t(\mathbf{x}_1) = t\mathbf{x}_1; \\ \boldsymbol{\sigma}_t(\mathbf{x}_1) = 1-t. \end{cases} \Rightarrow \begin{cases} p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \cdot \mathbf{I}); \\ \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0. \end{cases}$$



Conical gaussian paths

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \mathbf{I}); \quad \mathbf{x} = t\mathbf{x}_1 + (1-t)\mathbf{x}_0.$$

Conditional vector field

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, \mathbf{x}_1, t) = \mu'_t(\mathbf{x}_1) + \frac{\sigma'_t(\mathbf{x}_1)}{\sigma_t(\mathbf{x}_1)} \odot (\mathbf{x} - \mu_t(\mathbf{x}_1))$$

$$\begin{aligned}\mathbf{f}(\mathbf{x}, \mathbf{x}_1, t) &= \mathbf{x}_1 - \frac{1}{1-t} \cdot (\mathbf{x} - t\mathbf{x}_1) = \frac{\mathbf{x}_1 - \mathbf{x}}{1-t} = \\ &= \frac{\mathbf{x}_1 - t\mathbf{x}_1 - (1-t)\mathbf{x}_0}{1-t} = \mathbf{x}_1 - \mathbf{x}_0\end{aligned}$$

Conditional Flow Matching

$$\begin{aligned}\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 &= \\ \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x}_1 \sim \pi(\mathbf{x})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{x}_1)} \left\| \left(\frac{\mathbf{x}_1 - \mathbf{x}}{1-t} \right) - \mathbf{f}_\theta(\mathbf{x}, t) \right\|^2 &= \\ \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x}_1 \sim \pi(\mathbf{x})} \mathbb{E}_{\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})} \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_\theta(t\mathbf{x}_1 + (1-t)\mathbf{x}_0, t)\|^2\end{aligned}$$

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x}_1 \sim \pi(\mathbf{x})} \mathbb{E}_{\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})} \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

We fit straight lines between noise distribution $p(\mathbf{x})$ and the data distribution $\pi(\mathbf{x})$.

Training

1. Get the sample $\mathbf{x}_1 \sim \pi(\mathbf{x})$.
2. Sample timestamp $t \sim U[0, 1]$ and $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$.
3. Get noisy image $\mathbf{x}_t = t\mathbf{x}_1 + (1 - t)\mathbf{x}_0$.
4. Compute loss $\mathcal{L} = \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2$.

Sampling

1. Sample $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$.
2. Solve the ODE to get \mathbf{x}_1 :

$$\mathbf{x}_1 = \text{ODESolve}_f(\mathbf{x}_0, \theta, t_0 = 0, t_1 = 1).$$

Flow Matching

$$\mathbf{x}_t = t\mathbf{x}_1 + (1 - t)\mathbf{x}_0$$

- ▶ The conditional probability path $p_t(\mathbf{x}|\mathbf{z})$ is an **optimal transport path** from $p_0(\mathbf{x}|\mathbf{z})$ to $p_1(\mathbf{x}|\mathbf{z})$ (in terms of the conditional trajectories straightness).
- ▶ The marginal path $p_t(\mathbf{x})$ is not in general an optimal transport path from the standard normal $p_0(\mathbf{x})$ to the data distribution $p_1(\mathbf{x})$.



image credit: <https://mlg.eng.cam.ac.uk/blog/2024/01/20/flow-matching.html>

Outline

1. Conditional Flow Matching

Conical gaussian paths

Linear interpolation

Link with score-based models

Pair conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditioning latent variable

Let choose $\mathbf{z} = (\mathbf{x}_0, \mathbf{x}_1)$. Then $p(\mathbf{z}) = p(\mathbf{x}_0, \mathbf{x}_1) = p_0(\mathbf{x}_0)p_1(\mathbf{x}_1)$.

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) p_0(\mathbf{x}_0) p_1(\mathbf{x}_1) d\mathbf{x}_0 d\mathbf{x}_1$$

We need to ensure boundary constraints:

$$\begin{cases} p(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); (= \mathcal{N}(0, \mathbf{I})) \\ \pi(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}). \end{cases} \Rightarrow \begin{cases} p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0); \\ p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1). \end{cases}$$

Linear interpolation


$$p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0); \quad p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1).$$

Gaussian conditional probability path

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}(\mu_t(\mathbf{x}_0, \mathbf{x}_1), \sigma_t^2(\mathbf{x}_0, \mathbf{x}_1)); \quad \mathbf{x}_t = \mu_t(\mathbf{x}_0, \mathbf{x}_1) + \sigma_t^2(\mathbf{x}_0, \mathbf{x}_1) \odot \mathbf{x}_0.$$

Let consider straight conditional paths

$$\begin{cases} \mu_t(\mathbf{x}_1) = t\mathbf{x}_1 + (1-t)\mathbf{x}_0; \\ \sigma_t(\mathbf{x}_1) = \epsilon. \end{cases} \Rightarrow \begin{cases} p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0); \\ p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1). \end{cases}$$



Flow Matching: conical paths vs linear interpolation

$$z = x_1$$

$$p_t(x|x_1) = \mathcal{N}(tx_1, (1-t)^2 I)$$


$$x_t = tx_1 + (1-t)x_0.$$

$$z = (x_0, x_1)$$


$$p_t(x|x_0, x_1) = \mathcal{N}(tx_1 + (1-t)x_0, \epsilon^2 I)$$

$$x_t = tx_1 + (1-t)x_0.$$

Flow Matching (Lipman et al.)



Conditional Flow Matching



Linear interpolation

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}\left(t\mathbf{x}_1 + (1-t)\mathbf{x}_0, \epsilon^2 \mathbf{I}\right); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0.$$

Conditional vector field

$$\begin{aligned} \frac{d\mathbf{x}}{dt} &= \mathbf{f}(\mathbf{x}, \mathbf{x}_0, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_0, \mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_0, \mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1)} \odot (\mathbf{x} - \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1)) \\ \mathbf{f}(\mathbf{x}, \mathbf{x}_0, \mathbf{x}_1, t) &= \mathbf{x}_1 - \mathbf{x}_0 \end{aligned}$$

Conditional Flow Matching

$$\begin{aligned} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 &= \\ \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{(\mathbf{x}_0, \mathbf{x}_1) \sim p(\mathbf{x}_0, \mathbf{x}_1)} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1)} \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_\theta(\mathbf{x}_t, t)\|^2 & \end{aligned}$$

- ▶ We got the same procedure as for conical paths!
- ▶ Now we do not have the constraint that $p_0(\mathbf{x})$ should be $\mathcal{N}(0, \mathbf{I})$.

Conditional Flow Matching

- ▶ We could use this conditioning for transferring any distribution $p_0(\mathbf{x})$ to any distribution $p_1(\mathbf{x})$.
- ▶ It is possible to use this approach for paired problems (style transfer).

Training


1. Get the sample $(\mathbf{x}_0, \mathbf{x}_1) \sim p(\mathbf{x}_0, \mathbf{x}_1)$.
2. Sample timestamp $t \sim U[0, 1]$.
3. Get noisy image $\mathbf{x}_t = t\mathbf{x}_1 + (1 - t)\mathbf{x}_0$.
4. Compute loss $\mathcal{L} = \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_{\theta}(\mathbf{x}, t)\|^2$.

Sampling

1. Sample $\mathbf{x}_0 \sim p_0(\mathbf{x})$.
2. Solve the ODE to get \mathbf{x}_1 :

$$\mathbf{x}_1 = \text{ODESolve}_f(\mathbf{x}_0, \boldsymbol{\theta}, t_0 = 0, t_1 = 1).$$

Stable Diffusion 3: scalable flow matching



Outline

1. Conditional Flow Matching

Conical gaussian paths

Linear interpolation

Link with score-based models

Score-based generative models through SDEs

Training

$$\mathbb{E}_{\pi(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{q(\mathbf{x}(t)|\mathbf{x}(0))} \| \mathbf{s}_\theta(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) \|_2^2$$

Variance Exploding SDE (NCSN)

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}(\mathbf{x}(0), [\sigma^2(t) - \sigma^2(0)] \cdot \mathbf{I}), \quad \sigma(0) = 0.$$

Variance Preserving SDE (DDPM)

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}(\mathbf{x}(0)\alpha(t), (1 - \alpha(t)^2) \cdot \mathbf{I}); \quad \alpha(t) = e^{-\frac{1}{2} \int_0^t \beta(s) ds}$$

Flow matching uses the reverse time direction.

$$\textbf{NCSN: } p(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(\mathbf{x}_1, \sigma_{1-t}^2 \cdot \mathbf{I})$$

$$\textbf{DDPM: } p(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(\alpha_{1-t}\mathbf{x}_1, (1 - \alpha_{1-t}^2) \cdot \mathbf{I})$$

Flow matching vs score-based SDE models

Flow matching probability path

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N} \left(t\mathbf{x}_1, (1-t)^2 \mathbf{I} \right); \quad \mathbf{f}(\mathbf{x}, \mathbf{x}_1, t) = \frac{\mathbf{x}_1 - \mathbf{x}}{1-t}$$

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_1)} \odot (\mathbf{x} - \boldsymbol{\mu}_t(\mathbf{x}_1))$$

Variance Exploding SDE probability path

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N} \left(\mathbf{x}_1, \sigma_{1-t}^2 \mathbf{I} \right) \quad \Rightarrow \quad \mathbf{f}(\mathbf{x}, \mathbf{x}_1, t) = -\frac{\boldsymbol{\sigma}'_{1-t}}{\boldsymbol{\sigma}_{1-t}} \cdot (\mathbf{x} - \mathbf{x}_1)$$


Variance Preserving SDE probability path

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N} \left(\alpha_{1-t} \mathbf{x}_1, (1 - \alpha_{1-t}^2) \mathbf{I} \right) \Rightarrow \mathbf{f}(\mathbf{x}, \mathbf{x}_1, t) = \frac{\alpha'_{1-t}}{1 - \alpha_{1-t}^2} \cdot (\alpha_{1-t} \mathbf{x} - \mathbf{x}_1)$$

Flow matching vs score-based SDE models


SDE vector field

$$\mathbf{f}(\mathbf{x}, \mathbf{x}_1, t) = \frac{\alpha'_{1-t}}{1 - \alpha_{1-t}^2} \cdot (\alpha_{1-t} \mathbf{x} - \mathbf{x}_1)$$




FM vector field

$$\mathbf{f}(\mathbf{x}, \mathbf{x}_1, t) = \frac{\mathbf{x}_1 - \mathbf{x}}{1 - t}$$




Flow matching vs score-based SDE models


Trajectories




Diffusion



OT



Score matching w/ Diffusion



Flow Matching w/ OT

Summary

- ▶ Conditional flow matching allows to make the FM objective tractable.
- ▶ Conical gaussian paths is the example of the effective FM technique.
- ▶ Pair conditioning gives the same procedure, but it is more general.
- ▶ Diffusion and score-based model are the special case of flow matching approach with curved trajectories.