

Deep Generative Models

Lecture 12

Roman Isachenko



2026, Spring

Recap of Previous Lecture

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

Variance Exploding SDE (NCSN)

$$d\mathbf{x} = \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}, \quad \mathbf{f}(\mathbf{x}, t) = 0, \quad g(t) = \sqrt{\frac{d[\sigma^2(t)]}{dt}}$$

The variance grows since $\sigma(t)$ is a monotonically increasing function.

Variance Preserving SDE (DDPM)

$$d\mathbf{x} = -\frac{1}{2}\beta(t)\mathbf{x}(t)dt + \sqrt{\beta(t)} \cdot d\mathbf{w}$$

$$\mathbf{f}(\mathbf{x}, t) = -\frac{1}{2}\beta(t)\mathbf{x}(t), \quad g(t) = \sqrt{\beta(t)}$$

The variance is preserved if $\mathbf{x}(0)$ has unit variance.

Recap of Previous Lecture

Discrete-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}_0)} \mathbb{E}_{t \sim U\{1, T\}} \mathbb{E}_{q(\mathbf{x}_t | \mathbf{x}_0)} \| \mathbf{s}_{\theta, t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t | \mathbf{x}_0) \|_2^2$$

Continuous-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0, 1]} \mathbb{E}_{q(\mathbf{x}(t) | \mathbf{x}(0))} \| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t) | \mathbf{x}(0)) \|_2^2$$

NCSN

$$q(\mathbf{x}(t) | \mathbf{x}(0)) = \mathcal{N}(\mathbf{x}(0), [\sigma^2(t) - \sigma^2(0)] \cdot \mathbf{I})$$


DDPM

$$q(\mathbf{x}(t) | \mathbf{x}(0)) = \mathcal{N}\left(\mathbf{x}(0) e^{-\frac{1}{2} \int_0^t \beta(s) ds}, \left(1 - e^{-\int_0^t \beta(s) ds}\right) \cdot \mathbf{I}\right)$$

Recap of Previous Lecture

Sampling

To sample, solve the reverse SDE using numerical solvers (SDESolve).



- ▶ Discretizing the reverse SDE gives us ancestral sampling.
- ▶ If we use the probability flow ODE instead, then the reverse ODE yields DDIM sampling.

Recap of Previous Lecture

Consider ODE dynamics $\mathbf{x}(t)$ in the interval $t \in [0, 1]$ with $\mathbf{x}_0 \sim p_0(\mathbf{x}) = p(\mathbf{x})$, $\mathbf{x}_1 \sim p_1(\mathbf{x}) = p_{\text{data}}(\mathbf{x})$.

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t), \quad \text{with initial condition } \mathbf{x}(0) = \mathbf{x}_0.$$

KFP Theorem (Continuity Equation)

$$\frac{\partial p_t(\mathbf{x})}{\partial t} = -\text{div}(\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x})) \Leftrightarrow \frac{d \log p_t(\mathbf{x}(t))}{dt} = -\text{tr}\left(\frac{\partial \mathbf{f}(\mathbf{x}(t), t)}{\partial \mathbf{x}(t)}\right)$$

Solving the continuity equation using the adjoint method is complicated and unstable.

Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{f}(\mathbf{x}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_\theta$$

Recap of Previous Lecture

Introduce the latent variable \mathbf{z} :

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{z})p(\mathbf{z})d\mathbf{z}$$
$$\frac{\partial p_t(\mathbf{x}|\mathbf{z})}{\partial t} = -\text{div}(\mathbf{f}(\mathbf{x}, \mathbf{z}, t)p_t(\mathbf{x}|\mathbf{z})).$$

- ▶ $p_t(\mathbf{x}|\mathbf{z})$ is a **conditional probability path**
- ▶ $\mathbf{f}(\mathbf{x}, \mathbf{z}, t)$ is a **conditional vector field**

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t) \quad \Rightarrow \quad \frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, \mathbf{z}, t)$$

Theorem

The following vector field generates the probability path $p_t(\mathbf{x})$:

$$\mathbf{f}(\mathbf{x}, t) = \mathbb{E}_{p_t(\mathbf{z}|\mathbf{x})}\mathbf{f}(\mathbf{x}, \mathbf{z}, t) = \int \mathbf{f}(\mathbf{x}, \mathbf{z}, t) \frac{p_t(\mathbf{x}|\mathbf{z})p(\mathbf{z})}{p_t(\mathbf{x})} d\mathbf{z}$$

Recap of Previous Lecture

Flow Matching (FM)


$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{f}(\mathbf{x}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditional Flow Matching (CFM)

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Theorem

If $\text{supp}(p_t(\mathbf{x})) = \mathbb{R}^m$, then the optimal value of the FM objective equals the optimum for CFM.



Outline

1. Conditional Flow Matching

2. Conical Gaussian Paths

3. Linear Interpolation

Outline

1. Conditional Flow Matching

2. Conical Gaussian Paths

3. Linear Interpolation

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \| \mathbf{f}(\mathbf{x}, t) - \mathbf{f}_\theta(\mathbf{x}, t) \|^2 = \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \| \mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t) \|^2 \end{aligned}$$

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{x \sim p_t(x)} \|f(x, t) - f_{\theta}(x, t)\|^2 = \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_{\theta}(x, t)\|^2 \end{aligned}$$

Proof

$$\begin{aligned} \mathbb{E}_{x \sim p_t(x)} \|f(x, t) - f_{\theta}(x, t)\|^2 = \\ = \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} [\|f_{\theta}(x, t)\|^2 - 2f_{\theta}^{\top}(x, t)f(x, t)] + \text{const}(\theta) \end{aligned}$$

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{x \sim p_t(x)} \|f(x, t) - f_{\theta}(x, t)\|^2 = \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_{\theta}(x, t)\|^2 \end{aligned}$$

Proof

$$\begin{aligned} \mathbb{E}_{x \sim p_t(x)} \|f(x, t) - f_{\theta}(x, t)\|^2 = \\ = \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} [\|f_{\theta}(x, t)\|^2 - 2f_{\theta}^{\top}(x, t)f(x, t)] + \text{const}(\theta) \\ \mathbb{E}_{p_t(x)} [f_{\theta}^{\top}(x, t)f(x, t)] = \int p_t(x) \left[f_{\theta}^{\top}(x, t) \left(\int p_t(z|x) f(x, z, t) dz \right) \right] dx \end{aligned}$$

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{x \sim p_t(x)} \|f(x, t) - f_{\theta}(x, t)\|^2 = \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_{\theta}(x, t)\|^2 \end{aligned}$$

Proof

$$\begin{aligned} \mathbb{E}_{x \sim p_t(x)} \|f(x, t) - f_{\theta}(x, t)\|^2 = \\ = \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} [\|f_{\theta}(x, t)\|^2 - 2f_{\theta}^{\top}(x, t)f(x, t)] + \text{const}(\theta) \\ \mathbb{E}_{p_t(x)} [f_{\theta}^{\top}(x, t)f(x, t)] = \int p_t(x) \left[f_{\theta}^{\top}(x, t) \left(\int p_t(z|x)f(x, z, t)dz \right) \right] dx = \\ = \int \int p_t(x)p_t(z|x) [f_{\theta}^{\top}(x, t)f(x, z, t)] dz dx \end{aligned}$$

Conditional Flow Matching


Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{x \sim p_t(x)} \|f(x, t) - f_{\theta}(x, t)\|^2 = \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_{\theta}(x, t)\|^2 \end{aligned}$$


Proof

$$\begin{aligned} \mathbb{E}_{x \sim p_t(x)} \|f(x, t) - f_{\theta}(x, t)\|^2 = \\ = \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} [\|f_{\theta}(x, t)\|^2 - 2f_{\theta}^{\top}(x, t)f(x, t)] + \text{const}(\theta) \\ \mathbb{E}_{p_t(x)} [f_{\theta}^{\top}(x, t)f(x, t)] = \int p_t(x) \left[f_{\theta}^{\top}(x, t) \left(\int p_t(z|x)f(x, z, t)dz \right) \right] dx = \\ = \int \int p_t(x)p_t(z|x) [f_{\theta}^{\top}(x, t)f(x, z, t)] dz dx = \\ = \mathbb{E}_{p(z)} \mathbb{E}_{p_t(x|z)} [f_{\theta}^{\top}(x, t)f(x, z, t)] \end{aligned}$$

Conditional Flow Matching




Conditional Flow Matching



- ▶ We don't want to directly model $p_t(\mathbf{x})$, since it's complex.
- ▶ We've shown it's possible to solve the CFM task instead of the FM task.

Conditional Flow Matching



- ▶ We don't want to directly model $p_t(\mathbf{x})$, since it's complex.
- ▶ We've shown it's possible to solve the CFM task instead of the FM task.
- ▶ Let's choose a convenient conditioning latent variable \mathbf{z} .
- ▶ We'll parametrize $p_t(\mathbf{x}|\mathbf{z})$ instead of $p_t(\mathbf{x})$. It should satisfy the following constraints:

$$p(\mathbf{x}) = \mathcal{N}(0, \mathbf{I}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); \quad p_{\text{data}}(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}).$$

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \| \mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t) \|^2 \rightarrow \min_{\theta}$$

$$p(\mathbf{x}) = \mathcal{N}(0, \mathbf{I}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); \quad p_{\text{data}}(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}).$$

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \| \mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t) \|^2 \rightarrow \min_{\theta}$$

$$p(\mathbf{x}) = \mathcal{N}(0, \mathbf{I}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); \quad p_{\text{data}}(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}).$$

Training

1. Sample a time $t \sim U[0, 1]$ and $\mathbf{z} \sim p(\mathbf{z})$.
2. Draw $\mathbf{x}_t \sim p_t(\mathbf{x}|\mathbf{z})$.
3. Compute the loss $\mathcal{L} = \| \mathbf{f}(\mathbf{x}_t, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}_t, t) \|^2$.

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

$$p(\mathbf{x}) = \mathcal{N}(0, \mathbf{I}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); \quad p_{\text{data}}(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}).$$

Training

1. Sample a time $t \sim U[0, 1]$ and $\mathbf{z} \sim p(\mathbf{z})$.
2. Draw $\mathbf{x}_t \sim p_t(\mathbf{x}|\mathbf{z})$.
3. Compute the loss $\mathcal{L} = \|\mathbf{f}(\mathbf{x}_t, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}_t, t)\|^2$.

Sampling

1. Sample $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$.
2. Solve the ODE to obtain \mathbf{x}_1 :

$$\mathbf{x}_1 = \text{ODESolve}_f(\mathbf{x}_0, \theta, t_0 = 0, t_1 = 1).$$

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_\theta(x, t)\|^2 \rightarrow \min_{\theta}$$

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_\theta(x, t)\|^2 \rightarrow \min_{\theta}$$

Open Questions

- Q1 How should we choose the conditioning latent variable z ?
- Q2 How can we define $p_t(x|z)$ to enforce the constraints?

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_\theta(x, t)\|^2 \rightarrow \min_{\theta}$$

Open Questions

- Q1 How should we choose the conditioning latent variable z ?
- Q2 How can we define $p_t(x|z)$ to enforce the constraints?

Gaussian Conditional Probability Path [Q2]

$$p_t(x|z) = \mathcal{N}(\mu_t(z), \sigma_t^2(z))$$

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_\theta(x, t)\|^2 \rightarrow \min_{\theta}$$

Open Questions

- Q1 How should we choose the conditioning latent variable z ?
- Q2 How can we define $p_t(x|z)$ to enforce the constraints?

Gaussian Conditional Probability Path [Q2]

$$p_t(x|z) = \mathcal{N}(\mu_t(z), \sigma_t^2(z))$$

- ▶ There are infinitely many vector fields that generate a particular probability path.
- ▶ Let's consider the following dynamics:

$$x_t = \mu_t(z) + \sigma_t(z) \odot \epsilon, \quad \text{with fixed } \epsilon \sim \mathcal{N}(0, I)$$

Flow Matching

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{z}) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{z}), \boldsymbol{\sigma}_t^2(\mathbf{z})) ; \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{z}) + \boldsymbol{\sigma}_t(\mathbf{z}) \odot \boldsymbol{\epsilon}$$

Is it possible to derive the expression for $\mathbf{f}(\mathbf{x}_t, \mathbf{z}, t)$?

Flow Matching

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{z}) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{z}), \boldsymbol{\sigma}_t^2(\mathbf{z})) ; \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{z}) + \boldsymbol{\sigma}_t(\mathbf{z}) \odot \boldsymbol{\epsilon}$$

Is it possible to derive the expression for $\mathbf{f}(\mathbf{x}_t, \mathbf{z}, t)$?

Statement

$$\mathbf{f}(\mathbf{x}_t, \mathbf{z}, t) = \boldsymbol{\mu}'_t(\mathbf{z}) + \frac{\boldsymbol{\sigma}'_t(\mathbf{z})}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{z}))$$

Flow Matching

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{z}) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{z}), \boldsymbol{\sigma}_t^2(\mathbf{z})) ; \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{z}) + \boldsymbol{\sigma}_t(\mathbf{z}) \odot \boldsymbol{\epsilon}$$

Is it possible to derive the expression for $\mathbf{f}(\mathbf{x}_t, \mathbf{z}, t)$?

Statement

$$\mathbf{f}(\mathbf{x}_t, \mathbf{z}, t) = \boldsymbol{\mu}'_t(\mathbf{z}) + \frac{\boldsymbol{\sigma}'_t(\mathbf{z})}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{z}))$$

Proof

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{z}, t); \quad \boldsymbol{\epsilon} = \frac{1}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{z}))$$

Flow Matching

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{z}) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{z}), \boldsymbol{\sigma}_t^2(\mathbf{z})) ; \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{z}) + \boldsymbol{\sigma}_t(\mathbf{z}) \odot \boldsymbol{\epsilon}$$

Is it possible to derive the expression for $\mathbf{f}(\mathbf{x}_t, \mathbf{z}, t)$?

Statement

$$\mathbf{f}(\mathbf{x}_t, \mathbf{z}, t) = \boldsymbol{\mu}'_t(\mathbf{z}) + \frac{\boldsymbol{\sigma}'_t(\mathbf{z})}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{z}))$$

Proof

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{z}, t); \quad \boldsymbol{\epsilon} = \frac{1}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{z}))$$

$$\frac{d\mathbf{x}_t}{dt} = \boldsymbol{\mu}'_t(\mathbf{z}) + \boldsymbol{\sigma}'_t(\mathbf{z}) \odot \boldsymbol{\epsilon} = \boldsymbol{\mu}'_t(\mathbf{z}) + \frac{\boldsymbol{\sigma}'_t(\mathbf{z})}{\boldsymbol{\sigma}_t(\mathbf{z})} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{z}))$$

Outline

1. Conditional Flow Matching

2. Conical Gaussian Paths

3. Linear Interpolation

Endpoint Conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_\theta(x, t)\|^2 \rightarrow \min_{\theta}$$

Let's define our latent variable z .

Endpoint Conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Let's define our latent variable \mathbf{z} .

Conditioning Latent Variable [Q1]

Let us choose $\mathbf{z} = \mathbf{x}_1$. Then $p(\mathbf{z}) = p_1(\mathbf{x}_1)$.

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{x}_1) p_1(\mathbf{x}_1) d\mathbf{x}_1$$

Endpoint Conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Let's define our latent variable \mathbf{z} .

Conditioning Latent Variable [Q1]

Let us choose $\mathbf{z} = \mathbf{x}_1$. Then $p(\mathbf{z}) = p_1(\mathbf{x}_1)$.

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{x}_1) p_1(\mathbf{x}_1) d\mathbf{x}_1$$

We need to ensure the boundary constraints:

$$\begin{cases} p(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); (= \mathcal{N}(0, \mathbf{I})) \\ p_{\text{data}}(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}) \end{cases}$$

Endpoint Conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_\theta(x, t)\|^2 \rightarrow \min_{\theta}$$

Let define our latent variable z .

Conditioning Latent Variable [Q1]

Let us choose $z = x_1$. Then $p(z) = p_1(x_1)$.

$$p_t(x) = \int p_t(x|x_1)p_1(x_1)dx_1$$

We need to ensure the boundary constraints:

$$\begin{cases} p(x) = \mathbb{E}_{p(z)} p_0(x|z); (= \mathcal{N}(0, I)) \\ p_{\text{data}}(x) = \mathbb{E}_{p(z)} p_1(x|z) \end{cases} \Rightarrow \begin{cases} p_0(x|x_1) = \mathcal{N}(0, I); \\ p_1(x|x_1) = \delta(x - x_1) \end{cases}$$

Conical Gaussian Paths

$$p_0(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(0, \mathbf{I}); \quad p_1(\mathbf{x}|\mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1)$$

Gaussian Conditional Probability Path


$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_1) \odot \boldsymbol{\epsilon}$$

Conical Gaussian Paths

$$p_0(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(0, \mathbf{I}); \quad p_1(\mathbf{x}|\mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1)$$

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_1) \odot \boldsymbol{\epsilon}$$




Conical Gaussian Paths

$$p_0(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(0, \mathbf{I}); \quad p_1(\mathbf{x}|\mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1)$$

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_1) \odot \epsilon$$



Let's consider straight conditional paths:


$$\begin{cases} \boldsymbol{\mu}_t(\mathbf{x}_1) = t\mathbf{x}_1 \\ \boldsymbol{\sigma}_t(\mathbf{x}_1) = 1 - t \end{cases}$$

Conical Gaussian Paths

$$p_0(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(0, \mathbf{I}); \quad p_1(\mathbf{x}|\mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1)$$

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_1) \odot \epsilon$$



Let's consider straight conditional paths:

$$\begin{cases} \boldsymbol{\mu}_t(\mathbf{x}_1) = t\mathbf{x}_1 \\ \boldsymbol{\sigma}_t(\mathbf{x}_1) = 1 - t \end{cases} \Rightarrow \begin{cases} p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \cdot \mathbf{I}) \\ \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0 \end{cases}$$

Conical Gaussian Paths

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \mathbf{I}); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_1))$$

Conical Gaussian Paths

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \mathbf{I}); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_1))$$

$$\mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) = \mathbf{x}_1 - \frac{1}{1-t} \cdot (\mathbf{x}_t - t\mathbf{x}_1) = \frac{\mathbf{x}_1 - \textcolor{teal}{\mathbf{x}_t}}{1-t}$$

Conical Gaussian Paths

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \mathbf{I}); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_1))$$

$$\begin{aligned}\mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) &= \mathbf{x}_1 - \frac{1}{1-t} \cdot (\mathbf{x}_t - t\mathbf{x}_1) = \frac{\mathbf{x}_1 - \textcolor{teal}{\mathbf{x}_t}}{1-t} = \\ &= \frac{\mathbf{x}_1 - \textcolor{teal}{t\mathbf{x}_1} - (1-t)\mathbf{x}_0}{1-t}\end{aligned}$$

Conical Gaussian Paths

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \mathbf{I}); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_1))$$

$$\begin{aligned}\mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) &= \mathbf{x}_1 - \frac{1}{1-t} \cdot (\mathbf{x}_t - t\mathbf{x}_1) = \frac{\mathbf{x}_1 - \textcolor{teal}{\mathbf{x}_t}}{1-t} = \\ &= \frac{\mathbf{x}_1 - \textcolor{teal}{t\mathbf{x}_1} - (1-t)\mathbf{x}_0}{1-t} = \mathbf{x}_1 - \mathbf{x}_0\end{aligned}$$


Conical Gaussian Paths

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \mathbf{I}); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_1))$$

$$\begin{aligned}\mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t) &= \mathbf{x}_1 - \frac{1}{1-t} \cdot (\mathbf{x}_t - t\mathbf{x}_1) = \frac{\mathbf{x}_1 - \mathbf{x}_t}{1-t} = \\ &= \frac{\mathbf{x}_1 - t\mathbf{x}_1 - (1-t)\mathbf{x}_0}{1-t} = \mathbf{x}_1 - \mathbf{x}_0\end{aligned}$$



The conditional vector field $\mathbf{f}(\mathbf{x}_t, \mathbf{x}_1, t)$ defines straight lines between $p_{\text{data}}(\mathbf{x})$ and $\mathcal{N}(0, \mathbf{I})$.

Conical Gaussian Paths

$$\begin{aligned} & \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \| \mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t) \|^2 = \\ &= \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x}_1 \sim p_{\text{data}}(\mathbf{x})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{x}_1)} \left\| \frac{\mathbf{x}_1 - \mathbf{x}}{1-t} - \mathbf{f}_\theta(\mathbf{x}, t) \right\|^2 \end{aligned}$$

Conical Gaussian Paths

$$\begin{aligned} & \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \| \mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t) \|^2 = \\ &= \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x}_1 \sim p_{\text{data}}(\mathbf{x})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{x}_1)} \left\| \frac{\mathbf{x}_1 - \mathbf{x}}{1-t} - \mathbf{f}_\theta(\mathbf{x}, t) \right\|^2 = \\ &= \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x}_1 \sim p_{\text{data}}(\mathbf{x})} \mathbb{E}_{\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})} \| (\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_\theta(t\mathbf{x}_1 + (1-t)\mathbf{x}_0, t) \|^2 \end{aligned}$$

Conical Gaussian Paths

$$\begin{aligned} & \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(\textcolor{brown}{x}, z, t) - f_\theta(x, t)\|^2 = \\ &= \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{x_1 \sim p_{\text{data}}(x)} \mathbb{E}_{x \sim p_t(x|x_1)} \left\| \frac{x_1 - x}{1-t} - f_\theta(\textcolor{teal}{x}, t) \right\|^2 = \\ &= \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{x_1 \sim p_{\text{data}}(x)} \mathbb{E}_{x_0 \sim \mathcal{N}(0, I)} \| (x_1 - x_0) - f_\theta(tx_1 + (1-t)x_0, t) \|^2 \end{aligned}$$

- ▶ We fit straight lines between the noise distribution $p(x)$ and the data distribution $p_{\text{data}}(x)$.
- ▶ The **marginal** path $p_t(x)$ does not give straight lines.

Conical Gaussian Paths

$$\begin{aligned} & \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{z \sim p(z)} \mathbb{E}_{x \sim p_t(x|z)} \|f(x, z, t) - f_\theta(x, t)\|^2 = \\ &= \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{x_1 \sim p_{\text{data}}(x)} \mathbb{E}_{x \sim p_t(x|x_1)} \left\| \frac{x_1 - x}{1-t} - f_\theta(x, t) \right\|^2 = \\ &= \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{x_1 \sim p_{\text{data}}(x)} \mathbb{E}_{x_0 \sim \mathcal{N}(0, I)} \| (x_1 - x_0) - f_\theta(tx_1 + (1-t)x_0, t) \|^2 \end{aligned}$$

- ▶ We fit straight lines between the noise distribution $p(x)$ and the data distribution $p_{\text{data}}(x)$.
- ▶ The **marginal** path $p_t(x)$ does not give straight lines.



Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x}_1 \sim p_{\text{data}}(\mathbf{x})} \mathbb{E}_{\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})} \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_{\theta}(\mathbf{x}_t, t)\|^2 \rightarrow \min_{\theta}$$

Training

1. Sample $\mathbf{x}_1 \sim p_{\text{data}}(\mathbf{x})$.
2. Sample time $t \sim U[0, 1]$ and $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$.
3. Obtain the noisy image $\mathbf{x}_t = t\mathbf{x}_1 + (1 - t)\mathbf{x}_0$.
4. Compute the loss $\mathcal{L} = \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_{\theta}(\mathbf{x}_t, t)\|^2$.

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x}_1 \sim p_{\text{data}}(\mathbf{x})} \mathbb{E}_{\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})} \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_{\theta}(\mathbf{x}_t, t)\|^2 \rightarrow \min_{\theta}$$

Training


1. Sample $\mathbf{x}_1 \sim p_{\text{data}}(\mathbf{x})$.
2. Sample time $t \sim U[0, 1]$ and $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$.
3. Obtain the noisy image $\mathbf{x}_t = t\mathbf{x}_1 + (1 - t)\mathbf{x}_0$.
4. Compute the loss $\mathcal{L} = \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_{\theta}(\mathbf{x}_t, t)\|^2$.

Sampling


1. Sample $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$.
2. Solve the ODE to obtain \mathbf{x}_1 :

$$\mathbf{x}_1 = \text{ODESolve}_f(\mathbf{x}_0, \theta, t_0 = 0, t_1 = 1)$$

Conical Gaussian Paths



Conical Gaussian Paths



- ▶ Conical gaussian paths give us the way to construct generative model.
- ▶ Now we extend it to image-to-image formulation (mapping between two distinct $p_{\text{data}_1}(\mathbf{x})$ and $p_{\text{data}_2}(\mathbf{x})$).

Outline

1. Conditional Flow Matching

2. Conical Gaussian Paths

3. Linear Interpolation

Pair Conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditioning Latent Variable [Q1]

Let us choose $\mathbf{z} = (\mathbf{x}_0, \mathbf{x}_1)$. Then $p(\mathbf{z}) = p(\mathbf{x}_0, \mathbf{x}_1) = p_0(\mathbf{x}_0)p_1(\mathbf{x}_1)$.

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) p_0(\mathbf{x}_0) p_1(\mathbf{x}_1) d\mathbf{x}_0 d\mathbf{x}_1$$

Pair Conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditioning Latent Variable [Q1]

Let us choose $\mathbf{z} = (\mathbf{x}_0, \mathbf{x}_1)$. Then $p(\mathbf{z}) = p(\mathbf{x}_0, \mathbf{x}_1) = p_0(\mathbf{x}_0)p_1(\mathbf{x}_1)$.

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) p_0(\mathbf{x}_0) p_1(\mathbf{x}_1) d\mathbf{x}_0 d\mathbf{x}_1$$

We must enforce boundary constraints:

$$\begin{cases} p_0(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); \\ p_1(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}) \end{cases}$$

Pair Conditioning

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditioning Latent Variable [Q1]

Let us choose $\mathbf{z} = (\mathbf{x}_0, \mathbf{x}_1)$. Then $p(\mathbf{z}) = p(\mathbf{x}_0, \mathbf{x}_1) = p_0(\mathbf{x}_0)p_1(\mathbf{x}_1)$.

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) p_0(\mathbf{x}_0) p_1(\mathbf{x}_1) d\mathbf{x}_0 d\mathbf{x}_1$$

We must enforce boundary constraints:

$$\begin{cases} p_0(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}); \\ p_1(\mathbf{x}) = \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}) \end{cases} \Rightarrow \begin{cases} p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0) \\ p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1) \end{cases}$$

Linear Interpolation

$$p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0); \quad p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1)$$

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_0, \mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1) \odot \boldsymbol{\epsilon}$$

Linear Interpolation

$$p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0); \quad p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1)$$

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_0, \mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1) \odot \boldsymbol{\epsilon}$$

Let's consider straight conditional paths:

$$\begin{cases} \boldsymbol{\mu}_t(\mathbf{x}_1) = t\mathbf{x}_1 + (1 - t)\mathbf{x}_0 \\ \boldsymbol{\sigma}_t(\mathbf{x}_1) = \boldsymbol{\epsilon} \end{cases}$$

Linear Interpolation

$$p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0); \quad p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1)$$

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_0, \mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1) \odot \boldsymbol{\epsilon}$$

Let's consider straight conditional paths:

$$\begin{cases} \boldsymbol{\mu}_t(\mathbf{x}_1) = t\mathbf{x}_1 + (1-t)\mathbf{x}_0 \\ \boldsymbol{\sigma}_t(\mathbf{x}_1) = \boldsymbol{\epsilon} \end{cases} \Rightarrow \begin{cases} p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0) \\ p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1) \end{cases}$$

Linear Interpolation




$$p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0); \quad p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1)$$

Gaussian Conditional Probability Path

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}(\boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1), \boldsymbol{\sigma}_t^2(\mathbf{x}_0, \mathbf{x}_1)); \quad \mathbf{x}_t = \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1) + \boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1) \odot \boldsymbol{\epsilon}$$

Let's consider straight conditional paths:

$$\begin{cases} \boldsymbol{\mu}_t(\mathbf{x}_1) = t\mathbf{x}_1 + (1-t)\mathbf{x}_0 \\ \boldsymbol{\sigma}_t(\mathbf{x}_1) = \boldsymbol{\epsilon} \end{cases} \Rightarrow \begin{cases} p_0(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_0) \\ p_1(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \delta(\mathbf{x} - \mathbf{x}_1) \end{cases}$$



Flow Matching: Conical Paths vs. Linear Interpolation

$$\mathbf{z} = \mathbf{x}_1$$

$$\mathbf{z} = (\mathbf{x}_0, \mathbf{x}_1)$$

$$p_t(\mathbf{x}|\mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1, (1-t)^2 \mathbf{I})$$

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}(t\mathbf{x}_1 + (1-t)\mathbf{x}_0, \epsilon^2 \mathbf{I})$$

$$\mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

$$\mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Flow Matching: Conical Paths vs. Linear Interpolation

$$z = x_1$$

$$p_t(x|x_1) = \mathcal{N}(tx_1, (1-t)^2 I)$$


$$x_t = tx_1 + (1-t)x_0$$

$$z = (x_0, x_1)$$


$$p_t(x|x_0, x_1) = \mathcal{N}(tx_1 + (1-t)x_0, \epsilon^2 I)$$

$$x_t = tx_1 + (1-t)x_0$$

Flow Matching (Lipman et al.)



Conditional Flow Matching



Linear Interpolation

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}\left(t\mathbf{x}_1 + (1-t)\mathbf{x}_0, \epsilon^2 \mathbf{I}\right); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_0, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_0, \mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_0, \mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1))$$

Linear Interpolation

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}\left(t\mathbf{x}_1 + (1-t)\mathbf{x}_0, \epsilon^2 \mathbf{I}\right); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_0, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_0, \mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_0, \mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1))$$
$$\mathbf{f}(\mathbf{x}_t, \mathbf{x}_0, \mathbf{x}_1, t) = \mathbf{x}_1 - \mathbf{x}_0$$

Linear Interpolation

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}\left(t\mathbf{x}_1 + (1-t)\mathbf{x}_0, \epsilon^2 \mathbf{I}\right); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_0, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_0, \mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_0, \mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1))$$
$$\mathbf{f}(\mathbf{x}_t, \mathbf{x}_0, \mathbf{x}_1, t) = \mathbf{x}_1 - \mathbf{x}_0$$

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 =$$
$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{(\mathbf{x}_0, \mathbf{x}_1) \sim p(\mathbf{x}_0, \mathbf{x}_1)} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1)} \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2$$

Linear Interpolation

$$p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1) = \mathcal{N}\left(t\mathbf{x}_1 + (1-t)\mathbf{x}_0, \epsilon^2 \mathbf{I}\right); \quad \mathbf{x}_t = t\mathbf{x}_1 + (1-t)\mathbf{x}_0$$

Conditional Vector Field

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{f}(\mathbf{x}_t, \mathbf{x}_0, \mathbf{x}_1, t) = \boldsymbol{\mu}'_t(\mathbf{x}_0, \mathbf{x}_1) + \frac{\boldsymbol{\sigma}'_t(\mathbf{x}_0, \mathbf{x}_1)}{\boldsymbol{\sigma}_t(\mathbf{x}_0, \mathbf{x}_1)} \odot (\mathbf{x}_t - \boldsymbol{\mu}_t(\mathbf{x}_0, \mathbf{x}_1))$$
$$\mathbf{f}(\mathbf{x}_t, \mathbf{x}_0, \mathbf{x}_1, t) = \mathbf{x}_1 - \mathbf{x}_0$$

Conditional Flow Matching

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{f}(\mathbf{x}, \mathbf{z}, t) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2 =$$
$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{(\mathbf{x}_0, \mathbf{x}_1) \sim p(\mathbf{x}_0, \mathbf{x}_1)} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{x}_0, \mathbf{x}_1)} \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_\theta(\mathbf{x}, t)\|^2$$

- ▶ This yields the same procedure as for conical paths!
- ▶ Now, we do not require that $p_0(\mathbf{x})$ is necessarily $\mathcal{N}(0, \mathbf{I})$.

Conditional Flow Matching

- ▶ This conditioning allows us to transport any distribution $p_0(\mathbf{x})$ to any distribution $p_1(\mathbf{x})$.
- ▶ It's possible to apply this approach to paired tasks, e.g., style transfer.

Conditional Flow Matching

- ▶ This conditioning allows us to transport any distribution $p_0(\mathbf{x})$ to any distribution $p_1(\mathbf{x})$.
- ▶ It's possible to apply this approach to paired tasks, e.g., style transfer.

Training Procedure

1. Sample $(\mathbf{x}_0, \mathbf{x}_1) \sim p(\mathbf{x}_0, \mathbf{x}_1)$.
2. Sample time $t \sim U[0, 1]$.
3. Compute the noisy image $\mathbf{x}_t = t\mathbf{x}_1 + (1 - t)\mathbf{x}_0$.
4. Compute the loss $\mathcal{L} = \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_{\theta}(\mathbf{x}_t, t)\|^2$.

Conditional Flow Matching

- ▶ This conditioning allows us to transport any distribution $p_0(\mathbf{x})$ to any distribution $p_1(\mathbf{x})$.
- ▶ It's possible to apply this approach to paired tasks, e.g., style transfer.

Training Procedure


1. Sample $(\mathbf{x}_0, \mathbf{x}_1) \sim p(\mathbf{x}_0, \mathbf{x}_1)$.
2. Sample time $t \sim U[0, 1]$.
3. Compute the noisy image $\mathbf{x}_t = t\mathbf{x}_1 + (1 - t)\mathbf{x}_0$.
4. Compute the loss $\mathcal{L} = \|(\mathbf{x}_1 - \mathbf{x}_0) - \mathbf{f}_{\theta}(\mathbf{x}_t, t)\|^2$.

Sampling

1. Sample $\mathbf{x}_0 \sim p_0(\mathbf{x})$.
2. Solve the ODE to obtain \mathbf{x}_1 :

$$\mathbf{x}_1 = \text{ODESolve}_f(\mathbf{x}_0, \theta, t_0 = 0, t_1 = 1)$$

Stable Diffusion 3: Scalable Flow Matching



Summary

- ▶ Conditional flow matching makes the FM objective tractable.
- ▶ Conical Gaussian paths use endpoint conditioning $\mathbf{z} = \mathbf{x}_1$ and serve as an effective FM technique.
- ▶ Linear Interpolation paths use pair conditioning $\mathbf{z} = (\mathbf{x}_0, \mathbf{x}_1)$ and yields the same procedure, but is more general (suitable for paired tasks).