

Deep Generative Models

Lecture 10

Roman Isachenko



AI Masters

2026, Spring

Recap of Previous Lecture

Continuous-Time Dynamics (Neural ODE)

$$\frac{d\mathbf{x}(t)}{dt} = \mathbf{v}_{\theta}(\mathbf{x}(t), t); \quad \text{where } \mathbf{x}(t_0) = \mathbf{x}_0.$$

$$\psi_{t_1}(\mathbf{x}_0) = \int_{t_0}^{t_1} \mathbf{v}_{\theta}(\mathbf{x}(t), t) dt + \mathbf{x}_0 \approx \text{ODESolve}_v(\mathbf{x}_0, \theta, t_0, t_1).$$

- ▶ $\mathbf{v}_{\theta} : \mathbb{R}^m \times [t_0, t_1] \rightarrow \mathbb{R}^m$ is a vector field.
- ▶ $\psi : \mathbb{R}^m \times [t_0, t_1] \rightarrow \mathbb{R}^m$ is a flow (the solution of ODE):

Euler Update Step (ODESolve)


$$\mathbf{x}(t+h) = \mathbf{x}(t) + h \cdot \mathbf{v}_{\theta}(\mathbf{x}(t), t)$$

More advanced numerical methods (such as Runge-Kutta) are often used instead of unstable Euler update step.

Recap of Previous Lecture

$$\frac{d\mathbf{x}(t)}{dt} = \mathbf{v}_\theta(\mathbf{x}(t), t); \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

- ▶ Suppose $\mathbf{x}(0)$ is a random variable with density $p_0(\mathbf{x})$. Then, $\mathbf{x}(t)$ is a random variable with density $p_t(\mathbf{x})$.
- ▶ $p_t(\mathbf{x}) = p(\mathbf{x}, t)$ describes the **probability path** between $p_0(\mathbf{x})$ and $p_1(\mathbf{x})$.



Recap of Previous Lecture

Theorem (Picard)

If $\mathbf{v}$ is continuously differentiable in $\mathbf{x}$ (with bounded derivative) and continuous in t , then the ODE has a **unique** solution.

This means the ODE can be **uniquely inverted**:

$$\begin{aligned}\mathbf{x}(1) &= \mathbf{x}(0) + \int_0^1 \mathbf{v}_\theta(\mathbf{x}(t), t) dt \\ \mathbf{x}(0) &= \mathbf{x}(1) + \int_1^0 \mathbf{v}_\theta(\mathbf{x}(t), t) dt\end{aligned}$$

Note: Unlike discrete-time NF, $\mathbf{v}$ need not be invertible (uniqueness ensures bijectivity).

How can we compute $p_t(\mathbf{x})$ for any t ?

Recap of Previous Lecture

$$\frac{d\mathbf{x}(t)}{dt} = \mathbf{v}_\theta(\mathbf{x}(t), t); \quad \text{with initial condition } \mathbf{x}(t_0) = \mathbf{x}_0$$

Theorem (Continuity Equation)

If $\mathbf{v}$ is continuously differentiable in $\mathbf{x}$ (with bounded derivative) and continuous in t , and $p_t(\mathbf{x}) > 0$, then

$$\frac{d \log p_t(\mathbf{x}(t))}{dt} = -\text{tr} \left(\frac{\partial \mathbf{v}(\mathbf{x}(t), t)}{\partial \mathbf{x}(t)} \right)$$

Solution of the Continuity Equation

$$\log p_1(\mathbf{x}(1)) = \log p_0(\mathbf{x}(0)) - \int_0^1 \text{tr} \left(\frac{\partial \mathbf{v}(\mathbf{x}(t), t)}{\partial \mathbf{x}(t)} \right) dt.$$

- ▶ This solution gives us the density **along the trajectory**.
- ▶ However, it's difficult to efficiently estimate **the last term**.

Recap of Previous Lecture

SDE Basics

Let's define a stochastic process $\mathbf{x}(t)$ with initial condition $\mathbf{x}(0) \sim p_0(\mathbf{x})$:

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w},$$

where $\mathbf{w}(t)$ is the standard Wiener process (Brownian motion):

$$\mathbf{w}(t) - \mathbf{w}(s) \sim \mathcal{N}(0, (t-s)\mathbf{I}), \quad d\mathbf{w} = \epsilon \cdot \sqrt{dt}, \text{ where } \epsilon \sim \mathcal{N}(0, \mathbf{I}).$$

Discretization of SDE (Euler Method) - SDEsolve

$$\mathbf{x}(t + dt) = \mathbf{x}(t) + \mathbf{f}(\mathbf{x}(t), t) \cdot dt + g(t) \cdot \epsilon \cdot \sqrt{dt}$$

- ▶ At each time t , we have the density $p_t(\mathbf{x}) = p(\mathbf{x}, t)$.
- ▶ $p : \mathbb{R}^m \times [0, 1] \rightarrow \mathbb{R}_+$ is a **probability path** between $p_0(\mathbf{x})$ and $p_1(\mathbf{x})$.

Recap of Previous Lecture

Theorem (Kolmogorov-Fokker-Planck)

If $p_t(\mathbf{x}) \in C^{1,2}$ (i.e., C^1 in t and C^2 in $\mathbf{x}$), then

$$\frac{\partial p_t(\mathbf{x})}{\partial t} = -\operatorname{div}(\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x})) + \frac{1}{2}g^2(t)\Delta_{\mathbf{x}}p_t(\mathbf{x})$$

Langevin SDE (Special Case)

$$d\mathbf{x} = \frac{1}{2} \frac{\partial}{\partial \mathbf{x}} \log p_{\theta}(\mathbf{x}) dt + \mathbf{1} \cdot d\mathbf{w}$$

The density $p_{\theta}(\mathbf{x})$ is a **stationary** distribution for the SDE.

Langevin Dynamics

Samples from the following dynamics will come from $p_{\theta}(\mathbf{x})$ under mild regularity conditions for a small enough η :

$$\mathbf{x}_{t+1} = \mathbf{x}_t + \frac{\eta}{2} \nabla_{\mathbf{x}_t} \log p_{\theta}(\mathbf{x}_t) + \sqrt{\eta} \cdot \epsilon, \quad \epsilon \sim \mathcal{N}(0, \mathbf{I}).$$

Recap of Previous Lecture

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

Variance Exploding SDE (NCSN)

$$d\mathbf{x} = \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}, \quad \mathbf{f}(\mathbf{x}, t) = 0, \quad g(t) = \sqrt{\frac{d[\sigma^2(t)]}{dt}}$$

The variance grows since $\sigma(t)$ is a monotonically increasing function.

Variance Preserving SDE (DDPM)

$$d\mathbf{x} = -\frac{1}{2}\beta(t)\mathbf{x}(t)dt + \sqrt{\beta(t)} \cdot d\mathbf{w}$$

$$\mathbf{f}(\mathbf{x}, t) = -\frac{1}{2}\beta(t)\mathbf{x}(t), \quad g(t) = \sqrt{\beta(t)}$$

The variance is preserved if $\mathbf{x}(0)$ has unit variance.

Song Y., et al. Score-Based Generative Modeling through Stochastic Differential Equations, 2020

Outline

1. Probability Flow ODE
2. Reverse SDE
3. Score-Based Generative Models Through SDEs

Outline

1. Probability Flow ODE
2. Reverse SDE
3. Score-Based Generative Models Through SDEs

Probability Flow ODE

ODE and Continuity Equation

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt$$

$$\frac{d \log p_t(\mathbf{x}(t))}{dt} = -\text{tr} \left(\frac{\partial \mathbf{v}(\mathbf{x}(t), t)}{\partial \mathbf{x}(t)} \right) \Leftrightarrow \frac{\partial p_t(\mathbf{x})}{\partial t} = -\text{div}(\mathbf{v}(\mathbf{x}, t)p_t(\mathbf{x}))$$

The only source of randomness is the initial distribution $p_0(\mathbf{x})$.

Probability Flow ODE

ODE and Continuity Equation

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt$$

$$\frac{d \log p_t(\mathbf{x}(t))}{dt} = -\text{tr} \left(\frac{\partial \mathbf{v}(\mathbf{x}(t), t)}{\partial \mathbf{x}(t)} \right) \Leftrightarrow \frac{\partial p_t(\mathbf{x})}{\partial t} = -\text{div}(\mathbf{v}(\mathbf{x}, t)p_t(\mathbf{x}))$$

The only source of randomness is the initial distribution $p_0(\mathbf{x})$.

SDE and KFP Equation

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

$$\frac{\partial p_t(\mathbf{x})}{\partial t} = -\text{div}(\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x})) + \frac{1}{2}g^2(t)\Delta_{\mathbf{x}}p_t(\mathbf{x})$$

Now there are two sources of randomness: the initial distribution $p_0(\mathbf{x})$ and the Wiener process $\mathbf{w}(t)$.

Probability Flow ODE

Theorem

Suppose the SDE $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ induces the probability path $p_t(\mathbf{x})$ with $p_t(\mathbf{x}) > 0$. Then, there exists an ODE with the same probability path $p_t(\mathbf{x})$, given by


$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

Probability Flow ODE

Theorem

Suppose the SDE $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ induces the probability path $p_t(\mathbf{x})$ with $p_t(\mathbf{x}) > 0$. Then, there exists an ODE with the same probability path $p_t(\mathbf{x})$, given by

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$



Song Y., et al. *Score-Based Generative Modeling through Stochastic Differential Equations*, 2020

Probability Flow ODE

Theorem

Suppose the SDE $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ induces the probability path $p_t(\mathbf{x})$ with $p_t(\mathbf{x}) > 0$. Then, there exists an ODE with the same probability path $p_t(\mathbf{x})$, given by

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

Proof

$$\frac{\partial p_t(\mathbf{x})}{\partial t} = \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} [\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x})] + \frac{1}{2}g^2(t)\frac{\partial^2 p_t(\mathbf{x})}{\partial \mathbf{x}^2} \right)$$

Probability Flow ODE

Theorem

Suppose the SDE $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ induces the probability path $p_t(\mathbf{x})$ with $p_t(\mathbf{x}) > 0$. Then, there exists an ODE with the same probability path $p_t(\mathbf{x})$, given by

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

Proof

$$\begin{aligned} \frac{\partial p_t(\mathbf{x})}{\partial t} &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} [\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x})] + \frac{1}{2}g^2(t)\frac{\partial^2 p_t(\mathbf{x})}{\partial \mathbf{x}^2} \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x}) - \frac{1}{2}g^2(t)\frac{\partial p_t(\mathbf{x})}{\partial \mathbf{x}} \right] \right) \end{aligned}$$

Probability Flow ODE

Theorem

Suppose the SDE $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ induces the probability path $p_t(\mathbf{x})$ with $p_t(\mathbf{x}) > 0$. Then, there exists an ODE with the same probability path $p_t(\mathbf{x})$, given by

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

Proof

$$\begin{aligned} \frac{\partial p_t(\mathbf{x})}{\partial t} &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} [\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x})] + \frac{1}{2}g^2(t)\frac{\partial^2 p_t(\mathbf{x})}{\partial \mathbf{x}^2} \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x}) - \frac{1}{2}g^2(t)\frac{\partial p_t(\mathbf{x})}{\partial \mathbf{x}} \right] \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x}) - \frac{1}{2}g^2(t)p_t(\mathbf{x})\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right] \right) \end{aligned}$$

Probability Flow ODE

Theorem

Suppose the SDE $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ induces the probability path $p_t(\mathbf{x})$ with $p_t(\mathbf{x}) > 0$. Then, there exists an ODE with the same probability path $p_t(\mathbf{x})$, given by

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

Proof

$$\begin{aligned} \frac{\partial p_t(\mathbf{x})}{\partial t} &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} [\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x})] + \frac{1}{2}g^2(t)\frac{\partial^2 p_t(\mathbf{x})}{\partial \mathbf{x}^2} \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x}) - \frac{1}{2}g^2(t)\frac{\partial p_t(\mathbf{x})}{\partial \mathbf{x}} \right] \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\mathbf{f}(\mathbf{x}, t)p_t(\mathbf{x}) - \frac{1}{2}g^2(t)p_t(\mathbf{x})\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right] \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) p_t(\mathbf{x}) \right] \right) \end{aligned}$$

Probability Flow ODE

Proof (Continued)

$$\frac{\partial p_t(\mathbf{x})}{\partial t} = \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2} g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) p_t(\mathbf{x}) \right] \right)$$

Probability Flow ODE

Proof (Continued)

$$\begin{aligned}\frac{\partial p_t(\mathbf{x})}{\partial t} &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2} g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) p_t(\mathbf{x}) \right] \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} [\mathbf{v}(\mathbf{x}, t) p_t(\mathbf{x})] \right) = -\text{div}(\mathbf{v}(\mathbf{x}, t) p_t(\mathbf{x}))\end{aligned}$$

Probability Flow ODE

Proof (Continued)

$$\begin{aligned}\frac{\partial p_t(\mathbf{x})}{\partial t} &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} \left[\left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) p_t(\mathbf{x}) \right] \right) = \\ &= \text{tr} \left(-\frac{\partial}{\partial \mathbf{x}} [\mathbf{v}(\mathbf{x}, t)p_t(\mathbf{x})] \right) = -\text{div}(\mathbf{v}(\mathbf{x}, t)p_t(\mathbf{x}))\end{aligned}$$

$$\mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}); \quad \tilde{g}(t) = 0$$

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt + 0 \cdot d\mathbf{w} = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

$$\frac{\partial p_t(\mathbf{x})}{\partial t} = -\text{div}(\mathbf{v}(\mathbf{x}, t)p_t(\mathbf{x})) + \frac{1}{2}\tilde{g}^2(t)\Delta_{\mathbf{x}}p_t(\mathbf{x})$$

Probability Flow ODE

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w} \quad - \text{SDE}$$

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt \quad - \text{Probability Flow ODE}$$

$$\mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x})$$

Probability Flow ODE

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w} \quad - \text{SDE}$$

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt \quad - \text{Probability Flow ODE}$$

$$\mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x})$$

- $\mathbf{s}(\mathbf{x}, t) = \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x})$ is the continuous-time score function.


Probability Flow ODE

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w} \quad - \text{SDE}$$

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt \quad - \text{Probability Flow ODE}$$

$$\mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x})$$

- ▶ $\mathbf{s}(\mathbf{x}, t) = \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x})$ is the continuous-time score function.
- ▶ The ODE produces more stable trajectories.



Song Y., et al. *Score-Based Generative Modeling through Stochastic Differential Equations*, 2020

Why do we need to convert SDEs to PF-ODEs?

- ▶ We could use `SDEsolve` to discretize the SDE.
- ▶ However, this is not efficient (trajectories are not smooth).

Why do we need to convert SDEs to PF-ODEs?

- ▶ We could use `SDEsolve` to discretize the SDE.
- ▶ However, this is not efficient (trajectories are not smooth).

Efficient Solvers

- ▶ Converting SDEs to PF-ODEs yields more efficient inference.
- ▶ We can apply any `ODEsolve` procedure to reduce the number of inference steps.
- ▶ For example, we can use Heun's / Runge-Kutta methods to solve the PF-ODE instead of Euler's method.
- ▶ In practice, this reduces the number of steps from 100–1000 to 20–50.

Outline

1. Probability Flow ODE
2. Reverse SDE
3. Score-Based Generative Models Through SDEs

Reverse SDE

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt, \quad \mathbf{x}(t + dt) = \mathbf{x}(t) + \mathbf{v}(\mathbf{x}, t)dt$$

Here dt can be > 0 or < 0 .

Reverse SDE

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt, \quad \mathbf{x}(t + dt) = \mathbf{x}(t) + \mathbf{v}(\mathbf{x}, t)dt$$

Here dt can be > 0 or < 0 .

Reverse ODE

Let $\tau = 1 - t$ ($d\tau = -dt$).

$$d\mathbf{x} = -\mathbf{v}(\mathbf{x}, 1 - \tau)d\tau$$

Reverse SDE

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt, \quad \mathbf{x}(t + dt) = \mathbf{x}(t) + \mathbf{v}(\mathbf{x}, t)dt$$

Here dt can be > 0 or < 0 .

Reverse ODE

Let $\tau = 1 - t$ ($d\tau = -dt$).

$$d\mathbf{x} = -\mathbf{v}(\mathbf{x}, 1 - \tau)d\tau$$

- ▶ How do we reverse the SDE $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$?
- ▶ The Wiener process introduces randomness that must be reversed.

Reverse SDE

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt, \quad \mathbf{x}(t + dt) = \mathbf{x}(t) + \mathbf{v}(\mathbf{x}, t)dt$$

Here dt can be > 0 or < 0 .

Reverse ODE

Let $\tau = 1 - t$ ($d\tau = -dt$).

$$d\mathbf{x} = -\mathbf{v}(\mathbf{x}, 1 - \tau)d\tau$$

- ▶ How do we reverse the SDE $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$?
- ▶ The Wiener process introduces randomness that must be reversed.

Theorem

There exists a reverse SDE for $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ (assume $p_t(\mathbf{x}) > 0$), given by:

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt + g(t)d\bar{\mathbf{w}}$$

where $dt < 0$ and $\bar{\mathbf{w}}$ is a reverse-time Wiener process.

Song Y., et al. Score-Based Generative Modeling through Stochastic Differential Equations, 2020

Reverse SDE

Theorem

There exists a reverse SDE for $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ (assume $p_t(\mathbf{x}) > 0$), given by:

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt + g(t) d\bar{\mathbf{w}}$$

where $dt < 0$ and $\bar{\mathbf{w}}(t)$ is a reverse-time Wiener process.

Reverse SDE

Theorem

There exists a reverse SDE for $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ (assume $p_t(\mathbf{x}) > 0$), given by:

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt + g(t) d\bar{\mathbf{w}}$$

where $dt < 0$ and $\bar{\mathbf{w}}(t)$ is a reverse-time Wiener process.

- ▶ Here the score function appears: $\mathbf{s}(\mathbf{x}, t) = \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x})$.
- ▶ Since $\mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\mathbf{s}(\mathbf{x}, t)$ (PF-ODE), the reverse drift equals $\mathbf{f} - g^2\mathbf{s} = \mathbf{v} - \frac{1}{2}g^2\mathbf{s}$.

Reverse SDE

Theorem

There exists a reverse SDE for $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ (assume $p_t(\mathbf{x}) > 0$), given by:

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt + g(t) d\bar{\mathbf{w}}$$

where $dt < 0$ and $\bar{\mathbf{w}}(t)$ is a reverse-time Wiener process.

- ▶ Here the score function appears: $\mathbf{s}(\mathbf{x}, t) = \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x})$.
- ▶ Since $\mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\mathbf{s}(\mathbf{x}, t)$ (PF-ODE), the reverse drift equals $\mathbf{f} - g^2\mathbf{s} = \mathbf{v} - \frac{1}{2}g^2\mathbf{s}$.

Proof Sketch

- ▶ Convert the initial SDE to a probability flow ODE.
- ▶ Reverse the probability flow ODE.
- ▶ Convert the reversed probability flow ODE back to an SDE.

Song Y., et al. Score-Based Generative Modeling through Stochastic Differential Equations, 2020

Proof of Reverse SDE

- Convert the initial SDE to a probability flow ODE:

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

Proof of Reverse SDE

- Convert the initial SDE to a probability flow ODE:

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

- Reverse the probability flow ODE:

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

$$d\mathbf{x} = \left(-\mathbf{f}(\mathbf{x}, 1 - \tau) + \frac{1}{2}g^2(1 - \tau)\frac{\partial}{\partial \mathbf{x}} \log p_{1-\tau}(\mathbf{x}) \right) d\tau$$

Proof of Reverse SDE

- Convert the initial SDE to a probability flow ODE:

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

- Reverse the probability flow ODE:

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt$$

$$d\mathbf{x} = \left(-\mathbf{f}(\mathbf{x}, 1 - \tau) + \frac{1}{2}g^2(1 - \tau)\frac{\partial}{\partial \mathbf{x}} \log p_{1-\tau}(\mathbf{x}) \right) d\tau$$

- Convert the reversed probability flow ODE back to an SDE:

$$d\mathbf{x} = \left(-\mathbf{f}(\mathbf{x}, 1 - \tau) + \frac{1}{2}g^2(1 - \tau)\frac{\partial}{\partial \mathbf{x}} \log p_{1-\tau}(\mathbf{x}) \right) d\tau$$

$$d\mathbf{x} = \left(-\mathbf{f}(\mathbf{x}, 1 - \tau) + g^2(1 - \tau)\frac{\partial}{\partial \mathbf{x}} \log p_{1-\tau}(\mathbf{x}) \right) d\tau + g(1 - \tau)d\mathbf{w}$$

Here $\mathbf{w}(\tau)$ is a Wiener process in forward τ -time.

Reverse SDE

Theorem

There exists a reverse SDE for $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ (assume $p_t(\mathbf{x}) > 0$), given by:

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt + g(t) d\bar{\mathbf{w}}$$

where $dt < 0$ and $\bar{\mathbf{w}}(t)$ is a reverse-time Wiener process.

Proof (Continued)

$$d\mathbf{x} = \left(-\mathbf{f}(\mathbf{x}, 1 - \tau) + g^2(1 - \tau) \frac{\partial}{\partial \mathbf{x}} \log p_{1-\tau}(\mathbf{x}) \right) d\tau + g(1 - \tau) d\mathbf{w}$$

Reverse SDE

Theorem

There exists a reverse SDE for $d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$ (assume $p_t(\mathbf{x}) > 0$), given by:

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt + g(t) d\bar{\mathbf{w}}$$

where $dt < 0$ and $\bar{\mathbf{w}}(t)$ is a reverse-time Wiener process.

Proof (Continued)

$$d\mathbf{x} = \left(-\mathbf{f}(\mathbf{x}, 1 - \tau) + g^2(1 - \tau) \frac{\partial}{\partial \mathbf{x}} \log p_{1-\tau}(\mathbf{x}) \right) d\tau + g(1 - \tau) d\mathbf{w}$$

$$d\mathbf{x} = \left(\mathbf{f}(\mathbf{x}, t) - g^2(t) \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt + g(t) d\bar{\mathbf{w}}$$

Here $d\tau > 0$, $dt < 0$, and $\bar{\mathbf{w}}$ is reverse-time Wiener process.

Reverse SDE

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

$$\mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\mathbf{s}(\mathbf{x}, t), \quad \text{where} \quad \mathbf{s}(\mathbf{x}, t) = \nabla_{\mathbf{x}} \log p_t(\mathbf{x})$$

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt \quad - \text{Probability Flow ODE}$$

$$d\mathbf{x} = \left(\mathbf{v}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\mathbf{s}(\mathbf{x}, t)\right)dt + g(t)d\bar{\mathbf{w}} \quad - \text{Reverse SDE}$$


Reverse SDE

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}$$

$$\mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\mathbf{s}(\mathbf{x}, t), \quad \text{where} \quad \mathbf{s}(\mathbf{x}, t) = \nabla_{\mathbf{x}} \log p_t(\mathbf{x})$$

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt \quad \text{-- Probability Flow ODE}$$

$$d\mathbf{x} = \left(\mathbf{v}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\mathbf{s}(\mathbf{x}, t) \right)dt + g(t)d\bar{\mathbf{w}} \quad \text{-- Reverse SDE}$$



Song Y., et al. *Score-Based Generative Modeling through Stochastic Differential Equations*, 2020

Score Matching SDE

$$\mathbf{x}(t) = \mathbf{x}(t - dt) + \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}$$

Score Matching SDE

$$\mathbf{x}(t) = \mathbf{x}(t - dt) + \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}$$

Variance Exploding SDE

$$d\mathbf{x} = \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}$$

$\sigma(t)$ is a monotonically increasing function.

Score Matching SDE

$$\mathbf{x}(t) = \mathbf{x}(t - dt) + \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}$$

Variance Exploding SDE

$$d\mathbf{x} = \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}$$

$\sigma(t)$ is a monotonically increasing function.

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}, \quad \mathbf{f}(\mathbf{x}, t) = 0, \quad g(t) = \sqrt{\frac{d[\sigma^2(t)]}{dt}}$$

Score Matching SDE

$$\mathbf{x}(t) = \mathbf{x}(t - dt) + \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}$$

Variance Exploding SDE

$$d\mathbf{x} = \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w}$$

$\sigma(t)$ is a monotonically increasing function.

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w}, \quad \mathbf{f}(\mathbf{x}, t) = 0, \quad g(t) = \sqrt{\frac{d[\sigma^2(t)]}{dt}}$$

$$d\mathbf{x} = \left(-\frac{1}{2} \frac{d[\sigma^2(t)]}{dt} \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt \quad (\text{probability flow ODE})$$

$$d\mathbf{x} = \left(-\frac{d[\sigma^2(t)]}{dt} \frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x}) \right) dt + \sqrt{\frac{d[\sigma^2(t)]}{dt}} d\bar{\mathbf{w}} \quad (\text{reverse SDE})$$

Diffusion SDE

Variance Preserving SDE

$$d\mathbf{x} = -\frac{1}{2}\beta(t)\mathbf{x}(t)dt + \sqrt{\beta(t)} \cdot d\mathbf{w}$$

$$\mathbf{f}(\mathbf{x}, t) = -\frac{1}{2}\beta(t)\mathbf{x}(t), \quad g(t) = \sqrt{\beta(t)}$$

Variance is preserved as long as $\mathbf{x}(0)$ has unit variance.

Diffusion SDE

Variance Preserving SDE

$$d\mathbf{x} = -\frac{1}{2}\beta(t)\mathbf{x}(t)dt + \sqrt{\beta(t)} \cdot d\mathbf{w}$$

$$\mathbf{f}(\mathbf{x}, t) = -\frac{1}{2}\beta(t)\mathbf{x}(t), \quad g(t) = \sqrt{\beta(t)}$$

Variance is preserved as long as $\mathbf{x}(0)$ has unit variance.

$$d\mathbf{x} = \left(-\frac{1}{2}\beta(t)\mathbf{x}(t) - \frac{1}{2}\beta(t)\frac{\partial}{\partial\mathbf{x}} \log p_t(\mathbf{x}) \right) dt \quad (\text{probability flow ODE})$$

$$d\mathbf{x} = \left(-\frac{1}{2}\beta(t)\mathbf{x}(t) - \beta(t)\frac{\partial}{\partial\mathbf{x}} \log p_t(\mathbf{x}) \right) dt + \sqrt{\beta(t)}d\bar{\mathbf{w}} \quad (\text{reverse SDE})$$

Outline

1. Probability Flow ODE
2. Reverse SDE
3. Score-Based Generative Models Through SDEs

Score-Based Generative Models Through SDEs

Discrete-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}_0)} \mathbb{E}_{t \sim U\{1, T\}} \mathbb{E}_{q(\mathbf{x}_t | \mathbf{x}_0)} \left\| \mathbf{s}_{\theta, t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t | \mathbf{x}_0) \right\|_2^2$$

Is it possible to train score-based diffusion models in continuous time?

Score-Based Generative Models Through SDEs

Discrete-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}_0)} \mathbb{E}_{t \sim U\{1, T\}} \mathbb{E}_{q(\mathbf{x}_t | \mathbf{x}_0)} \left\| \mathbf{s}_{\theta, t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t | \mathbf{x}_0) \right\|_2^2$$

Is it possible to train score-based diffusion models in continuous time?

Continuous-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0, 1]} \mathbb{E}_{q(\mathbf{x}(t) | \mathbf{x}(0))} \left\| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t) | \mathbf{x}(0)) \right\|_2^2$$

Score-Based Generative Models Through SDEs


Discrete-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}_0)} \mathbb{E}_{t \sim U\{1, T\}} \mathbb{E}_{q(\mathbf{x}_t | \mathbf{x}_0)} \left\| \mathbf{s}_{\theta, t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t | \mathbf{x}_0) \right\|_2^2$$

Is it possible to train score-based diffusion models in continuous time?


Continuous-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0, 1]} \mathbb{E}_{q(\mathbf{x}(t) | \mathbf{x}(0))} \left\| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t) | \mathbf{x}(0)) \right\|_2^2$$



Song Y., et al. *Score-Based Generative Modeling through Stochastic Differential Equations*, 2020

Generative Models Taxonomy



Score-Based Generative Models Through SDEs

Continuous-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{q(\mathbf{x}(t)|\mathbf{x}(0))} \left\| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) \right\|_2^2$$

Score-Based Generative Models Through SDEs

Continuous-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{q(\mathbf{x}(t)|\mathbf{x}(0))} \left\| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) \right\|_2^2$$

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

$$\nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) = -\frac{1}{\sigma^2(t)}(\mathbf{x}(t) - \boldsymbol{\mu})$$

Note: Normality holds for $\mathbf{f}(\mathbf{x}, t)$ affine in $\mathbf{x}$ ($\mathbf{f}(\mathbf{x}, t) = f(t)\mathbf{x}$).

Score-Based Generative Models Through SDEs

Continuous-Time Objective

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{q(\mathbf{x}(t)|\mathbf{x}(0))} \left\| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) \right\|_2^2$$

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

$$\nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) = -\frac{1}{\sigma^2(t)}(\mathbf{x}(t) - \boldsymbol{\mu})$$

Note: Normality holds for $\mathbf{f}(\mathbf{x}, t)$ affine in $\mathbf{x}$ ($\mathbf{f}(\mathbf{x}, t) = f(t)\mathbf{x}$).

$$d\mathbf{x} = \sqrt{\frac{d[\sigma^2(t)]}{dt}} \cdot d\mathbf{w} \quad (\text{Variance Exploding SDE})$$

$$d\mathbf{x} = -\frac{1}{2}\beta(t)\mathbf{x}(t)dt + \sqrt{\beta(t)} \cdot d\mathbf{w} \quad (\text{Variance Preserving SDE})$$

Is it possible to explicitly derive $\boldsymbol{\mu}(\mathbf{x}(0), t)$ and $\sigma^2(t)$ for VE-SDE and VP-SDE?

Score-Based Generative Models Through SDEs

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

Theorem

The moments of the SDE $d\mathbf{x} = f(t)\mathbf{x}dt + g(t)d\mathbf{w}$ satisfy:

$$\frac{d\boldsymbol{\mu}(\mathbf{x}(0), t)}{dt} = f(t)\boldsymbol{\mu}(\mathbf{x}(0), t)$$

$$\frac{d\sigma^2(t)}{dt} = 2f(t)\sigma^2(t) + g^2(t)$$

Score-Based Generative Models Through SDEs

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

Theorem

The moments of the SDE $d\mathbf{x} = f(t)\mathbf{x}dt + g(t)d\mathbf{w}$ satisfy:

$$\frac{d\boldsymbol{\mu}(\mathbf{x}(0), t)}{dt} = f(t)\boldsymbol{\mu}(\mathbf{x}(0), t)$$

$$\frac{d\sigma^2(t)}{dt} = 2f(t)\sigma^2(t) + g^2(t)$$

Proof

$$\mathbb{E}[d\mathbf{x}|\mathbf{x}(0)] = \mathbb{E}[f(t)\mathbf{x}dt|\mathbf{x}(0)] + \mathbb{E}[g(t)d\mathbf{w}|\mathbf{x}(0)]$$

Score-Based Generative Models Through SDEs

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

Theorem

The moments of the SDE $d\mathbf{x} = f(t)\mathbf{x}dt + g(t)d\mathbf{w}$ satisfy:

$$\frac{d\boldsymbol{\mu}(\mathbf{x}(0), t)}{dt} = f(t)\boldsymbol{\mu}(\mathbf{x}(0), t)$$

$$\frac{d\sigma^2(t)}{dt} = 2f(t)\sigma^2(t) + g^2(t)$$

Proof

$$\begin{aligned}\mathbb{E}[d\mathbf{x}|\mathbf{x}(0)] &= \mathbb{E}[f(t)\mathbf{x}dt|\mathbf{x}(0)] + \mathbb{E}[g(t)d\mathbf{w}|\mathbf{x}(0)] \\ &= f(t)\mathbb{E}[\mathbf{x}|\mathbf{x}(0)] dt + g(t)\mathbb{E}[d\mathbf{w}|\mathbf{x}(0)]\end{aligned}$$

Score-Based Generative Models Through SDEs

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

Theorem

The moments of the SDE $d\mathbf{x} = f(t)\mathbf{x}dt + g(t)d\mathbf{w}$ satisfy:

$$\frac{d\boldsymbol{\mu}(\mathbf{x}(0), t)}{dt} = f(t)\boldsymbol{\mu}(\mathbf{x}(0), t)$$

$$\frac{d\sigma^2(t)}{dt} = 2f(t)\sigma^2(t) + g^2(t)$$

Proof

$$\begin{aligned}\mathbb{E}[d\mathbf{x}|\mathbf{x}(0)] &= \mathbb{E}[f(t)\mathbf{x}dt|\mathbf{x}(0)] + \mathbb{E}[g(t)d\mathbf{w}|\mathbf{x}(0)] \\ &= f(t)\mathbb{E}[\mathbf{x}|\mathbf{x}(0)]dt + g(t)\mathbb{E}[d\mathbf{w}|\mathbf{x}(0)] \\ &= f(t)\boldsymbol{\mu}(\mathbf{x}(0), t)dt\end{aligned}$$

Score-Based Generative Models Through SDEs

Theorem

$$\frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Score-Based Generative Models Through SDEs

Theorem

$$\frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Proof (Continued)

$$\mathbb{E}[d\mathbf{x}|\mathbf{x}(0)] = f(t)\mu(\mathbf{x}(0), t)dt$$

Score-Based Generative Models Through SDEs

Theorem

$$\frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Proof (Continued)

$$\mathbb{E}[d\mathbf{x}|\mathbf{x}(0)] = f(t)\mu(\mathbf{x}(0), t)dt$$

$$\frac{d\mathbb{E}[\mathbf{x}(t)|\mathbf{x}(0)]}{dt} = \frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Score-Based Generative Models Through SDEs

Theorem

$$\frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Proof (Continued)

$$\mathbb{E}[d\mathbf{x}|\mathbf{x}(0)] = f(t)\mu(\mathbf{x}(0), t)dt$$

$$\frac{d\mathbb{E}[\mathbf{x}(t)|\mathbf{x}(0)]}{dt} = \frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Examples

$$\text{NCSN: } f(t) = 0 \quad \Rightarrow \quad \mu = \mathbf{x}(0)$$

Score-Based Generative Models Through SDEs

Theorem

$$\frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Proof (Continued)

$$\mathbb{E}[d\mathbf{x}|\mathbf{x}(0)] = f(t)\mu(\mathbf{x}(0), t)dt$$

$$\frac{d\mathbb{E}[\mathbf{x}(t)|\mathbf{x}(0)]}{dt} = \frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Examples

$$\text{NCSN: } f(t) = 0 \quad \Rightarrow \quad \mu = \mathbf{x}(0)$$

$$\text{DDPM: } f(t) = -\frac{1}{2}\beta(t) \quad \Rightarrow \quad \frac{d\mu}{dt} = -\frac{1}{2}\beta(t)\mu$$

Score-Based Generative Models Through SDEs

Theorem

$$\frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Proof (Continued)

$$\mathbb{E}[d\mathbf{x}|\mathbf{x}(0)] = f(t)\mu(\mathbf{x}(0), t)dt$$

$$\frac{d\mathbb{E}[\mathbf{x}(t)|\mathbf{x}(0)]}{dt} = \frac{d\mu(\mathbf{x}(0), t)}{dt} = f(t)\mu(\mathbf{x}(0), t)$$

Examples

$$\textbf{NCSN: } f(t) = 0 \quad \Rightarrow \quad \mu = \mathbf{x}(0)$$

$$\textbf{DDPM: } f(t) = -\frac{1}{2}\beta(t) \quad \Rightarrow \quad \frac{d\mu}{dt} = -\frac{1}{2}\beta(t)\mu$$

$$\mu = \mathbf{x}(0) \exp\left(-\frac{1}{2} \int_0^t \beta(s) ds\right)$$

Score-Based Generative Models Through SDEs

Training

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{q(\mathbf{x}(t)|\mathbf{x}(0))} \left\| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) \right\|_2^2$$

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

Score-Based Generative Models Through SDEs

Training

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{q(\mathbf{x}(t)|\mathbf{x}(0))} \left\| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) \right\|_2^2$$

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

NCSN

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\mathbf{x}(0), [\sigma^2(t) - \sigma^2(0)] \cdot \mathbf{I}\right)$$

Score-Based Generative Models Through SDEs

Training

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x}(0))} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{q(\mathbf{x}(t)|\mathbf{x}(0))} \left\| \mathbf{s}_{\theta}(\mathbf{x}(t), t) - \nabla_{\mathbf{x}(t)} \log q(\mathbf{x}(t)|\mathbf{x}(0)) \right\|_2^2$$

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\boldsymbol{\mu}(\mathbf{x}(0), t), \sigma^2(t) \cdot \mathbf{I}\right)$$

NCSN

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\mathbf{x}(0), [\sigma^2(t) - \sigma^2(0)] \cdot \mathbf{I}\right)$$

DDPM

$$q(\mathbf{x}(t)|\mathbf{x}(0)) = \mathcal{N}\left(\mathbf{x}(0)e^{-\frac{1}{2} \int_0^t \beta(s) ds}, \left(1 - e^{-\int_0^t \beta(s) ds}\right) \cdot \mathbf{I}\right)$$


Here we omit the derivations of the variance.

Song Y., et al. Score-Based Generative Modeling through Stochastic Differential Equations, 2020

Score-Based Generative Models Through SDEs

Sampling


Solve the reverse SDE using numerical solvers (SDEsolve).



Score-Based Generative Models Through SDEs

Sampling

Solve the reverse SDE using numerical solvers (SDEsolve).



- ▶ Discretizing the reverse SDE provides ancestral sampling.
- ▶ Discretizing the probability flow ODE yields deterministic sampling.

Summary

- ▶ Every SDE admits a corresponding probability flow ODE that follows the same probability path $p_t(\mathbf{x})$.
- ▶ Converting SDEs to probability flow ODEs enables efficient numerical solvers, reducing the number of inference steps from 100–1000 to 20–50.
- ▶ SDEs can be reversed in time using the score function $\nabla_{\mathbf{x}} \log p_t(\mathbf{x})$.
- ▶ The continuous-time score matching objective generalizes DDPM and NCSN to continuous time.
- ▶ For affine-drift SDEs (VE-SDE and VP-SDE), the transition kernel $q(\mathbf{x}(t)|\mathbf{x}(0))$ is Gaussian, making the training objective tractable.
- ▶ Discretizing the reverse SDE yields ancestral sampling, while discretizing the probability flow ODE yields deterministic sampling.