

Deep Generative Models

Lecture 11

Roman Isachenko



AI Masters

2026, Spring


Recap of Previous Lecture

$$d\mathbf{x} = \mathbf{f}(\mathbf{x}, t)dt + g(t)d\mathbf{w} \quad (\text{SDE with the probability path } p_t(\mathbf{x}))$$

Probability Flow ODE

There exists an ODE with the identical probability path $p_t(\mathbf{x})$ of the form:

$$d\mathbf{x} = \mathbf{v}(\mathbf{x}, t)dt, \quad \mathbf{v}(\mathbf{x}, t) = \mathbf{f}(\mathbf{x}, t) - \frac{1}{2}g^2(t)\frac{\partial}{\partial \mathbf{x}} \log p_t(\mathbf{x})$$



Song Y., et al. *Score-Based Generative Modeling through Stochastic Differential Equations*, 2020

Outline

1. Conditional Flow Matching

Outline

1. Conditional Flow Matching

Flow Matching

Let's introduce the latent variable $\mathbf{z}$:

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{z})p(\mathbf{z})d\mathbf{z}$$

Here, $p_t(\mathbf{x}|\mathbf{z})$ is a **conditional probability path**.

Flow Matching

Let's introduce the latent variable $\mathbf{z}$:

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{z})p(\mathbf{z})d\mathbf{z}$$

Here, $p_t(\mathbf{x}|\mathbf{z})$ is a **conditional probability path**. The conditional probability path $p_t(\mathbf{x}|\mathbf{z})$ satisfies the KFP theorem:

$$\frac{\partial p_t(\mathbf{x}|\mathbf{z})}{\partial t} = -\operatorname{div}(\mathbf{v}(\mathbf{x}, \mathbf{z}, t)p_t(\mathbf{x}|\mathbf{z})),$$

where $\mathbf{v}(\mathbf{x}, \mathbf{z}, t)$ is a **conditional vector field**:

Flow Matching

Let's introduce the latent variable $\mathbf{z}$:

$$p_t(\mathbf{x}) = \int p_t(\mathbf{x}|\mathbf{z})p(\mathbf{z})d\mathbf{z}$$

Here, $p_t(\mathbf{x}|\mathbf{z})$ is a **conditional probability path**. The conditional probability path $p_t(\mathbf{x}|\mathbf{z})$ satisfies the KFP theorem:

$$\frac{\partial p_t(\mathbf{x}|\mathbf{z})}{\partial t} = -\operatorname{div}(\mathbf{v}(\mathbf{x}, \mathbf{z}, t)p_t(\mathbf{x}|\mathbf{z})),$$

where $\mathbf{v}(\mathbf{x}, \mathbf{z}, t)$ is a **conditional vector field**:

$$\frac{d\mathbf{x}_t}{dt} = \mathbf{v}(\mathbf{x}_t, t) \quad \Rightarrow \quad \frac{d\mathbf{x}_t}{dt} = \mathbf{v}(\mathbf{x}_t, \mathbf{z}, t)$$

Flow Matching

Flow Matching (FM)

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditional Flow Matching (CFM)

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Flow Matching

Flow Matching (FM)

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditional Flow Matching (CFM)

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_\theta(\mathbf{x}, \mathbf{z}, t)\|^2 \rightarrow \min_{\theta}$$

Theorem

If $\text{supp}(p_t(\mathbf{x})) = \mathbb{R}^m$, then the optimal value of the FM objective equals the optimal value of the CFM objective.

Flow Matching

Flow Matching (FM)

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

Conditional Flow Matching (CFM)

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_\theta(\mathbf{x}, \mathbf{z}, t)\|^2 \rightarrow \min_{\theta}$$

Theorem

If $\text{supp}(p_t(\mathbf{x})) = \mathbb{R}^m$, then the optimal value of the FM objective equals the optimal value of the CFM objective.

Proof

This can be proved in a similar way as in the denoising score matching theorem.

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 \end{aligned}$$

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 \end{aligned}$$

Proof

$$\begin{aligned} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} [\|\mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 - 2\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t)\mathbf{v}(\mathbf{x}, t)] + \text{const}(\theta) \end{aligned}$$

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 \end{aligned}$$

Proof

$$\begin{aligned} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} [\|\mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 - 2\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t)\mathbf{v}(\mathbf{x}, t)] + \text{const}(\theta) \\ \mathbb{E}_{p_t(\mathbf{x})} [\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t)\mathbf{v}(\mathbf{x}, t)] &= \int p_t(\mathbf{x}) \left[\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t) \left(\int p_t(\mathbf{z}|\mathbf{x}) \mathbf{v}(\mathbf{x}, \mathbf{z}, t) d\mathbf{z} \right) \right] d\mathbf{x} \end{aligned}$$

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 \end{aligned}$$

Proof

$$\begin{aligned} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} [\|\mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 - 2\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t)\mathbf{v}(\mathbf{x}, t)] + \text{const}(\theta) \\ \mathbb{E}_{p_t(\mathbf{x})} [\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t)\mathbf{v}(\mathbf{x}, t)] &= \int p_t(\mathbf{x}) \left[\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t) \left(\int p_t(\mathbf{z}|\mathbf{x}) \mathbf{v}(\mathbf{x}, \mathbf{z}, t) d\mathbf{z} \right) \right] d\mathbf{x} = \\ &= \int \int p_t(\mathbf{x}) p_t(\mathbf{z}|\mathbf{x}) [\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t) \mathbf{v}(\mathbf{x}, \mathbf{z}, t)] d\mathbf{z} d\mathbf{x} \end{aligned}$$

Conditional Flow Matching

Theorem

$$\begin{aligned} \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \arg \min_{\theta} \mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 \end{aligned}$$

Proof

$$\begin{aligned} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x})} \|\mathbf{v}(\mathbf{x}, t) - \mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 &= \\ = \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} [\|\mathbf{v}_{\theta}(\mathbf{x}, t)\|^2 - 2\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t)\mathbf{v}(\mathbf{x}, t)] + \text{const}(\theta) \\ \mathbb{E}_{p_t(\mathbf{x})} [\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t)\mathbf{v}(\mathbf{x}, t)] &= \int p_t(\mathbf{x}) \left[\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t) \left(\int p_t(\mathbf{z}|\mathbf{x}) \mathbf{v}(\mathbf{x}, \mathbf{z}, t) d\mathbf{z} \right) \right] d\mathbf{x} = \\ &= \int \int p_t(\mathbf{x}) p_t(\mathbf{z}|\mathbf{x}) [\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t) \mathbf{v}(\mathbf{x}, \mathbf{z}, t)] d\mathbf{z} d\mathbf{x} = \\ &= \mathbb{E}_{p(\mathbf{z})} \mathbb{E}_{p_t(\mathbf{x}|\mathbf{z})} [\mathbf{v}_{\theta}^{\top}(\mathbf{x}, t) \mathbf{v}(\mathbf{x}, \mathbf{z}, t)] \end{aligned}$$

Conditional Flow Matching: Training and Sampling

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

$$\mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}) = \mathcal{N}(\mathbf{0}, \mathbf{I}); \quad \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}) = p_{\text{data}}(\mathbf{x}).$$

Conditional Flow Matching: Training and Sampling

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

$$\mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}) = \mathcal{N}(0, \mathbf{I}); \quad \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}) = p_{\text{data}}(\mathbf{x}).$$

Training

1. Sample a time $t \sim U[0, 1]$ and $\mathbf{z} \sim p(\mathbf{z})$.
2. Draw $\mathbf{x}_t \sim p_t(\mathbf{x}|\mathbf{z})$.
3. Compute the loss $\mathcal{L} = \|\mathbf{v}(\mathbf{x}_t, \mathbf{z}, t) - \mathbf{v}_\theta(\mathbf{x}_t, t)\|^2$.

Conditional Flow Matching: Training and Sampling

$$\mathbb{E}_{t \sim U[0,1]} \mathbb{E}_{\mathbf{z} \sim p(\mathbf{z})} \mathbb{E}_{\mathbf{x} \sim p_t(\mathbf{x}|\mathbf{z})} \|\mathbf{v}(\mathbf{x}, \mathbf{z}, t) - \mathbf{v}_\theta(\mathbf{x}, t)\|^2 \rightarrow \min_{\theta}$$

$$\mathbb{E}_{p(\mathbf{z})} p_0(\mathbf{x}|\mathbf{z}) = \mathcal{N}(0, \mathbf{I}); \quad \mathbb{E}_{p(\mathbf{z})} p_1(\mathbf{x}|\mathbf{z}) = p_{\text{data}}(\mathbf{x}).$$

Training

1. Sample a time $t \sim U[0, 1]$ and $\mathbf{z} \sim p(\mathbf{z})$.
2. Draw $\mathbf{x}_t \sim p_t(\mathbf{x}|\mathbf{z})$.
3. Compute the loss $\mathcal{L} = \|\mathbf{v}(\mathbf{x}_t, \mathbf{z}, t) - \mathbf{v}_\theta(\mathbf{x}_t, t)\|^2$.

Sampling

1. Sample $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$.
2. Solve the ODE to obtain $\mathbf{x}_1$:

$$\mathbf{x}_1 = \psi_1(\mathbf{x}_0) = \text{ODESolve}_v(\mathbf{x}_0, \theta, t_0 = 0, t_1 = 1).$$

Summary

- ▶ Flow matching suggests fitting the vector field directly.
- ▶ Conditional flow matching introduces the latent variable $\mathbf{z}$, reformulating the initial task in terms of conditional dynamics.