

Deep Generative Models

Lecture 3

Roman Isachenko

Moscow Institute of Physics and Technology
Yandex School of Data Analysis

2026, Autumn

Recap of Previous Lecture

Jacobian Matrix

Let $\mathbf{f} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ be a differentiable function.

$$\mathbf{z} = \mathbf{f}(\mathbf{x}), \quad \mathbf{J} = \frac{\partial \mathbf{z}}{\partial \mathbf{x}} = \begin{pmatrix} \frac{\partial z_1}{\partial x_1} & \cdots & \frac{\partial z_1}{\partial x_m} \\ \vdots & \ddots & \vdots \\ \frac{\partial z_m}{\partial x_1} & \cdots & \frac{\partial z_m}{\partial x_m} \end{pmatrix} \in \mathbb{R}^{m \times m}$$

Change of Variables Theorem (CoV)


Let $\mathbf{x} \in \mathbb{R}^m$ be a random vector with density $p(\mathbf{x})$, and let $\mathbf{f} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ be a C^1 -diffeomorphism ($\mathbf{f}$ and $\mathbf{f}^{-1}$ are continuously differentiable mappings). If $\mathbf{z} = \mathbf{f}(\mathbf{x})$, then

$$p(\mathbf{x}) = p(\mathbf{z}) |\det(\mathbf{J}_{\mathbf{f}})| = p(\mathbf{z}) \left| \det \left(\frac{\partial \mathbf{z}}{\partial \mathbf{x}} \right) \right| = p(\mathbf{f}(\mathbf{x})) \left| \det \left(\frac{\partial \mathbf{f}(\mathbf{x})}{\partial \mathbf{x}} \right) \right|$$
$$p(\mathbf{z}) = p(\mathbf{x}) |\det(\mathbf{J}_{\mathbf{f}^{-1}})| = p(\mathbf{x}) \left| \det \left(\frac{\partial \mathbf{x}}{\partial \mathbf{z}} \right) \right| = p(\mathbf{f}^{-1}(\mathbf{z})) \left| \det \left(\frac{\partial \mathbf{f}^{-1}(\mathbf{z})}{\partial \mathbf{z}} \right) \right|$$

Recap of Previous Lecture

Definition

A normalizing flow is a C^1 -diffeomorphism that transforms data $\mathbf{x}$ to noise $\mathbf{z}$.



Log-Likelihood

$$\log p_{\theta}(\mathbf{x}) = \log p(\mathbf{f}_K \circ \cdots \circ \mathbf{f}_1(\mathbf{x})) + \sum_{k=1}^K \log |\det(\mathbf{J}_{\mathbf{f}_k})|$$

Recap of Previous Lecture

Flow Log-Likelihood

$$\log p_{\theta}(\mathbf{x}) = \log p(\mathbf{f}_{\theta}(\mathbf{x})) + \log |\det(\mathbf{J}_{\mathbf{f}})|$$

One significant challenge is efficiently computing the Jacobian determinant.

Linear Flows

$$\mathbf{z} = \mathbf{f}_{\theta}(\mathbf{x}) = \mathbf{W}\mathbf{x}, \quad \mathbf{W} \in \mathbb{R}^{m \times m}, \quad \theta = \mathbf{W}, \quad \mathbf{J}_{\mathbf{f}} = \mathbf{W}^{\top}$$

- ▶ LU Decomposition:

$$\mathbf{W} = \mathbf{P}\mathbf{L}\mathbf{U}.$$

- ▶ QR Decomposition:

$$\mathbf{W} = \mathbf{Q}\mathbf{R}.$$

Decomposition is performed only once during initialization. Then the decomposed matrices (**P**, **L**, **U** or **Q**, **R**) are optimized.

Recap of Previous Lecture

Consider an autoregressive model:

$$p_{\theta}(\mathbf{x}) = \prod_{j=1}^m p_{\theta}(x_j | \mathbf{x}_{1:j-1}), \quad p_{\theta}(x_j | \mathbf{x}_{1:j-1}) = \mathcal{N}(\mu_{j,\theta}(\mathbf{x}_{1:j-1}), \sigma_{j,\theta}^2(\mathbf{x}_{1:j-1})).$$

Gaussian Autoregressive Normalizing Flow

$$\mathbf{x} = \mathbf{f}_{\theta}^{-1}(\mathbf{z}) \quad \Rightarrow \quad x_j = \sigma_{j,\theta}(\mathbf{x}_{1:j-1}) \cdot z_j + \mu_{j,\theta}(\mathbf{x}_{1:j-1}).$$

$$\mathbf{z} = \mathbf{f}_{\theta}(\mathbf{x}) \quad \Rightarrow \quad z_j = (x_j - \mu_{j,\theta}(\mathbf{x}_{1:j-1})) \cdot \frac{1}{\sigma_{j,\theta}(\mathbf{x}_{1:j-1})}.$$

- ▶ This transformation is both C^1 -**diffeomorphism**, mapping $p(\mathbf{z})$ to $p_{\theta}(\mathbf{x})$.
- ▶ The Jacobian matrix for this transformation is triangular.

The generative function $\mathbf{f}_{\theta}^{-1}(\mathbf{z})$ is **sequential**, while the inference function $\mathbf{f}_{\theta}(\mathbf{x})$ is **not sequential**.

Recap of Previous Lecture

Let us partition $\mathbf{x}$ and $\mathbf{z}$ into two groups:

$$\mathbf{x} = [\mathbf{x}_1, \mathbf{x}_2] = [\mathbf{x}_{1:d}, \mathbf{x}_{d+1:m}]; \quad \mathbf{z} = [\mathbf{z}_1, \mathbf{z}_2] = [\mathbf{z}_{1:d}, \mathbf{z}_{d+1:m}].$$

Coupling Layer

$$\begin{cases} \mathbf{x}_1 = \mathbf{z}_1; \\ \mathbf{x}_2 = \mathbf{z}_2 \odot \sigma_{\theta}(\mathbf{z}_1) + \mu_{\theta}(\mathbf{z}_1). \end{cases} \quad \begin{cases} \mathbf{z}_1 = \mathbf{x}_1; \\ \mathbf{z}_2 = (\mathbf{x}_2 - \mu_{\theta}(\mathbf{x}_1)) \odot \frac{1}{\sigma_{\theta}(\mathbf{x}_1)}. \end{cases}$$

Both density estimation and sampling require just a single pass!

Jacobian

$$\det \left(\frac{\partial \mathbf{z}}{\partial \mathbf{x}} \right) = \det \begin{pmatrix} \mathbf{I}_d & \mathbf{0}_{d \times (m-d)} \\ \frac{\partial \mathbf{z}_2}{\partial \mathbf{x}_1} & \frac{\partial \mathbf{z}_2}{\partial \mathbf{x}_2} \end{pmatrix} = \prod_{j=1}^{m-d} \frac{1}{\sigma_{j,\theta}(\mathbf{x}_1)}.$$

A coupling layer is a special instance of an gaussian autoregressive normalizing flow.

Recap of Previous Lecture

Maximum Likelihood Estimation (MLE) Problem

$$\theta^* = \arg \max_{\theta} p_{\theta}(\mathbf{X}) = \arg \max_{\theta} \prod_{i=1}^n p_{\theta}(\mathbf{x}_i) = \arg \max_{\theta} \sum_{i=1}^n \log p_{\theta}(\mathbf{x}_i).$$

The distribution $p_{\theta}(\mathbf{x})$ should be highly complex (just like the true data distribution $p_{\text{data}}(\mathbf{x})$).

Latent Variable

Introduce a latent variable $\mathbf{z}$ for each observed sample $\mathbf{x}$:

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p(\mathbf{z}); \quad \log p_{\theta}(\mathbf{x}, \mathbf{z}) = \log p_{\theta}(\mathbf{x}|\mathbf{z}) + \log p(\mathbf{z}).$$

Recap of Previous Lecture

Maximum Likelihood Estimation (MLE) Problem

$$\theta^* = \arg \max_{\theta} p_{\theta}(\mathbf{X}) = \arg \max_{\theta} \prod_{i=1}^n p_{\theta}(\mathbf{x}_i) = \arg \max_{\theta} \sum_{i=1}^n \log p_{\theta}(\mathbf{x}_i).$$

The distribution $p_{\theta}(\mathbf{x})$ should be highly complex (just like the true data distribution $p_{\text{data}}(\mathbf{x})$).

Latent Variable

Introduce a latent variable $\mathbf{z}$ for each observed sample $\mathbf{x}$:

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p(\mathbf{z}); \quad \log p_{\theta}(\mathbf{x}, \mathbf{z}) = \log p_{\theta}(\mathbf{x}|\mathbf{z}) + \log p(\mathbf{z}).$$

$$p_{\theta}(\mathbf{x}) = \int p_{\theta}(\mathbf{x}, \mathbf{z}) d\mathbf{z} = \int p_{\theta}(\mathbf{x}|\mathbf{z})p(\mathbf{z}) d\mathbf{z}.$$

- ▶ $p(\mathbf{z})$ is prior distribution over the latent variable.
- ▶ $p_{\theta}(\mathbf{x}|\mathbf{z})$ is **decoder** distribution.
- ▶ Both $p_{\theta}(\mathbf{x}|\mathbf{z})$ and $p(\mathbf{z})$ are usually much simpler than $p_{\theta}(\mathbf{x})$.

Outline

1. Latent Variable Models (LVM) (continued)
2. Variational Evidence Lower Bound (ELBO)
3. Amortized Inference
4. ELBO Gradients, Reparametrization Trick

Outline


1. Latent Variable Models (LVM) (continued)
2. Variational Evidence Lower Bound (ELBO)
3. Amortized Inference
4. ELBO Gradients, Reparametrization Trick

MLE for LVM

$$\sum_{i=1}^n \log p_{\theta}(\mathbf{x}_i) = \sum_{i=1}^n \log \int p_{\theta}(\mathbf{x}_i | \mathbf{z}_i) p(\mathbf{z}_i) d\mathbf{z}_i \rightarrow \max_{\theta}.$$


MLE for LVM

$$\sum_{i=1}^n \log p_{\theta}(\mathbf{x}_i) = \sum_{i=1}^n \log \int p_{\theta}(\mathbf{x}_i | \mathbf{z}_i) p(\mathbf{z}_i) d\mathbf{z}_i \rightarrow \max_{\theta}.$$



MLE for LVM

$$\sum_{i=1}^n \log p_{\theta}(\mathbf{x}_i) = \sum_{i=1}^n \log \int p_{\theta}(\mathbf{x}_i | \mathbf{z}_i) p(\mathbf{z}_i) d\mathbf{z}_i \rightarrow \max_{\theta}.$$




Naive Monte Carlo Estimation

$$\log p_{\theta}(\mathbf{x}) = \log \mathbb{E}_{p(\mathbf{z})} p_{\theta}(\mathbf{x} | \mathbf{z}) \geq \mathbb{E}_{p(\mathbf{z})} \log p_{\theta}(\mathbf{x} | \mathbf{z}) \approx \frac{1}{K} \sum_{k=1}^K \log p_{\theta}(\mathbf{x} | \mathbf{z}_k),$$

where $\mathbf{z}_k \sim p(\mathbf{z})$.

MLE for LVM

$$\sum_{i=1}^n \log p_{\theta}(\mathbf{x}_i) = \sum_{i=1}^n \log \int p_{\theta}(\mathbf{x}_i | \mathbf{z}_i) p(\mathbf{z}_i) d\mathbf{z}_i \rightarrow \max_{\theta}.$$



Naive Monte Carlo Estimation

$$\log p_{\theta}(\mathbf{x}) = \log \mathbb{E}_{p(\mathbf{z})} p_{\theta}(\mathbf{x}|\mathbf{z}) \geq \mathbb{E}_{p(\mathbf{z})} \log p_{\theta}(\mathbf{x}|\mathbf{z}) \approx \frac{1}{K} \sum_{k=1}^K \log p_{\theta}(\mathbf{x}|\mathbf{z}_k),$$

where $\mathbf{z}_k \sim p(\mathbf{z})$.

Challenge: As the dimensionality of $\mathbf{z}$ increases, the number of samples needed to adequately cover the latent space grows exponentially.

Outline

1. Latent Variable Models (LVM) (continued)
2. Variational Evidence Lower Bound (ELBO)
3. Amortized Inference
4. ELBO Gradients, Reparametrization Trick

ELBO Derivation I

Bayes theorem

$$p_{\theta}(\mathbf{z}|\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{x})} \quad \Rightarrow \quad p_{\theta}(\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})}$$

ELBO Derivation I

Bayes theorem

$$p_{\theta}(\mathbf{z}|\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{x})} \quad \Rightarrow \quad p_{\theta}(\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})}$$

Log likelihood

$$\log p_{\theta}(\mathbf{x}) = \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}) d\mathbf{z}$$

ELBO Derivation I

Bayes theorem

$$p_{\theta}(\mathbf{z}|\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{x})} \quad \Rightarrow \quad p_{\theta}(\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})}$$

Log likelihood

$$\log p_{\theta}(\mathbf{x}) = \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}) d\mathbf{z} = \mathbb{E}_q \log \left[\frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} \right]$$

ELBO Derivation I

Bayes theorem

$$p_{\theta}(\mathbf{z}|\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{x})} \quad \Rightarrow \quad p_{\theta}(\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})}$$

Log likelihood

$$\begin{aligned} \log p_{\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}) d\mathbf{z} = \mathbb{E}_q \log \left[\frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} \right] = \\ &= \mathbb{E}_q \log \left[\frac{p_{\theta}(\mathbf{x}, \mathbf{z}) q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x}) q(\mathbf{z})} \right] \end{aligned}$$

ELBO Derivation I

Bayes theorem

$$p_{\theta}(\mathbf{z}|\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{x})} \quad \Rightarrow \quad p_{\theta}(\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})}$$

Log likelihood

$$\begin{aligned} \log p_{\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}) d\mathbf{z} = \mathbb{E}_q \log \left[\frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} \right] = \\ &= \mathbb{E}_q \log \left[\frac{p_{\theta}(\mathbf{x}, \mathbf{z}) q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x}) q(\mathbf{z})} \right] = \\ &= \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} \end{aligned}$$

ELBO Derivation I

Bayes theorem

$$p_{\theta}(\mathbf{z}|\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{x})} \quad \Rightarrow \quad p_{\theta}(\mathbf{x}) = \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})}$$

Log likelihood

$$\begin{aligned} \log p_{\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}) d\mathbf{z} = \mathbb{E}_q \log \left[\frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} \right] = \\ &= \mathbb{E}_q \log \left[\frac{p_{\theta}(\mathbf{x}, \mathbf{z}) q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x}) q(\mathbf{z})} \right] = \\ &= \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} \end{aligned}$$

- ▶ Here, $q(\mathbf{z})$ is any distribution such that $\int q(\mathbf{z}) d\mathbf{z} = 1$.
- ▶ We assume that $\text{supp}(q(\mathbf{z})) = \text{supp}(p_{\theta}(\mathbf{z}|\mathbf{x})) = \mathbb{R}^d$.

ELBO Derivation II

Variational Decomposition

$$\log p_{\theta}(\mathbf{x}) = \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})}$$

ELBO Derivation II

Variational Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} = \\ &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x}))\end{aligned}$$

ELBO Derivation II

Variational Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} = \\ &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})) \geq \mathcal{L}_{q,\theta}(\mathbf{x})\end{aligned}$$

ELBO Derivation II

Variational Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} = \\ &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})) \geq \mathcal{L}_{q,\theta}(\mathbf{x})\end{aligned}$$

Variational Evidence Lower Bound (ELBO)

$$\mathcal{L}_{q,\theta}(\mathbf{x}) = \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} \leq \log p_{\theta}(\mathbf{x})$$

ELBO Derivation II

Variational Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} = \\ &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})) \geq \mathcal{L}_{q,\theta}(\mathbf{x})\end{aligned}$$

Variational Evidence Lower Bound (ELBO)

$$\mathcal{L}_{q,\theta}(\mathbf{x}) = \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} \leq \log p_{\theta}(\mathbf{x})$$

- This inequality holds for any choice of $q(\mathbf{z})$.

ELBO Derivation II

Variational Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} = \\ &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})) \geq \mathcal{L}_{q,\theta}(\mathbf{x})\end{aligned}$$

Variational Evidence Lower Bound (ELBO)

$$\mathcal{L}_{q,\theta}(\mathbf{x}) = \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} \leq \log p_{\theta}(\mathbf{x})$$

- ▶ This inequality holds for any choice of $q(\mathbf{z})$.
- ▶ Instead of maximizing the likelihood, maximize the ELBO:

$$\max_{\theta} p_{\theta}(\mathbf{x}) \quad \rightarrow \quad \max_{q,\theta} \mathcal{L}_{q,\theta}(\mathbf{x})$$

ELBO Derivation II

Variational Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} + \mathbb{E}_q \log \frac{q(\mathbf{z})}{p_{\theta}(\mathbf{z}|\mathbf{x})} = \\ &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})) \geq \mathcal{L}_{q,\theta}(\mathbf{x})\end{aligned}$$

Variational Evidence Lower Bound (ELBO)

$$\mathcal{L}_{q,\theta}(\mathbf{x}) = \mathbb{E}_q \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} \leq \log p_{\theta}(\mathbf{x})$$

- ▶ This inequality holds for any choice of $q(\mathbf{z})$.
- ▶ Instead of maximizing the likelihood, maximize the ELBO:

$$\max_{\theta} p_{\theta}(\mathbf{x}) \quad \rightarrow \quad \max_{q,\theta} \mathcal{L}_{q,\theta}(\mathbf{x})$$

- ▶ Distribution $q(\mathbf{z})$ is treated as **variational** parameter.

Variational Evidence Lower Bound (ELBO)

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z}) \| p_{\theta}(\mathbf{z}|\mathbf{x}))$$

What is the optimal distribution $q^*(\mathbf{z})$ given fixed θ^* ?

Variational Evidence Lower Bound (ELBO)

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x}))$$

What is the optimal distribution $q^*(\mathbf{z})$ given fixed θ^* ?

$$q^*(\mathbf{z}) = \arg \max_q \mathcal{L}_{q,\theta^*}(\mathbf{x})$$

Variational Evidence Lower Bound (ELBO)

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x}))$$

What is the optimal distribution $q^*(\mathbf{z})$ given fixed θ^* ?

$$\begin{aligned} q^*(\mathbf{z}) &= \arg \max_q \mathcal{L}_{q,\theta^*}(\mathbf{x}) = \\ &= \arg \min_q D_{\text{KL}}(q(\mathbf{z})\|p_{\theta^*}(\mathbf{z}|\mathbf{x})) \end{aligned} .$$

Variational Evidence Lower Bound (ELBO)

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x}))$$

What is the optimal distribution $q^*(\mathbf{z})$ given fixed θ^* ?

$$\begin{aligned} q^*(\mathbf{z}) &= \arg \max_q \mathcal{L}_{q,\theta^*}(\mathbf{x}) = \\ &= \arg \min_q D_{\text{KL}}(q(\mathbf{z})\|p_{\theta^*}(\mathbf{z}|\mathbf{x})) = p_{\theta^*}(\mathbf{z}|\mathbf{x}). \end{aligned}$$

Variational Evidence Lower Bound (ELBO)

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x}))$$

What is the optimal distribution $q^*(\mathbf{z})$ given fixed θ^* ?

$$\begin{aligned} q^*(\mathbf{z}) &= \arg \max_q \mathcal{L}_{q,\theta^*}(\mathbf{x}) = \\ &= \arg \min_q D_{\text{KL}}(q(\mathbf{z})\|p_{\theta^*}(\mathbf{z}|\mathbf{x})) = p_{\theta^*}(\mathbf{z}|\mathbf{x}). \end{aligned}$$

Here we got the intuition about variational distribution $q(\mathbf{z})$ – it estimates the posterior $p_{\theta^*}(\mathbf{z}|\mathbf{x})$.


Variational Evidence Lower Bound (ELBO)

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z}) \| p_{\theta}(\mathbf{z}|\mathbf{x}))$$

What is the optimal distribution $q^*(\mathbf{z})$ given fixed θ^* ?

$$\begin{aligned} q^*(\mathbf{z}) &= \arg \max_q \mathcal{L}_{q,\theta^*}(\mathbf{x}) = \\ &= \arg \min_q D_{\text{KL}}(q(\mathbf{z}) \| p_{\theta^*}(\mathbf{z}|\mathbf{x})) = p_{\theta^*}(\mathbf{z}|\mathbf{x}). \end{aligned}$$

Here we got the intuition about variational distribution $q(\mathbf{z})$ – it estimates the posterior $p_{\theta^*}(\mathbf{z}|\mathbf{x})$.



Variational Evidence Lower Bound (ELBO)

$$\mathcal{L}_{q,\theta}(\mathbf{x}) = \int q(\mathbf{z}) \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} d\mathbf{z}$$

Variational Evidence Lower Bound (ELBO)

$$\begin{aligned}\mathcal{L}_{q,\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} + \int q(\mathbf{z}) \log \frac{p(\mathbf{z})}{q(\mathbf{z})} d\mathbf{z}\end{aligned}$$

Variational Evidence Lower Bound (ELBO)

$$\begin{aligned}\mathcal{L}_{q,\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} + \int q(\mathbf{z}) \log \frac{p(\mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z}))\end{aligned}$$

Variational Evidence Lower Bound (ELBO)

$$\begin{aligned}\mathcal{L}_{q,\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} + \int q(\mathbf{z}) \log \frac{p(\mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z}))\end{aligned}$$

Log-Likelihood Decomposition

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x}))$$

Variational Evidence Lower Bound (ELBO)

$$\begin{aligned}\mathcal{L}_{q,\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} + \int q(\mathbf{z}) \log \frac{p(\mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z}))\end{aligned}$$

Log-Likelihood Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})) = \\ &= \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z})) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})).\end{aligned}$$

Variational Evidence Lower Bound (ELBO)

$$\begin{aligned}\mathcal{L}_{q,\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} + \int q(\mathbf{z}) \log \frac{p(\mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z}))\end{aligned}$$

Log-Likelihood Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})) = \\ &= \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z})) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})).\end{aligned}$$

- What do we get if we drop $D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z}))$ term from ELBO?

Variational Evidence Lower Bound (ELBO)

$$\begin{aligned}\mathcal{L}_{q,\theta}(\mathbf{x}) &= \int q(\mathbf{z}) \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \int q(\mathbf{z}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} + \int q(\mathbf{z}) \log \frac{p(\mathbf{z})}{q(\mathbf{z})} d\mathbf{z} \\ &= \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z}))\end{aligned}$$

Log-Likelihood Decomposition

$$\begin{aligned}\log p_{\theta}(\mathbf{x}) &= \mathcal{L}_{q,\theta}(\mathbf{x}) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})) = \\ &= \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z})) + D_{\text{KL}}(q(\mathbf{z})\|p_{\theta}(\mathbf{z}|\mathbf{x})).\end{aligned}$$

- ▶ What do we get if we drop $D_{\text{KL}}(q(\mathbf{z})\|p(\mathbf{z}))$ term from ELBO?
- ▶ What do we have in the term $\mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z})$ in the case of Normal distribution?

Outline

1. Latent Variable Models (LVM) (continued)
2. Variational Evidence Lower Bound (ELBO)
3. Amortized Inference
4. ELBO Gradients, Reparametrization Trick

Parametric Variable Posterior

Variational Posterior

$$q(\mathbf{z}) = \arg \max_q \mathcal{L}_{q, \theta^*}(\mathbf{x}) = \arg \min_q D_{\text{KL}}(q \| p) = p_{\theta^*}(\mathbf{z} | \mathbf{x}).$$

Parametric Variable Posterior

Variational Posterior

$$q(\mathbf{z}) = \arg \max_q \mathcal{L}_{q, \theta^*}(\mathbf{x}) = \arg \min_q D_{\text{KL}}(q \| p) = p_{\theta^*}(\mathbf{z} | \mathbf{x}).$$

- ▶ $p_{\theta^*}(\mathbf{z} | \mathbf{x})$ may be **intractable**;
- ▶ $q(\mathbf{z})$ is individual for each data point $\mathbf{x}$.

Parametric Variable Posterior

Variational Posterior

$$q(\mathbf{z}) = \arg \max_q \mathcal{L}_{q, \theta^*}(\mathbf{x}) = \arg \min_q D_{\text{KL}}(q \| p) = p_{\theta^*}(\mathbf{z} | \mathbf{x}).$$

- ▶ $p_{\theta^*}(\mathbf{z} | \mathbf{x})$ may be **intractable**;
- ▶ $q(\mathbf{z})$ is individual for each data point $\mathbf{x}$.

Amortized Variational Inference

We restrict the family of possible distributions $q(\mathbf{z})$ to a parametric class $q_\phi(\mathbf{z} | \mathbf{x})$, **conditioned on data $\mathbf{x}$** and **parameterized by ϕ** .

Parametric Variable Posterior

Variational Posterior

$$q(\mathbf{z}) = \arg \max_q \mathcal{L}_{q, \theta^*}(\mathbf{x}) = \arg \min_q D_{\text{KL}}(q \| p) = p_{\theta^*}(\mathbf{z} | \mathbf{x}).$$

- ▶ $p_{\theta^*}(\mathbf{z} | \mathbf{x})$ may be **intractable**;
- ▶ $q(\mathbf{z})$ is individual for each data point $\mathbf{x}$.

Amortized Variational Inference

We restrict the family of possible distributions $q(\mathbf{z})$ to a parametric class $q_\phi(\mathbf{z} | \mathbf{x})$, **conditioned on data $\mathbf{x}$** and **parameterized by ϕ** .


Gradient Update

$$\begin{bmatrix} \phi_k \\ \theta_k \end{bmatrix} = \left[\begin{bmatrix} \phi_{k-1} + \eta \cdot \nabla_\phi \mathcal{L}_{\phi, \theta}(\mathbf{x}) \\ \theta_{k-1} + \eta \cdot \nabla_\theta \mathcal{L}_{\phi, \theta}(\mathbf{x}) \end{bmatrix} \right]_{(\phi_{k-1}, \theta_{k-1})}$$

ELBO Optimization

Gradient Update

$$\begin{bmatrix} \phi_k \\ \theta_k \end{bmatrix} = \begin{bmatrix} \phi_{k-1} + \eta \cdot \nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) \\ \theta_{k-1} + \eta \cdot \nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x}) \end{bmatrix} \Big|_{(\phi_{k-1}, \theta_{k-1})}$$



ELBO Optimization

ELBO

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{\phi, \theta}(\mathbf{x}) + D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p_{\theta}(\mathbf{z}|\mathbf{x})) \geq \mathcal{L}_{\phi, \theta}(\mathbf{x}).$$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

ELBO Optimization

ELBO

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{\phi, \theta}(\mathbf{x}) + D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p_{\theta}(\mathbf{z}|\mathbf{x})) \geq \mathcal{L}_{\phi, \theta}(\mathbf{x}).$$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Gradient Update

$$\begin{bmatrix} \phi_k \\ \theta_k \end{bmatrix} = \left[\begin{bmatrix} \phi_{k-1} + \eta \cdot \nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) \\ \theta_{k-1} + \eta \cdot \nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x}) \end{bmatrix} \right]_{(\phi_{k-1}, \theta_{k-1})}$$

- ▶ ϕ denotes the parameters of the variational posterior $q_{\phi}(\mathbf{z}|\mathbf{x})$.
- ▶ θ represents the parameters of the generative model $p_{\theta}(\mathbf{x}|\mathbf{z})$.

ELBO Optimization

ELBO

$$\log p_{\theta}(\mathbf{x}) = \mathcal{L}_{\phi, \theta}(\mathbf{x}) + D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p_{\theta}(\mathbf{z}|\mathbf{x})) \geq \mathcal{L}_{\phi, \theta}(\mathbf{x}).$$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Gradient Update

$$\begin{bmatrix} \phi_k \\ \theta_k \end{bmatrix} = \left[\begin{bmatrix} \phi_{k-1} + \eta \cdot \nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) \\ \theta_{k-1} + \eta \cdot \nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x}) \end{bmatrix} \right]_{(\phi_{k-1}, \theta_{k-1})}$$

- ▶ ϕ denotes the parameters of the variational posterior $q_{\phi}(\mathbf{z}|\mathbf{x})$.
- ▶ θ represents the parameters of the generative model $p_{\theta}(\mathbf{x}|\mathbf{z})$.

The remaining step is to obtain **unbiased** Monte Carlo estimates of the gradients: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$ and $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$.

Outline

1. Latent Variable Models (LVM) (continued)
2. Variational Evidence Lower Bound (ELBO)
3. Amortized Inference
4. ELBO Gradients, Reparametrization Trick

ELBO Gradients: $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

ELBO Gradients: $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Gradient $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x}) = \nabla_{\theta} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z}$$

ELBO Gradients: $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Gradient $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\begin{aligned}\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \nabla_{\theta} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} \\ &= \int q_{\phi}(\mathbf{z}|\mathbf{x}) \nabla_{\theta} \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z}\end{aligned}$$

ELBO Gradients: $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Gradient $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\begin{aligned}\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \nabla_{\theta} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} \\ &= \int q_{\phi}(\mathbf{z}|\mathbf{x}) \nabla_{\theta} \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} \\ &\approx \nabla_{\theta} \log p_{\theta}(\mathbf{x}|\mathbf{z}^*), \quad \mathbf{z}^* \sim q_{\phi}(\mathbf{z}|\mathbf{x}).\end{aligned}$$

ELBO Gradients: $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Gradient $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\begin{aligned}\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \nabla_{\theta} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} \\ &= \int q_{\phi}(\mathbf{z}|\mathbf{x}) \nabla_{\theta} \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} \\ &\approx \nabla_{\theta} \log p_{\theta}(\mathbf{x}|\mathbf{z}^*), \quad \mathbf{z}^* \sim q_{\phi}(\mathbf{z}|\mathbf{x}).\end{aligned}$$

Naive Monte Carlo Estimation

$$\log p_{\theta}(\mathbf{x}) \geq \int \log p_{\theta}(\mathbf{x}|\mathbf{z}) p(\mathbf{z}) d\mathbf{z} \approx \frac{1}{K} \sum_{k=1}^K \log p_{\theta}(\mathbf{x}|\mathbf{z}_k), \quad \mathbf{z}_k \sim p(\mathbf{z}).$$

ELBO Gradients: $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_q \log p_{\theta}(\mathbf{x}|\mathbf{z}) - D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Gradient $\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\begin{aligned}\nabla_{\theta} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \nabla_{\theta} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} \\ &= \int q_{\phi}(\mathbf{z}|\mathbf{x}) \nabla_{\theta} \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} \\ &\approx \nabla_{\theta} \log p_{\theta}(\mathbf{x}|\mathbf{z}^*), \quad \mathbf{z}^* \sim q_{\phi}(\mathbf{z}|\mathbf{x}).\end{aligned}$$

Naive Monte Carlo Estimation

$$\log p_{\theta}(\mathbf{x}) \geq \int \log p_{\theta}(\mathbf{x}|\mathbf{z}) p(\mathbf{z}) d\mathbf{z} \approx \frac{1}{K} \sum_{k=1}^K \log p_{\theta}(\mathbf{x}|\mathbf{z}_k), \quad \mathbf{z}_k \sim p(\mathbf{z}).$$

The variational posterior $q_{\phi}(\mathbf{z}|\mathbf{x})$ typically concentrates more probability mass in a much smaller region than the prior $p(\mathbf{z})$.

ELBO Gradients: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Gradient $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Unlike the θ -gradient, the density $q_{\phi}(\mathbf{z}|\mathbf{x})$ now depends on ϕ , so standard Monte Carlo estimation can't be applied:

$$\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) = \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

ELBO Gradients: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Gradient $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Unlike the θ -gradient, the density $q_{\phi}(\mathbf{z}|\mathbf{x})$ now depends on ϕ , so standard Monte Carlo estimation can't be applied:

$$\begin{aligned}\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z})) \\ &\neq \int q_{\phi}(\mathbf{z}|\mathbf{x}) \nabla_{\phi} \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))\end{aligned}$$

ELBO Gradients: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Gradient $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Unlike the θ -gradient, the density $q_{\phi}(\mathbf{z}|\mathbf{x})$ now depends on ϕ , so standard Monte Carlo estimation can't be applied:

$$\begin{aligned}\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z})) \\ &\neq \int q_{\phi}(\mathbf{z}|\mathbf{x}) \nabla_{\phi} \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))\end{aligned}$$

Reparametrization Trick (LOTUS Trick)

Assume $\mathbf{z} \sim q_{\phi}(\mathbf{z}|\mathbf{x})$ is generated by a random variable $\epsilon \sim p(\epsilon)$ via a deterministic mapping $\mathbf{z} = \mathbf{g}_{\phi}(\mathbf{x}, \epsilon)$. Then,

$$\mathbb{E}_{\mathbf{z} \sim q_{\phi}(\mathbf{z}|\mathbf{x})} \mathbf{f}(\mathbf{z}) = \mathbb{E}_{\epsilon \sim p(\epsilon)} \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon))$$

ELBO Gradients: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Gradient $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Unlike the θ -gradient, the density $q_{\phi}(\mathbf{z}|\mathbf{x})$ now depends on ϕ , so standard Monte Carlo estimation can't be applied:

$$\begin{aligned}\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z})) \\ &\neq \int q_{\phi}(\mathbf{z}|\mathbf{x}) \nabla_{\phi} \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))\end{aligned}$$

Reparametrization Trick (LOTUS Trick)

Assume $\mathbf{z} \sim q_{\phi}(\mathbf{z}|\mathbf{x})$ is generated by a random variable $\epsilon \sim p(\epsilon)$ via a deterministic mapping $\mathbf{z} = \mathbf{g}_{\phi}(\mathbf{x}, \epsilon)$. Then,

$$\mathbb{E}_{\mathbf{z} \sim q_{\phi}(\mathbf{z}|\mathbf{x})} \mathbf{f}(\mathbf{z}) = \mathbb{E}_{\epsilon \sim p(\epsilon)} \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon))$$

Note: The LHS expectation is with respect to the parametric distribution $q_{\phi}(\mathbf{z}|\mathbf{x})$, while the RHS is for the non-parametric $p(\epsilon)$.

Kingma D.P., Welling M. Auto-Encoding Variational Bayes, 2013

ELBO Gradients: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Reparametrization Trick (LOTUS Trick)

$$\nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \mathbf{f}(\mathbf{z}) d\mathbf{z} = \nabla_{\phi} \int p(\epsilon) \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon$$

,

ELBO Gradients: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Reparametrization Trick (LOTUS Trick)

$$\begin{aligned}\nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \mathbf{f}(\mathbf{z}) d\mathbf{z} &= \nabla_{\phi} \int p(\epsilon) \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon = \\ &= \int p(\epsilon) \nabla_{\phi} \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon \approx \nabla_{\phi} \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon^*)),\end{aligned}$$

where $\epsilon^* \sim p(\epsilon)$.

ELBO Gradients: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

Reparametrization Trick (LOTUS Trick)

$$\begin{aligned}\nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \mathbf{f}(\mathbf{z}) d\mathbf{z} &= \nabla_{\phi} \int p(\epsilon) \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon = \\ &= \int p(\epsilon) \nabla_{\phi} \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon \approx \nabla_{\phi} \mathbf{f}(\mathbf{g}_{\phi}(\mathbf{x}, \epsilon^*)),\end{aligned}$$

where $\epsilon^* \sim p(\epsilon)$.

Variational Assumption

$$p(\epsilon) = \mathcal{N}(0, \mathbf{I}); \quad \mathbf{z} = \mathbf{g}_{\phi}(\mathbf{x}, \epsilon) = \sigma_{\phi}(\mathbf{x}) \odot \epsilon + \mu_{\phi}(\mathbf{x});$$

$$q_{\phi}(\mathbf{z}|\mathbf{x}) = \mathcal{N}(\mu_{\phi}(\mathbf{x}), \sigma_{\phi}^2(\mathbf{x})).$$

Here, $\mu_{\phi}(\cdot)$ and $\sigma_{\phi}(\cdot)$ are parameterized functions (outputs of a neural network).

Thus, we can write $q_{\phi}(\mathbf{z}|\mathbf{x}) = \text{NN}_{e, \phi}(\mathbf{x})$, the **encoder**.

ELBO Gradient: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) = \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

ELBO Gradient: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) = \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Reconstruction Term

$$\begin{aligned} \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} &= \int p(\epsilon) \nabla_{\phi} \log p_{\theta}(\mathbf{x} | \mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon \approx \\ &\approx \nabla_{\phi} \log p_{\theta}(\mathbf{x} | \sigma_{\phi}(\mathbf{x}) \odot \epsilon^* + \mu_{\phi}(\mathbf{x})), \quad \text{where } \epsilon^* \sim \mathcal{N}(0, \mathbf{I}) \end{aligned}$$

ELBO Gradient: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) = \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Reconstruction Term

$$\begin{aligned} \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} &= \int p(\epsilon) \nabla_{\phi} \log p_{\theta}(\mathbf{x} | \mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon \approx \\ &\approx \nabla_{\phi} \log p_{\theta}(\mathbf{x} | \sigma_{\phi}(\mathbf{x}) \odot \epsilon^* + \mu_{\phi}(\mathbf{x})), \quad \text{where } \epsilon^* \sim \mathcal{N}(0, \mathbf{I}) \end{aligned}$$

The generative distribution $p_{\theta}(\mathbf{x}|\mathbf{z})$ can be implemented as a neural network.

We may write $p_{\theta}(\mathbf{x}|\mathbf{z}) = \text{NN}_{d, \theta}(\mathbf{z})$, called the **decoder**.

ELBO Gradient: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) = \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Reconstruction Term

$$\begin{aligned} \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} &= \int p(\epsilon) \nabla_{\phi} \log p_{\theta}(\mathbf{x} | \mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon \approx \\ &\approx \nabla_{\phi} \log p_{\theta}(\mathbf{x} | \sigma_{\phi}(\mathbf{x}) \odot \epsilon^* + \mu_{\phi}(\mathbf{x})), \quad \text{where } \epsilon^* \sim \mathcal{N}(0, \mathbf{I}) \end{aligned}$$

The generative distribution $p_{\theta}(\mathbf{x}|\mathbf{z})$ can be implemented as a neural network.

We may write $p_{\theta}(\mathbf{x}|\mathbf{z}) = \text{NN}_{d, \theta}(\mathbf{z})$, called the **decoder**.

KL Term

$p(\mathbf{z})$ is the prior over latents $\mathbf{z}$, typically $p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I})$.

$$\nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z})) = \nabla_{\phi} D_{\text{KL}}(\mathcal{N}(\mu_{\phi}(\mathbf{x}), \sigma_{\phi}^2(\mathbf{x})) \| \mathcal{N}(0, \mathbf{I}))$$

ELBO Gradient: $\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x})$

$$\nabla_{\phi} \mathcal{L}_{\phi, \theta}(\mathbf{x}) = \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} - \nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z}))$$

Reconstruction Term

$$\begin{aligned} \nabla_{\phi} \int q_{\phi}(\mathbf{z}|\mathbf{x}) \log p_{\theta}(\mathbf{x}|\mathbf{z}) d\mathbf{z} &= \int p(\epsilon) \nabla_{\phi} \log p_{\theta}(\mathbf{x} | \mathbf{g}_{\phi}(\mathbf{x}, \epsilon)) d\epsilon \approx \\ &\approx \nabla_{\phi} \log p_{\theta}(\mathbf{x} | \sigma_{\phi}(\mathbf{x}) \odot \epsilon^* + \mu_{\phi}(\mathbf{x})), \quad \text{where } \epsilon^* \sim \mathcal{N}(0, \mathbf{I}) \end{aligned}$$

The generative distribution $p_{\theta}(\mathbf{x}|\mathbf{z})$ can be implemented as a neural network.

We may write $p_{\theta}(\mathbf{x}|\mathbf{z}) = \text{NN}_{d, \theta}(\mathbf{z})$, called the **decoder**.

KL Term

$p(\mathbf{z})$ is the prior over latents $\mathbf{z}$, typically $p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I})$.

$$\nabla_{\phi} D_{\text{KL}}(q_{\phi}(\mathbf{z}|\mathbf{x}) \| p(\mathbf{z})) = \nabla_{\phi} D_{\text{KL}}(\mathcal{N}(\mu_{\phi}(\mathbf{x}), \sigma_{\phi}^2(\mathbf{x})) \| \mathcal{N}(0, \mathbf{I}))$$

This expression admits a closed-form analytic solution.

Summary

- ▶ LVMs maximize the variational evidence lower bound (ELBO) to obtain maximum likelihood estimates for the parameters.
- ▶ Parametric posterior distribution $q_{\phi}(\mathbf{z}|\mathbf{x})$ makes the method scalable.
- ▶ The reparametrization trick provides unbiased gradients with respect to the variational posterior $q_{\phi}(\mathbf{z}|\mathbf{x})$.