

Deep Generative Models

Lecture 7

Roman Isachenko

Moscow Institute of Physics and Technology
Yandex School of Data Analysis

2026, Autumn

Recap of Previous Lecture

Theorem (Informal)

Let $\mathbf{x}_0$ be a random vector. Under mild regularity conditions, samples from the following dynamics will eventually follow $p_{\theta}(\mathbf{x})$ (for sufficiently small η and large l):

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2} \cdot \nabla_{\mathbf{x}_l} \log p_{\theta}(\mathbf{x}_l) + \sqrt{\eta} \cdot \epsilon_l, \quad \epsilon_l \sim \mathcal{N}(0, \mathbf{I}).$$

- ▶ The density $p_{\theta}(\mathbf{x})$ is the **stationary** distribution of the Markov chain.
- ▶ The gradient is taken with respect to $\mathbf{x}$, not θ .
- ▶ $\nabla_{\mathbf{x}} \log p_{\theta}(\mathbf{x})$ defines a vector field.

Fisher Divergence

$$D_F(p_{\text{data}}, p_{\theta}) = \frac{1}{2} \mathbb{E}_{p_{\text{data}}} \|\nabla_{\mathbf{x}} \log p_{\theta}(\mathbf{x}) - \nabla_{\mathbf{x}} \log p_{\text{data}}(\mathbf{x})\|_2^2 \rightarrow \min_{\theta}$$

Lai C. H. et al. The principles of diffusion models, 2025.

Song Y. Generative Modeling by Estimating Gradients of the Data Distribution, blog post, 2021

Recap of Previous Lecture


Define **score function** $\mathbf{s}_\theta(\mathbf{x}) = \nabla_{\mathbf{x}} \log p_\theta(\mathbf{x})$.

Fisher Divergence

$$D_F(p_{\text{data}}, p_\theta) = \frac{1}{2} \mathbb{E}_{p_{\text{data}}} \|\mathbf{s}_\theta(\mathbf{x}) - \nabla_{\mathbf{x}} \log p_{\text{data}}(\mathbf{x})\|_2^2 \rightarrow \min_{\theta}$$

Langevin Dynamics

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2} \cdot \mathbf{s}_\theta(\mathbf{x}_l) + \sqrt{\eta} \cdot \epsilon_l, \quad \epsilon_l \sim \mathcal{N}(0, \mathbf{I})$$



Song Y. *Generative Modeling by Estimating Gradients of the Data Distribution*, blog post, 2021

Recap of Previous Lecture

Let us perturb the original data with Gaussian noise

$$q(\mathbf{x}_\sigma|\mathbf{x}) = \mathcal{N}(\mathbf{x}, \sigma^2 \cdot \mathbf{I}).$$

$$q(\mathbf{x}_\sigma) = \int q(\mathbf{x}_\sigma|\mathbf{x})p_{\text{data}}(\mathbf{x})d\mathbf{x}.$$

Assumption

The solution to

$$\frac{1}{2}\mathbb{E}_{q(\mathbf{x}_\sigma)}\|\mathbf{s}_{\theta,\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma} \log q(\mathbf{x}_\sigma)\|_2^2 \rightarrow \min_{\theta}$$

satisfies $\mathbf{s}_{\theta,\sigma}(\mathbf{x}_\sigma) \approx \mathbf{s}_{\theta,0}(\mathbf{x}_0) = \mathbf{s}_{\theta}(\mathbf{x})$ if σ is sufficiently small.

Theorem

$$\begin{aligned} \mathbb{E}_{q(\mathbf{x}_\sigma)}\|\mathbf{s}_{\theta,\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma} \log q(\mathbf{x}_\sigma)\|_2^2 &= \\ &= \mathbb{E}_{p_{\text{data}}(\mathbf{x})}\mathbb{E}_{q(\mathbf{x}_\sigma|\mathbf{x})}\|\mathbf{s}_{\theta,\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma} \log q(\mathbf{x}_\sigma|\mathbf{x})\|_2^2 + \text{const}(\theta) \end{aligned}$$

Here, $\nabla_{\mathbf{x}_\sigma} \log q(\mathbf{x}_\sigma|\mathbf{x}) = -\frac{\mathbf{x}_\sigma - \mathbf{x}}{\sigma^2} = -\frac{\epsilon}{\sigma}$. $\mathbf{s}_{\theta,\sigma}(\mathbf{x}_\sigma)$ attempts to **denoise** a corrupted sample.


Vincent P. A Connection Between Score Matching and Denoising Autoencoders, 2010

Recap of Previous Lecture

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x})} \mathbb{E}_{\mathcal{N}(0, \mathbf{I})} \left\| \mathbf{s}_{\theta, \sigma}(\mathbf{x} + \sigma \boldsymbol{\epsilon}) + \frac{\boldsymbol{\epsilon}}{\sigma} \right\|_2^2 \rightarrow \min_{\theta}$$

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2} \cdot \mathbf{s}_{\theta, \sigma}(\mathbf{x}_l) + \sqrt{\eta} \cdot \boldsymbol{\epsilon}_l$$

- ▶ For **small** σ , $\mathbf{s}_{\theta, \sigma}(\mathbf{x})$ becomes inaccurate and Langevin dynamics fails to traverse modes
- ▶ For **large** σ , robustness in low-density regions is achieved, but the model learns a distribution that is overly corrupted





Recap of Previous Lecture

Noise-Conditioned Score Network (NCSN)

- ▶ Define a sequence of noise levels: $\sigma_1 < \sigma_2 < \dots < \sigma_T$.
- ▶ Train the denoising score function $\mathbf{s}_{\theta, \sigma_t}(\mathbf{x}_t)$ for each noise level:

$$\sum_{t=1}^T \sigma_t^2 \cdot \mathbb{E}_{p_{\text{data}}(\mathbf{x})} \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x})} \left\| \mathbf{s}_{\theta, \sigma_t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}) \right\|_2^2 \rightarrow \min_{\theta}$$

- ▶ Sample using **annealed** Langevin dynamics (for $t = T, \dots, 1$).



Song Y. et al. *Generative Modeling by Estimating Gradients of the Data Distribution*, 2019

Recap of Previous Lecture

Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $t \sim U\{1, T\}$, $\epsilon \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \mathbf{x}_0 + \sigma_t \epsilon$.
3. Compute loss $\mathcal{L} = \sigma_t^2 \left\| \mathbf{s}_{\theta, \sigma_t}(\mathbf{x}_t) + \frac{\epsilon}{\sigma_t} \right\|^2$.
4. Update θ via stochastic gradient descent.

Sampling (Annealed Langevin Dynamics)

1. Sample $\mathbf{x}_T^0 \sim \mathcal{N}(0, \sigma_T^2 \mathbf{I}) \approx q(\mathbf{x}_T)$.
2. Update $\mathbf{x}_t^l$ via L steps of Langevin dynamics at each noise level σ_t :
$$\mathbf{x}_t^l = \mathbf{x}_t^{l-1} + \frac{\eta_t}{2} \mathbf{s}_{\theta, \sigma_t}(\mathbf{x}_t^{l-1}) + \sqrt{\eta_t} \epsilon_t^l.$$
3. Update $\mathbf{x}_{t-1}^0 := \mathbf{x}_t^L$ and proceed to the next noise level σ_{t-1} .

Song Y. et al. *Generative Modeling by Estimating Gradients of the Data Distribution*, 2019

Song Y. et al. *Improved Techniques for Training Score-Based Generative Models*, 2020


Outline

1. Forward Gaussian Diffusion Process
2. Reverse Gaussian Diffusion Process
3. Gaussian Diffusion Model as VAE
4. Diffusion ELBO Derivation

Outline

1. Forward Gaussian Diffusion Process
2. Reverse Gaussian Diffusion Process
3. Gaussian Diffusion Model as VAE
4. Diffusion ELBO Derivation

Generative Models Taxonomy



Forward Gaussian Diffusion Process

Let $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$, $\beta_t \ll 1$. Define a Markov chain:

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Forward Gaussian Diffusion Process

Let $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$, $\beta_t \ll 1$. Define a Markov chain:

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Forward Gaussian Diffusion Process

Let $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$, $\beta_t \ll 1$. Define a Markov chain:

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Langevin Dynamics

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2} \cdot \nabla_{\mathbf{x}_l} \log p_{\theta}(\mathbf{x}_l) + \sqrt{\eta} \boldsymbol{\epsilon}_l, \quad \boldsymbol{\epsilon}_l \sim \mathcal{N}(0, \mathbf{I})$$

Forward Gaussian Diffusion Process

Let $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$, $\beta_t \ll 1$. Define a Markov chain:

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Langevin Dynamics

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2} \cdot \nabla_{\mathbf{x}_l} \log p_{\theta}(\mathbf{x}_l) + \sqrt{\eta} \boldsymbol{\epsilon}_l, \quad \boldsymbol{\epsilon}_l \sim \mathcal{N}(0, \mathbf{I})$$

$$\begin{aligned} \mathbf{x}_t &= \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t \approx \left(1 - \frac{\beta_t}{2}\right) \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t = \\ &= \mathbf{x}_{t-1} + \frac{\beta_t}{2} (-\mathbf{x}_{t-1}) + \sqrt{\beta_t} \boldsymbol{\epsilon}_t \end{aligned}$$

Forward Gaussian Diffusion Process

Let $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$, $\beta_t \ll 1$. Define a Markov chain:

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Langevin Dynamics

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2} \cdot \nabla_{\mathbf{x}_l} \log p_{\theta}(\mathbf{x}_l) + \sqrt{\eta} \boldsymbol{\epsilon}_l, \quad \boldsymbol{\epsilon}_l \sim \mathcal{N}(0, \mathbf{I})$$

$$\begin{aligned} \mathbf{x}_t &= \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t \approx \left(1 - \frac{\beta_t}{2}\right) \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t = \\ &= \mathbf{x}_{t-1} + \frac{\beta_t}{2} (-\mathbf{x}_{t-1}) + \sqrt{\beta_t} \boldsymbol{\epsilon}_t \end{aligned}$$

► $\beta_t = \eta$

► $\nabla_{\mathbf{x}_{t-1}} \log p_{\theta}(\mathbf{x}_{t-1}) = -\mathbf{x}_{t-1} = \nabla_{\mathbf{x}_{t-1}} \log \mathcal{N}(0, \mathbf{I})$

Forward Gaussian Diffusion Process

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Forward Gaussian Diffusion Process

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Statement 1

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^t \alpha_s = \prod_{s=1}^t (1 - \beta_s)$. Then

$$q(\mathbf{x}_t | \mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \mathbf{x}_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

Forward Gaussian Diffusion Process

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Statement 1

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^t \alpha_s = \prod_{s=1}^t (1 - \beta_s)$. Then

$$q(\mathbf{x}_t | \mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \mathbf{x}_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

Thus, samples at any timestep t can be generated directly from $\mathbf{x}_0$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \boldsymbol{\epsilon}_t$$

Forward Gaussian Diffusion Process

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Statement 1

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^t \alpha_s = \prod_{s=1}^t (1 - \beta_s)$. Then

$$q(\mathbf{x}_t | \mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \mathbf{x}_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

Thus, samples at any timestep t can be generated directly from $\mathbf{x}_0$

$$\begin{aligned} \mathbf{x}_t &= \sqrt{\alpha_t} \mathbf{x}_{t-1} + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t = \\ &= \sqrt{\alpha_t} (\sqrt{\alpha_{t-1}} \mathbf{x}_{t-2} + \sqrt{1 - \alpha_{t-1}} \boldsymbol{\epsilon}_{t-1}) + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t \end{aligned}$$

Forward Gaussian Diffusion Process

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Statement 1

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^t \alpha_s = \prod_{s=1}^t (1 - \beta_s)$. Then

$$q(\mathbf{x}_t | \mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \mathbf{x}_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

Thus, samples at any timestep t can be generated directly from $\mathbf{x}_0$

$$\begin{aligned} \mathbf{x}_t &= \sqrt{\alpha_t} \mathbf{x}_{t-1} + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t = \\ &= \sqrt{\alpha_t} (\sqrt{\alpha_{t-1}} \mathbf{x}_{t-2} + \sqrt{1 - \alpha_{t-1}} \boldsymbol{\epsilon}_{t-1}) + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t = \\ &= \sqrt{\alpha_t \alpha_{t-1}} \mathbf{x}_{t-2} + (\sqrt{\alpha_t (1 - \alpha_{t-1})} \boldsymbol{\epsilon}_{t-1} + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t) \end{aligned}$$

Forward Gaussian Diffusion Process

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Statement 1

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^t \alpha_s = \prod_{s=1}^t (1 - \beta_s)$. Then

$$q(\mathbf{x}_t | \mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \mathbf{x}_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

Thus, samples at any timestep t can be generated directly from $\mathbf{x}_0$

$$\begin{aligned} \mathbf{x}_t &= \sqrt{\alpha_t} \mathbf{x}_{t-1} + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t = \\ &= \sqrt{\alpha_t} (\sqrt{\alpha_{t-1}} \mathbf{x}_{t-2} + \sqrt{1 - \alpha_{t-1}} \boldsymbol{\epsilon}_{t-1}) + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t = \\ &= \sqrt{\alpha_t \alpha_{t-1}} \mathbf{x}_{t-2} + (\sqrt{\alpha_t (1 - \alpha_{t-1})} \boldsymbol{\epsilon}_{t-1} + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t) = \\ &= \sqrt{\alpha_t \alpha_{t-1}} \mathbf{x}_{t-2} + \sqrt{1 - \alpha_t \alpha_{t-1}} \boldsymbol{\epsilon}'_t \end{aligned}$$

Forward Gaussian Diffusion Process

$$\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1 - \beta_t} \mathbf{x}_{t-1}, \beta_t \mathbf{I})$$

Statement 1

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^t \alpha_s = \prod_{s=1}^t (1 - \beta_s)$. Then


$$q(\mathbf{x}_t | \mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \mathbf{x}_0, (1 - \bar{\alpha}_t) \mathbf{I})$$

Thus, samples at any timestep t can be generated directly from $\mathbf{x}_0$

$$\begin{aligned} \mathbf{x}_t &= \sqrt{\alpha_t} \mathbf{x}_{t-1} + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t = \\ &= \sqrt{\alpha_t} (\sqrt{\alpha_{t-1}} \mathbf{x}_{t-2} + \sqrt{1 - \alpha_{t-1}} \boldsymbol{\epsilon}_{t-1}) + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t = \\ &= \sqrt{\alpha_t \alpha_{t-1}} \mathbf{x}_{t-2} + (\sqrt{\alpha_t (1 - \alpha_{t-1})} \boldsymbol{\epsilon}_{t-1} + \sqrt{1 - \alpha_t} \boldsymbol{\epsilon}_t) = \\ &= \sqrt{\alpha_t \alpha_{t-1}} \mathbf{x}_{t-2} + \sqrt{1 - \alpha_t \alpha_{t-1}} \boldsymbol{\epsilon}'_t = \\ &= \dots = \sqrt{\bar{\alpha}_t} \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \boldsymbol{\epsilon}, \quad \boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}) \end{aligned}$$


Forward Gaussian Diffusion Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right); \quad q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}\left(\sqrt{\bar{\alpha}_t}\mathbf{x}_0, (1-\bar{\alpha}_t)\mathbf{I}\right)$$



Forward Gaussian Diffusion Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right); \quad q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}\left(\sqrt{\bar{\alpha}_t}\mathbf{x}_0, (1-\bar{\alpha}_t)\mathbf{I}\right)$$




Statement 2

Applying the Markov chain to any distribution $p_{\text{data}}(\mathbf{x})$ yields $\mathbf{x}_\infty \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$, the **stationary** (limiting) distribution:

$$p_\infty(\mathbf{x}) = \int q(\mathbf{x}|\mathbf{x}')p_\infty(\mathbf{x}')d\mathbf{x}'$$

Forward Gaussian Diffusion Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right); \quad q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}\left(\sqrt{\bar{\alpha}_t}\mathbf{x}_0, (1-\bar{\alpha}_t)\mathbf{I}\right)$$



Statement 2


Applying the Markov chain to any distribution $p_{\text{data}}(\mathbf{x})$ yields $\mathbf{x}_\infty \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$, the **stationary** (limiting) distribution:

$$p_\infty(\mathbf{x}) = \int q(\mathbf{x}|\mathbf{x}')p_\infty(\mathbf{x}')d\mathbf{x}'$$

$$p_\infty(\mathbf{x}) = \int q(\mathbf{x}_\infty|\mathbf{x}_0)p_{\text{data}}(\mathbf{x}_0)d\mathbf{x}_0 \approx \mathcal{N}(0, \mathbf{I}) \int p_{\text{data}}(\mathbf{x}_0)d\mathbf{x}_0 = \mathcal{N}(0, \mathbf{I})$$

Forward Gaussian Diffusion Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right); \quad q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}\left(\sqrt{\bar{\alpha}_t}\mathbf{x}_0, (1-\bar{\alpha}_t)\mathbf{I}\right)$$



Statement 2

Applying the Markov chain to any distribution $p_{\text{data}}(\mathbf{x})$ yields $\mathbf{x}_\infty \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$, the **stationary** (limiting) distribution:


$$p_\infty(\mathbf{x}) = \int q(\mathbf{x}|\mathbf{x}')p_\infty(\mathbf{x}')d\mathbf{x}'$$

$$p_\infty(\mathbf{x}) = \int q(\mathbf{x}_\infty|\mathbf{x}_0)p_{\text{data}}(\mathbf{x}_0)d\mathbf{x}_0 \approx \mathcal{N}(0, \mathbf{I}) \int p_{\text{data}}(\mathbf{x}_0)d\mathbf{x}_0 = \mathcal{N}(0, \mathbf{I})$$

Note: This holds iff $\bar{\alpha}_t \rightarrow 0$, i.e., $\sum_{t=1}^{\infty} \beta_t = +\infty$.


Forward Gaussian Diffusion Process

Diffusion describes the migration of particles from regions of high density to those of low density.



Forward Gaussian Diffusion Process


Diffusion describes the migration of particles from regions of high density to those of low density.



1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \epsilon_t, \epsilon_t \sim \mathcal{N}(0, \mathbf{I}), t \geq 1$
3. After $T \gg 1$ steps: $\mathbf{x}_T \sim p_{\infty}(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

Forward Gaussian Diffusion Process

Diffusion describes the migration of particles from regions of high density to those of low density.



1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \epsilon_t, \epsilon_t \sim \mathcal{N}(0, \mathbf{I}), t \geq 1$
3. After $T \gg 1$ steps: $\mathbf{x}_T \sim p_{\infty}(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

If this process can be reversed, we can sample from $p_{\text{data}}(\mathbf{x})$ by starting from noise $p_{\infty}(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.

Our goal now becomes inverting this diffusion.

Denoising Score Matching

NCSN

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\mathbf{x}_0, \sigma_t^2 \mathbf{I}), \quad q(\mathbf{x}_1) \approx p_{\text{data}}(\mathbf{x}), \quad q(\mathbf{x}_T) \approx \mathcal{N}(0, \sigma_T^2 \mathbf{I})$$

$$\nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}) = -\frac{\mathbf{x}_t - \mathbf{x}}{\sigma_t^2}$$

Denoising Score Matching

NCSN

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\mathbf{x}_0, \sigma_t^2 \mathbf{I}), \quad q(\mathbf{x}_1) \approx p_{\text{data}}(\mathbf{x}), \quad q(\mathbf{x}_T) \approx \mathcal{N}(0, \sigma_T^2 \mathbf{I})$$

$$\nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}) = -\frac{\mathbf{x}_t - \mathbf{x}}{\sigma_t^2}$$

Gaussian Diffusion

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t}\mathbf{x}_0, (1-\bar{\alpha}_t)\mathbf{I}), \quad q(\mathbf{x}_1) \approx p_{\text{data}}(\mathbf{x}), \quad q(\mathbf{x}_T) \approx \mathcal{N}(0, \mathbf{I})$$

$$\nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}_0) = -\frac{\mathbf{x}_t - \sqrt{\bar{\alpha}_t}\mathbf{x}_0}{1 - \bar{\alpha}_t}$$

Denoising Score Matching

NCSN

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\mathbf{x}_0, \sigma_t^2 \mathbf{I}), \quad q(\mathbf{x}_1) \approx p_{\text{data}}(\mathbf{x}), \quad q(\mathbf{x}_T) \approx \mathcal{N}(0, \sigma_T^2 \mathbf{I})$$

$$\nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}) = -\frac{\mathbf{x}_t - \mathbf{x}}{\sigma_t^2}$$

Gaussian Diffusion

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t}\mathbf{x}_0, (1-\bar{\alpha}_t)\mathbf{I}), \quad q(\mathbf{x}_1) \approx p_{\text{data}}(\mathbf{x}), \quad q(\mathbf{x}_T) \approx \mathcal{N}(0, \mathbf{I})$$

$$\nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}_0) = -\frac{\mathbf{x}_t - \sqrt{\bar{\alpha}_t}\mathbf{x}_0}{1 - \bar{\alpha}_t}$$

Theorem (Denoising Score Matching)

$$\begin{aligned} \mathbb{E}_{q(\mathbf{x}_t)} \|\mathbf{s}_{\theta,t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t)\|_2^2 &= \\ &= \mathbb{E}_{p_{\text{data}}(\mathbf{x})} \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x})} \|\mathbf{s}_{\theta,t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x})\|_2^2 + \text{const}(\theta) \end{aligned}$$

Denoising Score Matching

NCSN

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\mathbf{x}_0, \sigma_t^2 \mathbf{I}), \quad q(\mathbf{x}_1) \approx p_{\text{data}}(\mathbf{x}), \quad q(\mathbf{x}_T) \approx \mathcal{N}(0, \sigma_T^2 \mathbf{I})$$

$$\nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}) = -\frac{\mathbf{x}_t - \mathbf{x}}{\sigma_t^2}$$

Gaussian Diffusion

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \mathbf{x}_0, (1 - \bar{\alpha}_t) \mathbf{I}), \quad q(\mathbf{x}_1) \approx p_{\text{data}}(\mathbf{x}), \quad q(\mathbf{x}_T) \approx \mathcal{N}(0, \mathbf{I})$$

$$\nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}_0) = -\frac{\mathbf{x}_t - \sqrt{\bar{\alpha}_t} \mathbf{x}_0}{1 - \bar{\alpha}_t}$$

Theorem (Denoising Score Matching)


$$\begin{aligned} \mathbb{E}_{q(\mathbf{x}_t)} \|\mathbf{s}_{\theta,t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t)\|_2^2 &= \\ &= \mathbb{E}_{p_{\text{data}}(\mathbf{x})} \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x})} \|\mathbf{s}_{\theta,t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x})\|_2^2 + \text{const}(\theta) \end{aligned}$$

Note: This enables applying the NCSN approach with annealed Langevin dynamics to diffusion-based denoising models.


Outline

1. Forward Gaussian Diffusion Process
2. Reverse Gaussian Diffusion Process
3. Gaussian Diffusion Model as VAE
4. Diffusion ELBO Derivation

Reverse Gaussian Diffusion Process




Reverse Gaussian Diffusion Process



Forward Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1 - \beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right)$$

Reverse Gaussian Diffusion Process




Forward Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right)$$

Reverse Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx p_\theta(\mathbf{x}_{t-1}|\mathbf{x}_t)$$

Reverse Gaussian Diffusion Process



Forward Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right)$$

Reverse Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx p_\theta(\mathbf{x}_{t-1}|\mathbf{x}_t)$$

$q(\mathbf{x}_{t-1})$ and $q(\mathbf{x}_t)$ are intractable:

$$q(\mathbf{x}_t) = \int q(\mathbf{x}_t|\mathbf{x}_0)p_{\text{data}}(\mathbf{x}_0)d\mathbf{x}_0$$

Reverse Gaussian Diffusion Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Theorem (Feller, 1949)

If β_t is sufficiently small, $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ is Gaussian (thus, diffusion requires $T \approx 1000$ steps for convergence)

Feller W. On the theory of stochastic processes, with particular reference to applications, 1949


Xiao Z., Kreis K., Vahdat A. Tackling the generative learning trilemma with denoising diffusion GANs, 2021

Reverse Gaussian Diffusion Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Theorem (Feller, 1949)


If β_t is sufficiently small, $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ is Gaussian (thus, diffusion requires $T \approx 1000$ steps for convergence)



Feller W. *On the theory of stochastic processes, with particular reference to applications*, 1949

Xiao Z., Kreis K., Vahdat A. *Tackling the generative learning trilemma with denoising diffusion GANs*, 2021

Reverse Gaussian Diffusion Process (Ancestral Sampling)




Define the reverse process as:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) \approx p_\theta(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}(\mu_{\theta,t}(\mathbf{x}_t), \sigma_{\theta,t}^2(\mathbf{x}_t))$$

Feller's theorem justifies this Gaussian assumption.

Reverse Gaussian Diffusion Process (Ancestral Sampling)



Define the reverse process as:


$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) \approx p_\theta(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}(\boldsymbol{\mu}_{\theta,t}(\mathbf{x}_t), \boldsymbol{\sigma}_{\theta,t}^2(\mathbf{x}_t))$$

Feller's theorem justifies this Gaussian assumption.

Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t$
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

Reverse Gaussian Diffusion Process (Ancestral Sampling)



Define the reverse process as:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) \approx p_\theta(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}(\mu_{\theta,t}(\mathbf{x}_t), \sigma_{\theta,t}^2(\mathbf{x}_t))$$

Feller's theorem justifies this Gaussian assumption.


Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \epsilon_t$
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

Reverse Process

1. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$
2. $\mathbf{x}_{t-1} = \sigma_{\theta,t}(\mathbf{x}_t) \epsilon + \mu_{\theta,t}(\mathbf{x}_t)$
3. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$

Reverse Gaussian Diffusion Process (Ancestral Sampling)



Define the reverse process as:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) \approx p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}(\boldsymbol{\mu}_{\theta,t}(\mathbf{x}_t), \boldsymbol{\sigma}_{\theta,t}^2(\mathbf{x}_t))$$

Feller's theorem justifies this Gaussian assumption.

Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \mathbf{x}_{t-1} + \sqrt{\beta_t} \boldsymbol{\epsilon}_t$
3. $\mathbf{x}_T \sim p_{\infty}(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

Reverse Process

1. $\mathbf{x}_T \sim p_{\infty}(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$
2. $\mathbf{x}_{t-1} = \boldsymbol{\sigma}_{\theta,t}(\mathbf{x}_t) \boldsymbol{\epsilon} + \boldsymbol{\mu}_{\theta,t}(\mathbf{x}_t)$
3. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$

Note: The forward process is non-learnable, i.e., it does not involve trainable parameters

Conditioned Reverse Distribution

Reverse Kernel (**Intractable**)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Conditioned Reverse Distribution

Reverse Kernel (**Intractable**)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Conditioned Reverse Kernel (**Tractable**)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)q(\mathbf{x}_{t-1}|\mathbf{x}_0)}{q(\mathbf{x}_t|\mathbf{x}_0)}$$

Conditioned Reverse Distribution

Reverse Kernel (**Intractable**)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Conditioned Reverse Kernel (**Tractable**)

$$\begin{aligned} q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) &= \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)q(\mathbf{x}_{t-1}|\mathbf{x}_0)}{q(\mathbf{x}_t|\mathbf{x}_0)} \\ &= \frac{\mathcal{N}(\sqrt{1-\beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \mathbf{I}) \cdot \mathcal{N}(\sqrt{\bar{\alpha}_{t-1}} \cdot \mathbf{x}_0, (1-\bar{\alpha}_{t-1}) \cdot \mathbf{I})}{\mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1-\bar{\alpha}_t) \cdot \mathbf{I})} \end{aligned}$$

Conditioned Reverse Distribution

Reverse Kernel (**Intractable**)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Conditioned Reverse Kernel (**Tractable**)

$$\begin{aligned} q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) &= \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)q(\mathbf{x}_{t-1}|\mathbf{x}_0)}{q(\mathbf{x}_t|\mathbf{x}_0)} \\ &= \frac{\mathcal{N}(\sqrt{1-\beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \mathbf{I}) \cdot \mathcal{N}(\sqrt{\bar{\alpha}_{t-1}} \cdot \mathbf{x}_0, (1-\bar{\alpha}_{t-1}) \cdot \mathbf{I})}{\mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1-\bar{\alpha}_t) \cdot \mathbf{I})} \\ &= \mathcal{N}(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \cdot \mathbf{I}) \end{aligned}$$

Here,

$$\begin{aligned} \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) &= \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t} \cdot \mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t} \cdot \mathbf{x}_0; \\ \tilde{\beta}_t &= \frac{(1-\alpha_t)(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t} = \text{const.} \end{aligned}$$

Distribution Summary

Forward process maps any distribution $p_{\text{data}}(\mathbf{x})$ to $\mathcal{N}(0, \mathbf{I})$ by injection of noise:

$$\begin{aligned}q(\mathbf{x}_t | \mathbf{x}_{t-1}) &= \mathcal{N}(\sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \cdot \mathbf{I}); \\q(\mathbf{x}_t | \mathbf{x}_0) &= \mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_t) \cdot \mathbf{I}).\end{aligned}$$

Distribution Summary

Forward process maps any distribution $p_{\text{data}}(\mathbf{x})$ to $\mathcal{N}(0, \mathbf{I})$ by injection of noise:

$$\begin{aligned}q(\mathbf{x}_t|\mathbf{x}_{t-1}) &= \mathcal{N}(\sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \cdot \mathbf{I}); \\q(\mathbf{x}_t|\mathbf{x}_0) &= \mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_t) \cdot \mathbf{I}).\end{aligned}$$

Reverse process refers to an intractable distribution that can be approximated by a normal distribution (with unknown parameters) for small β_t :

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx \mathcal{N}(\mu_{\theta,t}(\mathbf{x}_t), \sigma_{\theta,t}^2(\mathbf{x}_t))$$

Distribution Summary

Forward process maps any distribution $p_{\text{data}}(\mathbf{x})$ to $\mathcal{N}(0, \mathbf{I})$ by injection of noise:

$$\begin{aligned}q(\mathbf{x}_t|\mathbf{x}_{t-1}) &= \mathcal{N}(\sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \cdot \mathbf{I}); \\q(\mathbf{x}_t|\mathbf{x}_0) &= \mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_t) \cdot \mathbf{I}).\end{aligned}$$

Reverse process refers to an intractable distribution that can be approximated by a normal distribution (with unknown parameters) for small β_t :

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx \mathcal{N}(\boldsymbol{\mu}_{\theta,t}(\mathbf{x}_t), \boldsymbol{\sigma}_{\theta,t}^2(\mathbf{x}_t))$$

Conditioned reverse process is a normal distribution with known parameters, describing how to denoise a noisy image $\mathbf{x}_t$ when we know the clean image $\mathbf{x}_0$.

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\boldsymbol{\beta}}_t \cdot \mathbf{I})$$

Outline

1. Forward Gaussian Diffusion Process
2. Reverse Gaussian Diffusion Process
3. Gaussian Diffusion Model as VAE
4. Diffusion ELBO Derivation

Gaussian Diffusion Model as VAE

Let's treat $\mathbf{z} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$ as a latent variable (**note:** each $\mathbf{x}_t$ has the same dimension), and $\mathbf{x} = \mathbf{x}_0$ as the observed variable.

Latent Variable Model


$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$

Gaussian Diffusion Model as VAE

Let's treat $\mathbf{z} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$ as a latent variable (**note:** each $\mathbf{x}_t$ has the same dimension), and $\mathbf{x} = \mathbf{x}_0$ as the observed variable.

Latent Variable Model

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$




Gaussian Diffusion Model as VAE

Let's treat $\mathbf{z} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$ as a latent variable (**note:** each $\mathbf{x}_t$ has the same dimension), and $\mathbf{x} = \mathbf{x}_0$ as the observed variable.

Latent Variable Model

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$



Forward Diffusion

- Variational posterior distribution (encoder)

$$q(\mathbf{z}|\mathbf{x}) = q(\mathbf{x}_1, \dots, \mathbf{x}_T|\mathbf{x}_0) = \prod_{t=1}^T q(\mathbf{x}_t|\mathbf{x}_{t-1}).$$


- **Note:** there are no learnable parameters.

Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$


Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$



Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$




Reverse Diffusion

- Generative distribution (decoder)

$$p_{\theta}(\mathbf{x}|\mathbf{z}) = p_{\theta}(\mathbf{x}_0|\mathbf{x}_1).$$

Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$



Reverse Diffusion

- Generative distribution (decoder)


$$p_{\theta}(\mathbf{x}|\mathbf{z}) = p_{\theta}(\mathbf{x}_0|\mathbf{x}_1).$$

- Prior distribution

$$p_{\theta}(\mathbf{z}) = p_{\theta}(\mathbf{x}_1, \dots, \mathbf{x}_T) = \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t) \cdot p(\mathbf{x}_T).$$

Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$



Reverse Diffusion

- Generative distribution (decoder)

$$p_{\theta}(\mathbf{x}|\mathbf{z}) = p_{\theta}(\mathbf{x}_0|\mathbf{x}_1).$$

- Prior distribution

$$p_{\theta}(\mathbf{z}) = p_{\theta}(\mathbf{x}_1, \dots, \mathbf{x}_T) = \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t) \cdot p(\mathbf{x}_T).$$

Note: This differs from the vanilla VAE due to the simple decoder $p_{\theta}(\mathbf{x}|\mathbf{z})$ and the complex prior $p_{\theta}(\mathbf{z})$ (Markov chain).

Das A. An Introduction to Diffusion Probabilistic Models, blog post, 2021

Outline

1. Forward Gaussian Diffusion Process
2. Reverse Gaussian Diffusion Process
3. Gaussian Diffusion Model as VAE
4. Diffusion ELBO Derivation

ELBO for Gaussian Diffusion Model

Standard ELBO

$$\log p_{\theta}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\phi, \theta}(\mathbf{x}) \rightarrow \max_{\phi, \theta}$$

ELBO for Gaussian Diffusion Model

Standard ELBO

$$\log p_{\theta}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\phi, \theta}(\mathbf{x}) \rightarrow \max_{\phi, \theta}$$

Derivation

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p_{\theta}(\mathbf{x}_0, \mathbf{x}_{1:T})}{q(\mathbf{x}_{1:T}|\mathbf{x}_0)}$$

ELBO for Gaussian Diffusion Model

Standard ELBO

$$\log p_{\theta}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\phi, \theta}(\mathbf{x}) \rightarrow \max_{\phi, \theta}$$

Derivation

$$\begin{aligned} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p_{\theta}(\mathbf{x}_0, \mathbf{x}_{1:T})}{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t|\mathbf{x}_{t-1})} \end{aligned}$$

ELBO for Gaussian Diffusion Model

Standard ELBO

$$\log p_{\theta}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\theta}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\phi, \theta}(\mathbf{x}) \rightarrow \max_{\phi, \theta}$$

Derivation

$$\begin{aligned} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p_{\theta}(\mathbf{x}_0, \mathbf{x}_{1:T})}{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t|\mathbf{x}_{t-1})} \end{aligned}$$

- ▶ Let's try to decompose the ELBO into individual KL divergence terms.
- ▶ We need to replace $q(\mathbf{x}_t|\mathbf{x}_{t-1})$ with $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ in the denominator.
- ▶ Let's condition on $\mathbf{x}_0$ to make the reverse $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ tractable.

ELBO for Gaussian Diffusion Model

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) q(\mathbf{x}_t | \mathbf{x}_0)}{q(\mathbf{x}_{t-1} | \mathbf{x}_0)}$$

ELBO for Gaussian Diffusion Model

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) q(\mathbf{x}_t | \mathbf{x}_0)}{q(\mathbf{x}_{t-1} | \mathbf{x}_0)}$$

Derivation (continued)

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1})}$$

ELBO for Gaussian Diffusion Model

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) q(\mathbf{x}_t | \mathbf{x}_0)}{q(\mathbf{x}_{t-1} | \mathbf{x}_0)}$$

Derivation (continued)

$$\begin{aligned} \mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1})} \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0)} \end{aligned}$$

ELBO for Gaussian Diffusion Model

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) q(\mathbf{x}_t | \mathbf{x}_0)}{q(\mathbf{x}_{t-1} | \mathbf{x}_0)}$$

Derivation (continued)

$$\begin{aligned}\mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1})} \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0)} \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) p_{\theta}(\mathbf{x}_0 | \mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{q(\mathbf{x}_1 | \mathbf{x}_0) \prod_{t=2}^T q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0)}\end{aligned}$$

ELBO for Gaussian Diffusion Model

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) q(\mathbf{x}_t | \mathbf{x}_0)}{q(\mathbf{x}_{t-1} | \mathbf{x}_0)}$$

Derivation (continued)

$$\begin{aligned}\mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1})} \\&= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0)} \\&= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) p_{\theta}(\mathbf{x}_0 | \mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{q(\mathbf{x}_1 | \mathbf{x}_0) \prod_{t=2}^T q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0)} \\&= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) p_{\theta}(\mathbf{x}_0 | \mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{q(\mathbf{x}_1 | \mathbf{x}_0) \prod_{t=2}^T \frac{q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) q(\mathbf{x}_t | \mathbf{x}_0)}{q(\mathbf{x}_{t-1} | \mathbf{x}_0)}}\end{aligned}$$

ELBO for Gaussian Diffusion Model

$$q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) q(\mathbf{x}_t | \mathbf{x}_0)}{q(\mathbf{x}_{t-1} | \mathbf{x}_0)}$$

Derivation (continued)

$$\begin{aligned}\mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1})} \\&= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{\prod_{t=1}^T q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0)} \\&= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) p_{\theta}(\mathbf{x}_0 | \mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{q(\mathbf{x}_1 | \mathbf{x}_0) \prod_{t=2}^T q(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{x}_0)} \\&= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) p_{\theta}(\mathbf{x}_0 | \mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{q(\mathbf{x}_1 | \mathbf{x}_0) \prod_{t=2}^T \frac{q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0) q(\mathbf{x}_t | \mathbf{x}_0)}{q(\mathbf{x}_{t-1} | \mathbf{x}_0)}} \\&= \mathbb{E}_{q(\mathbf{x}_{1:T} | \mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) p_{\theta}(\mathbf{x}_0 | \mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1} | \mathbf{x}_t)}{q(\mathbf{x}_T | \mathbf{x}_0) \prod_{t=2}^T q(\mathbf{x}_{t-1} | \mathbf{x}_t, \mathbf{x}_0)}\end{aligned}$$

ELBO for Gaussian Diffusion Model

Derivation (continued)

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_T|\mathbf{x}_0) \prod_{t=2}^T q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)}$$

ELBO for Gaussian Diffusion Model

Derivation (continued)

$$\begin{aligned}\mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_T|\mathbf{x}_0) \prod_{t=2}^T q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)} = \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \left[\log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) + \log \frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^T \log \left(\frac{p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)} \right) \right]\end{aligned}$$

ELBO for Gaussian Diffusion Model

Derivation (continued)

$$\begin{aligned}\mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_T|\mathbf{x}_0) \prod_{t=2}^T q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)} = \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \left[\log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) + \log \frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^T \log \left(\frac{p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)} \right) \right] = \\ &= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) + \mathbb{E}_{q(\mathbf{x}_T|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \\ &\quad + \sum_{t=2}^T \mathbb{E}_{q(\mathbf{x}_{t-1}, \mathbf{x}_t|\mathbf{x}_0)} \log \left(\frac{p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)} \right)\end{aligned}$$

ELBO for Gaussian Diffusion Model

Derivation (continued)

$$\begin{aligned}\mathcal{L}_{\phi, \theta}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) \prod_{t=2}^T p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_T|\mathbf{x}_0) \prod_{t=2}^T q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)} = \\&= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \left[\log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) + \log \frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^T \log \left(\frac{p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)} \right) \right] = \\&= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) + \mathbb{E}_{q(\mathbf{x}_T|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \\&\quad + \sum_{t=2}^T \mathbb{E}_{q(\mathbf{x}_{t-1}, \mathbf{x}_t|\mathbf{x}_0)} \log \left(\frac{p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)} \right) = \\&= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) - D_{\text{KL}}(q(\mathbf{x}_T|\mathbf{x}_0) \| p(\mathbf{x}_T)) - \\&\quad - \sum_{t=2}^T \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\text{KL}}(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t))}_{\mathcal{L}_t}\end{aligned}$$

ELBO for Gaussian Diffusion Model

$$\begin{aligned}\mathcal{L}_{\phi, \theta}(\mathbf{x}) = & \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) - D_{\text{KL}}(q(\mathbf{x}_T|\mathbf{x}_0) \| p(\mathbf{x}_T)) - \\ & - \sum_{t=2}^T \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\text{KL}}(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t))}_{\mathcal{L}_t}\end{aligned}$$

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) - D_{\text{KL}}(q(\mathbf{x}_T|\mathbf{x}_0) \| p(\mathbf{x}_T)) - \underbrace{\sum_{t=2}^T \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\text{KL}}(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t))}_{\mathcal{L}_t}$$

- **First term** is the decoder distribution

$$\log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) = \log \mathcal{N}(\mathbf{x}_0 | \boldsymbol{\mu}_{\theta,1}(\mathbf{x}_1), \boldsymbol{\sigma}_{\theta,1}^2(\mathbf{x}_1)),$$

with $\mathbf{x}_1 \sim q(\mathbf{x}_1|\mathbf{x}_0)$.

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) - D_{\text{KL}}(q(\mathbf{x}_T|\mathbf{x}_0) \| p(\mathbf{x}_T)) - \underbrace{\sum_{t=2}^T \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\text{KL}}(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t))}_{\mathcal{L}_t}$$

- ▶ **First term** is the decoder distribution

$$\log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) = \log \mathcal{N}(\mathbf{x}_0 | \boldsymbol{\mu}_{\theta,1}(\mathbf{x}_1), \boldsymbol{\sigma}_{\theta,1}^2(\mathbf{x}_1)),$$

with $\mathbf{x}_1 \sim q(\mathbf{x}_1|\mathbf{x}_0)$.

- ▶ **Second term** is constant:
 - ▶ $p(\mathbf{x}_T) = \mathcal{N}(0, \mathbf{I})$;
 - ▶ $q(\mathbf{x}_T|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_T} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_T) \cdot \mathbf{I})$.

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_{\phi, \theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) - D_{\text{KL}}(q(\mathbf{x}_T|\mathbf{x}_0) \| p(\mathbf{x}_T)) - \underbrace{\sum_{t=2}^T \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\text{KL}}(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t))}_{\mathcal{L}_t}$$

- ▶ **First term** is the decoder distribution

$$\log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) = \log \mathcal{N}(\mathbf{x}_0 | \boldsymbol{\mu}_{\theta,1}(\mathbf{x}_1), \boldsymbol{\sigma}_{\theta,1}^2(\mathbf{x}_1)),$$

with $\mathbf{x}_1 \sim q(\mathbf{x}_1|\mathbf{x}_0)$.

- ▶ **Second term** is constant:
 - ▶ $p(\mathbf{x}_T) = \mathcal{N}(0, \mathbf{I})$;
 - ▶ $q(\mathbf{x}_T|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_T} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_T) \cdot \mathbf{I})$.
- ▶ **Third term** is the main contributor to the ELBO.

Summary

- ▶ The Gaussian diffusion process is a Markov chain that incrementally corrupts data with specific Gaussian noise.
- ▶ Denoising score matching, together with Langevin dynamics, can be applied to the Gaussian diffusion process.
- ▶ The reverse process reconstructs data from noise samples, although its precise form is intractable.
- ▶ We approximate the reverse process using normality assumptions.
- ▶ Gaussian diffusion model can be interpreted as a VAE with a hierarchy of latent variables.
- ▶ The ELBO for Gaussian diffusion model may be formulated as a sum over many KL divergence terms.