JOINS OF LOCALISATIONS IN THE LATTICE OF TWO-OUT-OF-THREE CLASSES: THE FINITE-ITERATION CRITERION FOR CLOSURE OPERATORS ON EXACT CATEGORIES, WITH APPLICATIONS TO TOPOSES, TORSION THEORIES AND FUZZY SETS

RYOTA SHIBUSAWA

2020 *Mathematics Subject Classification.* 18E08 (primary), 18F10, 18E40, 18B25, 06D22, 03E72 (secondary).

Key words and phrases. closure operator, exact category, Lawvere–Tierney topology, hereditary torsion theory, Gabriel localisation, nucleus, sheafification, two-out-of-three, fuzzy set, join of localisations, finite iteration.

Corresponding author. E-mail: `r-shibusawa@daiichi-koudai.ac.jp`.

Abstract. A closure operator c on the subobjects of an exact category, stable under pullback and preserving finite intersections, determines the class W_c of *c-bidense* maps, those whose image is c-dense and whose kernel pair lies in the closure of the diagonal. For a Lawvere–Tierney topology on a topos this is the class of maps inverted by sheafification; for a hereditary torsion theory on a Grothendieck abelian category, whose closure operators are exactly the hereditary torsion theories, it is the class of maps with torsion kernel and cokernel, inverted by the Gabriel quotient; for a nucleus on a frame viewed as a category it is the class of inequalities identified by the nucleus. These classes are elements of the complete lattice $\mathrm{Same}(\mathcal{C})$ of classes of morphisms closed under composition and two-out-of-three. We prove that for two closure operators c, c' the join $W_c \vee W_{c'}$ is the class of maps whose image and kernel pair are reached by a bounded number of alternating applications of c and c', and that every such map is a composite of maps of $W_c \cup W_{c'}$, so that the join is the closure of the union under composition alone. In a Grothendieck topos and in a Grothendieck abelian category the least closure operator $\bar{c}$ above c and c' is the transfinite iterate of cc', and $W_c \vee W_{c'} = W_{\bar{c}}$ if and only if $(cc')^k = \bar{c}$ for some k: for subtoposes this is the identity $(jj')^k = j \vee j'$ on the subobject classifier, for torsion theories T, T' it says that every object of the join torsion class has an alternating T'/T-filtration of bounded length. For persistence modules, representations of the ordinal ω, with the torsion classes of modules supported on the even and on the odd vertices, the join torsion class is everything but the interval modules need filtrations of unbounded length, so the join of the classes of maps that are isomorphisms at the odd vertices and at the even vertices is a proper class, the linear analogue of the known failure for sheaves on $\mathbb{N}$. In a frame viewed as a poset the witnesses that force uniformity are absent and the join is governed by the pointwise identity, as it is for isomorphisms up to a nucleus of fuzzy sets by a monomorphism version of the same argument; the paper thus places the joins of sheaf-equivalence classes, of torsion-theoretic localisations, of nuclei on a frame and of isomorphisms up to a nucleus of fuzzy sets under one theorem.

1. Introduction

Let $\mathcal{C}$ be a category. A *sameness* on $\mathcal{C}$ is a class of morphisms containing the identities and closed under composition and two-out-of-three; samenesses form a complete lattice $\mathrm{Same}(\mathcal{C})$ under inclusion, meets being intersections and the join of two classes the least sameness containing both [11]. Every localisation of $\mathcal{C}$, in particular every reflective subcategory, contributes the class W of maps it inverts. For a family of localisations closed under joins, the question studied in [13, 14, 20] is whether W sends the join of two localisations to the join of the two classes: whether every map inverted by the join localisation is obtained from maps inverted by the two localisations by finitely many compositions and cancellations. For subtoposes of a presheaf topos the answer was known to be yes on every finite

category [14] and no on the poset $\mathbb{N}$ [13], and [20] gave the exact criterion for subtoposes of a Grothendieck topos: $W_j \vee W_{j'} = W_{j\vee j'}$ iff $(jj')^k = j \vee j'$ for some k, an identity between endomorphisms of the subobject classifier. The same identity, $(j_2 j_1)^N = j_1 \vee j_2$ for nuclei on a Heyting algebra, is the finite-iteration axiom of the lax logic of the join of two nuclei [16], and it is the exact condition for the join of the classes of isomorphisms up to the two nuclei in the sameness lattice of fuzzy sets to be the class of the join nucleus [18], whereas in the sameness lattice of the Heyting algebra itself the pointwise identity suffices.

This paper proves one theorem behind these results and applies it to a setting not covered by them, the Gabriel localisations of abelian categories. The common structure is a *closure operator* c on the subobjects of an exact category $\mathcal{C}$: a pullback-stable, intersection-preserving closure on each $\mathrm{Sub}(X)$. It determines the sameness W_c of *c-bidense* maps, the maps $f\colon X \to Y$ with $c(\mathrm{im} f) = Y$ and $\ker f \subseteq c(\Delta_X)$. For two closure operators c, c' let U_k be the class of maps with

$$(cc')^k(\mathrm{im} f) = Y \qquad \text{and} \qquad \ker f \subseteq (cc')^k(\Delta_X),$$

the maps that are bidense *uniformly* for the k-fold alternation of the two closures.

Theorem 1.1. *Let c, c' be closure operators on an exact category $\mathcal{C}$. Then*

$$W_c \vee W_{c'} = \bigcup_k U_k$$

in $\mathrm{Same}(\mathcal{C})$*; U_0 is the class of isomorphisms, and for $k \geq 1$ every map in U_k is a composite of $2k$ regular epimorphisms followed by $2k$ monomorphisms lying alternately in $W_{c'}$ and W_c. In particular the join is the closure of $W_c \cup W_{c'}$ under composition.*

Theorem 1.2. *Let $\mathcal{C}$ be a Grothendieck topos or a Grothendieck abelian category, and c, c' closure operators on $\mathcal{C}$. Then the least closure operator $\bar{c}$ above c and c' is the transfinite iterate of cc', and $W_c \vee W_{c'} = W_{\bar{c}}$ if and only if $(cc')^k = \bar{c}$ for some k.*

Theorem 1.2 is proved under axioms (Section 4) that isolate what is used: chains of subobjects have pullback-stable unions, and the *fold map* of two copies of an object glued along a subobject S witnesses whether S is reached in k steps; the two classes of categories named in the statement are the ones in which we verify the axioms, by different arguments.

In a Grothendieck topos the closure operators are the Lawvere–Tierney topologies and W_c is the class of maps inverted by sheafification; this is the main theorem of [20], whose proofs are reproduced here in the generality in which they were written, and whose consequences for presheaf toposes, ideal topologies, locales and subspaces we summarise (Section 5.1, 5.2). In a Grothendieck abelian category $\mathcal{A}$ the closure operators are exactly the hereditary torsion theories T (Proposition 5.5): $c_T(A)$ is the preimage of

the torsion part of X/A, W_{c_T} is the class of maps with kernel and cokernel in T, the maps inverted by the quotient functor $\mathcal{A} \to \mathcal{A}/T$, and $\bar{c}$ is the closure operator of the join torsion theory $T \vee T'$. Theorems 1.1 and 1.2 read:

Corollary 1.3. *Let T, T' be hereditary torsion theories on a Grothendieck abelian category $\mathcal{A}$ and let E_k be the class of objects having a filtration $0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_{2k} = M$ with $M_{2i+1}/M_{2i} \in T'$ and $M_{2i+2}/M_{2i+1} \in T$. Then $W_T \vee W_{T'}$ is the class of maps whose kernel and cokernel lie in $\bigcup_k E_k$, the closure of $W_T \cup W_{T'}$ under composition; and $W_T \vee W_{T'} = W_{T \vee T'}$ iff $T \vee T' = E_k$ for some k, i.e. iff the objects of the join torsion class have alternating T'/T-filtrations of bounded length.*

For persistence modules, i.e. representations $\cdots \to M_2 \to M_1 \to M_0$ of the ordinal ω in vector spaces, with T and T' the modules supported on the even and on the odd vertices, W_T is the class of maps that are isomorphisms at the odd vertices, $T \vee T'$ is everything, and the interval module $k[0, N]$ needs an alternating filtration of length $N+2$; so $W_T \vee W_{T'}$ is a proper class, the class of maps whose kernel and cokernel are "bounded in alternation" (Example 5.6). This is the linear analogue of the failure for sheaves on $\mathbb{N}$ [13], and, as far as we know, the first statement of this kind about Gabriel localisations.

The frame L itself, viewed as a poset category, is an exact category in which every morphism is a monomorphism, with closure operators $c_n(x \leq y) = n(x) \wedge y$ for nuclei n; Theorem 1.1 applies and gives $W_n \vee W_{n'} = \{x \leq y : (nn')^k x \geq y \text{ for some } k\}$, so that the join is the class of the join nucleus iff the identity $(nn')^k x = (n \vee n')x$ holds *pointwise*; the witnesses of Theorem 1.2 do not exist in a poset, and indeed the uniform identity is not needed there (Section 5.3). Fuzzy sets are handled by a monomorphism version of Theorem 1.1 (Proposition 3.5), since the maps inverted by the reflection onto j-closed fuzzy sets are bijections; this recovers the description of their joins by bounded alternating chains [18] (Section 5.4).

The proofs are elementary and use only the calculus of closure operators on subobjects. The inclusion $U_k \subseteq W_c \vee W_{c'}$ is a peeling argument: a regular epimorphism whose kernel pair lies in the k-fold alternating closure of the diagonal is divided by the equivalence relation "kernel pair intersected with the c'-closure of the diagonal", which is a c'-bidense map, and the induced map needs one closure fewer; dually a monomorphism is enlarged to the c'-closure of its image. The failure direction of Theorem 1.2 uses, for a $\bar{c}$-dense subobject S of Y that is not reached in k steps, the fold map of two copies of Y glued along S; the coproduct of these maps over all k is $\bar{c}$-bidense but lies in no U_k. The criterion is uniform in the subobject, and it is the iteration of the closure operators, not of the reflectors, that matters: on a monoid of order 8 the alternating sheafification tower never stabilises although the join is preserved (Remark 5.2).

The quotient-closed regime is different. For subvarieties R, R' of a variety of algebras, $W_R \vee W_{R'} = W_{R \cap R'}$ always holds, because the composite of the two reflectors is the reflector onto $R \cap R'$, while the meet $W_R \cap W_{R'}$ is the class of the join variety only under a fracture condition [15]; the natural closure operators there are not pullback-stable, and we explain in Section 6 why no common mechanism exists.

What is known. Samenesses and the join theorem for a composite reflector are from [11, 12]; the meet theorem, the reduction lemma and the $\mathbb{N}$ counterexample from [13]; the finite case from [14]; the criterion for subtoposes of Grothendieck toposes, with the presheaf, ideal-type, object-generated, localic and subspace cases, from [20]. Closure operators on categories are the subject of [3]; universal closure operators on a topos, their correspondence with Lawvere–Tierney topologies and the bidense description of the maps inverted by sheafification are in [10, V.1, V.3], [4, A4.3–A4.4]. Hereditary torsion theories, Gabriel's quotient categories and the characterisation of the maps they invert are in [5, 19]; the correspondence between hereditary torsion theories and closure operators on module categories is discussed in [3], and its extension to normal and homological categories in [2]; the join of hereditary torsion theories and its description by transfinite extensions go back to [6]. Joins of nuclei as transfinite iterates are in [7]; the weak topologies of [9] compute the join of two Lawvere–Tierney topologies as such an iterate. The lax logic of the join of two nuclei and its finite-iteration axiom are in [16]; the sameness lattice of fuzzy sets and the uniform and pointwise criteria in [18]; the quotient-closed regime in [15].

What is new. The main new result is Corollary 1.3: for hereditary torsion theories T, T' on a Grothendieck abelian category, the join of the classes of maps inverted by the two Gabriel quotients is the class of maps whose kernel and cokernel have finite alternating T'/T-filtrations, it is generated by composition alone, and it is the class of the join torsion theory exactly when the lengths are bounded; Example 5.6 shows that the bound can fail even when the join torsion theory is the whole category. The description of $T \vee T'$ by transfinite alternating filtrations and the dictionary between closure operators and hereditary torsion theories (Proposition 5.5) are classical; the statement about the join of the inverted classes, and its failure, are new as far as we know. The general Theorems 1.1 and 1.2, in exact categories with closure operators and with the witness axioms verified in toposes and in abelian categories, are the tool; the topos-theoretic, localic, frame and fuzzy-set statements of [20, 18] are recovered as instances, with the pointwise/uniform difference explained by the presence or absence of witnesses (Section 5), and the quotient-closed regime is contrasted in Section 6.

2. Closure operators on exact categories

Throughout, $\mathcal{C}$ is an *exact* category in the sense of Barr [1]: it has finite limits, every morphism factors as a regular epimorphism followed by a monomorphism, regular epimorphisms are stable under pullback, and every equivalence relation is the kernel pair of its coequaliser. We write $\mathrm{im} f \subseteq Y$ for the image of $f\colon X \to Y$, $\ker f \subseteq X \times X$ for its kernel pair, $\Delta_X \subseteq X \times X$ for the diagonal, $g^{-1}(B)$ for the pullback of a subobject and $g(A)$ for the image of a subobject. We use freely the calculus of subobjects in a regular category: $\mathrm{im}(gh) = g(\mathrm{im} h)$; $\ker(gh) = (h \times h)^{-1}(\ker g)$; for a regular epimorphism e, $e(e^{-1}(B)) = B$; products of regular epimorphisms are regular epimorphisms; and for $g\colon Y \to Z$ and $A \subseteq Y$,

$$g^{-1}(g(A)) = \pi_1\big(\ker g \cap (Y \times A)\big),$$

as both are the image of $Y \times_Z A \to Y$.

Definition 2.1. A *closure operator* on $\mathcal{C}$ assigns to each object X a map $A \mapsto c(A)$ on $\mathrm{Sub}(X)$ that is inflationary, monotone, idempotent, preserves binary intersections, and is stable under pullback: $g^{-1}(c(B)) = c(g^{-1}(B))$ for every $g\colon X \to Y$ and $B \subseteq Y$. Closure operators are ordered pointwise. A subobject $A \subseteq X$ is *c-dense* if $c(A) = X$.

These are idempotent closure operators in the sense of [3], with the additional requirements that they preserve binary intersections and commute with pullbacks along all morphisms (the hereditary property of [3] concerns pullbacks along monomorphisms only). On a topos they are exactly the universal closure operators, which correspond to the Lawvere–Tierney topologies [10, V.1]; on a Grothendieck abelian category they are exactly the hereditary torsion theories (Proposition 5.5). A *word* is a composite $w = c_1 c_2 \cdots c_n$ of closure operators, c_n applied first; words are inflationary, monotone, pullback-stable and preserve binary intersections, but need not be idempotent.

Lemma 2.2. *Let w be a word, $g\colon X \to Y$, $A, K \subseteq X$, $B \subseteq Y$.*

(i) $g(w(A)) \subseteq w(g(A))$.
(ii) $w(A) \cap K \subseteq w(A \cap K)$.
(iii) $w(A \times B) = w(A) \times w(B)$ *in* $\mathrm{Sub}(X \times Y)$.
(iv) *For a closure operator c, $c(\Delta_X)$ is an equivalence relation on X.*

Proof. It suffices to treat a single closure operator c in (i)–(iii) and to iterate. (i) $A \subseteq g^{-1}(g(A))$, so $c(A) \subseteq c(g^{-1}(g(A))) = g^{-1}(c(g(A)))$. (ii) Let $m\colon K \hookrightarrow X$; then $A \cap K = m(m^{-1}(A))$ and $c(A) \cap K = m(m^{-1}(c(A))) = m(c_K(m^{-1}(A))) \subseteq c(m(m^{-1}(A)))$ by (i), where c_K is the closure on $\mathrm{Sub}(K)$, the pullback of c. (iii) $A \times B = \pi_1^{-1}(A) \cap \pi_2^{-1}(B)$, and c preserves intersections and pullbacks. (iv) Reflexivity and symmetry are clear (the twist map preserves Δ_X and commutes with c). The composite of the relation $c(\Delta_X)$ with itself is $\pi_{13}\big(\pi_{12}^{-1}c(\Delta_X) \cap \pi_{23}^{-1}c(\Delta_X)\big) = \pi_{13}\big(c(\pi_{12}^{-1}\Delta_X \cap \pi_{23}^{-1}\Delta_X)\big) \subseteq$

$c(\pi_{13}(\delta)) = c(\Delta_X)$ by (i), where $\delta \subseteq X^3$ is the image of the triple diagonal. □

Definition 2.3. A map $f\colon X \to Y$ is *c-bidense* if $\mathrm{im}f$ is c-dense in Y and $\ker f \subseteq c(\Delta_X)$. Write W_c for the class of c-bidense maps.

A monomorphism is c-bidense iff it is c-dense, and a regular epimorphism iff its kernel pair lies in $c(\Delta_X)$. On a topos with the closure operator of a Lawvere–Tierney topology j, W_c is the class of maps inverted by the sheafification a_j [4, A4.3]; in general W_c is defined by the closure operator alone, and Lemma 3.2 below (with $c' = c$) shows that it is a sameness.

3. Uniform bidensity and the join

Fix closure operators c, c' on $\mathcal{C}$ and write $c^{(k)} := (cc')^k$, a word of length $2k$ ending with c', $c^{(0)}$ being the identity; since closures are inflationary, $c^{(k)} \le c^{(k+1)}$ and $c, c' \le c^{(1)}$. Write $V := W_c \vee W_{c'}$.

Definition 3.1. For $k \ge 0$ let U_k be the class of maps $f\colon X \to Y$ with $c^{(k)}(\mathrm{im}f) = Y$ and $\ker f \subseteq c^{(k)}(\Delta_X)$. Put $U = \bigcup_k U_k$.

U_0 is the class of isomorphisms: $\mathrm{im}f = Y$ means that the monic part of f is an isomorphism, and $\ker f = \Delta_X$ that f is a monomorphism, so f is a monic regular epimorphism.

Lemma 3.2. *$U_k \subseteq U_{k+1}$, $W_c \cup W_{c'} \subseteq U_1$, and U is a sameness. Hence $V \subseteq U$. With $c' = c$ the same proof shows that W_c is a sameness.*

Proof. The first two assertions follow from $c^{(k)} \le c^{(k+1)}$ and $c, c' \le c^{(1)}$. Identities lie in U_0. Let $h\colon X \to Y$ and $g\colon Y \to Z$; we use Lemma 2.2 for the words $c^{(k)}$.

Composition. Let $h \in U_k$, $g \in U_l$. Then $\mathrm{im}(gh) = g(\mathrm{im}h)$ and $c^{(k+l)}(\mathrm{im}gh) \supseteq c^{(l)}\big(g(c^{(k)}(\mathrm{im}h))\big) = c^{(l)}(g(Y)) = c^{(l)}(\mathrm{im}g) = Z$ by (i). And $\ker(gh) = (h\times h)^{-1}(\ker g) \subseteq (h\times h)^{-1}(c^{(l)}(\Delta_Y)) = c^{(l)}((h\times h)^{-1}\Delta_Y) = c^{(l)}(\ker h) \subseteq c^{(l+k)}(\Delta_X)$.

Cancellation of g. Let $gh \in U_k$ and $g \in U_k$. Then $\ker h \subseteq \ker(gh) \subseteq c^{(k)}(\Delta_X)$. Let $B := g^{-1}(\mathrm{im}gh) \subseteq Y$; then $c^{(k)}(B) = g^{-1}(c^{(k)}(\mathrm{im}gh)) = g^{-1}(Z) = Y$. Moreover $B = \pi_1\big(\ker g \cap (Y \times \mathrm{im}h)\big)$, and

$$\begin{aligned}\pi_1\big(\ker g \cap (Y\times \mathrm{im}h)\big) &\subseteq \pi_1\big(c^{(k)}(\Delta_Y) \cap (Y\times \mathrm{im}h)\big)\\ &\subseteq \pi_1\big(c^{(k)}(\Delta_Y \cap (Y\times \mathrm{im}h))\big)\\ &\subseteq c^{(k)}\big(\pi_1(\Delta_Y \cap (Y\times \mathrm{im}h))\big) = c^{(k)}(\mathrm{im}h)\end{aligned}$$

by (ii) and (i). Hence $Y = c^{(k)}(B) \subseteq c^{(2k)}(\mathrm{im}h)$.

Cancellation of h. Let $gh \in U_k$ and $h \in U_k$. Then $\mathrm{im}g \supseteq \mathrm{im}(gh)$, so $c^{(k)}(\mathrm{im}g) = Z$. Since $h\times h\colon X\times X \to \mathrm{im}h \times \mathrm{im}h$ is a regular epimorphism, $\ker g \cap (\mathrm{im}h \times \mathrm{im}h) = (h\times h)(\ker gh) \subseteq (h\times h)(c^{(k)}(\Delta_X)) \subseteq c^{(k)}(\Delta_Y)$ by (i);

and $c^{(k)}(\mathrm{im}h \times \mathrm{im}h) = Y \times Y$ by (iii). So $\ker g = \ker g \cap c^{(k)}(\mathrm{im}h \times \mathrm{im}h) \subseteq c^{(k)}\big(\ker g \cap (\mathrm{im}h \times \mathrm{im}h)\big) \subseteq c^{(2k)}(\Delta_Y)$ by (ii).

For $c' = c$ one has $c^{(k)} = c$ for all $k \geq 1$, so $U_k = W_c$ and the argument shows that W_c is closed under composition and cancellation. □

Lemma 3.3 (peeling)**.** *Let $w = c_1 c_2 \cdots c_n$ be a word in closure operators.*

(a) *If $f\colon X \to Y$ is a regular epimorphism with $\ker f \subseteq w(\Delta_X)$, then $f = f_1 \circ \cdots \circ f_n \circ u$ with u an isomorphism and f_i a regular epimorphism in W_{c_i}.*

(b) *If $f\colon X \to Y$ is a monomorphism with $w(X) = Y$, then $f = f_1 \circ \cdots \circ f_n \circ u$ with u an isomorphism and f_i a monomorphism in W_{c_i}.*

For $n \geq 1$ the isomorphism u can be absorbed into f_n.

Proof. Induction on n; for $n = 0$ the hypothesis says that f is an isomorphism. Write $w = w'c_n$. (a) Let $R := \ker f \cap c_n(\Delta_X)$, an equivalence relation on X by Lemma 2.2(iv), and let $q\colon X \twoheadrightarrow X/R$ be its coequaliser, which has $\ker q = R$ by exactness; so $\ker q \subseteq c_n(\Delta_X)$ and $q \in W_{c_n}$. Since $R \subseteq \ker f$, f coequalises the two projections of R and $f = \bar{f} \circ q$ with $\bar{f}$ a regular epimorphism, and $\ker \bar{f} = (q \times q)(\ker f)$ because $q \times q$ is a regular epimorphism and $\ker f = (q \times q)^{-1}(\ker \bar{f})$. Now

$$\begin{aligned}(q \times q)(\ker f) &= (q \times q)\big(w'(c_n\Delta_X) \cap \ker f\big) \subseteq (q \times q)\big(w'(c_n\Delta_X \cap \ker f)\big) \\ &= (q \times q)(w'(R)) \subseteq w'\big((q \times q)(R)\big) = w'(\Delta_{X/R})\end{aligned}$$

by Lemma 2.2(ii) and (i), as $(q \times q)(\ker q) = \Delta_{X/R}$. Induction applied to $\bar{f}$ and w' gives the factorisation. (b) Regard X as a subobject of Y and let $Z := c_n(X) \subseteq Y$. The inclusion $X \hookrightarrow Z$ is c_n-dense (the closure of X in Z is $c_n(X) \cap Z = Z$ by pullback stability), hence in W_{c_n}, and $w'(Z) = w(X) = Y$. Induction applied to $Z \hookrightarrow Y$ and w' gives the factorisation. □

Proof of Theorem 1.1. Let C be the closure of $W_c \cup W_{c'}$ under composition. Then $C \subseteq V \subseteq U$ by Lemma 3.2. Let $f \in U_k$, $k \geq 1$, and factor $f = m \circ e$ with $e\colon X \twoheadrightarrow \mathrm{im}f$ a regular epimorphism and $m\colon \mathrm{im}f \hookrightarrow Y$ a monomorphism. Then $\ker e = \ker f \subseteq c^{(k)}(\Delta_X)$ and $c^{(k)}(\mathrm{im}f) = Y$, and $c^{(k)}$ is a word of length $2k$ ending with c'. By Lemma 3.3, e is a composite of $2k$ regular epimorphisms and m a composite of $2k$ monomorphisms, alternately in $W_{c'}$ and W_c, beginning with $W_{c'}$. So $U \subseteq C$, and $C = V = U$. □

Remark 3.4. In general the join of two samenesses requires the two-out-of-three saturation of the compositional closure [11]: on the poset $0 < 1 < 2$ viewed as a category, the classes $W_1 = \{0 \to 1\}$ and $W_2 = \{0 \to 2\}$ (with the identities) are samenesses, their compositional closure is $\{0 \to 1, 0 \to 2\}$, and two-out-of-three forces $1 \to 2$ into the join, which is therefore the class of all morphisms. For two bidense classes no cancellation is needed. The number of steps is controlled in both directions: a composite of n maps of $W_c \cup W_{c'}$ lies in U_n, and a map of U_k is a composite of $4k$ such maps.




FIGURE 1. Theorem 1.1: a map in U_k is $2k$ quotients by "kernel pair intersected with the closure of the diagonal" followed by $2k$ alternating closures of its image, each step bidense for one of the two closure operators.

Proposition 3.5 (monomorphisms)**.** *Let $\mathcal{C}$ be a category with finite limits and c, c' closure operators on $\mathcal{C}$ (pullback-stable, intersection-preserving closures on the subobject lattices); exactness is not assumed. For monomorphisms $m\colon X \hookrightarrow Y$ let M_k be the class of those with $c^{(k)}(X) = Y$ and $M = \bigcup_k M_k$. Then M contains the c-dense and the c'-dense monomorphisms, is closed under composition, and under cancellation among monomorphisms: if $gh \in M_k$ then $h \in M_k$ and $g \in M_k$; and every $m \in M_k$ is a composite of $2k$ monomorphisms alternately c'-dense and c-dense. Consequently, if B is a sameness consisting of monomorphisms such that for $m\colon X \hookrightarrow Y$ in B the inclusions $X \hookrightarrow c(X)$ and $X \hookrightarrow c'(X)$ are in B, then the join in* $\mathrm{Same}(\mathcal{C})$ *of the c-dense and the c'-dense members of B is $M \cap B$.*

Proof. For $h\colon X \hookrightarrow Y$ in M_k and $g\colon Y \hookrightarrow Z$ in M_l, pullback stability along $Y \hookrightarrow Z$ gives $c_Z^{(k)}(X) \cap Y = c_Y^{(k)}(X) = Y$, so $c_Z^{(k)}(X) \supseteq Y$ and $c_Z^{(k+l)}(X) \supseteq c_Z^{(l)}(Y) = Z$. If $gh \in M_k$, then $c_Y^{(k)}(X) = c_Z^{(k)}(X) \cap Y = Z \cap Y = Y$, so $h \in M_k$, and $c_Z^{(k)}(Y) \supseteq c_Z^{(k)}(X) = Z$, so $g \in M_k$; neither cancellation needs the other factor to be in M. The factorisation is Lemma 3.3(b), whose proof uses only pullback stability. For the last assertion, $M \cap B$ is a sameness containing the two classes, and each $m \in M_k \cap B$ is a composite of its peeling factors, which lie in B by hypothesis and by two-out-of-three in B. □

4. The finite-iteration criterion

In this section $\mathcal{C}$ satisfies

(H1) every $\mathrm{Sub}(X)$ is a set and has unions of chains, these are stable under pullback, and $A \cap \bigcup_\alpha B_\alpha = \bigcup_\alpha (A \cap B_\alpha)$ for chains (B_α).

Let c, c' be closure operators and define the transfinite iterates $c^{(0)}(A) = A$, $c^{(\alpha+1)}(A) = c^{(1)}(c^{(\alpha)}(A))$ and $c^{(\lambda)}(A) = \bigcup_{\alpha<\lambda} c^{(\alpha)}(A)$ for limit λ; they increase with α, are constant from some ordinal γ_A on since $\mathrm{Sub}(X)$ is a set, commute with pullback by (H1), and $c^{(\alpha)}(c^{(\beta)}(A)) = c^{(\beta+\alpha)}(A)$.

Lemma 4.1. *Under (H1), $\bar{c}(A) := \bigcup_\alpha c^{(\alpha)}(A)$ is a closure operator, the least one above c and c'.*

Proof. $\bar{c}$ is inflationary and monotone; $\bar{c}(A) = c^{(\gamma)}(A)$ for $\gamma = \gamma_A$ is a fixed point of $c^{(1)}$, hence of c and c' (as $c'(B) \subseteq c(c'(B)) = B$ for a fixed point B) and of $\bar{c}$; it commutes with pullback because each $c^{(\alpha)}$ does and pullback preserves unions of chains; and it preserves binary intersections because each $c^{(\alpha)}$ does and, by (H1), $\bigcup_\alpha c^{(\alpha)}(A) \cap \bigcup_\beta c^{(\beta)}(B) = \bigcup_{\alpha,\beta} c^{(\alpha)}(A) \cap c^{(\beta)}(B) = \bigcup_\gamma c^{(\gamma)}(A \cap B)$. It lies above c and c', and if d is a closure operator above c and c' then $c^{(\alpha)}(A) \subseteq d(A)$ for all α by induction, d being monotone and idempotent, with unions at limits. □

A map in U_k is $\bar{c}$-bidense, since $c^{(k)} \leq \bar{c}$; so $V \subseteq U \subseteq W_{\bar{c}}$. The converse inclusion $W_{\bar{c}} \subseteq U$ is what the criterion is about, and its failure is detected by *witnesses*.

Definition 4.2. A *witness* for a subobject $S \subseteq Y$ is a regular epimorphism $f_S \colon X_S \to Y$ such that, for every word w and for $w = \bar{c}$, $\ker f_S \subseteq w(\Delta_{X_S})$ iff $w(S) = Y$. A *sum* of a sequence of maps $f_k \colon X_k \to Y_k$ is a map $F \colon X \to Y$ with monomorphisms $\sigma_k \colon X_k \hookrightarrow X$, $\tau_k \colon Y_k \hookrightarrow Y$, $F\sigma_k = \tau_k f_k$, such that $(\sigma_k \times \sigma_k)^{-1}(\ker F) = \ker f_k$ and $\tau_k^{-1}(\mathrm{im} F) = \mathrm{im} f_k$ for every k, and such that F is $\bar{c}$-bidense whenever every f_k is.

Theorem 4.3. *Assume (H1), that every subobject has a witness, and that every sequence of witnesses has a sum. Then $W_c \vee W_{c'} = W_{\bar{c}}$ iff $c^{(k)} = \bar{c}$ for some k.*

Proof. If $c^{(k)} = \bar{c}$, a $\bar{c}$-bidense map lies in $U_k \subseteq V$, and $V \subseteq W_{\bar{c}}$ always. If $c^{(k)} \neq \bar{c}$ for every k, choose for each k a subobject $S_k \subseteq Y_k$ with $c^{(k)}(S_k) \neq \bar{c}(S_k)$; replacing Y_k by $\bar{c}(S_k)$ we may assume that S_k is $\bar{c}$-dense in Y_k but $c^{(k)}(S_k) \neq Y_k$, as $c^{(k)} \leq \bar{c}$ and both commute with pullback along $\bar{c}(S_k) \hookrightarrow Y_k$. Let f_k be a witness for S_k and F a sum of the f_k. Each f_k is $\bar{c}$-bidense, being a regular epimorphism with $\ker f_k \subseteq \bar{c}(\Delta)$, so $F \in W_{\bar{c}}$. But $F \notin U_k$ for every k: pulling back along σ_k and τ_k, the conditions $c^{(k)}(\mathrm{im} F) = Y$ and $\ker F \subseteq c^{(k)}(\Delta_X)$ restrict, by pullback stability of $c^{(k)}$ and the definition of a sum, to $c^{(k)}(\mathrm{im} f_k) = Y_k$ and $\ker f_k \subseteq c^{(k)}(\Delta_{X_k})$, i.e. to $f_k \in U_k$, which fails since $c^{(k)}(S_k) \neq Y_k$. So $F \in W_{\bar{c}} \setminus U = W_{\bar{c}} \setminus V$ by Theorem 1.1. □

It remains to produce witnesses and sums. The witness is always the *fold map* $f_S \colon Y \amalg_S Y \to Y$ of the pushout of $Y \hookleftarrow S \hookrightarrow Y$; what has to be checked is the description of its kernel pair, and this is where the two classes of categories differ.

Coherent categories with effective unions. Assume

(T) finite unions of subobjects exist, are stable under pullback and are *effective*: $A \cup B$ is the pushout of $A \hookleftarrow A \cap B \hookrightarrow B$; pushouts of monomorphisms along monomorphisms exist, are pullback squares and have monic injections; and countable coproducts exist and are disjoint and universal.

Every Grothendieck topos satisfies (T) [4, A1.4, A2.4, C2.2], and (H1) as well, subobject lattices being frames. The category of groups does not: pushouts of monomorphisms are amalgamated free products, which are pullbacks with monic injections, but the union of the two factors in $\mathbb{Z} * \mathbb{Z}$ is not the whole group, and coproducts are not universal.

Proposition 4.4. *Assume (T). For $S \subseteq Y$ the fold map $f_S \colon X_S := Y \amalg_S Y \to Y$ is a witness, and the coproduct $\coprod_k f_k$ of a sequence of witnesses is a sum.*

Proof. Let $\iota_1, \iota_2 \colon Y \hookrightarrow X_S$ be the injections, monomorphisms with intersection S, which is also the equaliser of ι_1, ι_2. The union $\iota_1 Y \cup \iota_2 Y \subseteq X_S$ is the pushout of $\iota_1 Y \hookleftarrow S \hookrightarrow \iota_2 Y$ by effectiveness, hence the inclusion $\iota_1 Y \cup \iota_2 Y \hookrightarrow X_S$ is an isomorphism: the two injections *cover* X_S. The map f_S is a regular epimorphism, being split by ι_1. Since unions are pullback-stable, subobject lattices are distributive and $X_S \times X_S = \bigcup_{a,b} \iota_a Y \times \iota_b Y$; hence $\ker f_S = \bigcup_{a,b} \ker f_S \cap (\iota_a Y \times \iota_b Y)$, and $\ker f_S \cap (\iota_a Y \times \iota_b Y) = (\iota_a \times \iota_b)\big((\iota_a \times \iota_b)^{-1} \ker f_S\big) = (\iota_a \times \iota_b)(\Delta_Y)$, as $f_S \iota_a = f_S \iota_b = \mathrm{id}_Y$. For $a = b$ this is contained in Δ_{X_S}; for $a \neq b$ it is the image of (ι_1, ι_2) or (ι_2, ι_1). Now $(\iota_1, \iota_2)^{-1}(\Delta_{X_S}) = S$, so by pullback stability $(\iota_1, \iota_2)^{-1}(w(\Delta_{X_S})) = w(S)$, and the image of (ι_1, ι_2) lies in $w(\Delta_{X_S})$ iff $w(S) = Y$; the same holds for (ι_2, ι_1) by symmetry. So $\ker f_S \subseteq w(\Delta_{X_S})$ iff $w(S) = Y$.

For the sum, let $F = \coprod_k f_k \colon \coprod X_k \to \coprod Y_k$ with the coproduct injections σ_k, τ_k, monomorphisms since coproducts are disjoint. Since coproducts are universal, $(\coprod X_k)^2 = \coprod_{k,l} X_k \times X_l$, the pullback of $\ker F$ along $\sigma_k \times \sigma_k$ is $\ker f_k$ and the pullback of $\mathrm{im} F$ along τ_k is $\mathrm{im} f_k$ (images are stable under pullback). If every f_k is $\bar{c}$-bidense then F is a regular epimorphism, and $\ker F$ is the union of the images of the $\ker f_k$ (countable unions being images of coproducts), which lie in the images of $\bar{c}(\Delta_{X_k})$, hence in $\bar{c}(\Delta_{\coprod X_k})$ by Lemma 2.2(i), as $\Delta_{\coprod X_k}$ is the union of the images of the Δ_{X_k}. □

Abelian categories. Let $\mathcal{A}$ be a Grothendieck abelian category; it is exact, satisfies (H1) (subobject lattices are sets, unions of chains are directed colimits, exact and pullback-stable by AB5, and distribute over finite intersections), and has countable coproducts, but these are not disjoint and pushouts of monomorphisms are not unions in the sense of (T). For a map $f \colon X \to Y$ write $d \colon X \times X \to X$ for $(x, x') \mapsto x - x'$, a split epimorphism with $d^{-1}(0) = \Delta_X$ and $d^{-1}(K) = \ker f$ for the kernel object $K = \mathrm{Ker}\, f \subseteq X$. For a word w, pullback stability gives $w(\Delta_X) = d^{-1}(w(0))$, and since d is a

split epimorphism, $d^{-1}(K) \subseteq d^{-1}(w(0))$ iff $K \subseteq w(0)$; so

$$(1)\quad \ker f \subseteq w(\Delta_X) \quad \text{iff} \quad \operatorname{Ker} f \subseteq w(0 \subseteq X) \quad \text{iff} \quad w(0 \subseteq \operatorname{Ker} f) = \operatorname{Ker} f,$$

the last step by pullback stability along $\operatorname{Ker} f \hookrightarrow X$. Call an object M *w-torsion* if $w(0 \subseteq M) = M$; by pullback stability along a quotient map, $w(S) = Y$ iff Y/S is w-torsion.

Proposition 4.5. *In a Grothendieck abelian category, for $S \subseteq Y$ the fold map $f_S \colon Y \amalg_S Y \to Y$ is a witness, and the direct sum $\bigoplus_k f_k$ of a sequence of witnesses is a sum.*

Proof. $X_S = (Y \oplus Y)/\{(s,-s) : s \in S\}$ and $f_S[(y_1,y_2)] = y_1 + y_2$ is an epimorphism with kernel object $\{[(y,-y)]\} \cong Y/S$. By (1), $\ker f_S \subseteq w(\Delta_{X_S})$ iff Y/S is w-torsion iff $w(S) = Y$. For the sum, $F = \bigoplus f_k$ has $\operatorname{Ker} F = \bigoplus \operatorname{Ker} f_k$ and $\operatorname{Coker} F = \bigoplus \operatorname{Coker} f_k$; the pullbacks of $\ker F$ and $\operatorname{im} F$ along the summand inclusions are $\ker f_k$ and $\operatorname{im} f_k$ by (1) and the corresponding statement for images. If every f_k is $\bar{c}$-bidense, then F is an epimorphism and $\operatorname{Ker} F = \bigoplus \operatorname{Ker} f_k$ is $\bar{c}$-torsion, because $\bar{c}(0 \subseteq \bigoplus K_k) \supseteq \bigoplus \bar{c}(0 \subseteq K_k)$ by Lemma 2.2(i) applied to the summand inclusions and the fact that unions of subobjects are sums; so $F \in W_{\bar{c}}$ by (1). □

Proof of Theorem 1.2. Theorem 4.3 with Lemma 4.1 and Proposition 4.4, respectively 4.5. □




FIGURE 2. The witness: two copies of Y glued along S; the kernel pair of the fold map lies in $w(\Delta)$ exactly when $w(S) = Y$. In a topos the two copies cover X_S; in an abelian category the kernel of the fold map is Y/S.

Remark 4.6. Theorem 1.1 needs only exactness, and Proposition 3.5 only finite limits. In Theorem 4.3, (H1) serves to identify $\bar{c}$ with the transfinite iterate, witnesses to detect a single failure and sums to combine infinitely many; without sums one still has, for each fixed k, that $W_{\bar{c}} \subseteq U_k$ iff $c^{(k)} = \bar{c}$, provided $\bar{c}$ exists and witnesses do.

5. Instances

5.1. **Toposes.** Let $\mathcal{E}$ be a Grothendieck topos. Closure operators on $\mathcal{E}$ are the universal closure operators, in bijection with Lawvere–Tierney topologies $j\colon \Omega \to \Omega$ (the closure of $A \subseteq X$ is classified by $j \circ \chi_A$) and with subtoposes, and W_{c_j} is the class of maps inverted by the sheafification a_j [10, V.1], [4, A4.3]. The hypotheses (H1) and (T) hold, and the identity $c^{(k)} = \bar{c}$ on all subobject lattices is the identity $(jj')^k = j \vee j'$ on Ω (apply both sides to the generic subobject). Theorems 1.1 and 1.2 specialise to [20, Thms. 1.1, 1.2]:

Corollary 5.1. *For Lawvere–Tierney topologies j, j' on a Grothendieck topos, $W_j \vee W_{j'}$ is the class of maps whose image and kernel pair are reached by a bounded number of alternating closures, the closure of $W_j \cup W_{j'}$ under composition, and $W_j \vee W_{j'} = W_{j\vee j'}$ iff $(jj')^k = j \vee j'$ for some k.*

For a presheaf topos $\mathrm{Presh}(S)$ and Grothendieck topologies J, J' on the small category S the conditions are conditions on sieves: a map f is in U_k iff its image sieves $\{b : y \cdot b \in \mathrm{im}f\}$ and its equaliser sieves $\{b : x \cdot b = x' \cdot b\}$, $f(x) = f(x')$, are sent to the maximal sieve by $(jj')^k$. For topologies of ideal type, J_D and J_E for idempotent two-sided ideals D, E of S, this is local bijectivity along the ideal $(DE)^k$, and the criterion is the stabilisation of the chain $DE \supseteq DEDE \supseteq \cdots$, which holds on every finite category and fails on $(\mathbb{N}, \leq)$ for the ideals of the even and of the odd points [20, §5]. For ideals generated by two sets of objects A, B, the join of "bijective at the objects of A" and "bijective at the objects of B" is "bijective at $A \cap B$" iff every morphism through a long enough alternating zigzag of B and A factors through $A \cap B$; on $\mathbb{N}$ with the even and the odd points the join is the class of maps of towers bijective up to a uniform shift, a proper subclass of the isomorphisms of pro-sets [20, §6].

Remark 5.2. The classical sufficient condition for $W_j \vee W_{j'} = W_{j\vee j'}$ is that the alternating tower of sheafifications $(a_{j'}a_j)^k X$ reach the $(j \vee j')$-sheaves for one k and all X [12, Thm. 4.1]. The criterion of Theorem 1.2 is the iteration of the closure operators, and the two differ: on the contracted monoid of order 8 of [14, §7] the chain stabilises, so the join is preserved, while the strictly alternating plus tower never reaches the sheaves uniformly [14, Prop. 7.1], and a computation on the right M-sets with at most three elements shows that no composite of at most eight plus constructions, in particular no $(a_E a_D)^k$ with $k \leq 2$, is a sheaf for all of them [20, §8].

5.2. **Locales.** Let L be a frame and $\mathrm{Sh}(L)$ its topos of sheaves. Closure operators on $\mathrm{Sh}(L)$ correspond to the nuclei on L (sublocales), ordered as the nuclei [8, II.2], [4, A4.5]; the subobjects of the representable $y(U)$ are the $V \leq U$; and for the nucleus n with sublocale inclusion i, the sheafification is i_*i^* with $i^*(y(V)) = y(n(V))$, so $V \hookrightarrow W$ is dense iff $n(V) = n(W)$, and the closure of V in $y(U)$ is $n(V) \wedge U$, the largest $W \leq U$ with $n(W) = n(V)$. By

induction, using $U \le n(U), n'(U)$, $(cc')^k(V) = (nn')^k(V) \wedge U$ and $\bar{c}(V) = (n \vee n')(V) \wedge U$ [20, Lemma 7.1].

Corollary 5.3. *For nuclei n, n' on a frame L, $W_n \vee W_{n'} = W_{n\vee n'}$ in* $\mathrm{Same}(\mathrm{Sh}(L))$ *iff $(nn')^k = n \vee n'$ in the frame of nuclei for some k.*

Proof. Theorem 1.2 and the formula above with $U = \top$. □

Open and closed nuclei satisfy the condition with $k = 1$: $o(U)c(U')$, $V \mapsto U \to (U' \vee V)$, is idempotent and hence the join. For a topological space X and subspaces A, B, with nuclei $n_A(V) = \mathrm{int}(V \cup (X \setminus A))$, the class W_{n_A} consists of the maps of sheaves whose inverse image to A is an isomorphism, and the condition reads: the alternating derivation $T(F) = \mathrm{cl}(\mathrm{cl}(F \cap B) \cap A)$ on closed sets reaches its transfinite limit in a uniform number of steps [20, Prop. 7.5]. It holds for closed and for open subspaces with $k = 1$ and fails for the rationals and the irrationals in $\mathbb{R}$: for a countable closed set C whose points of even Cantor–Bendixson rank are rational and of odd rank irrational, $T^j(C)$ is the $2j$-th derived set, nonempty for $2j$ below the rank, while the transfinite limit is empty; so in $\mathrm{Same}(\mathrm{Sh}(\mathbb{R}))$ the join of "isomorphism over $\mathbb{Q}$" and "isomorphism over the irrationals" is strictly smaller than the class of the meet subtopos, although the two sublocales meet in a nonempty dense sublocale [20, Ex. 7.6].

5.3. **Frames as categories: the pointwise criterion.** Let L be a frame, or just a meet-semilattice with top, viewed as a category. It is exact: every morphism $x \le y$ is a monomorphism, the only regular epimorphisms are the identities, and the (regular epi, mono) factorisation of $x \le y$ is $x \le y$ itself; kernel pairs are diagonals and equivalence relations trivial. $\mathrm{Sub}(y) = {\downarrow}y$, and the pullback of $z \le y$ along $x \le y$ is $z \wedge x$. For a nucleus n on L put $c_n(z \le y) := n(z) \wedge y$; this is a closure operator: inflationary, monotone, idempotent $(n(n(z) \wedge y) \wedge y = n(z) \wedge n(y) \wedge y = n(z) \wedge y)$, intersection-preserving, and pullback-stable $(c_n(z \wedge x) = n(z) \wedge n(x) \wedge x = c_n(z) \wedge x)$. The c_n-bidense maps are the $x \le y$ with $n(x) \wedge y = y$, i.e. $n(x) \ge y$, i.e. $n(x) = n(y)$: W_{c_n} is the class $W_n^L = \{x \le y : nx = ny\}$ of [18, Thm. 1.1].

Corollary 5.4. *For nuclei n, n' on L, in* $\mathrm{Same}(L)$*,*

$$W_n^L \vee W_{n'}^L = \{x \le y : \ (nn')^k x \ge y \text{ for some } k\},$$

every such $x \le y$ being the composite of the chain $x \le n'x \wedge y \le nn'x \wedge y \le \cdots \le y$ of inequalities alternately in $W_{n'}^L$ and W_n^L; and $W_n^L \vee W_{n'}^L = W_{n\vee n'}^L$ iff for every x some iterate $(nn')^k x$ equals $(n \vee n')x$.

Proof. Theorem 1.1: $x \le y$ is in U_k iff $c^{(k)}(x) = (nn')^k x \wedge y = y$, and the peeling factorisation is the displayed chain. For the last assertion, $W_{n\vee n'}^L = \{x \le y : (n \vee n')x \ge y\}$; if the pointwise identity holds, then $(n \vee n')x \ge y$ gives $(nn')^k x \ge y$ for the k of x; conversely take $y = (n \vee n')x$. □

Here Theorem 4.3 does not apply: in a poset the pushout of $y \hookleftarrow s \hookrightarrow y$ is y with both injections the identity, its fold map is id_y, and no witnesses

exist (the only regular epimorphisms are identities, whose kernel pair is the diagonal). Accordingly the criterion is pointwise, as in [18, Thm. 1.1], and the examples there of nuclei on the Gödel chain $[0,1]$ that stabilise pointwise but not uniformly separate $\mathrm{Same}(L)$ from $\mathrm{Same}(\mathrm{Sh}(L))$: with $F = \{0,1\} \cup \{1/2k\}$ and $G = \{0,1\} \cup \{1/(2k+1)\}$ as fixed sets, $W_{n_F} \vee W_{n_G} = W_{n_F \vee n_G}$ in $\mathrm{Same}(L)$ by Corollary 5.4 but not in $\mathrm{Same}(\mathrm{Sh}([0,1]))$ by Corollary 5.3 [20, Ex. 7.4].

5.4. **Fuzzy sets.** Let L be a complete lattice and $\mathbf{Set}(L)$ Goguen's category of L-fuzzy sets: objects (X,μ) with $\mu\colon X \to L$, morphisms $f\colon (X,\mu) \to (Y,\nu)$ maps with $\mu \le \nu f$. It has finite limits (computed on underlying sets with the initial structure) but is not exact: for the three-element chain $L = \{0 < a < 1\}$ and $X = \{x_0, x_1\}$ with $\mu \equiv 1$, the relation on $X \times X$ with degree 1 on the diagonal and a off it is an internal equivalence relation that is the kernel pair of no map, as kernel pairs of maps out of (X,μ) have degree 1 wherever they are nonempty. For a nucleus j on L preserving binary meets, the j-closed fuzzy sets ($\mu = j\mu$) form a reflective subcategory with reflector $(X,\mu) \mapsto (X, j\mu)$, and the class W_j of maps inverted by it consists of the bijections f with $j\mu = j\nu f$ [18]. Subobjects of (Y,ν) are the (A,α) with $A \subseteq Y$ and $\alpha \le \nu|_A$, pulled back along f to $(f^{-1}A, \alpha f \wedge \mu)$; put

$$c_j(A,\alpha) := (Y,\ j\bar{\alpha} \wedge \nu),$$

where $\bar{\alpha}$ extends α by the bottom element outside A. This is inflationary, monotone, idempotent, preserves binary intersections (as j preserves binary meets) and is pullback-stable: both $f^{-1}(c_j(A,\alpha))$ and $c_j(f^{-1}(A,\alpha))$ are $(X, j(\bar{\alpha}f) \wedge \mu)$, using $\mu \le \nu f$ and $\mu \le j\mu$. A bijection $f\colon (X,\mu) \to (Y,\nu)$ is a monomorphism with image $(Y, \mu f^{-1})$, so it is c_j-dense iff $j(\mu f^{-1}) \ge \nu$ iff $j\mu = j\nu f$: on bijections, c_j-density is membership in W_j, and $f \in M_k$ iff $(jj')^k\mu \ge \nu f$. Bijections form a sameness B, and for a bijection f the inclusions $\mathrm{im}f \hookrightarrow c_j(\mathrm{im}f)$ are bijections, since $c_j(\mathrm{im}f)$ has underlying set Y. Hence Proposition 3.5 gives

$$W_j \vee W_{j'} = \{f \text{ bijective} :\ (jj')^k\mu \ge \nu f \text{ for some } k\},$$

the description of [18, Lemma 4.1, Thm. 4.3] by uniformly bounded alternating chains, each such f being a composite of bijections alternately in $W_{j'}$ and W_j. The uniform criterion $(j'j)^k = j \vee j'$ of [18, Thm. 1.1(ii)] is proved there with witnesses that glue fuzzy sets along a common part and sums of such witnesses, in the pattern of Theorem 4.3; we do not claim $\mathbf{Set}(L)$ as an instance of Theorem 1.2. (For non-bijective maps the class of c_j-bidense maps is larger than W_j: the fold map of two points of degree $\bot$ onto one is c_j-bidense for every j. The join of two classes of bijections is computed among bijections, so this does not affect the statement.)

5.5. **Abelian categories and torsion theories.** Let $\mathcal{A}$ be a Grothendieck abelian category. A *hereditary torsion theory* is given by a class T of objects closed under subobjects, quotients, extensions and coproducts; equivalently

by its radical $t\colon \mathcal{A} \to \mathcal{A}$, the subfunctor of the identity with $t(M)$ the largest subobject of M in T, which is left exact ($t(N) = N \cap t(M)$ for $N \subseteq M$) and idempotent ($t(M/t(M)) = 0$) [5, 19].

Proposition 5.5. *Closure operators on $\mathcal{A}$ correspond bijectively to hereditary torsion theories (this is classical, cf.* [3, 19]*; we include the proof to fix the dictionary): T gives $c_T(A) := \pi^{-1}(t(X/A))$ for $A \subseteq X$ with $\pi\colon X \to X/A$, and c gives $T_c = \{M : c(0 \subseteq M) = M\}$ with radical $t_c(M) = c(0 \subseteq M)$. Under this correspondence $c_T \leq c_{T'}$ iff $T \subseteq T'$, W_{c_T} is the class of maps with kernel and cokernel in T, the maps inverted by the quotient functor $\mathcal{A} \to \mathcal{A}/T$, and the least closure operator above c_T and $c_{T'}$ is $c_{T \vee T'}$ for the least hereditary torsion theory $T \vee T'$ containing T and T'.*

Proof. c_T is inflationary, monotone (the image of $t(X/A)$ in X/B lies in $t(X/B)$ for $A \subseteq B$) and idempotent ($X/c_T(A) = (X/A)/t(X/A)$ is torsion-free). It preserves intersections: $X/(A \cap B)$ embeds in $X/A \oplus X/B$, and as t is left exact and additive, $x + (A \cap B)$ is torsion iff $x + A$ and $x + B$ are. It is pullback-stable: for $f\colon X' \to X$, $X'/f^{-1}(A)$ embeds in X/A, so $t(X'/f^{-1}(A))$ is the preimage of $t(X/A)$ by left exactness, i.e. $c_T(f^{-1}A) = f^{-1}(c_T A)$. Conversely, for a closure operator c, $t_c(M) = c(0)$ is a subfunctor (by Lemma 2.2(i)), left exact (pullback stability along monomorphisms), and idempotent ($c(0 \subseteq M/c(0))$ pulls back to $c(c(0)) = c(0)$), so T_c is a hereditary torsion theory, and $c(A) = \pi^{-1}(c(0 \subseteq X/A)) = c_{T_c}(A)$ by pullback stability along π; and $t_{c_T} = t$. Order: $c_T \leq c_{T'}$ iff $t \leq t'$ iff $T \subseteq T'$. A map $f\colon X \to Y$ has $c_T(\mathrm{im} f) = Y$ iff $Y/\mathrm{im} f = \mathrm{Coker}\, f$ is torsion, and $\ker f \subseteq c_T(\Delta_X)$ iff $\mathrm{Ker}\, f \subseteq t(X)$ iff $\mathrm{Ker}\, f \in T$ by (1); these are the maps inverted by the quotient functor [5, III]. The last assertion follows from the order isomorphism. □

For a word w in $c_T, c_{T'}$ the w-torsion objects are those with a filtration by the canonical subobjects $0 \subseteq t'(M) \subseteq \pi^{-1}t(M/t'(M)) \subseteq \cdots$; so an object is $c^{(k)}$-torsion iff it lies in the class E_k of Corollary 1.3, any alternating filtration being dominated by the canonical one, and E_k is closed under subobjects and quotients. By (1) and pullback stability along $Y \to Y/\mathrm{im} f$, a map is in U_k iff its kernel and cokernel lie in E_k.

Proof of Corollary 1.3. Theorems 1.1 and 1.2 with Proposition 5.5 and the description of U_k; the identity $c^{(k)} = \bar{c}$ says that every $c_{T \vee T'}$-torsion object, i.e. every object of $T \vee T'$, is $c^{(k)}$-torsion. □

By Lemma 4.1, $T \vee T'$ is the class of objects with a transfinite alternating filtration, the classical description of the join of two hereditary torsion theories; the corollary says that the join of the two classes of maps inverted by the Gabriel quotients is the class of the join torsion theory exactly when a finite bound on the length exists. When T and T' commute in the sense that $t(t'(M)) = t'(t(M))$ and the join is the class of extensions of a T-object by a T'-object, $k = 1$: for instance the p-torsion and the q-torsion abelian groups for distinct primes.

Example 5.6. Let $\mathcal{A}$ be the category of persistence modules over a field k: representations $\cdots \to M_2 \to M_1 \to M_0$ of the ordinal ω with arrows $n+1 \to n$, a Grothendieck abelian category. Let T and T' be the classes of modules supported on the even, respectively on the odd, vertices; each is closed under subobjects, quotients, extensions and direct sums, hence a hereditary torsion theory, with $t(M)_n = 0$ for n odd and $t(M)_n = \mathrm{Ker}(M_n \to M_{n-1})$ for n even ($t(M)_0 = M_0$), and symmetrically t'. A map f has kernel and cokernel in T iff f_n is an isomorphism for every odd n, so W_T is the class of maps that are isomorphisms at the odd vertices and $W_{T'}$ at the even vertices, the linear analogue of the classes of [13, Thm. 7.1]. The join torsion theory is everything: a module that is torsion-free for both has $M_0 = 0$ and all maps $M_n \to M_{n-1}$ injective, hence is 0. The interval module $I = k[0, N]$, with $I_n = k$ for $n \leq N$, identities as maps and $I_n = 0$ for $n > N$, has $t'(I) = 0$, $t(I) = k[0, 0]$, then $t'(I/k[0, 0]) = k[1, 1]$, and so on: its canonical alternating filtration is $0 \subseteq 0 \subseteq k[0, 0] \subseteq k[0, 1] \subseteq \cdots \subseteq k[0, N]$, of length $N + 2$, and $I \in E_k$ iff $2k \geq N + 2$. So $E_k \neq T \vee T'$ for every k, and by Corollary 1.3 the join of "isomorphism at the odd vertices" and "isomorphism at the even vertices" is a proper class of maps, although the join localisation is the zero category and $W_{T\vee T'}$ is the class of all maps; it is the class of maps whose kernel and cokernel have alternating filtrations of bounded length, for instance it contains $0 \to k[0, N]$ for each N but not $0 \to \bigoplus_N k[0, N]$. The witness of Proposition 4.5 for $S = 0 \subseteq k[0, N]$ is the fold map $k[0, N] \oplus k[0, N] \to k[0, N]$, in $W_{T\vee T'}$ but not in U_k for $2k < N+2$, and the direct sum of these fold maps over N lies in no U_k.

5.6. **Summary.** The settings are instances of one theorem, and the identity $(cc')^k = \bar{c}$ that governs them is, for nuclei, the finite-iteration axiom of the lax logic of the join of two nuclei [16]: uniform in the subobject in toposes, locales, abelian categories and fuzzy sets, where witnesses and sums exist, and pointwise in the frame itself, where they do not. The pointwise condition is also the one that characterises $\Box_C \vee \Box_{C'} = \Box_{C\cap C'}$ for relativized necessities on the frame of up-sets of a poset (absence of infinite alternating chains [17]), and a disjoint union of finite alternating chains of unbounded length satisfies it but not the uniform one.

6. The dual regime: quotient-closed reflections

Let $\mathcal{V}$ be a variety of algebras and $R \subseteq \mathcal{V}$ a subvariety, with reflector $a_R(A) = A/\rho_R(A)$, $\rho_R(A)$ the least congruence with quotient in R. A homomorphism $f\colon A \to B$ is inverted by a_R iff $\mathrm{im}f$ meets every $\rho_R(B)$-class and $(f \times f)^{-1}(\rho_R(B)) \subseteq \rho_R(A)$. This looks like bidensity, but the candidate closure operators are not pullback-stable: for R the abelian groups, $\rho_R(A)$ is the commutator congruence, and the inclusion $A_3 \hookrightarrow S_3$ has $\rho_R(A_3)$ trivial while $\rho_R(S_3)$ restricted to A_3 is everything; likewise the $\rho_R(B)$-saturation of a subalgebra is not stable under intersection with subalgebras (in S_3, with $K = \{e, (13)\}$ and $A = \{e, (12)\}$, the saturation of A contains (13) but $A\cap K$

is trivial). Accordingly the mechanism is different: because R is closed under quotients, the composite $a_{R'}a_R$ is already the reflector onto $R \cap R'$, and the join theorem of [12, Thm. 4.1] gives $W_R \vee W_{R'} = W_{R\cap R'}$ for all subvarieties, with no condition [15, Thm. 1.1]: at the level of reflectors, $k = 1$. The hard direction there is the meet, $W_R \cap W_{R'} = W_{R\vee R'}$, which holds iff a fracture condition is satisfied (subdirect representation and absence of dense proper subalgebras [15, Thm. 1.2]). The two regimes are thus dual: for closure-type reflections (subtoposes) the meet of classes is automatic and the join is governed by finite iteration of closures; for quotient-closed reflections (subvarieties) the join is automatic and the meet is governed by fracture. Remark 5.2 shows that in the closure-type regime the reflector-level condition ($k = 1$ or any uniform k for the tower of reflectors) is not the right one.

7. Questions

Question 7.1. Theorem 1.2 reduces join preservation to the identity $(cc')^k = \bar{c}$. For subspaces this is the uniform termination of the alternating derivation on closed sets, for ideal topologies the stabilisation of the ideal chain. Is there a structural description of the pairs of nuclei, or of subtoposes, satisfying it, for instance in terms of the lattice of sublocales?

Question 7.2. Does Theorem 1.2 hold in an elementary topos, where (H1) and the coproducts in (T) may be unavailable? By Remark 4.6 the obstacle is the existence of $\bar{c}$ and the combination of infinitely many witnesses. Likewise, which exact categories beyond toposes and abelian categories have witnesses and sums?

Question 7.3. Is there a common generalisation of the closure-type and the quotient-closed regimes, for reflective subcategories of an exact category whose reflector is determined by a closure operator that is only *weakly* hereditary in the sense of [3]?

Funding

No funding was received for this work.

Declarations

The author has no competing interests to declare.

References

[1] M. Barr, *Exact categories*, in: Exact Categories and Categories of Sheaves, Lecture Notes in Math. 236, Springer, 1971, 1–120.
[2] M. M. Clementino, D. Dikranjan, W. Tholen, *Torsion theories and radicals in normal categories*, J. Algebra **305** (2006), 98–129.
[3] D. Dikranjan, W. Tholen, *Categorical Structure of Closure Operators*, Mathematics and its Applications 346, Kluwer, 1995.

[4] P. T. Johnstone, *Sketches of an Elephant: A Topos Theory Compendium*, Oxford Logic Guides 43–44, Oxford University Press, 2002.
[5] P. Gabriel, *Des catégories abéliennes*, Bull. Soc. Math. France **90** (1962), 323–448.
[6] S. E. Dickson, *A torsion theory for abelian categories*, Trans. Amer. Math. Soc. **121** (1966), 223–235.
[7] M. H. Escardó, *Joins in the frame of nuclei*, Appl. Categ. Structures **11** (2003), 117–124.
[8] P. T. Johnstone, *Stone Spaces*, Cambridge Studies in Advanced Mathematics 3, Cambridge University Press, 1982.
[9] Z. Khanjanzadeh, A. Madanshekaf, *Weak topologies on toposes*, Bull. Iranian Math. Soc. **47** (2021), 461–486; arXiv:1808.06104.
[10] S. Mac Lane, I. Moerdijk, *Sheaves in Geometry and Logic*, Springer, 1992.
[11] R. Shibusawa, *The sameness lattice of a category: weak-equivalence structures, Galois functoriality, and comparison intervals*, preprint (2026), archived at DOI 10.5281/zenodo.22823772.
[12] R. Shibusawa, *Postnikov truncations as joins of sheaf and homotopy equivalences*, preprint (2026), archived at DOI 10.5281/zenodo.22956181.
[13] R. Shibusawa, *Meets and joins of Grothendieck topologies in the sameness lattice*, preprint (2026), archived at DOI 10.5281/zenodo.22978446.
[14] R. Shibusawa, *Grothendieck topologies on a finite category form a sublattice of the lattice of two-out-of-three classes*, preprint (2026), archived at DOI 10.5281/zenodo.23204943.
[15] R. Shibusawa, *Lattices of subvarieties and of logics as sublattices of the lattice of two-out-of-three classes*, preprint (2026), archived at DOI 10.5281/zenodo.23407364.
[16] R. Shibusawa, *The join of two nuclei as a lax modality: axiomatization, finite models, and inexpressibility*, preprint (2026), archived at DOI 10.5281/zenodo.23330576.
[17] R. Shibusawa, *Joins of relativized necessities on a poset: an alternating-chain criterion and the induction scheme of lax logic*, preprint (2026), archived at DOI 10.5281/zenodo.23411203.
[18] R. Shibusawa, *Nuclei on the algebra of truth values and the sameness lattice of fuzzy sets: meets, joins and the finite-iteration axiom*, preprint (2026), archived at DOI 10.5281/zenodo.23413073.
[19] B. Stenström, *Rings of Quotients*, Grundlehren der mathematischen Wissenschaften 217, Springer, 1975.
[20] R. Shibusawa, *Joins of Lawvere–Tierney topologies in the lattice of two-out-of-three classes: the finite-iteration criterion*, preprint (2026), archived at DOI 10.5281/zenodo.23487202.

Daiichi Institute of Technology
Email address: r-shibusawa@daiichi-koudai.ac.jp