

Mankiewicz Extension Theorem

An isometry equivalence from an open connected subset onto an open subset of real normed spaces extends uniquely to an affine isometry equivalence of the ambient spaces. The selected route also reaches convex-domain and ball specializations.

Exact source: MathlibAnnex v0.4.0 · 11 declaration tiles · 9 levels

Local distance information becomes a rigid affine structure. This Project exposes the reusable chain from reflection and midpoint geometry to global extension.

Real normed spaces and an isometry equivalence. The principal theorem requires an open connected source and an open target. The convex specialization requires a convex source set with nonempty ambient interior and exact preservation of ambient-interior membership; the target set is not separately assumed convex. Exact assumptions, including radius conditions for balls, remain in the linked exact declarations.

The eleven declaration explanations are reviewed for source-exposition correspondence. This does not claim a line-by-line human proof audit. Boundary Inputs include Lean Core, Mathlib, and omitted MathlibAnnex support declarations.

Eleven reviewed canonical Declaration Cards remain independent records. Project relations navigate within this page.

[Exact Project source](#)

[Exact source manifest](#)

Proof architecture

Reflection to midpoint rigidity — A bounded symmetric lens is reflection-invariant about the midpoint. Center rigidity sends that midpoint to the target midpoint, first on lenses and then on contained balls.

Local affinity through omitted support lemmas — Midpoint preservation yields line-map preservation. Radial-map support declarations, including private technical lemmas, connect this result to a local affine isometric extension. The path witnesses retain these declarations even though they are outside the selected Card scope.

Global assembly and specializations — Local extensions agree and assemble over an open connected domain. A convex source with nonempty ambient interior, together with exact preservation of ambient-interior membership, yields the convex specialization; open- and closed-ball corollaries follow under their exact radius conditions. The closed-ball route also uses reflection rigidity.

Boundary Inputs

961 exact Boundary Inputs: EXTERNAL_COMPILED_DECLARATION 896, OMITTED_INTERNAL_NATIVE_DECLARATION 65. Selected reachability is retained through every omitted internal declaration.

NormedAddCommGroup — Compiled external provider used by selected Project declarations.

NormedAddCommGroup.toSeminormedAddCommGroup — Compiled external provider used by selected Project declarations.

Real — Compiled external provider used by selected Project declarations.

Set — Compiled external provider used by selected Project declarations.

AddCommGroup.toDivisionAddCommMonoid — Compiled external provider used by selected Project declarations.

AddCommMonoidWithOne.toAddMonoidWithOne — Compiled external provider used by selected Project declarations.

AddGroup.toSubNegMonoid — Compiled external provider used by selected Project declarations.

AddGroupWithOne.toAddGroup — Compiled external provider used by selected Project declarations.

AddGroupWithOne.toAddMonoidWithOne — Compiled external provider used by selected Project declarations.

AddMonoid.toAddSemigroup — Compiled external provider used by selected Project declarations.

AddMonoidWithOne — Compiled external provider used by selected Project declarations.

AddMonoidWithOne.toNatCast — Compiled external provider used by selected Project declarations.

AddMonoidWithOne.toOne — Compiled external provider used by selected Project declarations.

AddSemigroup.toAdd — Compiled external provider used by selected Project declarations.

AffineIsometryEquiv — Compiled external provider used by selected Project declarations.

Dependency-first reading route

Level 0

Center transport under bounded point-reflection symmetry

MathlibAnnex.IsometryEquiv.map_center_of_mapsTo_pointReflection

f maps the source reflection center to the target reflection center.

Route / section: Theorem Project

Immediate prerequisites in this Project: None in this Project

Used by in this Project: Midpoint preservation on a symmetric lens

[Open Declaration Card](#)

[Read exact source](#)

Symmetric lens

MathlibAnnex.symmetricLens

The symmetric lens is the intersection of the closed radius- r balls centered at x and y . This definition needs no positivity assumption.

Route / section: Theorem Project

Immediate prerequisites in this Project: None in this Project

Used by in this Project: Midpoint preservation on a symmetric lens

[Open Declaration Card](#)

[Read exact source](#)

Level 1

Midpoint preservation on a symmetric lens

MathlibAnnex.IsometryEquiv.map_midpoint_of_symmetricLens

The isometry maps the midpoint of the source foci to the midpoint of the target foci.

Route / section: Theorem Project

Immediate prerequisites in this Project: Center transport under bounded point-reflection symmetry, Symmetric lens

Used by in this Project: Midpoint preservation under lens containment

[Open Declaration Card](#)

[Read exact source](#)

Level 2

Midpoint preservation under lens containment

`MathlibAnnex.IsometryEquiv.map_midpoint_of_symmetricLens_subset`

`f` preserves the midpoint of `x` and `y`. The target half-distance bound follows from the isometry.

Route / section: Theorem Project

Immediate prerequisites in this Project: Midpoint preservation on a symmetric lens

Used by in this Project: Midpoint preservation on a quarter ball

[Open Declaration Card](#)

[Read exact source](#)

Level 3

Midpoint preservation on a quarter ball

`MathlibAnnex.IsometryEquiv.map_midpoint_of_mem_ball`

`f` sends the midpoint of `x` and `y` to the midpoint of `f(x)` and `f(y)`.

Route / section: Theorem Project

Immediate prerequisites in this Project: Midpoint preservation under lens containment

Used by in this Project: Affine-segment preservation on a quarter ball

[Open Declaration Card](#)

[Read exact source](#)

Level 4

Affine-segment preservation on a quarter ball

`MathlibAnnex.IsometryEquiv.map_lineMap_of_mem_ball`

`f` preserves the affine combination $(1-a)x + ay$, with the exact subtype membership witnesses supplied in Lean.

Route / section: Theorem Project

Immediate prerequisites in this Project: Midpoint preservation on a quarter ball

Used by in this Project: Local affine-isometry chart on a smaller ball

[Open Declaration Card](#)

[Read exact source](#)

Level 5

Local affine-isometry chart on a smaller ball

`MathlibAnnex.IsometryEquiv.exists_affineExtension_eqOn_ball`

An ambient affine isometry equivalence agrees with f on the ball about c of radius $R/8$.

Route / section: Theorem Project

Immediate prerequisites in this Project: Affine-segment preservation on a quarter ball

Used by in this Project: Mankiewicz extension on open connected domains

[Open Declaration Card](#)

[Read exact source](#)

Level 6

Mankiewicz extension on open connected domains

`MathlibAnnex.IsometryEquiv.existsUnique_affineExtension`

There is a unique ambient real affine isometry equivalence whose restriction to s is f .

Route / section: Theorem Project

Immediate prerequisites in this Project: Local affine-isometry chart on a smaller ball

Used by in this Project: Affine extension from open balls, Convex-set extension via ambient interiors

[Open Declaration Card](#)

[Read exact source](#)

Level 7

Affine extension from open balls

`MathlibAnnex.IsometryEquiv.existsUnique_affineExtension_ball`

f extends uniquely to an ambient real affine isometry equivalence. The two positive radii need not be assumed equal.

Route / section: Theorem Project

Immediate prerequisites in this Project: Mankiewicz extension on open connected domains

Used by in this Project: None in this Project

[Open Declaration Card](#)

[Read exact source](#)

Convex-set extension via ambient interiors

`MathlibAnnex.IsometryEquiv.existsUnique_affineExtension_of_convex`

f extends uniquely to an ambient real affine isometry equivalence. No additional convexity assumption on t is stated.

Route / section: Theorem Project

Immediate prerequisites in this Project: Mankiewicz extension on open connected domains

Used by in this Project: Affine extension from equal-radius closed balls

[Open Declaration Card](#)

[Read exact source](#)

Level 8

Affine extension from equal-radius closed balls

MathlibAnnex.IsometryEquiv.existsUnique_affineExtension_closedBall

f extends uniquely to an ambient real affine isometry equivalence.

Route / section: Theorem Project

Immediate prerequisites in this Project: Convex-set extension via ambient interiors

Used by in this Project: None in this Project

[Open Declaration Card](#)

[Read exact source](#)