A Comparative Effectiveness Study of Intravitreal Aflibercept, Bevacizumab and Ranibizumab for Diabetic Macular Edema (T)

Intraocular Pressure Measurement Form

tblTIOP

PtID: **(tblPtRoster.PtID)**

Patient Initials:

Study Eye: **(tblPtRoster.StudyEye)**

IOP measurement is required in both eyes. Goldmann Tonometer should be used.

☐ **Not Done** **(IOPNotDone)**

If Required but not done, give reason in COMMENTS

If Done, please complete the following

GENERAL INFORMATION

1. IOP measurement date: __ __/__ __ __/__ __ __ __ **(IOPExamDtDaysFromEnroll)**
dd/MMM/yyyy

2. IOP Tester:_____

3. How was IOP measured? O Goldmann O Tonopen O Other _____ **(IOPMeasured)**

4. When was the IOP measured? O Pre-Dilation O Post-Dilation **(IOPDilation)**

INTRAOCULAR PRESSURE MEASUREMENT – RIGHT EYE (OD)

1. Intraocular Pressure (OD): _____mm Hg **(IOPOD)** ☐ **Not Done (if required, enter reason not done in comments)** **(IOPNotDoneOD)**

INTRAOCULAR PRESSURE MEASUREMENT – LEFT EYE (OS)

1. Intraocular Pressure (OS): _____mm Hg **(IOPOS)** ☐ **Not Done (if required, enter reason not done in comments)** **(IOPNotDoneOS)**

COMMENTS