A Comparative Effectiveness Study of Intravitreal Aflibercept, Bevacizumab and Ranibizumab for Diabetic Macular Edema (T)

Intravitreal Injection Form

tblTIntraInj

PtID: (tblPtRoster.PtID)

Patient Initials:

Study Eye: (tblPtRoster.StudyEye)

☐ **Intravitreal injection not done (InjNotDone)**

If not done, give reason: **(RandInjMissReas)**

- O Adverse Event
- O Patient Refuses
- O Other _____

If Done, please complete the following.

GENERAL INFORMATION

1. Treatment Date: ___ ___/___ ___ ___/___ ___ ___ ___ **(InjectDtDaysFromEnroll)**
dd/MMM/yyyy

2. Name of Investigator _____

3. Site Location: _____

TREATMENT CONFIRMATION

1. Was the correct study eye verified by two individuals using the study website or a printout from the study website AND was the eye marked (with a sticker or marking pen)? O Yes O No **(ConfirmEye)**

2. Was the correct treatment verified by two individuals using the study website or a printout from the study website? O Yes O No **(ConfirmTrt)**

ANESTHETIC USED

1. Record anesthetic used for the injection.

Check all that apply:

☐ Topical non-viscous anesthetic drops; record name(s) and concentration(s) (e.g. Ophthetic, Tetracaine) below **(AnesthTopical)**

☐ Topical viscous anesthetic drops or gel; record name(s) and concentration(s) (e.g. TetraVisc) below **(AnesthTopViscious)**

☐ Subconjunctival injection; record name(s) and concentration (s) below **(AnesthSubconj)**

☐ Other; describe below **(AnesthOth)**

1a. Describe:

_____

_____

PREP

1. Indicate the prep that was done.

*Check all that apply**:***

☐ 5% povidone iodine placed in lower fornix **(PovIodFornix)**

☐ 5% or 10% povidone iodine cotton-tipped applicator applied to upper and lower eyelids and eyelashes. **(PovIodLidLash)**

☐ 5% or 10% povidone iodine cotton-tipped applicator applied to the conjunctiva over and surrounding the injection site **(PovIodConj5)**

☐ 5% povidone-iodine topical drop(s) applied to the conjunctiva over and surrounding the injection site **(PovIodFlush)**

☐ 10% povidone iodine Swabstick applied to the conjunctiva over and surrounding the injection site **(PovIodConj10)**

☐ Other (OthPrep); describe and indicate reason for using an "other" prep_____ **(OthPrep)**

2. Was an eyelid speculum used? O Yes O No **(LidSpeculum)**

INJECTION

1. Eye Treated: O OD O OS **(TrtEye)**

2. Drug and dose administered:_____ **(InjType2)**

If intravitreal (i.e. not sham), complete the following:

2a. What gauge needle was used for the injection? O 27 O 30 **(NeedleGauge)**

2b. Location of injection: (InjectLoc)

- O inferior-temporal
- O inferior-nasal
- O superior-temporal
- O superior-nasal

2c. Was the injection performed 3.0-4.0 mm posterior to the limbus, via the pars plana as per the injection procedure? O Yes O No **(InjProc)**

i. If No, explain: _____
_____

2d. Did needle separation occur during priming and/or during intravitreal injection? (NeedleSep)

O Yes, during injection O Yes, during priming O No

i. If Yes, describe: _____
_____

2e. Did any other problem occur during injection? O Yes O No **(InjectOthProb)**

i. If Yes, describe: _____
_____

NUMBERED DRUG INFORMATION

1. Syringe/vial number used for the injection_____

POST INJECTION

1. Was indirect ophthalmoscopy performed to confirm the perfusion of the central retinal artery?
O Yes O No **(IndirectOphthal)**

2. Was vision checked to confirm that there is some perception of vision in the study eye?
O Yes O No **(VisionCheck)**

3. Intraocular pressure (last recorded measurement):
___ mm Hg **(IOP)** ☐ Not Measured **(IOPNotMeas)**

4. Did the subject receive treatment to lower the intraocular pressure? O Yes O No **(TrtLowIOP)**

If YES:

4a. Check all that apply,
☐ Topical treatment(s); describe below **(TrtLowIOPTopTrt)**
☐ Paracentesis **(Paracentesis)**
☐ Other; describe below **(TrtLowIOPOth)**

Describe:_____

An Adverse Event Form does NOT need to be completed for increased intraocular pressure noted on this form.

ADVERSE EVENTS

1. Did the subject experience any adverse events from the intravitreal injection (other than an increase in intraocular pressure) prior to leaving office on day of injection? O Yes O No **(InjAE)**
(If Yes, complete an Adverse Event Form)

ANTIBIOTIC USED

1. Was a pre-injection topical antibiotic used? (TopAntibioticUsed)
O Yes, prescribed for home use and given in clinic on day of injection
O Yes, prescribed for home use only
O Yes, given in clinic on day of injection only
O No

If Yes:

1a. Describe drug and regimen

_____
_____

2. Were post-injection topical antibiotics used? (PostInjAntiHomeUse)

- O Yes, given in clinic post-injection and prescribed for home use
- O Yes, given in clinic post-injection only
- O Yes, prescribed for home use only
- O No

(If Yes, record on the Concomitant Medication Form)

COMMENTS

Record any pertinent information about the procedure that is not covered above