A Comparative Effectiveness Study of Intravitreal Aflibercept, Bevacizumab and Ranibizumab for Diabetic Macular Edema (T)

Protocol Specific Exam Questions Follow-up

tblTFUExamQuestions

GENERAL

NOTE: If study drug may be given, female subjects should be questioned about the possibility of pregnancy. Investigator judgment is used to determine when a pregnancy test is needed prior to injection. Injections should be discontinued if a subject is pregnant or if a subject who indicates that pregnancy is possible refuses to have a pregnancy test done.

1. Is the study participant pregnant or attempting to become pregnant? (PtPreg)
O Yes O No or participant is male

Medical Conditions
Please record all pre-existing conditions on the Pre-existing Medical Conditions form. Please record all medications on the Concomitant Medications form.

1. Please indicate if the participant has had the listed condition:

) **Myocardial infarction (MI) (MICond)**
O Yes O No knowledge of having the condition

) **Coronary artery disease (without MI) (CADCond)**
O Yes O No knowledge of having the condition

) **Stroke (StrokeCond)**
O Yes O No knowledge of having the condition

) **Transient ischemic attacks (TIACond)**
O Yes O No knowledge of having the condition

) **Gastointestinal perforation (GasPerfCond)**
O Yes O No knowledge of having the condition

) **Renal disease (Known renal disease identified through microalbuminuria testing) (RenalDiseaseCond)**
O Yes O No knowledge of having the condition

) **Hypertension**
O Yes O No knowledge of having the condition

) **Severe Hemorrhage (includes hemoptysis, epistaxis, gastrointestinal, and vaginal bleeding)**
O Yes O No knowledge of having the condition **(SevHemorrhageCond)**

RIGHT EYE (OD)

1. Have <u>both</u> of the following criteria for "complete focal photocoagulation" been met in the right eye?

) **All leaking microaneurysms within areas of retinal thickening or contributing to the edema that is threatening the center of the macula have been directly treated at some time with laser burns directly over the microaneurysms**

) **All areas of current retinal thickening have been treated with laser burns (either focal or grid), such that the laser burns are "on average" within 100 microns of each other (range between 50 and 150 microns) in a grid pattern throughout the area of retinal thickening**

O **Yes** O **No** O **Cannot assess (i.e. no view)** O **Did not assess (CompleteLaserOD)**

2. Is edema "threatening the fovea" on clinical exam in the right eye?

) **Edema threatening the fovea is defined as edema within 500 microns of the foveal center or edema associated with lipid within 500 microns of the foveal center or ≥ 1 disc area of edema the posterior edge of which is within 1 disc diameter of the foveal center**

O **Yes** O **No** O **Cannot assess (i.e. no view)** O **Did not assess** **(EdemaThreatFoveaOD)**

3. Is DME present in the right eye that you believe is causing visual acuity loss?

O **Yes** O **No** O **Cannot assess (i.e. no view)** O **Did not assess** **(DMEPresCauseVALossOD)**

LEFT EYE (OS)

1. Have <u>both</u> of the following criteria for "complete focal photocoagulation" been met in the left eye?

) **All leaking microaneurysms within areas of retinal thickening or contributing to the edema that is threatening the center of the macula have been directly treated at some time with laser burns directly over the microaneurysms**

) **All areas of current retinal thickening have been treated with laser burns (either focal or grid), such that the laser burns are "on average" within 100 microns of each other (range between 50 and 150 microns) in a grid pattern throughout the area of retinal thickening**

O **Yes** O **No** O **Cannot assess (i.e. no view)** O **Did not assess** **(CompleteLaserOS)**

2. Is edema "threatening the fovea" on clinical exam in the left eye?

) **Edema threatening the fovea is defined as edema within 500 microns of the foveal center or edema associated with lipid within 500 microns of the foveal center or ≥ 1 disc area of edema the posterior edge of which is within 1 disc diameter of the foveal center**

O **Yes** O **No** O **Cannot assess (i.e. no view)** O **Did not assess** **(EdemaThreatFoveaOS)**

3 Is DME present in the left eye that you believe is causing visual acuity loss?

O **Yes** O **No** O **Cannot assess (i.e. no view)** O **Did not assess** **(DMEPresCauseVALossOS)**

<u>COMMENTS</u>