A Comparative Effectiveness Study of Intravitreal Aflibercept, Bevacizumab and Ranibizumab for Diabetic Macular Edema (T)

Visual Acuity Testing Form (Annual Follow-up)

tblTVAReVATest

Patient ID: **(tblPtRoster.PtID)**

Patient Initials:

Study Eye: **(tblPtRoster.StudyEye)**

Refraction followed by visual acuity testing is required in each eye. Visual acuity should be measured in each eye separately using the E-ETDRS program on the Electronic Visual Acuity Tester. If the EVA is unavailable, ETDRS charts should be used. Snellen or other charts should NOT be used. Both refraction and visual acuity testing should be performed by a certified masked examiner.

VISUAL ACUITY TESTING

Visual Acuity Testing: ○ Not Done ○ Right eye only ○ Left eye only ○ Both eyes **(VACompleted)**

Reason not done:_____

Note: If visual acuity is too poor for EVA or Chart testing please include the eye in the selection above and complete the form below as indicated.

GENERAL INFORMATION

1. Name of VA Tester: _____

2. Was the visual acuity tester masked to the study participant's treatment group? **(VAMasked)**

Complete If EVA Used

1. EVA Instrument # *(from label or palm):* _____

2. Calibration Checks

Verify the following:

☐ 2a. Testing distance = 3 meters (118 inches) from monitor screen to center of exam chair seat **(InstrMeasDist)**

☐ 2b. Brightness of screen within range on light meter **(InstrBright)**

☐ 2c. Size of EVA calibration square: horizontal = 114 mm and vertical = 114 mm **(InstrCalibSize)**

REFRACTION

Indicate below if a refraction was performed in either eye <u>prior</u> to the initial visual acuity testing.

1. Name of refractionist (if done):_____

2. Right Eye:_____ _____ @ _____ o OR ☐ **Not Done** **(VARefNotDoneOD)**
sph cyl axis
(VARefSphOD) (VARefCylOD) (VARefAxisOD)

3. Left Eye:_____ _____ @ _____ o OR ☐ **Not Done** **(VARefNotDoneOS)**
sph cyl axis
(VARefSphOS) (VARefCylOS) (VARefAxisOS)

VISUAL ACUITY TESTING – RIGHT EYE (OD)

1. How was visual acuity measured in the right eye? ○ **EVA-ETDRS** ○ **ETDRS Chart** ○ **Non-protocol testing** **(VAMethodOD)**

(Note: if visual acuity is too poor for EVA or Chart testing, please select EVA-ETDRS above and enter visual acuity letter score of 0 below.)

1a. If EVA-ETDRS was not used, reason: _____

If ETDRS charts were used, please fax acuity worksheet to the Jaeb Center 1-800-816-7601. See study participant data entry menu to print form.

2. Correction Used for Visual Acuity Testing:

☐ **Today's refraction recorded above** **(TodayVARefUsedOD)** OR 'Other' _____ _____ @ _____ o
sph cyl axis
(VACorSphOD) (VACorCylOD) (VACorAxisOD)

3. Visual Acuity Letter Score for Right Eye:_____ **(EtdrsOD)**

(If vision is too poor for EVA or Chart testing, please enter acuity score of 0 for that eye)

If any aspects of the acuity testing were not completed according to the protocol, please detail in COMMENTS.

3a. EVA Score Code OD: __ __ - __ __ __ (EVAValCodeOD) ☐ **Not Verified (EVAValCodeODNV)**

If visual acuity letter score is 0, answer the following:

3b. Which of the following best describes the vision:

O counting fingers O hand motion O light perception O no light perception **(VAZeroVisOD)**

3c. Please indicate the reason for poor vision: (VAScoreZeroCatOD)

- **O** Cataract
- **O** Corneal edema
- **O** Endophthalmitis
- **O** Macular edema
- **O** Macular non-perfusion
- **O** No cause identified
- **O** Optic nerve abnormality
- **O** Rhegmatogenous retinal detachment
- **O** Traction retinal detachment
- **O** Uveitis
- **O** Vitreomacular interface abnormality – epiretinal membrane
- **O** Vitreomacular interface abnormality - macular hole
- **O** Vitreomacular interface abnormality – other
- **O** Vitreous hemorrhage
- **O** Vitritis
- **O** Other

If other, describe _____

4. If Non-protocol testing was used OD, the Snellen acuity based on the standard office chart testing:
_____ **(SnellenOD)**

VISUAL ACUITY TESTING – LEFT EYE (OS)

1. How was visual acuity measured in the left eye? O **EVA-ETDRS** O **ETDRS Chart** O **Non-protocol testing** **(VAMethodOS)**

(Note: if visual acuity is too poor for EVA or Chart testing, please select EVA-ETDRS above and enter visual acuity letter score of 0 below.)

1a. If EVA-ETDRS was not used, reason: _____

If ETDRS charts were used, please fax acuity worksheet to the Jaeb Center 1-800-816-7601. See study participant data entry menu to print form.

2. Correction Used for Visual Acuity Testing:

☐ **Today's refraction recorded above (TodayVARefUsedOS)** OR 'Other' _____ (sph) _____ (cyl) @ _____ (axis) o
(VACorSphOS) (VACorCylOS) (VACorAxisOS)

3. Visual Acuity Letter Score for Left Eye: _____ **(EtdrsOS)**

(If vision is too poor for EVA or Chart testing, please enter acuity score of 0 for that eye)

If any aspects of the acuity testing were not completed according to the protocol, please detail in COMMENTS.

3a. EVA Score Code OS: __ __ - __ __ __ (EVAValCodeOS) ☐ **Not Verified (EVAValCodeOSNV)**

If visual acuity letter score is 0, answer the following:

3b. Which of the following best describes the vision:

O counting fingers O hand motion O light perception O no light perception **(VAZeroVisOS)**

3c. Please indicate the reason for poor vision: (VAScoreZeroCatOS)

- **O** Cataract
- **O** Corneal edema
- **O** Endophthalmitis
- **O** Macular edema
- **O** Macular non-perfusion
- **O** No cause identified
- **O** Optic nerve abnormality
- **O** Rhegmatogenous retinal detachment
- **O** Traction retinal detachment
- **O** Uveitis
- **O** Vitreomacular interface abnormality – epiretinal membrane
- **O** Vitreomacular interface abnormality - macular hole
- **O** Vitreomacular interface abnormality – other
- **O** Vitreous hemorrhage
- **O** Vitritis
- **O** Other

If other, describe _____

4. If Non-protocol testing was used OS, the Snellen acuity based on the standard office chart testing:
_____ **(SnellenOS)**

VISUAL ACUITY TESTING DATE

1. Visual acuity testing date:

___ ___ / ___ ___ ___ /___ ___ ___ ___ **(VAExamDt)**
dd/MMM/yyyy

1a. EVA date code:__ __ __ - __ __ __ __ (EVAValCode) ☐ **Not Verified (EVAValCodeNV)**

DECREASED VISUAL ACUITY-Right Eye (OD)

1. If visual acuity has decreased at least 10 letters since the prior visit please indicate the primary reason for the visual acuity decrease (select link above to view prior visual acuity): (DecreasedVAOD)

- O Cataract
- O Corneal edema
- O Endophthalmitis
- O Macular edema
- O Macular non-perfusion
- O No cause identified
- O Optic nerve abnormality
- O Rhegmatogenous retinal detachment
- O Traction retinal detachment
- O Uveitis
- O Vitreomacular interface abnormality – epiretinal membrane
- O Vitreomacular interface abnormality - macular hole
- O Vitreomacular interface abnormality – other
- O Vitreous hemorrhage
- O Vitritis
- O Other

1a. If other, describe _____

DECREASED VISUAL ACUITY-Left Eye (OS)

1. If visual acuity has decreased at least 10 letters since the prior visit please indicate the primary reason for the visual acuity decrease (select link above to view prior visual acuity): (DecreasedVAOS)

- O Cataract
- O Corneal edema
- O Endophthalmitis
- O Macular edema
- O Macular non-perfusion
- O No cause identified
- O Optic nerve abnormality
- O Rhegmatogenous retinal detachment
- O Traction retinal detachment
- O Uveitis
- O Vitreomacular interface abnormality – epiretinal membrane
- O Vitreomacular interface abnormality - macular hole
- O Vitreomacular interface abnormality – other
- O Vitreous hemorrhage
- O Vitritis
- O Other

1a. If other, describe _____

COMMENTS