

# Title

## Subtitle

Rebecca Wei

Northwestern University

Date/Event



A more categorical approach (or somehow otherwise motivate  $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

# A more categorical approach (or somehow otherwise motivate $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

Objects: algebra maps  $f : A \rightarrow A$

Morphisms:  $Hoch(A)(f, g) = (C^\bullet(A, {}_f A_g), {}_f \delta_g)$

# A more categorical approach (or somehow otherwise motivate $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

Objects: algebra maps  $f : A \rightarrow A$

Morphisms:  $Hoch(A)(f, g) = (C^\bullet(A, {}_f A_g), {}_f \delta_g)$

$$\begin{aligned} {}_f \delta_g(\phi)(a_1 \otimes \dots \otimes a_n) = & \epsilon_\phi \left( \textcolor{red}{f(a_1)} \cdot \phi(a_2, \dots, a_n) + \right. \\ & + \sum_{1 \leq i \leq n-1} (-1)^i \phi(a_1, \dots, a_i a_{i+1}, \dots, a_n) + \\ & \left. + (-1)^n \phi(a_1, \dots, a_{n-1}) \cdot \textcolor{red}{g(a_n)} \right) \\ \epsilon_\phi = & (-1)^{|\phi|+1} \end{aligned}$$

# A more categorical approach (or somehow otherwise motivate $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

Objects: algebra maps  $f : A \rightarrow A$

Morphisms:  $Hoch(A)(f, g) = (C^\bullet(A, {}_f A_g), {}_f \delta_g)$

Composition: cup product on cochains

# A more categorical approach (or somehow otherwise motivate $B(n)$ )


Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

Objects: algebra maps  $f : A \rightarrow A$

Morphisms:  $Hoch(A)(f, g) = (C^\bullet(A, {}_f A_g), {}_f \delta_g)$

Composition: cup product on cochains



A more categorical approach (or somehow otherwise motivate  $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

$$Bar(Hoch(A)) \otimes Bar(Hoch(A)) \xrightarrow{\bullet} Hoch(A)$$




# A more categorical approach (or somehow otherwise motivate $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

$$Bar(Hoch(A)) \otimes Bar(Hoch(A)) \xrightarrow{\bullet} Hoch(A)$$




A morphism from  $f_0$  to  $f_n$  in  $Bar(Hoch(A))$

# A more categorical approach (or somehow otherwise motivate $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

$$Bar(Hoch(A)) \otimes Bar(Hoch(A)) \xrightarrow{\bullet} Hoch(A)$$



# A more categorical approach (or somehow otherwise motivate $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

In this context, braces give multilinear maps:

$$\begin{array}{ccc} Bar(Hoch(A)) \otimes Bar(Hoch(A)) & \xrightarrow{\bullet} & Hoch(A) \\ & \searrow \bullet & \uparrow \\ & & Bar(Hoch(A)) \end{array}$$

Then,  $(Bar(Hoch(A)), \bullet)$  is an algebra in  $DGCocats$ .

# A more categorical approach (or somehow otherwise motivate $B(n)$ )

Fix an algebra,  $A$ .

Define a dg category,  $Hoch(A)$ :

In this context, braces give multilinear maps:

$$\begin{array}{ccc} Bar(Hoch(A)) \otimes Bar(Hoch(A)) & \xrightarrow{\bullet} & Hoch(A) \\ & \searrow \bullet & \uparrow \\ & & Bar(Hoch(A)) \end{array}$$

Then,  $(Bar(Hoch(A)), \bullet)$  is an algebra in  $DGCocats$ .

But we have more...

A more categorical approach, (or somehow otherwise motivate  $B(n)$ )

Fix algebras,  $A_0, A_1, \dots, A_n$ .

Define a dg cocategory  $B(A_0 \rightarrow A_1 \rightarrow \dots \rightarrow A_n \rightarrow A_0)$

where, for  $n=0$ ,  $B(A_0 \rightarrow A_0) := \text{Bar}(\text{Hoch}(A_0))$ .

# A sheafy-cyclic object in $\mathrm{DGCocat}$

Fact: We have a functor