



Thomas Lewiner

Constructing Discrete Morse Functions

MsC Thesis

Thesis presented to the Post-graduate Program in Applied Mathematics of the Matematica Department, PUC-Rio as partial fulfillment of the requirements for the degree of Master in Applied Mathematics

Adviser : Prof. Hélio Côrtes Vieira Lopes
Co-Adviser: Prof. Geovan Tavares dos Santos

Rio de Janeiro
July 2002

Thomas Lewiner

Constructing Discrete Morse Functions

Thesis presented to the Post-graduate Program in Applied Mathematics of the Mathematics Department, PUC-Rio as partial fulfillment of the requirements for the degree of Master in Applied Mathematics. Approved by the following commission:

Prof. Hélio Côrtes Vieira Lopes

Adviser

Department of Mathematics — PUC-Rio

Prof. Geovan Tavares dos Santos

Co-Adviser

Department of Mathematics — PUC-Rio

Prof. Luis Velho

Visgraf Laboratory — IMPA

Prof. Jorge Stolfi

Institute of Computing — UNICAMP

Prof. José Eugênio Leal

Head of the Science and Engineering Center — PUC-Rio

Rio de Janeiro — July 10, 2002

All rights reserved.

Thomas Lewiner

Thomas Lewiner graduated from the Ecole Polytechnique (Paris, France) in Algebra and Computer Science, and in Theoretical Physics. He then specialized at the Ecole Supérieure des Télécommunications (Paris, France) in Signal and Image Processing, and in Project Management, while working for Inventel in wireless telecommunication systems based on BlueTooth technology. He then obtained a Master degree at the PUC-Rio in computational topology, and actively participated on the department's work for Petrobras.

Bibliographic data

Lewiner, Thomas

Constructing Discrete Morse Functions / Thomas Lewiner; adviser: Hélio Côrtes Vieira Lopes; co-adviser: Geovan Tavares dos Santos. — Rio de Janeiro : PUC-Rio, Department of Mathematicc, 2002.

v., 17 f: il. ; 29,7 cm

1. MsC Thesis - Pontifícia Universidade Católica do Rio de Janeiro, Department of Mathematicc.

Bibliography included.

1. Mathematicc – Dissertation. 2. Morse Theory. 3. Forman Theory. 4. Computational Topology. 5. Computational Geometry. 6. Solid Modeling. 7. Discrete Mathematics. I. Lopes, Hélio Côrtes Vieira. II. Santos, Geovan Tavares dos. III. Pontifícia Universidade Católica do Rio de Janeiro. Department of Mathematicc. IV. Title.

CDD: 510

Acknowledgments

To my advisors Professors H lio Lopes and Geovan Tavares for the support, the everyday kindness and the incentive for the realisation of this work.

To the CNPq and the PUC Rio, for the financial support, without which this work would not have been realized.

To my grand mothers, who suffered the most of my expatriation. To my mother and father, sisters and all my family.

To my colleagues of the PUC Rio, who have me loved this place.

To the professors Marcos da Silvera, Jean Marie Nicolas and Anne Germa who offered me the opportunity of this cooperation.

To the people of the Mathematic departament for the constant help, particularly to Ana Cristina, Creuza and Sinesio.

Abstract

Lewiner, Thomas; Lopes, Hélio Côrtes Vieira; Santos, Geovan Tavares dos. **Constructing Discrete Morse Functions**. Rio de Janeiro, 2002. 17p. MsC Thesis — Department of Matematicc, Pontifícia Universidade Católica do Rio de Janeiro.

Morse theory has been considered a powerful tool in its applications to computational topology, computer graphics and geometric modeling. It was originally formulated for smooth manifolds. Recently, Robin Forman formulated a version of this theory for discrete structures such as cell complexes. It opens up several categories of interesting objects (particularly meshes) to applications of Morse theory.

Once a Morse function has been defined on a manifold, then information about its topology can be deduced from its critical elements. The purpose of this work is to design an algorithm to define optimal discrete Morse functions on general cell complex, where optimality entails having the least number of critical elements. This problem is proven here to be MAX-SNP hard. However, we provide a linear algorithm that, for the case of 2-manifolds, always reaches optimality.

Moreover, we proved various results on the structure of a discrete Morse function. In particular, we provide an equivalent representation by hyperforests. From this point of view, we designed a construction of discrete Morse functions for general cell complexes of arbitrary finite dimension. The resulting algorithm is quadratic in time and, although not guaranteed to be optimal, gives optimal answers in most of the practical cases.

Keywords

Morse Theory. Forman Theory. Computational Topology. Computational Geometry. Solid Modeling. Discrete Mathematics.

Contents

I	Future works	15
	Bibliography	17

List of Figures

I.1	Uma figura	16
-----	------------	----

Un abîme effrayant, une profusion de questions de toutes sortes où ma responsabilité était en jeu se présentaient à moi. Et la plus importante: qu'est-ce qui doit remplacer l'objet manquant? Le danger d'un art ornemental m'apparaissait clairement, la morte existence illusoire des formes stylisées ne pouvait que me rebuter.

C'est seulement après de nombreuses années d'un travail patient, d'une réflexion intense, d'essais nombreux et prudents où je développais toujours plus la capacité de vivre purement, abstraitement les formes picturales et de m'absorber toujours plus profondément dans ces profondeurs insondables, que j'arrivais à ces formes picturales avec lesquelles je travaille aujourd'hui et qui, comme je l'espère et le veux, se développeront bien plus encore.

Il a fallu beaucoup de temps avant que cette question: 'qu'est ce qui doit remplacer l'objet ?' trouve en moi une véritable réponse. Souvent je me retourne vers mon passé et je suis désespéré de voir combien de temps il m'a fallu pour arriver à cette solution.

Wassily Kandinsky, *Regards sur le passé.*

I

Future works

This work was focused on Forman’s discrete Morse theory. We analyzed the building blocs of this theory, and proved the layered structure of discrete Morse functions. We represented this layer structure by a collection of hyperforests and gave a complete characterization of the critical cells in terms of regular components of hyperforests. We used this analysis to introduce a scheme for constructing discrete Morse function on finite cell complexes of arbitrary dimension. This construction is quadratic in time in the worst cases, and is proven to be linear and optimal in the case of 2-manifolds. The experimental results showed our algorithm gave an optimal result in most of the cases. This opens the question of which conditions on the cell complex would ensure the optimality of the resulting function.

An important application of this work to computer graphics would be in the field of geometric compression. The algorithm Grow&Fold of A. Szymczak and J. Rossignac [Szy00] could be justified and enhanced by our algorithm to minimize the number of so-called “glue faces” in order to achieve a better encoding. This work has been done in an optimal way for the case of surfaces with handles in [Lop02].

We plan to continue this work in three directions. First, as mentioned above, apply Forman’s theory and the analysis of our algorithm for solid mesh compression. Second, we will try to apply discrete Morse theory to resolve singularities that arise from shape reconstruction. Finally, we look forward to produce a topologically consistent morphing based on mapping directly the discrete gradient field between two objects of the same homotopy type.




Figure I.1: Uma figura

Bibliography

- [Lop02] H. Lopes, J. Rossignac, A. Safanova, A. Szymczak and G. Tavares. **Edgebreaker: a simple compression for surfaces with handles**. In *7th ACM Siggraph Symposium on Solid Modeling and Application*, 2002. I
- [Szy00] A. Szymczak and J. Rossignac. **Grow & Fold: Compression of Tetrahedral Meshes**. In *Solid Modelling '99*, 1999. to appear. I