

HARMONIC FUNCTIONS AND THE POSITIVE MASS THEOREM FOR ASYMPTOTICALLY FLAT HALF-SPACES



Bachelor's Thesis in Mathematics
eingereicht an der Fakultät für Mathematik und Informatik
der Georg-August-Universität Göttingen
am 19. Juni 2023

von
Henry Ruben Fischer

Erstgutachter:
Prof. Dr. Thomas Schick

Zweitgutachter:
Prof. Dr. Max Wardetzki

Contents

1	Introduction	1
2	Prerequisites	1
3	Main basic identity	7
4	Existence and uniqueness of asymptotically linear harmonic functions	7
5	Proof of the Mass Lower Bound	9
A	Different Exhausting Sequences for Computation of the Mass	11

1 Introduction

TODO Add historical context and applications of the Positive Mass theorem.

Todo

In the following, we will explore the relatively new proof of the Positive Mass theorem using (spacetime) harmonic functions, and in particular consider the case of rigidity. We will then look at how these functions look in some simple example cases. Finally, we will apply this method to the Positive Mass theorem on asymptotically flat half spaces with connected (non-compact) boundary, acquiring an explicit lower bound for the mass in the process, i.e. we will prove the following theorem (all notation and concepts will be introduced later):

Theorem 1.1. *Let (M, g) be an asymptotically flat half-space (M, g) of dimension $n = 3$ with horizon boundary $\Sigma \subset M$, associated exterior region $M(\Sigma)$ and connected non-compact boundary ∂M . Let (x_1, x_2, x_3) be asymptotically flat coordinates such that outside of a compact set, M is diffeomorphic to $\{x \in \mathbb{R}_+^3 \mid |x| > 1\}$. Assume that the following three conditions hold*

- $R(g) \geq 0$ in $M(\Sigma)$.
- $H(\partial M, g) \geq 0$ on $M(\Sigma) \cap \partial M$.

Then there exists a unique harmonic function u on $M(\Sigma)$ asymptotic to x_3 fulfilling zero Dirichlet boundary conditions on $\partial M \cap M(\Sigma)$ and zero Neumann boundary conditions on Σ , and we have

$$m \geq \int_{M(\Sigma)} \left(\frac{|\nabla^2 u|^2}{|\nabla u|} + R(g)|\nabla u| \right) dV + \int_{\partial M \cap M(\Sigma)} H(\partial M, g)|\nabla u| dA \geq 0,$$

where m is the ADM mass of M . Equality $m = 0$ occurs if and only if (M, g) is isometric to (\mathbb{R}_+^3, δ) .

2 Prerequisites

In the following let M always be a 3-dimensional Riemannian manifold, equipped with the unique Levi-Civita-Connection.

Note that by ∇ we will always denote the covariant derivative / the Levi-Civita connection. In particular we will have $\nabla f = df$, and $\text{grad } f = (df)^\sharp$, such that in coordinates df will be given by $\nabla_i f = \partial_i f$ and $\text{grad } f$ will be given by $\nabla^i f$.

We will often be considering M as a three dimensional spacelike surface embedded in a larger $3 + 1$ -dimensional Lorentzian manifold \tilde{M} (for the signature of the metric

of \tilde{M} we choose the convention $(-, +, +, +)$, such that the induced metric on M is positive definite. M itself will also often have a boundary ∂M . Thus some basic facts about submanifolds will be helpful. Most of the following is from [Lee19, Chapter 2.1].

Let Σ^m be a submanifold of (pseudo-)Riemannian manifolds (M^n, g) (equipped with Levi-Civita connection ∇). We have an induced metric $\gamma = g|_{T\Sigma}$ (also called the *first fundamental form*).

Fact 2.1. Denoting the Levi-Civita connection of (Σ, γ) by $\hat{\nabla}$, we have for any $p \in \Sigma$, tangent vector $v \in T_p\Sigma$ and $Y \in \Gamma(T\Sigma)$,

$$\hat{\nabla}_X Y = (\nabla_X \tilde{Y})^\top,$$

where \tilde{Y} is any extension of Y to a vector field on M . Here $(-)^{\top}$ denotes the orthogonal projection from $T_p M$ to $T_p \Sigma$.

TODO This might be too similar to [Lee19, p. 2.1]

Todo

Thus Σ intrinsically contains information about tangential parts of tangential derivatives. But this information does not determine the orthogonal part! This motivates the following definition:

Definition 2.2 (Second fundamental form). The *second fundamental form* of Σ is a tensor $\mathbf{A} \in \Gamma(T^*\Sigma \otimes T^*\Sigma \otimes N\Sigma)$ such that

$$\mathbf{A}(X, Y) := (\nabla_X \tilde{Y})^\perp,$$

where $(-)^{\perp}$ denotes the orthogonal projection from $T_p M$ to $T_p \Sigma$.

Fact 2.3. $\mathbf{A}(X, Y) = \mathbf{A}(Y, X)$, i.e. \mathbf{A} is symmetric (since $\nabla_X Y - \nabla_Y X = [X, Y] \in T_\Sigma$).

Definition 2.4. The *mean curvature vector* \mathbf{H} is the trace of \mathbf{A} over $T_p \Sigma$, i.e. for an orthonormal basis e_1, \dots, e_n of $T_p \Sigma$ we define

$$\mathbf{H} := \sum_{i=1}^m \mathbf{A}(e_i, e_i).$$

If Σ is an orientable hypersurface of M , we can choose a normal direction ν (if Σ has an interior and exterior, we typically implicitly choose ν to be the outward normal). Then we define

$$A(X, Y) := g(\mathbf{A}(\mathbf{X}, \mathbf{Y}), -\nu) \quad H := g(\mathbf{H}, -\nu) = \text{tr}_\gamma(A).$$

We also call A the second fundamental form and H the mean curvature. Note that we have

$$A(X, Y) = g(\nabla_X Y, -\nu) = \underbrace{\nabla_X(g(Y, -\nu))}_{=0} - g(Y, \nabla_X(-\nu)) = g(Y, \nabla_X \nu).$$

The following (which is [Exercise 2.3 in Lee19]) is one of the main facts we will require:

Fact 2.5. *Given a hypersurface Σ in (M, g) and a smooth function f on M ,*

$$\Delta_M f = \Delta_\Sigma + \nabla_\nu \nabla_\nu f + H \nabla_\nu f.$$

Proof. Choose an orthonormal frame e_1, \dots, e_n of $T_p M$ and $e_1, \dots, e_{n-1} \in T_p \Sigma$ and $e_n = \nu$, then

$$\begin{aligned} \Delta_g f &= \sum_{i=1}^n (\nabla \nabla f)(e_i, e_i) \\ &= \nabla_M \nabla_\nu f + \sum_{i=1}^{n-1} g(\nabla_{e_i}(\text{grad}_M f), e_i) \\ &= \nabla_\nu \nabla_\nu f + \sum_{i=1}^{n-1} g(\nabla_{e_i}(\text{grad}_\Sigma f + \nu \cdot \nabla_\nu f), e_i) \\ &= \nabla_\nu \nabla_\nu f + \sum_{i=1}^{n-1} (\gamma(\hat{\nabla}_{e_i}(\text{grad}_\Sigma f), e_i) + \underbrace{g(\mathbf{A}(e_i, \text{grad } \Sigma), e_i)}_{=0}) \\ &\quad + \nabla_{e_i} \nabla_\nu f \cdot \underbrace{g(\nu, e_i)}_{=0} + \nabla_\nu f \cdot g(\nabla_{e_i} \nu, e_i) \\ &= \nabla_\nu \nabla_\nu f + \sum_{i=1}^{n-1} (\hat{\nabla} \hat{\nabla} f)(e_i, e_i) + \nabla_\nu f \cdot \sum_{i=1}^{n-1} A(e_i, e_i) \\ &= \nabla_\nu \nabla_\nu f + \Delta_\Sigma f + \nabla_\nu f \cdot H. \end{aligned}$$

■

We will now establish the necessary definitions (mostly adapted from [ABL16], [EK23] and [Bra+19]) to state the main result of this thesis.

Definition 2.6 (Asymptotically flat half-space). Let (M, g) be a connected, complete Riemannian manifold of dimension 3, with scalar curvature R and a non-compact

boundary ∂M with mean curvature H (computed as the divergence along ∂M of an outward pointing unit normal ν).

We call (M, g) an *asymptotically flat half-space* with decay rate $\tau > 0$ if there exists a compact subset K such that $M \setminus K$ consists of a finite number of connected components M_{end}^i called *ends*, such that for each of these ends there exists a diffeomorphism $\Phi: M_{\text{end}}^i \rightarrow \{x \in \mathbb{R}_+^3 \mid |x| > 1\}$ and such that in the coordinate system given by this diffeomorphism we have the following asymptotic as $r \rightarrow \infty$:

$$|\partial^l(g_{ij} - \delta_{ij})| = O(r^{-\tau-l}) \quad (1)$$

for $l = 0, 1, 2$. Here $r = |x|$ and δ is the Euclidean metric on $\mathbb{R}_+^3 = \{x \in \mathbb{R}^3 \mid x_3 \geq 0\}$.

In the following, we will often use the Einstein summation convention with the index ranges $i, j, \dots = 1, \dots, 3$ and $\alpha, \beta, \dots = 1, 2$. Note that, along ∂M , $\{\partial_\alpha\}_\alpha$ spans $T_{\partial M}$, while ∂_n points inwards.

Definition 2.7. If R and H are integrable over M and ∂M respectively and $\tau > 1/2$, then the *mass* of each end of M is well defined and given by

$$\begin{aligned} \mathfrak{m}_{(M_{\text{end}}^i, g)} &= \frac{1}{16\pi} \lim_{r \rightarrow \infty} \left(\int_{\cap \mathbb{S}_{r,+}^2} [g_{ij,j} - g_{jj,i}] \mu^i dA \right), \\ &\quad + \int_{\mathbb{S}_r^1} g_{\alpha 3} \theta^\alpha dl \end{aligned}$$

where the integrals are computed in the asymptotically flat chart, $\mathbb{S}_{r,+}^2(0) = \{\mathbb{R}\}_+^3 \cap \mathbb{S}_r^2(0)$ is a large upper coordinate hemisphere with outward unit normal μ , and θ is the outward pointing unit co-normal to $\mathbb{S}_r^1 = (\{\mathbb{R}\}^2 \times \{0\}) \cap \mathbb{S}_r^2(0) = \partial \mathbb{S}_{r,+}^2$, oriented as the boundary of $(\partial M)_r \subset \partial M$




Figure 1: An asymptotically flat half space with horizon boundary and a large coordinate sphere (from Definition 2.7)

Remark 2.8. In the definition above, the factor $1/(16\pi)$ is a normalization factor used also for the ADM mass of asymptotically flat manifolds with the full \mathbb{R}^n as a model space, where it ensures that we recover the mass of the Schwarzschild solution.

Note that thus in our case of asymptotically flat half-space, the mass of a *half Schwarzschild space* $M_m = \{x \in \mathbb{R}^3_+ \mid |x| > (m/2)^{1/(n-2)}\}$ with conformal metric

$$g_m = \left(1 + \frac{m}{2|x|}\right)^2 \delta, \quad m > 0$$

will be

$$\mathfrak{m}_{(M_m, g_m, =)} = \frac{m}{2},$$

which is half the ADM mass of the standard Schwarzschild space.

In [ABL16], Almaraz, Barbosa, and de Lima showed that this mass is well defined and a geometric invariant, and, in fact, non-negative under suitable energy conditions:

Theorem 2.9. *For (M, g) as above in Definition 2.7, if $R \geq 0$ and $H \geq 0$ on M and ∂M respectively, then*

$$\mathfrak{m}_{(M, g)} \geq 0,$$

with equality occurring if and only if (M, g) is isometric to (\mathbb{R}^3_+, δ) .

For the positive mass theorem on 3-dimensional asymptotically flat manifolds modeled on the full \mathbb{R}^3 , recently [Bra+19] a new method using harmonic functions has been used to achieve a relatively elementary proof of the above theorem and in particular an explicit lower bound for the mass. This thesis will attempt to establish an equivalent result for the case of asymptotically flat half spaces. We will need two further definitions adopted from [EK23] to understand the statement of our main result.

Definition 2.10. Let $\Sigma \subset M$ be a compact separating hypersurface satisfying $\partial\Sigma = \Sigma \cap \partial M$ with normal n_Σ pointing towards the closure $M(\Sigma)$ of the noncompact component of $M \setminus \Sigma$. We call a connected component Σ_0 of M *closed* if $\partial\Sigma_0 = \emptyset$ or a *free boundary hypersurface* if $\partial\Sigma_0 \neq \emptyset$ and $n_{\Sigma_0}(x) \in T_x \partial M$ for every $x \in \Sigma_0 \cap \partial M = \partial\Sigma_0$ (i.e. if Σ_0 meets ∂M orthogonally along its boundary).

We say that an (M, g) has horizon boundary Σ if Σ is a non-empty compact minimal (i.e. having zero mean curvature) hypersurface, whose connected components are all either closed or free boundary hypersurfaces such that $M(\Sigma) \setminus \Sigma$ contains no minimal closed or free boundary hypersurfaces. Σ is also called an *outer most minimal surface* and the region $M(\Sigma)$ outside Σ is called an *exterior region*.

Remark 2.11. By [Koe20, Lemma 2.3] if $H \geq 0$ on ∂M , then there either exists a unique horizon boundary $\Sigma \subset M$ or contains no compact hypersurfaces.

The main result of this thesis is then the following, which will prove Theorem 2.9 as a corollary:

Theorem 2.12. *For (M, g) as above in Definition 2.7, if $R \geq 0$ and $H \geq 0$ on M and ∂M respectively, then there exists a unique harmonic function u asymptotic to the linear function x_3 and satisfying zero Dirichlet boundary conditions on ∂M and zero Neumann boundary conditions on the horizon boundary Σ , and we have*

$$\mathfrak{m}_{(M, g)} \geq \frac{1}{16\pi} \int_{M(\Sigma)} \left(\frac{|\nabla^2 u|^2}{|\nabla u|} + R(g)|\nabla u| \right) dV + \frac{1}{16\pi} \int_{\partial M \cap M(\Sigma)} H(\partial M, g) |\nabla u| dA \geq 0.$$

Proposition 3.8 in „A positive mass theorem for asymptotically flat manifolds with a non-compact boundary“ is super important! We have nice harmonic functions which are themselves asymptotically flat coordinates. 3.9 then gives uniqueness, which we can use along with a doubling argument along the horizon boundary to show existence of asymptotically flat coordinates for the case with non empty horizon boundary.

3 Main basic identity

The basic identity underlying the method of harmonic functions is the following (this is a special case of [Proposition 3.2 in HKK21] and slightly more general than [Bra+19, Proposition 4.2 in]):

Proposition 3.1. *Let (Ω, g) be a compact 3-dimensional manifold with boundary $\partial\Omega$ (smooth almost everywhere), having outward unit normal n . Let $u: \Omega \rightarrow \mathbb{R}$ be a harmonic function (i.e. $\Delta_g u = 0$), and denote the open subset of $\partial\Omega$ on which $|\nabla u| \neq 0$ by $\partial_{\neq 0}\Omega$. If \bar{u} and \underline{u} denote the maximum and minimum of u and Σ_t are t -level sets of u , then*

$$\int_{\partial_{\neq 0}\Omega} \partial_n |\nabla u| dA \geq \int_{\underline{u}}^{\bar{u}} \int_{\Sigma_t} \left(\frac{1}{2} \frac{|\nabla^2 u|^2}{|\nabla u|^2} + R \right) dA dt.$$

4 Existence and uniqueness of asymptotically linear harmonic functions

To use Proposition 3.1, we will require harmonic functions with properties as in Theorem 2.12. More specifically we will require asymptotically linear harmonic coordinates on $M(\Sigma)$ (for Σ the horizon boundary of M) with certain boundary conditions, i.e. $\tilde{x}_1, \tilde{x}_2, \tilde{x}_3$ such that

$$\begin{aligned} \Delta \tilde{x}_i &= 0 & \partial_{n_\Sigma} \tilde{x}_i &= 0 \text{ on } \Sigma & \tilde{x}_i - x_i &\in C_{1-\tau+\varepsilon}^{2,\alpha} & i = 1, 2, 3 \\ \partial_\nu \tilde{x}_\alpha &= 0 \text{ on } \partial M \cap M(\Sigma), & \text{for } \alpha &= 1, 2 & \tilde{x}_3 &= 0 \text{ on } \partial M \cap M(\Sigma) \end{aligned}$$

for some $\varepsilon > 0$ and $0 < \alpha < 1$.

TODO Explain weighted Hölder spaces.

Todo

To prove the following proposition in generality, we will require a special case of it, but with multiple ends. For simplicity we will thus state the whole proposition for manifolds with multiple ends.

Proposition 4.1. *Suppose (M, g) is an asymptotically flat half-space with decay-rate $\tau > 1/2$, asymptotically flat coordinates $\{x_i\}^j$ in each end M_{ext}^j and horizon boundary Σ . Assume (e.g. by shrinking the ends a bit) that the closures of the ends M_{ext}^j are disjoint. Then there exist (up to constant shift) unique smooth functions $\{\tilde{x}_i\}: M(\Sigma) \rightarrow \mathbb{R}$ satisfying*

$$\begin{cases} \Delta \tilde{x}_\beta = 0 & \text{in } M(\Sigma), \\ \partial_\nu \tilde{x}_\beta = 0 & \text{on } \partial M \cap M(\Sigma), \\ \partial_{n_\Sigma} \tilde{x}_\beta = 0 & \text{on } \Sigma, \end{cases}$$

for $\beta = 1, 2$,

$$\begin{cases} \Delta \tilde{x}_3 = 0 & \text{in } M(\Sigma), \\ \tilde{x}_3 = 0 & \text{on } \partial M \cap M(\Sigma), \\ \partial_{n_\Sigma} \tilde{x}_3 = 0 & \text{on } \Sigma, \end{cases}$$

and

$$x_i^j - \tilde{x}_i \in C_{1-\tau+\varepsilon}^{2,\alpha} \quad \text{in } M_{\text{ext}}^j.$$

Moreover, for each end M_{ext}^j , the functions $\{\tilde{x}_i\}$ form an asymptotically flat coordinate system in a neighborhood of infinity.

TODO Maybe change the wording here to align with our wording elsewhere?

Todo

Proof. We first show existence and uniqueness for the case $\Sigma = \emptyset$, then extend to $\Sigma \neq \emptyset$ via a reflection argument along Σ .

Step 1. [ABL16, Proposition 3.8] proves existence for $\Sigma = \emptyset$ and one end. But by replacing the x_i in the proof with arbitrary smooth extensions of the x_i^j (which are defined on open sets with disjoint closures) with $x_i|_{M_{\text{ext}}^j} = x_i^j$, $x_3|_{\partial M} = 0$, we can easily generalise the statement to multiple ends.

We want to show uniqueness for the case $\Sigma = \emptyset$. Let $\{\tilde{x}_i\}$ and $\{\tilde{x}'_i\}$ be two harmonic coordinates fulfilling all the properties. By [ABL16, Proposition 3.9] (the proof of which extends without changes to the case with multiple ends), there exist an orthogonal matrix $(Q_i^j)_{i,j=1}^3$ and constants $\{a_i\}_{i=1}^3$, such that

$$\tilde{x}_i = Q_i^j \tilde{x}'_j + a_i.$$

We have

$$(\delta_i^j - Q_i^j) \tilde{x}_j - a_i = \tilde{x}_i - \tilde{x}'_i = o(r^{1/2}) \quad \text{as } r \rightarrow \infty,$$

and further

$$(\delta_i^j - Q_i^j)(x_j - \tilde{x}_j) - a_i = o(r^{1/2})$$

which implies

$$(\delta_i^j - Q_i^j)x_j = o(r^1)$$

and thus we must have $Q_i^j = \delta_i^j$ (since otherwise $(\delta_i^j - Q_i^j)x_j$ would be linear). Hence

$$\tilde{x}_i = \tilde{x}'_i + a_i.$$

Note that $a_3 = 0$, since $\tilde{x}_i = 0 = \tilde{x}'_i$ on ∂M .

Step 2. Consider now the case $\Sigma \neq \emptyset$. We adapt the proof of [EK23, Proposition 46].

Consider the differentiable manifold $(\hat{M} = M \times \{-1, +1\})/\sim$, where $(x, \pm 1) \sim (x, \mp 1)$ if and only if $x_1, x_2 \in \Sigma$ and $x_1 = x_2$ (i.e. \hat{M} is constructed by gluing two copies of M along Σ). We equip \hat{M} with the Riemannian metric $\hat{g}(\hat{x}) = \gamma(\pi(\hat{x}))$, where $\pi([(x, \pm 1)]) = x$.

Then by [EK23, Lemma 19], \hat{g} is of class C^2 away from $\pi^{-1}(\Sigma)$ and on $\pi^{-1}(\Sigma)$ the coefficients of $\Delta_{\hat{g}}$ are Lipschitz since Σ is minimal.

Note that \hat{M} has twice as many ends as M , where we set $\hat{x}_i^{j,\pm}(\hat{x}) = x_i^j(\pi(x))$ to be the asymptotic coordinates in these ends.

We can thus apply the result from Step 1 to \hat{M} (for which we don't consider any boundary conditions on horizon boundaries) to obtain asymptotically linear harmonic coordinates \hat{x}_i on \hat{M} with Dirichlet boundary condition on $\partial\hat{M}$. But note that $\hat{x}_i \circ \tau$ is another solution, where we let $\tau: \hat{M} \rightarrow \hat{M}$ be given by $\tau([(x, \pm 1)]) = [(x, \mp 1)]$. Then by the already established uniqueness for the case without horizon boundary, $\hat{x}_i \circ \tau = \hat{x}_i + a_i$ for some constants a_i . But since these must agree on $\pi^{-1}(\Sigma)$ (τ is the identity there), we have $a_i = 0$.

In particular we get on $\pi^{-1}(\Sigma)$

$$\partial_{n_\Sigma} \hat{x}_i = -\partial_{n_\Sigma}(\hat{x}_i \circ \tau) = -\partial_{n_\Sigma}(\hat{x}_i) = -\partial_{n_\Sigma}(\hat{x}_i)$$

and thus \hat{x} satisfies Neumann boundary conditions on $\pi^{-1}(\Sigma)$ (here we need to fix n_Σ , e.g. choose it to point towards $M \times \{+1\}$).

In particular we get a solution to our original problem on M by setting $\tilde{x}_i(x) = \hat{x}_i([(x, +1)])$.

The argument from [EK23, Proposition 3.9] extends straightforwardly to also show uniqueness (up to adding constants) for the \tilde{x}_i on $M(\Sigma)$.

■

5 Proof of the Mass Lower Bound

We proceed by constructing a proof parallel to [Bra+19, Section 6] and [HKK21, Section 6].

To this end, let (M, g) be an asymptotically flat half-space and horizon boundary Σ with asymptotically flat harmonic coordinates x_1, x_2, x_3 as in proposition 4.1. Note

that from now on we will again consider M to only have a single end M_{ext} and that although we call x_1, x_2, x_3 are defined on all of $M(\Sigma)$ and we call them harmonic coordinates, they are only guaranteed to form a coordinate system in M_{ext} .

TODO The above wording is very similar to [HKK21, Section 6].

Todo

By [ABL16, Proposition 3.7], we can compute the mass in these harmonic coordinates. For $L > 0$ define coordinate half-cylinders $C_L = D_L \cup T_L$ given by

$$\begin{aligned} D_L &= \{x \in M_{\text{ext}} \mid (x_1)^2 + (x_2)^2 \leq L^2, \ x_3 = L\} \\ T_L &= \{x \in M_{\text{ext}} \mid (x_1)^2 + (x_2)^2 = L^2, \ 0 \leq x_3 \leq L\}. \end{aligned}$$

Further define

$$\begin{aligned} \mathbb{S}_L^1 &= \{x \in M_{\text{ext}} \mid (x_1)^2 + (x_2)^2 = L^2, \ 0 = x_3\} = \partial C_L = C_L \cap \partial M \\ (\partial M)_L &= \{x \in \partial M \cap M(\Sigma) \mid (x_1)^2 + (x_2)^2 \leq L\} \end{aligned}$$

and let Ω_L be the compact component of $M(\Sigma) \setminus C_L$. By an argument similar to the proof of [Bar86, Proposition 4.1], we can compute the mass as

$$\mathfrak{m}_{(M,g)} = \lim_{L \rightarrow \infty} \left(\int_{C_L} (g_{ij,j} - g_{jj,i}) \mu^i dA + \int_{\mathbb{S}_L^1} g_{\alpha 3} \theta^\alpha dl \right)$$

where now μ is the outward unit normal to C_L and θ is as in *definition 2.7*. We delegate the details here to the Appendix, see proposition A.1.

A Different Exhausting Sequences for Computation of the Mass

Proposition A.1. *Suppose that (M, g) is an asymptotically flat half space with asymptotically flat coordinates x_1, x_2, x_3 . Let $\{D_k^3\}_1^\infty$ be an exhaustion of M by closed sets such that the sets $S_k = \partial D_k$ are connected 2-dimensional submanifolds (smooth almost everywhere) of M with $\partial S_k = \partial M \cap S_k$ such that*

$$R_k := \inf\{|x| \mid x \in S_k\} \rightarrow \infty \quad \text{as } k \rightarrow \infty,$$

$$R_k^2 \cdot |S_k| \text{ is bounded as } k \rightarrow \infty,$$

and $R_1 \geq R_0$, where $|S_k|$, the area of S_k , and $|x|$ are as usual calculated with respect to the euclidean background metric. Then

$$\mathfrak{m}_{(M,g)} = \lim_{k \rightarrow \infty} \int_{S_k} \int_{\mathbb{S}_{r,+}^2} [g_{ij,j} - g_{jj,i}] \mu^i dA + \int_{\partial S_k} g_{\alpha 3} \theta^\alpha dl$$

is independent of the sequence S_k , where as in definition 2.7 μ^i is the outward normal to S_k and θ^α the co-normal to ∂S_k oriented as the boundary of the compact component of $\partial M \setminus \partial S_k$.

Proof. **TODO** Prove this

■ Todo

References

- [ABL16] Sérgio Almaraz, Ezequiel Barbosa, and Levi Lopes de Lima. „A Positive Mass Theorem for Asymptotically Flat Manifolds with a Non-Compact Boundary“. In: *Communications in Analysis and Geometry* 24.4 (2016), pp. 673–715. ISSN: 10198385, 19449992. DOI: 10.4310/CAG.2016.v24.n4.a1. URL: <http://www.intlpress.com/site/pub/pages/journals/items/cag/content/vols/0024/0004/a001/> (visited on 03/22/2022).
- [Bar86] Robert Bartnik. „The Mass of an Asymptotically Flat Manifold“. In: *Communications on Pure and Applied Mathematics* 39.5 (Sept. 1986), pp. 661–693. ISSN: 00103640, 10970312. DOI: 10.1002/cpa.3160390505. URL: <https://onlinelibrary.wiley.com/doi/10.1002/cpa.3160390505> (visited on 04/24/2023).
- [Bra+19] Hubert L. Bray et al. „Harmonic Functions and The Mass of 3-Dimensional Asymptotically Flat Riemannian Manifolds“. Comment: 22 pages, 5 figures. Nov. 15, 2019. arXiv: 1911.06754 [gr-qc]. URL: <http://arxiv.org/abs/1911.06754> (visited on 03/22/2022).

-
- [EK23] Michael Eichmair and Thomas Koerber. „Doubling of Asymptotically Flat Half-spaces and the Riemannian Penrose Inequality“. In: *Communications in Mathematical Physics* (Jan. 31, 2023). ISSN: 0010-3616, 1432-0916. DOI: 10.1007/s00220-023-04635-7. URL: <https://link.springer.com/10.1007/s00220-023-04635-7> (visited on 04/30/2023).
- [HKK21] Sven Hirsch, Demetre Kazaras, and Marcus Khuri. „Spacetime Harmonic Functions and the Mass of 3-Dimensional Asymptotically Flat Initial Data for the Einstein Equations“. Comment: J. Differential Geom., to appear. Jan. 16, 2021. arXiv: 2002.01534 [gr-qc, physics:hep-th]. URL: <http://arxiv.org/abs/2002.01534> (visited on 03/22/2022).
- [Koe20] Thomas Koerber. *The Riemannian Penrose Inequality for Asymptotically Flat Manifolds with Non-Compact Boundary*. Comment: 42 pages, some minor changes and corrected typos, all comments welcome. Jan. 14, 2020. arXiv: 1909.13283 [math]. URL: <http://arxiv.org/abs/1909.13283> (visited on 06/09/2023). preprint.
- [Lee19] Dan A. Lee. *Geometric Relativity*. Graduate Studies in Mathematics volume 201. Providence, Rhode Island: American Mathematical Society, 2019. 361 pp. ISBN: 978-1-4704-5081-6.