

Note

Feng-Yang Hsieh

1 ATLAS latest study

The couplings of longitudinally polarized W and Z bosons to the Higgs boson prevent the divergence of tree-level Vector Boson Scattering (VBS) amplitudes at high energies when the two outgoing bosons are longitudinally polarized, restoring unitarity at the TeV scale. Thus, the VBS processes with longitudinal vector bosons provide a sensitive test of the electroweak (EW) symmetry-breaking mechanism.

The most recent ATLAS study is detailed in Ref. [1]. This study focuses on VBS. Specifically, it investigates the EW production of same-sign W boson pairs (EW $W^\pm W^\pm jj$), a process to which VBS provides a significant contribution. The key concepts and event selection criteria of this work are summarized below.

The object selection criteria are as follows:

- **Electrons:** Must have transverse momentum $p_T > 27$ GeV and pseudorapidity $|\eta| < 2.47$, excluding the calorimeter transition region of $1.37 < |\eta| < 1.52$.
- **Muons:** Must have $p_T > 27$ GeV and $|\eta| < 2.5$.
- **Jets:** At least two jets with $|\eta| < 4.5$ are required in each event. The leading jet must have $p_T > 65$ GeV, and the subleading jet must have $p_T > 35$ GeV.

The Signal Region (SR) event selection requires:

- A same-sign lepton pair with an invariant mass $m_{\ell\ell} > 20$ GeV.
- Missing transverse momentum magnitude, E_T^{miss} , greater than 30 GeV.
- The two highest- p_T jets must satisfy a dijet invariant mass $m_{jj} > 500$ GeV with an absolute rapidity difference $|\Delta y_{jj}| > 2.0$.

2 Monte Carlo simulation

2.1 Sample Generation

The signal processes are the electroweak (EW) production of $W^\pm W^\pm jj$ in various polarization states. We simulate the EW same-sign W boson pair production at a center-of-mass energy of $\sqrt{s} = 13$ TeV. The events are generated using **MadGraph 3.3.1** [2]. Both W bosons must decay leptonically. Parton showering and hadronization are simulated with **Pythia 8.306** [3], followed by a fast detector simulation using **Delphes 3.4.2** [4]. Jets are reconstructed via the anti- k_t algorithm [5] with a distance parameter of $R = 0.4$, implemented in **FastJet 3.3.2** [6]. Reconstructed jets are required to have a transverse momentum of $p_T > 25$ GeV.

The following **MadGraph** scripts are used to generate the Monte Carlo (MC) samples for these polarized processes:

Longitudinal mode: $pp \rightarrow W_L^\pm W_L^\pm jj$

```
generate p p > j j w+{0} w+{0} QCD=0, w+ > l+ vl  
add process p p > j j w-{0} w-{0} QCD=0, w- > l- vl~  
output EW_WWjj_LL  
launch EW_WWjj_LL
```

```
shower=Pythia8  
detector=Delphes  
analysis=OFF  
madspin=OFF  
done
```

Cards/delphes_card.dat

```
set run_card nevents 10000  
set run_card ebeam1 6500.0  
set run_card ebeam2 6500.0  
  
set run_card ptj 34.0  
set run_card ptl 26.0  
set run_card misset 29.0
```

```

set run_card etaj 4.6
set run_card etal 2.6

set run_card mmjj 480
set run_card deltaeta 1.9

set run_card ptj1min 64
set run_card ptj2min 34

set run_card use_syst False

done

```

Transverse mode: $pp \rightarrow W_T^\pm W_T^\pm jj$

```

generate p p > j j w+{T} w+{T} QCD=0, w+ > l+ vl
add process p p > j j w-{T} w-{T} QCD=0, w- > l- vl~
...

```

Mixed mode: $pp \rightarrow W_L^\pm W_T^\pm jj$

```

generate p p > j j w+{0} w+{T} QCD=0, w+ > l+ vl
add process p p > j j w-{0} w-{T} QCD=0, w- > l- vl~
...

```

2.2 Event Selection

The event selection criteria applied after the **Delphes** simulation are defined as follows:

- **Lepton cut:** The invariant mass of the same-sign lepton pair, $m_{\ell\ell}$, must be greater than 20 GeV.
- **MET cut:** The missing transverse momentum magnitude, E_T^{miss} , must be greater than 30 GeV.
- **Jet cut:** The two highest- p_T tagging jets must satisfy a dijet invariant mass of $m_{jj} > 500$ GeV with an absolute rapidity difference of $|\Delta y_{jj}| > 2.0$.

Table 1 summarizes the cutflow (number of expected events and the corresponding selection efficiencies) at each selection stage. The reconstruction requirements for electrons, muons, and jets are identical to those described in Section 1.

Table 1: Number of events and selection efficiencies (pass rates) for EW $W^\pm W^\pm jj$ production at different selection stages.

Cut	LL	pass rate	LT	pass rate	TT	pass rate
Total	100000	1.00	100000	1.00	100000	1.00
Lepton cut	26777	0.27	27270	0.27	29825	0.30
MET cut	24781	0.25	25518	0.26	28323	0.28
Jet cut	20005	0.20	21087	0.21	23298	0.23

Figure 1 displays the kinematic distributions for the various polarization states in the SR. Specifically, we present the absolute pseudorapidity difference between the leading and subleading leptons $|\Delta\eta_{\ell\ell}|$, the transverse mass of the dilepton and E_T^{miss} system m_T , and the azimuthal angle difference between the leading and subleading jets $\Delta\phi_{jj}$. The ratios of the individual process contributions to the total signal are shown in the bottom panels. For these combined distributions, we assume that the $W_L^\pm W_L^\pm jj$, $W_L^\pm W_T^\pm jj$, and $W_T^\pm W_T^\pm jj$ states contribute 10%, 30%, and 60% to the total signal, respectively.




Figure 1: Kinematic distributions of $|\Delta\eta_{\ell\ell}|$, m_T , and $\Delta\phi_{jj}$ in the signal region. The ratios of the contributions from different polarization states are shown in the bottom panels.

2.3 Absolute Cross-Sections

Table 2 summarizes the absolute cross-sections for the EW $W^\pm W^\pm jj$ production across different selection stages. To validate our simulation framework, the final cross-sections after

the jet cut are compared with the reference results from a previous thesis [7].

Table 2: Absolute cross-sections (in fb) for the EW $W^\pm W^\pm jj$ production at various selection stages. The reference values from the previous thesis and the relative fractions are provided at the final jet cut stage for comparison.

Cut	LL	LT	TT	Total
Total	0.443	2.514	3.824	6.781
Lepton cut	0.119	0.685	1.140	1.945
MET cut	0.110	0.641	1.083	1.834
Jet cut	0.089	0.530	0.891	1.510
Jet cut (Thesis)	0.177	1.030	1.718	2.925
Fraction	5.9%	35.1%	59.0%	100%
Fraction (Thesis)	6.1%	35.2%	58.7%	100%

As observed in Table 2, while the relative fractions of the polarization states are in excellent agreement with the reference thesis, the absolute cross-sections in our simulation are approximately a factor of two lower. Further investigation is needed.

2.4 WW Center-of-Mass Reference Frame

To properly study the scattering dynamics, we modify our **MadGraph** scripts to define the W boson polarizations in the $W^\pm W^\pm$ center-of-mass (CM) frame.

The following generation commands are used to produce the MC samples with this specific reference frame definition:

```

Longitudinal mode:  $pp \rightarrow W_L^\pm W_L^\pm jj$ 
generate p p > w+{0} w+{0} j j QCD=0, w+ > l+ vl
add process p p > w-{0} w-{0} j j QCD=0, w- > l- vl~
output MG5/EW_WWjj_LL-WW_cmf
launch MG5/EW_WWjj_LL-WW_cmf

shower=Pythia8
detector=Delphes
analysis=OFF
madspin=OFF
done

```

Cards/delphes_card.dat

```
set run_card nevents 10000
set run_card ebeam1 6500.0
set run_card ebeam2 6500.0
```

```
set run_card ptj 34.0
set run_card etaj 4.6
set run_card mmjj 480
set run_card deltaeta 1.9
```

```
set run_card etal 2.6
set run_card ptl 26.0
set run_card misset 29.0
```

```
set run_card ptj1min 64
set run_card ptj2min 34
```

```
set run_card me_frame 3, 4, 5, 6
```

```
set run_card use_syst False
```

done

Transverse mode: $pp \rightarrow W_T^\pm W_T^\pm jj$

```
generate p p > w+{T} w+{T} j j QCD=0, w+ > l+ vl
add process p p > w-{T} w-{T} j j QCD=0, w- > l- vl~
...
```

Mixed mode: $pp \rightarrow W_L^\pm W_T^\pm jj$

```
generate p p > w+{0} w+{T} j j QCD=0, w+ > l+ vl
add process p p > w-{0} w-{T} j j QCD=0, w- > l- vl~
...
```

The key modification lies in the `me_frame` setting. More details can be found in ref. [8]. It is crucial to note that when utilizing the in-line decay syntax, **MadGraph** expands the decay products directly into the final-state particle list. Consequently, the indices must be specified as 3, 4, 5, 6 to encompass all the decay products (leptons and neutrinos) required to reconstruct the WW system.

The remaining shower, detector, and run card settings for the transverse and mixed modes are identical to those of the longitudinal mode and are omitted here for brevity.

Table 3 summarizes the absolute cross-sections for the EW $W^\pm W^\pm jj$ production when the polarizations are defined in the $W^\pm W^\pm$ CM frame. The absolute cross-sections are still a factor of two lower.

Table 3: Absolute cross-sections for the EW $W^\pm W^\pm jj$ production at various selection stages, with W boson polarizations defined in the $W^\pm W^\pm$ center-of-mass frame. Reference values from the previous thesis and relative fractions are provided at the final jet cut stage for comparison.

Cut	LL [fb]	LT [fb]	TT [fb]	Total [fb]
Total	0.677	2.190	3.889	6.756
Lepton cut	0.190	0.577	1.143	1.910
MET cut	0.177	0.535	1.081	1.793
Jet cut	0.147	0.443	0.895	1.485
Jet cut (Thesis)	0.277	0.874	1.744	2.869
Fraction	9.4%	30.4%	60.2%	100%
Fraction (Thesis)	9.6%	30.6%	59.8%	100%

Figure 2 displays the kinematic distributions for the various polarization states in the SR within the WW center-of-mass frame. These results are consistent with Figure 2 in ref. [1].

2.5 Simulation with Sherpa

To investigate the cross-section discrepancy, we employ a different Monte Carlo generator, **Sherpa** 3 [9], which is also the primary generator utilized in the ATLAS analysis.

We simulate the EW same-sign W boson pair production at a center-of-mass energy of $\sqrt{s} = 13$ TeV. Both W bosons must decay leptonically. Parton showering and hadronization are handled internally by **Sherpa**, followed by a fast detector simulation using **Delphes** 3.4.2 [4]. Jets are reconstructed via the anti- k_t algorithm [5] with a distance parameter of $R = 0.4$,




Figure 2: Kinematic distributions of $|\Delta\eta_{\ell\ell}|$, m_T , and $\Delta\phi_{jj}$ in the signal region. The ratios of the contributions from different polarization states are shown in the bottom panels.

implemented in **FastJet** 3.3.2 [6]. The reconstructed jets are required to have a transverse momentum of $p_T > 25$ GeV.

Unlike **MadGraph**, which can generate pure polarized samples, **Sherpa** generates events and assigns polarization weights to each of them. These weights represent the probability of an event belonging to a specific polarization state. Due to technical constraints within the generator, the W^+W^+jj and W^-W^-jj processes must be generated as separate samples.

The following YAML configuration snippet illustrates the generation of the W^+W^+jj sample using **Sherpa**:

W^+W^+jj sample

EVENTS: 100

RANDOM_SEED: 1

BEAMS: 2212

BEAM_ENERGIES: 6500

PDF_SET: NNPDF30_nlo_as_0118

ME_GENERATORS:

- Comix

COMIX_DEFAULT_GAUGE: 0

scales


```

MEPS:
  CORE_SCALE: VAR{0.5*Abs2(p[2]+p[3])}

# physical constants and width settings
PARTICLE_DATA:
  24: {Width: 0} # W Boson
  23: {Width: 0} # Z Boson
WIDTH_SCHEME: Fixed

# decay and polarization settings
HARD_DECAYS:
  Enabled: true
  Channels:
    24,12,-11: {Status: 2} # W+ -> nu_e e+
    24,14,-13: {Status: 2} # W+ -> nu_mu mu+
  Pol_Cross_Section:
    Enabled: true
    Reference_System: [Lab, COM]

# process definition: same-sign W boson pair production with two jets
PROCESSES:
  - 93 93 -> 24 24 93 93:
    Order: {QCD: 0, EW: 4}

# final state particles cuts
SELECTORS:
  - [PT, 93, 34, 10000]
  - [Eta, 93, -4.6, 4.6]
  - [Mass, 93, 93, 480, 10000]
  - [DY, 93, 93, 1.9, 10000]

  - VariableSelector:
    Variable: PT
    Flavs: 93
    Ranges:
      - [60.0, 10000.0]

```

```

    Ordering: [PT_UP]

# output settings
EVENT_OUTPUT: HepMC3_GenEvent[EW_WpWpjj.hepmc]

W-W-jj sample
EVENTS: 100
RANDOM_SEED: 1

BEAMS: 2212
BEAM_ENERGIES: 6500

PDF_SET: NNPDF30_nlo_as_0118

ME_GENERATORS:
  - Comix
COMIX_DEFAULT_GAUGE: 0

# scales
MEPS:
  CORE_SCALE: VAR{0.5*Abs2(p[2]+p[3])}

# physical constants and width settings
PARTICLE_DATA:
  24: {Width: 0} # W Boson
  23: {Width: 0} # Z Boson
WIDTH_SCHEME: Fixed

# decay and polarization settings
HARD_DECAYS:
  Enabled: true
  Channels:
    -24,-12,11: {Status: 2} # W- -> nubar e-
    -24,-14,13: {Status: 2} # W- -> nubar mu-
  Pol_Cross_Section:

```

```

Enabled: true
Reference_System: [Lab, COM]

# process definition: same-sign W boson pair production with two jets
PROCESSES:
- 93 93 -> -24 -24 93 93:
    Order: {QCD: 0, EW: 4}

# final state particles cuts
SELECTORS:

- [PT, 93, 34, 10000]
- [Eta, 93, -4.6, 4.6]
- [Mass, 93, 93, 480, 10000]
- [DY, 93, 93, 1.9, 10000]

- VariableSelector:
    Variable: PT
    Flavs: 93
    Ranges:
        - [60.0, 10000.0]
    Ordering: [PT_UP]

# output settings
EVENT_OUTPUT: HepMC3_GenEvent[EW_WmWmjj.hepmc]

```

It is important to note that the decay channel specification in **Sherpa** is strictly sensitive. The particle ordering must match the internal definitions precisely; for instance, 24,12,-11: {Status: 2} is valid, whereas 24,-11,12: {Status: 2} will not be recognized.

Table 4 details the absolute cross-sections for the individual W^+W^+jj and W^-W^-jj samples at various selection stages, with the polarization states defined in the $W^\pm W^\pm$ center-of-mass frame.

The combined absolute cross-sections for the EW $W^\pm W^\pm jj$ production are summarized in table 5. While the relative fractions of the polarization states remain in excellent agreement with the reference thesis, the combined absolute cross-section generated by **Sherpa**

Table 4: Absolute cross-sections for the separate W^+W^+jj and W^-W^-jj samples at various selection stages simulated with **Sherpa**. Polarization states are defined in the $W^\pm W^\pm$ center-of-mass frame.

Stage	W^+W^+jj [fb]				W^-W^-jj [fb]			
	LL	LT	TT	Total	LL	LT	TT	Total
Total	0.527	1.633	2.949	5.108	0.162	0.475	0.850	1.487
Lepton cut	0.153	0.441	0.863	1.457	0.048	0.143	0.295	0.486
MET cut	0.132	0.394	0.776	1.302	0.040	0.122	0.250	0.412
Jet cut	0.106	0.319	0.633	1.057	0.030	0.094	0.199	0.323

(1.380 fb at the jet cut stage) is still approximately a factor of two lower than the reference value (2.933 fb).

Table 5: Combined absolute cross-sections for the EW $W^\pm W^\pm jj$ production simulated with **Sherpa**. Reference values from the previous thesis and relative fractions are provided at the final jet cut stage for comparison.

Cut	LL [fb]	LT [fb]	TT [fb]	Total [fb]
Total	0.688	2.108	3.799	6.595
Lepton cut	0.202	0.583	1.158	1.943
MET cut	0.172	0.516	1.026	1.714
Jet cut	0.136	0.413	0.831	1.380
Jet cut (Thesis)	0.285	0.909	1.778	2.933
Fraction	9.9%	29.9%	60.2%	100%
Fraction (Thesis)	9.6%	30.6%	59.8%	100%

2.6 Particle-Level Selection

Upon a detailed review of the reference thesis [7], we identified that their reported yields and cross-sections are evaluated at the particle level rather than the reconstructed level. To perform a fair comparison, we modified our event selection framework to operate on particle-level objects.

Specifically, the variables accessed in the **Delphes** analysis were updated as follows:

- **Leptons:** Reconstructed leptons (**Electron** and **Muon** collections) are replaced by generator-level stable final-state leptons from the **Particle** branch.

- **Missing Transverse Momentum:** The reconstructed missing transverse energy (`MissingET`) is replaced by the true missing transverse momentum (`GenMissingET`).
- **Jets:** Reconstructed jets (`Jet`) are replaced by truth-level jets (`GenJet`).

Furthermore, the kinematic selection criteria at the particle level are different from the reconstructed-level SR selection defined in Section 2.2. Based on the reference thesis, the particle-level selection is defined as follows:

- **Leptons:** Exactly two same-sign leptons with $p_T > 27$ GeV. Muons must satisfy $|\eta| < 2.5$. Electrons must satisfy $|\eta| < 2.47$, excluding the transition region $1.37 \leq |\eta| \leq 1.52$, with an additional restriction of $|\eta| < 1.37$ for the ee channel.
- **Dilepton Mass & Z -veto:** The dilepton invariant mass must be $m_{\ell\ell} > 20$ GeV. Additionally, a Z -boson veto is applied to the ee channel, requiring $|m_{ee} - m_Z| > 15$ GeV to suppress background contributions.
- **Missing Transverse Momentum:** $E_T^{\text{miss}} \geq 30$ GeV.
- **Jets:** At least two jets are required. The leading and subleading jets must satisfy $p_T > 65$ GeV and $p_T > 35$ GeV, respectively. The dijet invariant mass threshold is relaxed to $m_{jj} \geq 200$ GeV, and the rapidity difference must be $\Delta y_{jj} > 2$.

The PDF sets are also updated to the same as the reference thesis [7]. We use the NNPDF23_nlo_as_0119 PDF set [10] for MadGraph and PDF4LHC21_40_pdfas PDF set [11] for Sherpa.

The updated absolute cross-sections for the EW $W^\pm W^\pm jj$ production at the particle level are summarized in Table 6. Although the relative fractions of the polarization states remain in excellent agreement with the reference thesis, a systematic normalization discrepancy still exists in the absolute cross-sections.

The thesis also mentioned the overlapping removal strategies for the jet and lepton objects. However, it is not clear if they applied the overlapping removal and what kind of strategy they considered.

2.7 Background Processes

In this section, we focus on the primary background sources: the irreducible QCD $W^\pm W^\pm jj$ production and the reducible $W^\pm Zjj$ process.

Table 6: Combined absolute cross-sections for the EW $W^\pm W^\pm jj$ production simulated with MadGraph and Sherpa using **particle-level selection**. Reference values from the previous thesis and relative fractions are provided at the final jet cut stage for comparison.

Cut	MadGraph [fb]				Sherpa [fb]			
	LL	LT	TT	Total	LL	LT	TT	Total
Total	0.383	1.188	2.185	3.755	0.729	2.252	3.989	6.970
Lepton cut	0.267	0.850	1.665	2.782	0.291	0.906	1.746	2.943
MET cut	0.260	0.834	1.636	2.731	0.240	0.786	1.522	2.548
Jet cut	0.205	0.623	1.129	1.957	0.188	0.569	1.021	1.777
Jet cut (Thesis)	0.277	0.874	1.744	2.895	0.285	0.909	1.778	2.972
Fraction	10.5%	31.9%	57.7%	100%	10.6%	32.0%	57.5%	100%
Fraction (Thesis)	9.6%	30.2%	60.2%	100%	9.6%	30.6%	59.8%	100%

2.7.1 QCD $W^\pm W^\pm jj$ Production

The QCD-induced same-sign W boson pair production in association with two jets is an irreducible background, as its final state is identical to that of the signal. At tree level, this process is of order $\mathcal{O}(\alpha_S^2 \alpha^4)$.

The following MadGraph scripts are used to generate the MC samples:

```
generate p p > w+ w+ j j QCD<=2 QED<=2, w+ > l+ vl
add process p p > w- w- j j QCD<=2 QED<=2, w- > l- vl~
...

```

2.7.2 INT $W^\pm W^\pm jj$ Production

The interference (INT) $W^\pm W^\pm jj$ background arises from the quantum interference between the purely electroweak and the QCD-induced amplitudes. At tree level, this process is of order $\mathcal{O}(\alpha_S \alpha^5)$.

2.7.3 $W^\pm Z jj$ Production

The WZ production in association with jets, where $WZ \rightarrow \ell \nu \ell \ell$, is the most prominent reducible background. It mimics the same-sign WW topology when one of the leptons from the Z boson decay is “lost” (e.g., falls outside the detector acceptance or fails identification criteria).

The following MadGraph scripts are used to generate the MC samples:

```
generate p p > w+ z j j QCD=0, w+ > l+ vl, z > l+ l-
add process p p > w- z j j QCD=0, w- > l- vl~, z > l+ l-

...
```

2.7.4 Non-Prompt Leptons

Leptons originating from hadron decays and jets misidentified as leptons are referred to as non-prompt leptons. Since this background is estimated using data-driven methods in the experimental analysis, we omit its estimation in our current Monte Carlo simulation framework.

2.7.5 Full Process vs. Narrow Width Approximation (NWA)

The most rigorous approach to generating these events is to simulate the full $2 \rightarrow 6$ process. For instance, for the EW $W^\pm W^\pm jj$ signal, the full process command would be:

```
generate p p > j j l+ vl l+ vl QCD=0
add process p p > j j l- vl~ l- vl~ QCD=0
```

However, simulating the full process requires evaluating a vast number of Feynman diagrams and integrating over a highly complex phase space. This often leads to severe integration inefficiencies and divergences, making it prohibitive to generate a sufficient number of events.

We investigated the $W^\pm Zjj$ case and found that the full simulation and the Narrow Width Approximation (NWA) approach produce highly consistent absolute cross-sections after applying the SR cuts. Therefore, we temporarily adopt the NWA for our current generation setup to ensure efficiency.

2.7.6 Yields and Comparisons

To validate our simulation framework, the expected event yields are compared against the reference ATLAS results. Table 7 summarizes the expected signal and background yields in the SR. The “ATLAS” column represents the values reported in Table 3 of Ref. [12], the “Thesis” columns represent the values reported in Table 9.1 and Table G.2 of Ref. [7], while the remaining columns represent our simulation results. Here, we normalize our simulated yields to an integrated luminosity of $\mathcal{L} = 139 \text{ fb}^{-1}$.

Table 7: Expected yields in the signal region (SR). The ATLAS results are shown alongside Thesis values and our simulation estimates using different generator versions, assuming an integrated luminosity of 139 fb^{-1} .

Process	ATLAS	Thesis 9.1	Thesis G.2 ¹	MG 3.3.1	MG 3.7.0	Sh 3.0.3
$W^\pm W^\pm jj$ EW	235	206.52	211.29 ²	177.98	190.28	191.18
$W^\pm W^\pm jj$ QCD	24	24.05	24.27	14.38	12.96	19.12
$W^\pm Z jj$ EW	14.9	14.95	15.07	9.12	9.50	32.55
$W^\pm Z jj$ QCD	83	82.75 ³	77.0	18.47	16.30	80.15

2.8 Determination of the Cross-Section for Sherpa Samples

When utilizing **Sherpa** to generate MC samples, there are two primary stages. First, the generator performs a numerical integration over the defined phase space to calculate the total cross-section. Subsequently, it generates events and writes the kinematic information and local cross-section into the resulting **HepMC** file.

In our previous analyses, the cross-section was estimated by extracting the values recorded event-by-event in the **HepMC** file and computing their average. However, the values written to the event record are local estimates derived during the generating stage. Consequently, they are subject to large statistical fluctuations, which can introduce biases into the final normalization.

Conversely, the cross-section reported in the **Sherpa** log output that was evaluated during the initial phase-space integration stage is more suitable. Because this integration stage utilizes vast sampling points before generating events, it reduces statistical fluctuations and yields a more stable cross-section.

Table 8 presents a comparison of the absolute cross-sections (σ_{gen}) and the corresponding expected yields (N_{exp}) in the SR obtained via these two methods: the integration log output (New) and the averaged **HepMC** weights (Old).

As observed in the table, the **HepMC** average consistently overestimates the expected yields by roughly 4% to 9% across various processes. All subsequent analyses will utilize the cross-section obtained from the **Sherpa** generation log.

¹Values from the pre-fit event yields (Table G.2), where the three DNN score bins are merged.

²The sum of the LL, TL, and TT polarization processes.

³The predicted events appear to be scaled down (e.g., $82.75/2 \approx 41.38 \rightarrow 28.50$). The authors note in the caption that an additional normalization factor was applied to achieve a better agreement with the expected data. However, the exact derivation of this factor is not detailed.

Table 8: Comparison of the generation cross-sections (σ_{gen}) and expected yields (N_{exp}) in the SR using the **Sherpa** log output (New) versus the averaged **HepMC** values (Old). The percentage difference highlights the consistent overestimation bias introduced by the **HepMC** method.

Sample	$\sigma_{\text{gen}}^{\text{New}}$ [fb]	$\sigma_{\text{gen}}^{\text{Old}}$ [fb]	$N_{\text{exp}}^{\text{New}}$	$N_{\text{exp}}^{\text{Old}}$	Diff.
$W^\pm W^\pm jj$ EW	7.895	8.689	178.22	196.14	−9.13%
$W^\pm W^\pm jj$ QCD	2.245	2.441	17.88	19.44	−8.01%
$W^\pm Zjj$ EW	8.242	8.585	32.76	34.13	−4.00%
$W^\pm Zjj$ QCD	20.279	21.278	72.44	76.01	−4.70%

2.9 MadGraph Run Card Selection Criteria Issues

To increase the event generation efficiency for the SR phase space, we applied kinematic cuts at the generator level (as detailed in Section 2.4). While these cuts are appropriate for generating pure $W^\pm W^\pm jj$ signal samples, they introduce biases when extending the analysis to Control Regions (CR) or simulating reducible backgrounds such as $W^\pm Zjj$.

For the CR, the analysis investigates the low- m_{jj} region, specifically $200 \text{ GeV} < m_{jj} < 500 \text{ GeV}$. The previous generator-level cut of `set run_card mmjj 480` is problematic, as it truncates almost all valid events that would populate this CR.

Furthermore, for the $W^\pm Zjj \rightarrow \ell \nu \ell \ell$ background, the global lepton transverse momentum cut `set run_card ptl 26.0` causes an underestimation. The `ptl` parameter in **MadGraph** enforces a minimum p_T requirement on *all* generated leptons.

Example: Consider a $W^+ Z \rightarrow \ell^+ \nu \ell^+ \ell^-$ event where the two same-sign leptons have $p_T = 45 \text{ GeV}$ and 35 GeV , while the third lepton (ℓ^-) is soft with $p_T = 15 \text{ GeV}$. In the detector simulation, this soft ℓ^- might be “lost” (e.g., falling outside the acceptance or failing identification), leaving exactly two same-sign leptons that pass the SR cuts ($p_T > 27 \text{ GeV}$). However, because the third lepton’s p_T is lower than the 26.0 GeV threshold, **MadGraph** will discard this event at the generation level. Consequently, this topology is completely lost.

To resolve these issues, the **MadGraph** run card scripts must be updated to use looser, ordered cuts. Taking the QCD $W^\pm Zjj$ process as an example:

```
generate p p > w+ z j j QCD<=2 QED<=2, w+ > l+ vl, z > l+ l-
add process p p > w- z j j QCD<=2 QED<=2, w- > l- vl~, z > l+ l-
output MG5/WZjj_QCD-NWA-v370
launch MG5/WZjj_QCD-NWA-v370
```

```
shower=Pythia8
detector=Delphes
analysis=OFF
madspin=OFF
done
```

Cards/delphes_card.dat

```
set run_card nevents 10000
set run_card ebeam1 6500.0
set run_card ebeam2 6500.0
```

```
set run_card pdlabel lhpdf
set run_card lhaid 260000
```

```
set run_card ptj 20.0
set run_card etaj 5.0
set run_card mmjj 150.0
set run_card deltaeta 1.5
set run_card ptj1min 50.0
set run_card ptj2min 25.0
```

```
set run_card etal 2.7
set run_card ptl1min 20.0
set run_card ptl2min 20.0
```

```
set run_card misset 10.0
```

```
set run_card use_syst False
set run_card cut_decays True
```

```
done
```

The primary changes include relaxing the dijet invariant mass baseline to `mmjj 150.0`, and replacing the global `ptl` cut with ordered lepton cuts: `ptl1min 20.0` and `ptl2min 20.0`. This ensures that only the leading and subleading leptons are constrained, leaving the third

lepton unrestricted. Other kinematic variables are relaxed to safely encompass the full reconstructed-level phase space.

Table 9 presents the updated event yields in the SR. In this revised comparison, we utilized the cross-sections from the **Sherpa** log outputs (Section 2.8), and applied the relaxed, ordered generator-level cuts for the **MadGraph** 3.7.0 samples to prevent phase-space truncation (Section 2.9).

Table 9: Updated expected yields in the SR. The ATLAS results are shown alongside the reference Thesis values and our refined simulation estimates (MG 3.7.0 with relaxed cuts and Sherpa 3.0.3 with log-based cross-sections). All values assume an integrated luminosity of 139 fb^{-1} .

Process	ATLAS	Thesis 9.1	Thesis G.2	MG 3.7.0	Sherpa 3.0.3
$W^\pm W^\pm jj$ EW	235	206.52	211.29	204.08	178.22
$W^\pm W^\pm jj$ QCD	24	24.05	24.27	14.48	17.88
$W^\pm Z jj$ EW	14.9	14.95	15.07	22.94	32.76
$W^\pm Z jj$ QCD	83	82.75	77.0	50.23	72.44

3 Dense Neural Network

Because the kinematic relationships between the signal and background processes are complex, a single variable is insufficient for optimal separation. Therefore, multivariate analysis techniques, specifically Deep Neural Networks (DNNs), are employed to extract sensitive features to distinguish signal and background events [1]. We consider the Dense Neural Networks (Dense-NNs) as our first trial.

3.1 Input Features

The Dense-NN relies on a set of low-level and high-level kinematic variables. These input features are defined as follows:

- $(p_T^l, \eta^l, \text{flavor}^l)$: Transverse momentum, pseudorapidity, and flavor type of the leading and sub-leading leptons.
- (p_T^j, η^j) : Transverse momentum and pseudorapidity of the leading and sub-leading jets.
- E_T^{miss} : Missing transverse momentum magnitude.

- $\Delta\phi(x, \ell_1)$: Difference in azimuthal angle between the object x and the leading lepton. The object x can be the sub-leading lepton, leading jet, sub-leading jet, or the missing transverse momentum E_T^{miss} .
- Z_ℓ^* : Zeppenfeld variables [13] of the leading and sub-leading leptons. The pseudorapidities of the leptons and tagging jets are used to define this variable:

$$Z_\ell^* = \left| \frac{\eta_\ell - \frac{1}{2}(\eta_{j_1} + \eta_{j_2})}{\eta_{j_1} - \eta_{j_2}} \right|. \quad (1)$$

- $m_T(x, E_T^{\text{miss}})$: Transverse mass of the object x and the E_T^{miss} system. The object x can be the leading lepton, sub-leading lepton, or the di-lepton system. The standard definition for a two-object system is:

$$m_T(x, E_T^{\text{miss}}) = \sqrt{(E_T^x + E_T^{\text{miss}})^2 - |\mathbf{p}_T^x + \mathbf{p}_T^{\text{miss}}|^2}, \quad (2)$$

where the scalar transverse energy is evaluated as $E_T = \sqrt{m^2 + |\mathbf{p}_T|^2}$.

- m_T^0 : Projected transverse mass:

$$m_T^0 = \sqrt{(p_T^{\ell_1} + p_T^{\ell_2} + E_T^{\text{miss}})^2 - |\mathbf{p}_T^{\ell_1} + \mathbf{p}_T^{\ell_2} + \mathbf{p}_T^{\text{miss}}|^2}. \quad (3)$$

Because the rest mass of an electron or muon is negligible in this high-energy context ($m \approx 0$), the transverse energy of a single lepton simplifies to its transverse momentum ($E_T^\ell \approx p_T^\ell$). The key difference between the standard di-lepton transverse mass (m_T) and the projected transverse mass (m_T^0) lies in how the scalar energy sum is constructed. The standard m_T treats the two leptons as a single composite system, whereas m_T^0 treats the two leptons as separate massless entities and directly sums their individual scalar transverse momenta.

- $(\Delta R_{\ell\ell}, \Delta\eta_{\ell\ell}, m_{\ell\ell}, p_T^{\ell\ell})$: Angular distance ΔR , pseudorapidity difference, invariant mass, and transverse momentum of the di-lepton system.
- $(\Delta R_{jj}, \Delta y_{jj}, m_{jj}, \Delta\phi_{jj})$: Angular distance ΔR , rapidity difference, invariant mass, and azimuthal angle difference of the di-jet system.
- $\frac{p_T^{\ell_1} p_T^{\ell_2}}{p_T^{j_1} p_T^{j_2}}$: Ratio of the scalar transverse momenta products of the leptons to the jets.
- $\min(\Delta R_{\ell,j})$: Minimum angular distance ΔR between any of the signal leptons and jets.

3.2 Dataset Construction and Validation Strategy

To ensure the robustness of the neural network models and prevent over-training, a 5-fold cross-validation strategy is employed, which is used in the ATLAS analysis. The total dataset is partitioned into training, validation, and test sets with a fixed ratio of 3:1:1.

The assignment of each event to a specific subset is determined by its event number. For a given fold index $i_{\text{fold}} \in \{0, 1, 2, 3, 4\}$, the splitting rule is defined as follows:

- **Test Set:** Events satisfying $(\text{Event Number} - i_{\text{fold}}) \pmod{5} = 0$.
- **Validation Set:** Events satisfying $(\text{Event Number} - i_{\text{fold}} + 1) \pmod{5} = 0$.
- **Training Set:** All remaining events that do not fall into the test or validation sets.

The training processes are divided into two distinct stages: a primary inclusive network ($\text{DNN}_{W^\pm W^\pm}$) designed to separate the EW signal from all backgrounds, and secondary polarization-specific networks (DNN_{pol}) trained exclusively on the EW samples to distinguish specific polarization states.

Table 10 details the exact physics processes used for these neural networks and their corresponding class labels (Signal or Background) in the reference thesis [7]. For the polarization networks, the mixed states are grouped such that “LX” denotes the sum of LL and TL states, while “TX” denotes the sum of TL and TT states.

Each fold is used to train an individual Dense-NN, resulting in an ensemble of five distinct networks. Crucially, this 5-fold application strategy is essential to prevent evaluation bias when predicting the final scores.

In the reference thesis [7], the author states that “predictions obtained using testing datasets can be assumed to be identically distributed as the predictions on measured data” and that “the prediction is calculated using the DNN where the respective events were part of the testing dataset.” However, the thesis omits a direct procedural description of how to handle the experimental data, which possesses no true labels and never participates in the training, validation, or testing splits.

To bridge this methodological gap and ensure a statistically rigorous evaluation, we *infer* that the experimental data must be processed using the identical splitting rule applied to the Monte Carlo samples. Specifically, for any given event, the modulo operation $(\text{Event Number}) \pmod{5} = i_{\text{fold}}$ dictates which specific network in the ensemble (DNN_i) is exclusively used to score it. This method guarantees that the evaluation of the experimental data mirrors the independent evaluation of the respective MC test sets, thereby preserving the identically distributed assumption.

Table 10: Process usage and assigned labels across the different DNN training stages based on the reference thesis [7]. Labels are denoted as S = Signal, B = Background, and - = Not Used. The LX category represents the sum of the LL and TL states, whereas TX represents the sum of the TL and TT states.

Physics Process	Gen. Events	$\text{DNN}_{W^\pm W^\pm}$	DNN_{pol} (LL vs TX)	DNN_{pol} (LX vs TT)
$W^\pm W^\pm jj$ -EW LL (WW-cmf)	215,008	-	S	S
$W^\pm W^\pm jj$ -EW TL (WW-cmf)	170,387	-	B	S
$W^\pm W^\pm jj$ -EW TT (WW-cmf)	456,456	-	B	B
$W^\pm W^\pm jj$ -EW LL (pp-cmf)	19,673	-	-	-
$W^\pm W^\pm jj$ -EW TL (pp-cmf)	21,471	-	-	-
$W^\pm W^\pm jj$ -EW TT (pp-cmf)	46,320	-	-	-
$W^\pm W^\pm jj$ -EW (Unpol.)	201,488	S	-	-
$W^\pm W^\pm jj$ -QCD	25,375	B	-	-
$W^\pm W^\pm jj$ -Int.	79,805	B	-	-
$W^\pm Z$ -EW	8,273	B	-	-
$W^\pm Z$ -QCD (Comb.)	49,999	B	-	-
Top quark processes	2,192	B	-	-
ZZ production	2,880	B	-	-
Charge-flip	8,231	B	-	-

3.3 Kinematic Distributions

To illustrate the separation power of the input features, we present the normalized distributions of several key kinematic variables. The comparisons are divided into two categories, reflecting the two-stage strategy of the Dense-NN training: (1) separating the $W^\pm W^\pm jj$ -EW signal from various background processes, and (2) distinguishing between the different polarization states of the $W^\pm W^\pm jj$ electroweak production.

3.3.1 Signal vs Background Discrimination

In the first stage of the DNN, the goal is to isolate the inclusive $W^\pm W^\pm jj$ -EW signal from the dominant backgrounds, which include $W^\pm W^\pm jj$ -QCD, $W^\pm Z jj$ -EW, and $W^\pm Z jj$ -QCD.

Figure 3 shows the distributions of six features. The VBS signature is characterized by two forward jets with a large invariant mass (m_{jj}) and a large pseudorapidity gap ($|\Delta\eta_{jj}|$), making them powerful discriminants against QCD-induced processes. Furthermore, the Zepfenfeld variable ($Z_{\ell_1}^*$) quantifies the centrality of the lepton relative to the tagging jets; signal leptons tend to fall between the two forward jets. General kinematics, such as the leading jet transverse momentum ($p_T^{j_1}$), di-lepton invariant mass ($m_{\ell\ell}$), and missing transverse momentum (E_T^{miss}), provide additional discrimination power against the WZ and QCD backgrounds.

3.3.2 Polarization Discrimination

The second stage of the DNN focuses entirely on the $W^\pm W^\pm jj$ -EW process, aiming to separate the longitudinally polarized state (LL) from the transverse states (LT and TT).

Figure 4 illustrates six variables. The W_L bosons tend to emit leptons more centrally and with different spin correlations compared to W_T bosons, leading to distinct shapes in the angular separations $|\Delta\eta_{\ell\ell}|$ and $\Delta\phi_{\ell\ell}$, as well as the leading lepton transverse momentum ($p_T^{\ell_1}$). The transverse mass of the di-lepton system (m_T) and the projected transverse mass (m_T^0) are also sensitive to the different kinematic boundaries of the polarization states. Additionally, the azimuthal angle difference between the jets ($\Delta\phi_{jj}$) carries information about the intermediate vector boson interference in the VBS process.

3.4 Neural Network Architecture

The Dense-NN architecture utilized in the analysis is presented in Figure 5.




Figure 3: Normalized distributions of six key kinematic variables for the inclusive $W^\pm W^\pm jj$ -EW signal and the $W^\pm W^\pm jj$ -QCD, $W^\pm Zjj$ -EW, and $W^\pm Zjj$ -QCD backgrounds.




Figure 4: Normalized distributions of six polarization-sensitive variables for the LL, LT, and TT polarization states of the $W^\pm W^\pm jj$ -EW process.

The network begins with an input normalization layer. This layer applies the scaling to the input features first. Then, normalizing the resulting distribution to have a mean of 0 and a variance of 1.

Following the normalization layer, the architecture relies on a core block consisting of a fully-connected (dense) layer, a Batch Normalization layer, a Swish [14] activation function, and a Dropout layer for regularization. This composite block is repeated L times to form the deep hidden layers of the network.

Finally, the network terminates with an output dense layer.




Figure 5: Schematic representation of the feed-forward neural network architecture. The sequence begins with an input normalization layer. The core architecture consists of L repeated hidden layers, each comprising a dense layer with a constant number of neurons, followed by batch normalization, a Swish activation function, and a dropout layer for regularization. The final dense layer provides the binary classification output.

The Swish activation function utilized in the hidden blocks is defined as:

$$\text{swish}_{\beta}(x) = \frac{x}{1 + e^{-\beta x}} \quad (4)$$

where β is a learnable parameter or a fixed constant. Unlike ReLU, Swish is smooth and non-monotonic, characteristics that have been shown to improve gradient propagation and performance in deep networks.

3.5 Implementation and Preliminary Validation

The neural network architecture, data pipeline, and training loop are implemented using PyTorch 2.5.1 [15]. To verify the computational setup, we conducted a preliminary validation test on both the primary inclusive task $\text{DNN}_{W^{\pm}W^{\pm}}$ and the secondary polarization task DNN_{pol} .

For this initial test, we utilized the complete set of input features described in Section 3.1. The dataset comprised approximately 40,000 events for the $\text{DNN}_{W^{\pm}W^{\pm}}$ task and

73,000 events for the DNN_{pol} task. The models were trained for 10 epochs using the Adam optimizer with a learning rate of 10^{-3} and a batch size of 256. This configuration verified the 5-fold cross-validation as well as the Dense-NN forward and backward propagation.

Figure 6 illustrates the evolution of the loss and the Area Under the ROC Curve (AUC) across the initial training epochs. The quantitative performance of the models across the 5-fold ensemble is summarized in Table 11. The negligible discrepancy between the validation and test AUC scores indicates that the data splitting strategy effectively prevents data leakage and over-training, even at this early stage.

Table 11: Preliminary performance of the Dense-NN models after 10 epochs of training. The values represent the mean AUC score and its standard deviation across the 5 independent folds.

Classification Task	Validation AUC	Test AUC
$\text{DNN}_{W^\pm W^\pm}$	0.786 ± 0.078	0.788 ± 0.076
DNN_{pol} (LL vs. TX)	0.948 ± 0.021	0.947 ± 0.024


As anticipated for a preliminary test, the training curves indicate that the models have not yet converged. The next steps will involve generating larger training samples, increasing the number of epochs, and testing the optimized hyperparameters.

3.6 Feature Scaling, Normalization, and Full Training


In Section 3.5, the preliminary training results for the DNN_{pol} task yielded an unreasonably high AUC (AUC ≈ 0.948). Upon thorough investigation, it was a result of a subtle data leakage issue during the feature normalization process. Initially, the normalization procedure (calculating the mean and standard deviation) was applied to each physics process separately before merging them into a unified dataset. This encoded class-specific properties into the scaled variables, introducing a weak label leakage that allowed the network to artificially distinguish the processes.

This issue has been resolved. The scaling function is now strictly fitted on the completely merged *training* dataset. The derived parameters are subsequently applied to transform the training, validation, and test sets.


With the corrected data pipeline, we executed a full-scale training utilizing the complete set of input features described in Section 3.1. The dataset size was expanded to approximately 400,000 events for the inclusive $\text{DNN}_{W^\pm W^\pm}$ task, and 740,000 events for the DNN_{pol} task.




(a) Loss ($\text{DNN}_{W^\pm W^\pm}$)



(b) AUC ($\text{DNN}_{W^\pm W^\pm}$)



(c) Loss (DNN_{pol})



(d) AUC (DNN_{pol})

Figure 6: Preliminary training and validation curves (Loss and AUC) over 10 epochs for the inclusive and polarization DNN tasks.

The networks were trained using the Adam optimizer with a learning rate of 10^{-3} and a batch size of 256. To prevent overfitting during this extended run, an early stopping mechanism was implemented. The validation loss was monitored, and the training was terminated if no improvement was observed for 20 consecutive epochs. Table 12 summarizes the hyperparameter settings.

Table 12: Hyperparameters used in Dense-NN training.

Parameter	Value
Number of hidden layers	4
Number of nodes per layer	128
Dropout rate	0.3
Batch size	256
Learning rate	0.001
Early stopping patience	20

Figure 7 illustrates the evolution of the loss and the AUC across the training epochs. The quantitative performance of the fully trained models across the 5-fold ensemble is summarized in Table 13.


Table 13: Final performance of the Dense-NN models across the 5 independent folds after full training with early stopping. The table presents the mean AUC scores and their standard deviations. The corrected normalization strategy yields mathematically robust and realistic separation power.

Classification Task	Validation AUC	Test AUC
$\text{DNN}_{W^\pm W^\pm}$ (EW vs. Background)	0.806 ± 0.001	0.806 ± 0.001
DNN_{pol} (LL vs. TX)	0.839 ± 0.001	0.839 ± 0.001


The revised test AUC for the polarization task represents a much more realistic separation power. Furthermore, the small variance between the folds and the negligible discrepancy between the validation and test AUC scores confirm that the data splitting and normalization strategy effectively prevents overtraining.

3.7 DNN Output Score Distributions


To visually assess the discrimination power of the trained networks, we evaluate the DNN output score distributions for both the inclusive and polarization-specific classification




(a) Loss ($\text{DNN}_{W^\pm W^\pm}$)



(b) AUC ($\text{DNN}_{W^\pm W^\pm}$)



(c) Loss (DNN_{pol})



(d) AUC (DNN_{pol})

Figure 7: Full training and validation curves (Loss and AUC) for the inclusive and polarization Dense-NN tasks. The early stopping mechanism successfully prevents overtraining.

tasks. Figure 8 presents the normalized predicted probability scores evaluated on the test datasets.




Figure 8: Distributions of the DNN output scores for the test datasets. (a) The inclusive $\text{DNN}_{W^\pm W^\pm}$ separating the EW signal from backgrounds. (b) The DNN_{pol} distinguishing the LL signal from the TX states.

In an idealized binary classification scenario, the output scores would exhibit distinct peaks at 1.0 for the signal and 0.0 for the background. However, as observed in Figure 8, the distributions display a profile with noticeable overlapping regions.

Upon preliminary investigation, this behavior is attributed to the incorporation of physical event weights during the training process. However, we observed a discrepancy between our current score distributions and the reference plots presented in the thesis [7].

3.8 Training Weight Strategy

To properly account for the different production cross-sections and the imbalanced MC dataset sizes, we evaluated three distinct sample weighting strategies for the NN training loss function:

1. **Physical Yield Weighting:** Each event is assigned a weight corresponding to its expected physical yield in the SR.
2. **Inverse Generation Weighting:** Each event is weighted simply by the inverse of the total number of generated events for its respective process ($w \propto 1/N_{\text{gen}}$).

3. Hybrid Weighting: A combination of the physical yield and inverse event count methods.

The pure “Physical Yield” strategy reflects the true kinematic composition and imbalance among all physical processes, closely mimicking the exact conditions of real collision data. However, if used alone, it fails to account for the imbalance within the generated MC samples themselves. This is why the “Inverse Generation” and “Hybrid” approaches were introduced and evaluated.

Furthermore, a crucial **class-balancing** step is enforced. The event weights are globally normalized such that the sum of the weights in the signal category exactly equals the sum of the weights in the background category ($\sum w_{\text{sig}} = \sum w_{\text{bkg}}$). This global scaling preserves the relative physical ratios of the internal sub-processes while preventing the neural network from trivially predicting the majority class to minimize the loss.

The Hybrid weighting strategy combined with global class balancing provides the most reasonable learning strategy. Thus, we presented the training results of this strategy. Figure 9 illustrates the resulting Dense-NN output evaluated on the test datasets utilizing this optimized weighting scheme. The distributions satisfy our expectation that they exhibit peaks near 1 for signal and 0 for the background.




Figure 9: Distributions of the DNN output scores for the test datasets using the optimized hybrid weighting strategy. (a) The inclusive $\text{DNN}_{W^{\pm}W^{\pm}}$ separating the EW signal from backgrounds. (b) The DNN_{pol} distinguishing the LL signal from the TX states.

4 Statistical framework

This section outlines the statistical framework utilized to extract the final physics results. The ultimate goal of this analysis is to evaluate the Discovery Significance Z_0 and the Upper Limit constraints.

4.1 Profile Likelihood Fit

First, we need to compute the profile likelihood $\mathcal{L}$, which evaluates the goodness-of-fit between the theoretical model and the observed data across the 1D histograms. For each bin i , the expected number of events is defined as m_i , and the observed number of events is denoted as n_i .

The expectation m_i consists of the signal and background contributions, parameterized by the signal strength μ , and is modified by systematic uncertainties. These uncertainties are modeled using nuisance parameters (NPs), denoted as $\boldsymbol{\theta}$ for systematics, and $\boldsymbol{\gamma}$ for Monte Carlo (MC) statistical uncertainties:

$$m_i(\mu, \boldsymbol{\theta}, \boldsymbol{\gamma}) = \gamma_i (\mu s_i(\boldsymbol{\theta}) + b_i(\boldsymbol{\theta})), \quad (5)$$

where s_i and b_i are the nominal number of predicted signal and background events in bin i , respectively.

The complete likelihood function $\mathcal{L}$ is constructed as the product of Poisson probabilities for all bins, multiplied by constraint terms for the nuisance parameters:

$$\mathcal{L}(\mu, \boldsymbol{\theta}, \boldsymbol{\gamma}) = \prod_{i=1}^{N_{\text{bins}}} \text{Pois}(n_i | m_i) \text{Pois} \left(\frac{1}{(\delta_i^{\text{stat}})^2} \middle| \frac{\gamma_i}{(\delta_i^{\text{stat}})^2} \right) \prod_{j=1}^{N_{\text{nuis}}} f_{\text{Gauss}}(\theta_j). \quad (6)$$

Here, the Poisson probability is defined as $\text{Pois}(n_i | m_i) = \frac{m_i^{n_i}}{n_i!} e^{-m_i}$. The auxiliary functions $f_{\text{Gauss}}(\theta_j)$ are standard normal distributions constraining the systematic NPs $\boldsymbol{\theta}$.

4.1.1 Test Statistic and Profile Likelihood Ratio

To test specific hypotheses, the profile likelihood ratio $\lambda(\mu)$ is defined as:

$$\lambda(\mu) = \frac{\mathcal{L}(\mu, \hat{\boldsymbol{\theta}}, \hat{\boldsymbol{\gamma}})}{\mathcal{L}(\hat{\mu}, \hat{\boldsymbol{\theta}}, \hat{\boldsymbol{\gamma}})}. \quad (7)$$

In the denominator, $\hat{\mu}$, $\hat{\boldsymbol{\theta}}$, and $\hat{\boldsymbol{\gamma}}$ are the unconditional maximum likelihood estimators that globally maximize $\mathcal{L}$ across the entire parameter space. In the numerator, $\hat{\boldsymbol{\theta}}$ and $\hat{\boldsymbol{\gamma}}$ are the

conditional maximum likelihood estimators that maximize $\mathcal{L}$ for a specified, fixed value of μ .

Based on $\lambda(\mu)$, the test statistic q_μ is constructed:

$$q_\mu = -2 \ln \lambda(\mu). \quad (8)$$

4.1.2 Discovery Significance and Upper Limits

The presence of a new signal is tested by rejecting the background-only hypothesis ($\mu = 0$). The corresponding test statistic is $q_0 = -2 \ln \lambda(0)$. For a sufficiently large dataset, the discovery significance Z_0 can be approximated as:

$$Z_0 \approx \sqrt{q_0}. \quad (9)$$

In the absence of a significant excess, the analysis sets a 95% Confidence Level (CL) upper limit on the signal strength. This is derived using the CL_s method, which also utilizes the test statistic q_μ .

4.2 The histogram binning

In this analysis, the outputs of the Dense-NNs serve as the direct inputs for the profile likelihood fit. The outputs can be interpreted as classification tuple ($\text{DNN}_{\text{pol}}, \text{DNN}_{W^\pm W^\pm}$) on a two-dimensional score plane:

- Y-axis ($\text{DNN}_{W^\pm W^\pm}$): The inclusive network score, which separates the pure EW $W^\pm W^\pm jj$ signal from reducible and irreducible backgrounds (e.g., WZ , QCD WW).
- X-axis (DNN_{pol}): The polarization-specific network score, which evaluates the EW events to distinguish the targeted polarization state (e.g., LL signal) from the other polarization states (e.g., TX backgrounds).

This study uses a rebinning algorithm to maximize the significance and reduce the computation time. This procedure slices the 2D plane into a sequence of 1D bins, requiring a delicate balance between preserving the Dense-NN output shape information and avoiding statistical fluctuations. The algorithm is listed below:

- **Region Slicing:** The Y-axis ($\text{DNN}_{W^\pm W^\pm}$) is first divided into discrete purity intervals. Each interval is then evaluated along the X-axis (DNN_{pol}) and flattened into consecutive 1D strips.

- **Statistical Stability:** To prevent statistical uncertainties, the bin edges of each histogram are merged until there are at least ten events originating from background or transverse polarized contributions in each bin.
- **Significance Maximization:** Adjacent bins are merged if keeping them separate would not improve the significance.

Table 14 is the final binning from the thesis.

Table 14: The optimized splits in the $\text{DNN}_{W^\pm W^\pm}$ classification and the corresponding bin edges in polarization classification.

Polarization	Split in $\text{DNN}_{W^\pm W^\pm}$	Binning in polarization classification
LL in WW -cmf	[0.0, 0.2, 0.6, 1.0]	[0.0, 0.4, 0.7, 1.0] [0.0, 0.5, 0.8, 1.0] [0.0, 0.4, 0.6, 0.7, 0.8, 1.0]
LX in WW -cmf	[0.0, 0.3, 0.7, 1.0]	[0.0, 0.3, 0.7, 1.0] [0.0, 0.3, 0.5, 0.6, 0.7, 0.8, 1.0] [0.0, 0.15, 0.35, 0.45, 0.55, 0.7, 1.0]
LL in pp -cmf	[0.0, 0.15, 0.5, 1.0]	[0.0, 0.55, 0.8, 1.0] [0.0, 0.7, 0.8, 1.0] [0.0, 0.4, 0.6, 0.7, 0.8, 1.0]
LX in pp -cmf	[0.0, 0.25, 0.6, 1.0]	[0.0, 0.45, 0.6, 1.0] [0.0, 0.25, 0.4, 0.55, 0.75, 1.0] [0.0, 0.15, 0.3, 0.4, 0.5, 0.6, 0.7, 1.0]

4.3 Expected Results

The fit model must be validated using an expected dataset. This is achieved by utilizing an **Asimov dataset**, which is a representative pseudo-dataset where the “observed” event counts in each bin are set exactly equal to the nominal theoretical predictions (signal plus background), absent any statistical fluctuations.

Fitting the Asimov dataset serves a critical purpose: it determines the theoretical sensitivity of the analysis based on the current detector acceptance, background estimations, and the discrimination power of the DNN. This yields a baseline expected discovery significance (Z_0) for the targeted polarization states.

Table 15 summarizes the expected significances for the LL and LX signal states as reported in the reference thesis [7].

Table 15: Expected discovery significances for the LL and LX polarization states, evaluated using the Asimov dataset. The comparison highlights the impact of defining the polarization vectors in the WW center-of-mass frame versus the partonic frame.

Reference Frame	LL Significance (Z_0)	LX Significance (Z_0)
WW-cmf	1.512σ	4.557σ
pp-cmf	0.937σ	3.975σ

As shown in the table, defining the polarization in the WW center-of-mass frame yields notably higher expected sensitivities for both the LL and LX states compared to the initial-state partonic frame.

4.4 Results of the Statistical Fit

Using the profile likelihood fit framework and the unrolled 1D binning configuration described above, we evaluate the expected discovery significance Z_0 and the expected 95% CL upper limit on the signal strength parameter μ for both the LL mode and the LX mode. The fit is performed using the Asimov dataset constructed from our own simulated Monte Carlo samples in the WW center-of-mass frame.

Table 16 summarizes our obtained expected discovery significances and expected 95% CL upper limits on the signal strength parameters, alongside the reference results from the thesis [7] for comparison.

Table 16: Expected discovery significances (Z_0) and expected 95% CL upper limits on the signal strength parameters (μ) for the LL and LX polarization states in the WW center-of-mass frame. The reference values from the thesis are provided for comparison.

Signal State	Expected Significance (Z_0)		Expected 95% CL Upper Limit	
	Thesis	Our Result	Thesis	Our Result
LL (μ_{LL})	1.512σ	1.600σ	2.273	1.255
LX (μ_{LX})	4.557σ	4.852σ	—	0.356

As shown in Table 16, despite utilizing unoptimized Dense-NNs and slightly varied input features, our simulation successfully reproduces the overall statistical trend. The expected significances are in agreement with the reference values.

However, a discrepancy is observed in the expected 95% CL upper limit for the LL state, where our result is approximately half of the reference value. This deviation directly

highlights the impact of Nuisance Parameters (NPs). Our current fit considers only statistical uncertainties. In a full fit, systematic uncertainties flatten and widen the profile likelihood curve. Consequently, the absence of these systematic constraints results in a tighter upper limit.

5 Some research directions

We propose some research directions to further enhance the discovery significance and methodological robustness of the longitudinal W boson scattering study.

5.1 Enhance the discovery significance

The current analysis relies on a baseline Dense-NN, which flattens kinematic features and may lose underlying spatial correlations. We propose to utilize advanced architectures to boost classification accuracy then enhance the discovery significance:

- **Advanced Architectures:** Replace the Dense-NN with CNN, **ParticleNet** (Graph Neural Networks, GNNs) [16], or **Particle Transformer** (ParT) [17] to better capture the complex VBS topologies.
- **Multi-Task Learning (MTL):** Instead of training separate classifiers for inclusive WW signal and polarization states, implement a single MTL framework with a shared representation space and multiple classification heads. This can yield better generalization and higher separation power.
- **Multi-Class classification:** Instead of utilizing the step separation strategy, implement a network to distinguish LL, LT, TT, and other background classes directly.

5.2 Unsupervised / Self-supervised learning

Current supervised training heavily relies on MC simulations, which introduces theoretical uncertainties and potential biases when applied to measured data.

- **Weakly Supervised Learning & Topic Modeling:** The Classification Without Labels (CWoLa) paradigm [18] successfully operates on two mixed data regions for binary classification. We can extend this to multi-class scenarios using Topic Modeling [19]. This approach treats different mixed sample regions as “documents” and the underlying physics processes (e.g., pure LL, LT, TT states) as “topics,” enabling the extraction of pure kinematic distributions directly from collision data.

- **Self-Supervised Contrastive Learning:** Implementing self-supervised frameworks, such as JetCLR [20], allows the neural network to learn robust, physics-invariant representations without requiring explicit MC labels. By applying physical data augmentations (e.g., azimuthal rotations, longitudinal boosts, or momentum smearing), the model maps physically similar events closer together in a latent space. This latent representation can subsequently be evaluated or fine-tuned using minimal MC labels via linear probing.
- **Objective and Impact:** The ultimate goal is to exploit these data-driven techniques to construct optimal kinematic observables and isolate the polarization states with minimal dependence on pure MC labels.

5.3 Fully Hadronic Channel

While the baseline analysis focuses on the clean same-sign fully leptonic channel ($\ell^\pm \nu \ell^\pm \nu$), the fully hadronic decay mode ($WW \rightarrow qq\bar{q}\bar{q}$) represents a compelling yet highly challenging frontier for VBS polarization studies. To date, no experimental results for VBS exist in this channel.

References

- [1] G. Aad *et al.*, “Evidence for Longitudinally Polarized W Bosons in the Electroweak Production of Same-Sign W Boson Pairs in Association with Two Jets in pp Collisions at s=13 TeV with the ATLAS Detector,” *Phys. Rev. Lett.*, vol. 135, no. 11, p. 111802, 2025.
- [2] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations,” *JHEP*, vol. 07, p. 079, 2014.
- [3] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, “An introduction to PYTHIA 8.2,” *Comput. Phys. Commun.*, vol. 191, pp. 159–177, 2015.
- [4] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens, and M. Selvaggi, “DELPHES 3, A modular framework for fast simulation of a generic collider experiment,” *JHEP*, vol. 02, p. 057, 2014.

- [5] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- k_t jet clustering algorithm,” *JHEP*, vol. 04, p. 063, 2008.
- [6] M. Cacciari, G. P. Salam, and G. Soyez, “FastJet User Manual,” *Eur. Phys. J. C*, vol. 72, p. 1896, 2012.
- [7] M. Stange, M. Kobel, F. Siegert, and K. Potamianos, *Longitudinal $W^\pm$ Boson Polarization in Same-charged $W^\pm W^\pm$ Scattering in the ATLAS Experiment*. Verlagsname nicht ermittelbar, 2025.
- [8] D. Buarque Franzosi, O. Mattelaer, R. Ruiz, and S. Shil, “Automated predictions from polarized matrix elements,” *JHEP*, vol. 04, p. 082, 2020.
- [9] E. Bothmann *et al.*, “Event generation with Sherpa 3,” *JHEP*, vol. 12, p. 156, 2024.
- [10] R. D. Ball *et al.*, “Parton distributions with LHC data,” *Nucl. Phys. B*, vol. 867, pp. 244–289, 2013.
- [11] R. D. Ball *et al.*, “The PDF4LHC21 combination of global PDF fits for the LHC Run III,” *J. Phys. G*, vol. 49, no. 8, p. 080501, 2022.
- [12] G. Aad *et al.*, “Measurement and interpretation of same-sign W boson pair production in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” *JHEP*, vol. 04, p. 026, 2024.
- [13] D. L. Rainwater, R. Szalapski, and D. Zeppenfeld, “Probing color singlet exchange in $Z + \text{two jet}$ events at the CERN LHC,” *Phys. Rev. D*, vol. 54, pp. 6680–6689, 1996.
- [14] P. Ramachandran, B. Zoph, and Q. V. Le, “Searching for activation functions,” *CoRR*, vol. abs/1710.05941, 2017.
- [15] J. Ansel, E. Yang, H. He, N. Gimelshein, A. Jain, M. Voznesensky, B. Bao, P. Bell, D. Berard, E. Burovski, G. Chauhan, A. Chourdia, W. Constable, A. Desmaison, Z. DeVito, E. Ellison, W. Feng, J. Gong, M. Gschwind, B. Hirsh, S. Huang, K. Kalambarkar, L. Kirsch, M. Lazos, M. Lezcano, Y. Liang, J. Liang, Y. Lu, C. K. Luk, B. Maher, Y. Pan, C. Puhersch, M. Reso, M. Saroufim, M. Y. Siraichi, H. Suk, S. Zhang, M. Suo, P. Tillet, X. Zhao, E. Wang, K. Zhou, R. Zou, X. Wang, A. Mathews, W. Wen, G. Chanan, P. Wu, and S. Chintala, “Pytorch 2: Faster machine learning through dynamic python bytecode transformation and graph compilation,” in *Proceedings of the*

29th ACM International Conference on Architectural Support for Programming Languages and Operating Systems, Volume 2, ASPLOS '24, (New York, NY, USA), p. 929–947, Association for Computing Machinery, 2024.

- [16] H. Qu and L. Gouskos, “ParticleNet: Jet Tagging via Particle Clouds,” *Phys. Rev. D*, vol. 101, no. 5, p. 056019, 2020.
- [17] H. Qu, C. Li, and S. Qian, “Particle Transformer for Jet Tagging,” 2 2022.
- [18] E. M. Metodiev, B. Nachman, and J. Thaler, “Classification without labels: Learning from mixed samples in high energy physics,” *JHEP*, vol. 10, p. 174, 2017.
- [19] E. M. Metodiev and J. Thaler, “Jet Topics: Disentangling Quarks and Gluons at Colliders,” *Phys. Rev. Lett.*, vol. 120, no. 24, p. 241602, 2018.
- [20] B. M. Dillon, G. Kasieczka, H. Olschlager, T. Plehn, P. Sorrenson, and L. Vogel, “Symmetries, safety, and self-supervision,” *SciPost Phys.*, vol. 12, no. 6, p. 188, 2022.