

Cheatsheet

@r2cp

Page 3

$$\mathbb{E}[X \cdot Y] \neq \mathbb{E}[X] \cdot \mathbb{E}[Y]$$
$$\mathbb{E}[X \cdot Y] = \mathbb{E}[\mathbb{E}[Y \cdot X|Y]] = \mathbb{E}[Y \cdot \mathbb{E}[X|Y]]$$

Linearity of Expectation where a and c are given scalars:

$$\mathbb{E}[aX + cY] = a\mathbb{E}[X] + c\mathbb{E}[Y]$$

If Variance of X is known:

$$\mathbb{E}[X^2] = var(X) + \mathbb{E}[X]$$

15 Variance

Variance is the squared distance from the mean.

$$Var(X) = \mathbb{E}[(X - \mathbb{E}(X))^2]$$

$$Var(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$$

Variance of a product with constant a :

$$Var(aX) = a^2 Var(X)$$

Variance of sum of two dependent r.v.:

$$Var(X + Y) = Var(X) + Var(Y) + 2Cov(X, Y)$$

Variance of sum/difference of two independent r.v.:

$$Var(X + Y) = Var(X) + Var(Y)$$
$$Var(X - Y) = Var(X) + Var(Y)$$

16 Covariance

The Covariance is a measure of how much the values of each of two correlated random variables determine each other

$$Cov(X, Y) = \mathbb{E}[(X - \mu_X)(Y - \mu_Y)]$$

$$Cov(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$$

$$Cov(X, Y) = \mathbb{E}[(X)(Y - \mu_Y)]$$

Possible notations:

$$Cov(X, Y) = \sigma(X, Y) = \sigma_{(X, Y)}$$

Covariance is commutative:

$$Cov(X, Y) = Cov(Y, X)$$

Covariance with of r.v. with itself is variance:

$$Cov(X, X) = \mathbb{E}[(X - \mu_X)^2] = Var(X)$$

Useful properties:

$$Cov(aX + h, bY + c) = abCov(X, Y)$$

$$Cov(X, X + Y) = Var(X) + cov(X, Y)$$

$$Cov(aX + bY, Z) = aCov(X, Z) + bCov(Y, Z)$$

If $Cov(X, Y) = 0$, we say that X and Y are uncorrelated. If X and Y are independent, their Covariance is zero. The converse is not always true. It is only true if X and Y form a gaussian vector, ie. any linear combination $\alpha X + \beta Y$ is gaussian for all $(\alpha, \beta) \in \mathbb{R}^2$ without $[0, 0]$.

17 correlation coefficient

$$\rho(X, Y) = \frac{Cov(X, Y)}{\sqrt{Var(X)Var(Y)}}$$

18 Important probability distributions

Bernoulli

Parameter $p \in [0, 1]$, discrete

$$p_X(k) = \begin{cases} p, & \text{if } k = 1 \\ (1 - p), & \text{if } k = 0 \end{cases}$$

$$\mathbb{E}[X] = p$$

$$Var(X) = p(1 - p)$$

Likelihood n trials:

$$L_n(X_1, \dots, X_n, p) = p^{\sum_{i=1}^n X_i} (1 - p)^{n - \sum_{i=1}^n X_i}$$

Loglikelihood n trials:

$$\ell_n(p) = \ln(p) \sum_{i=1}^n X_i + (n - \sum_{i=1}^n X_i) \ln(1 - p)$$

MLE:

$$\hat{p}_{MLE} = \frac{\sum_{i=1}^n (X_i)}{n}$$

Fisher Information:

$$I(p) = \frac{1}{p(1 - p)}$$

Canonical exponential form:

$$f_\theta(y) = \exp(y\theta - \ln(1 + e^\theta) + \frac{0}{c(y, \phi)})$$

$$\theta = \ln\left(\frac{p}{1 - p}\right)$$
$$\phi = 1$$

Binomial

Parameters p and n , discrete. Describes the number of successes in n independent Bernoulli trials.

$$p_X(k) = \binom{n}{k} p^k (1 - p)^{n - k}, k = 0, \dots, n$$

$$\mathbb{E}[X] = np$$

$$Var(X) = np(1 - p)$$

Likelihood:

$$L_n(X_1, \dots, X_n, \theta) = \left(\prod_{i=1}^n \binom{n}{X_i}\right) \theta^{\sum_{i=1}^n X_i} (1 - \theta)^{nK - \sum_{i=1}^n X_i}$$

Loglikelihood:

$$\ell_n(\theta) = C + \left(\sum_{i=1}^n X_i\right) \log \theta + (nK - \sum_{i=1}^n X_i) \log(1 - \theta)$$

MLE:

Fisher Information:

$$I(p) = \frac{n}{p(1 - p)}$$

Canonical exponential form:

$$f_\theta(y) = \exp(y \underbrace{(\ln(p) - \ln(1 - p))}_{\theta} + n \underbrace{\ln(1 - p)}_{-b(\theta)} + \ln \underbrace{\left(\binom{n}{y}\right)}_{c(y, \phi)})(x) = \begin{cases} \lambda \exp(-\lambda x), & \text{if } x \geq 0 \\ 0, & \text{o.w.} \end{cases}$$

Geometric

Number of T trials up to (and including) the first success.

$$p_T(t) = (1 - p)^{t-1}, t = 1, 2, \dots$$

$$\mathbb{E}[T] = \frac{1}{p}$$

$$var(T) = \frac{1 - p}{p^2}$$

Pascal

The negative binomial or Pascal distribution is a generalization of the geometric distribution. It relates to the random experiment of repeated independent trials until observing m successes. I.e. the time of the k th arrival.

$$Y_k = T_1 + \dots + T_k$$

$$T_i \sim iid \text{Geometric}(p)$$

$$\mathbb{E}[Y_k] = \frac{k}{p}$$

$$Var(Y_k) = \frac{k(1 - p)}{p^2}$$

$$p_{Y_k}(t) = \binom{t-1}{k-1} p^k (1 - p)^{t-k}$$
$$t = k, k + 1, \dots$$

Poisson

Parameter λ , discrete, approximates the binomial PMF when n is large, p is small, and $\lambda = np$.

$$p_X(k) = \exp(-\lambda) \frac{\lambda^k}{k!} \text{ for } k = 0, 1, \dots,$$

$$\mathbb{E}[X] = \lambda$$

$$Var(X) = \lambda$$

Likelihood:

$\ell_n(\lambda) = -n\lambda + \log(\lambda) \left(\sum_{i=1}^n x_i\right) - \log\left(\prod_{i=1}^n x_i!\right)$

MLE:

$$\hat{\lambda}_{MLE} = \frac{1}{n} \sum_{i=1}^n (X_i)$$

Fisher Information:

$$I(\lambda) = \frac{1}{\lambda}$$

Canonical exponential form:

$$f_\theta(y) = \exp(y\theta - \underbrace{e^\theta}_{b(\theta)} - \underbrace{\ln y!}_{c(y, \phi)})$$

$$\theta = \ln \lambda$$
$$\phi = 1$$

Poisson process:

k arrivals in t slots

$$p_X(k, t) = \mathbb{P}(N_t = k) = e^{-\lambda t} \frac{(\lambda t)^k}{k!}$$

$$\mathbb{E}[N_t] = \lambda t$$

$$Var(N_t) = \lambda t$$

Exponential

Parameter λ , continuous

$$f_X(x) = \begin{cases} 1 - \exp(-\lambda x), & \text{if } x \geq 0 \\ 0, & \text{o.w.} \end{cases}$$

$$\mathbb{E}[X] = \frac{1}{\lambda}$$

$$\mathbb{E}[X^2] = \frac{2}{\lambda^2}$$

$$Var(X) = \frac{1}{\lambda^2}$$

Likelihood:

$$

MLE:

$$

Canonical exponential form:

$$

$$

$$

$$

Loglikelihood:

$$

$$

$$

CDF of standard gaussian:

$$

Loglikelihood:

$$$$

Canonical exponential form:

Gaussians are invariant under affine transformation:

$$

Standardization:

$$

Higher moments:

$$

$$

$$

$$

Loglikelihood:

Cauchy

continuous, parameter m ,

$$

$$

Chi squared

The χ_d^2 distribution with d degrees of freedom is given by the distribution of $Z_1^2 + Z_2^2 + \dots + Z_d^2$, where $Z_1, \dots, Z_d \stackrel{iid}{\sim} \mathcal{N}(0, 1)$

If $V \sim \chi_k^2$:

$$

Student's T Distribution

$T_n := \frac{Z}{\sqrt{V/n}}$ where $Z \sim \mathcal{N}(0, 1)$, and Z and V are independent

18.1 Useful to know

18.1.1 Min of iid exponential r.v

Let X_1, \dots, X_n be i.i.d. $Exp(\lambda)$ random variables.

Distribution of $\min_i(X_i)$

$$

18.1.2 Counting Committees

Out of $2n$ people, we want to choose a committee of n people, one of whom will be its chair. In how many different ways can this be done?"

$2n$ people, consisting of n boys and n girls, we want to select a committee of n people. In how many ways can this be done?"

$2n$ elements have?"

$$

The covariance matrix Σ is a $d \times d$ matrix. It is a table of the pairwise covariances of the elements of the random vector. Its diagonal elements are the variances of the elements of the random vector, the off-diagonal elements are its covariances. Note that the covariance is commutative e.g. $\sigma_{12} = \sigma_{21}$

Alternative forms:

$$

Every Covariance matrix is positive definite.

$\alpha^T X$ is (univariate) Gaussian or constant for any constant non-zero vector $\alpha \in \mathbb{R}^d$.

Multivariate Gaussians

The distribution of X the d -dimensional Gaussian or normal distribution, is completely specified by the vector mean $\mu = \mathbb{E}[X] = (\mathbb{E}[X^{(1)}], \dots, \mathbb{E}[X^{(d)}])^T$ and the $d \times d$ covariance matrix Σ . If Σ is invertible, then the pdf of X is:

$$

Where $\det(\Sigma)$ is the determinant of Σ , which is positive when Σ is invertible. If $\mu = 0$ and Σ is the identity matrix, then X is called a standard normal random vector.

If the covariant matrix Σ is diagonal, the pdf factors into pdfs of univariate Gaussians, and hence the components are independent.

The linear transform of a gaussian $X \sim N_d(\mu, \Sigma)$ with conformable matrices A and B is a gaussian:

$$

$$

$$

19 Random Vectors

A random vector $X = (X^{(1)}, \dots, X^{(d)})^T$ of dimension $d \times 1$ is a vector-valued function from a probability space ω to \mathbb{R}^d :

$$

where each $X^{(k)}$, is a (scalar) random variable on Ω .

PDF of X : joint distribution of its components $X^{(1)}, \dots, X^{(d)}$.

CDF of X :

$$

The sequence X_1, X_2, \dots converges in probability to X if and only if each component of the sequence $X_1^{(k)}, X_2^{(k)}, \dots$ converges in probability to $X^{(k)}$.

Expectation of a random vector

The expectation of a random vector is the elementwise expectation. Let X be a random vector of dimension $d \times 1$.

$$

Let X and Y be random matrices of the same dimension, and let A and B be conformable matrices of constants.

$$

Covariance Matrix

Let X be a random vector of dimension $d \times 1$ with expectation μ_X .

Matrix outer products!

$$

21 Matrix Algebra

$$\|Ax\|^2 = (Ax)^T(Ax) = x^T A^T Ax$$

22 Calculus

Differentiation under the integral sign

$$\frac{d}{dx} \left(\int_{a(x)}^{b(x)} f(x,t) dt \right) = f(x,b(x))b'(x) - f(x,a(x))a'(x) + \int_{a(x)}^{b(x)} f_x(x,t) dt.$$

Concavity in 1 dimension

If $g : I \rightarrow \mathbb{R}$ is twice differentiable in the interval I :

concave:

if and only if $g''(x) \leq 0$ for all $x \in I$

strictly concave:

if $g''(x) < 0$ for all $x \in I$

convex:

if and only if $g''(x) \geq 0$ for all $x \in I$

strictly convex if:

$g''(x) > 0$ for all $x \in I$

Multivariate Calculus

The Gradient ∇ of a twice differentiable function f is defined as:

$$\nabla f : \mathbb{R}^d \rightarrow \mathbb{R}^d$$

$$\theta = \begin{pmatrix} \theta_1 \\ \theta_2 \\ \vdots \\ \theta_d \end{pmatrix} \mapsto \begin{pmatrix} \frac{\partial f}{\partial \theta_1} \\ \frac{\partial f}{\partial \theta_2} \\ \vdots \\ \frac{\partial f}{\partial \theta_d} \end{pmatrix}$$

Hessian

The Hessian of f is a symmetric matrix of second partial derivatives of f

$$Hh(\theta) = \nabla^2 h(\theta) = \begin{pmatrix} \frac{\partial^2 h}{\partial \theta_1 \partial \theta_1}(\theta) & \cdots & \frac{\partial^2 h}{\partial \theta_1 \partial \theta_d}(\theta) \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 h}{\partial \theta_d \partial \theta_1}(\theta) & \cdots & \frac{\partial^2 h}{\partial \theta_d \partial \theta_d}(\theta) \end{pmatrix} \in \mathbb{R}^{d \times d}$$

A symmetric (real-valued) $d \times d$ matrix A is:

Positive semi-definite:

$x^T A x \geq 0$ for all $x \in \mathbb{R}^d$.

Positive definite:

$x^T A x > 0$ for all non-zero vectors $x \in \mathbb{R}^d$

Negative semi-definite (resp. negative definite):

$x^T A x$ is negative for all $x \in \mathbb{R}^d - \{0\}$.

Positive (or negative) definiteness implies positive (or negative) semi-definiteness.

If the Hessian is positive definite then f attains a local minimum at a (convex).

If the Hessian is negative definite at a , then f attains a local maximum at a (concave).

If the Hessian has both positive and negative eigenvalues then a is a saddle point for f .

23 Covariance Matrix

Let X be a random vector of dimension $d \times 1$ with expectation μ_X .

Matrix outer products!

$$\Sigma = \mathbb{E}[(X - \mu_X)(X - \mu_X)^T]$$
$$= \mathbb{E}[XX^T] - \mathbb{E}[X]\mathbb{E}[X]^T$$
$$= \mathbb{E}[XX^T] - \mu_X \mu_X^T$$

24 Probability

Unit 6: Derived distributions

$Z = X + Y$ (independent)

$p_Z(z) = \sum_x p_X(x)p_Y(z-x)$

$f_Z(z) = \int_{-\infty}^{\infty} f_X(x)f_Y(z-x)dx$

Sum of independent normals is normal.

Monotonic transformation: $Y = g(X)$

$F_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|, h(y) = g^{-1}(x)$

Unit 6: Deeper view of conditioning

Law of iterated expectations $\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X|Y]]$.

Law of total variance $Var(X) = \mathbb{E}[Var(X|Y)] + Var(\mathbb{E}[X|Y])$

Sum of a random number of independent r.v.'s: $Y = X_1 + \dots + X_N$

$\mathbb{E}[Y] = \mathbb{E}[N] \cdot \mathbb{E}[X]$

$Var(Y) = \mathbb{E}[N] Var(X) + (\mathbb{E}[X])^2 Var(N)$

Unit 8: Limit theorems and classical statistics

Markov inequality: $X \geq 0$ and $a > 0$, then $P(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$

Chebyshev inequality: $c > 0$, then $P(|X - \mathbb{E}[X]| \geq c) \leq \frac{Var(X)}{c^2}$

Convergence in probability: for every $\epsilon > 0$, $P(|X_n - a| \geq \epsilon) \rightarrow 0$

Weak law of large numbers: X_i (i.i.d.), $M_n = \frac{X_1 + \dots + X_n}{n} \rightarrow \mathbb{E}[X]$

Central limit theorem: X_i (i.i.d.), CDF of $\frac{X_1 + \dots + X_n - n\mathbb{E}[X]}{\sqrt{na} \sigma_X} \rightarrow$ standard normal CDF

Unit 9: The Bernoulli and Poisson processes

Bernoulli process

Number of successes/arrivals S in n time slots:

$S = X_1 + \dots + X_n$ (Binomial)

$P(S = k) = \binom{n}{k} p^k (1-p)^{n-k}, k = 0, \dots, n$

$\mathbb{E}[S] = np$

$Var(S) = np(1-p)$

Time until the first success/arrival T_1

$T_1 = \min\{i : X_i = 1\}$ (Geometric)

$P(T_1 = k) = (1-p)^{k-1}p, k = 1, 2, \dots$

$\mathbb{E}[T_1] = \frac{1}{p}$

$Var(T_1) = \frac{1-p}{p^2}$

Time of the k -th success/arrival $Y_k = T_1 + \dots + T_k$ (Pascal), T_i are iid Geometric(p)

$p_{Y_k}(t) = \binom{t-1}{k-1} p^k (1-p)^{t-k}, t = k, k+1, \dots$

$\mathbb{E}[Y_k] = \frac{k}{p}$

$Var(Y_k) = \frac{k(1-p)}{p^2}$

Merged process

$\text{Bernoulli}(p) + \text{Bernoulli}(q) = \text{Bernoulli}(p+q-pq)$

$P(\text{arrival in first process} | \text{arrival}) = \frac{p}{p+q-pq}$

Splitting process

Split arrivals into two streams, using independent coin flips of a coin with bias q :

Resulting streams are Bernoulli, success rates are pq and $p(1-q)$.

The two streams are not independent

Poisson Process

Number of arrivals in interval τ : N_τ (Poisson)

$P(N_\tau = k, \tau) = \frac{(\lambda \tau)^k e^{-\lambda \tau}}{k!}, k = 0, 1, \dots$

$\mathbb{E}[N_\tau] = \lambda \tau$

Time until the first success/arrival T_1

(Exponential(λ))

$T_1 = \min\{i : X_i = 1\}$ (Geometric)

$f_{T_1}(t) = \lambda e^{-\lambda t}, t \geq 0$

$\mathbb{E}[T_1] = \frac{1}{\lambda}$

$Var(T_1) = \frac{1}{\lambda^2}$

Time of the k -th success/arrival: $Y_k = T_1 + \dots + T_k$ (Erlang), T_i are iid Exponentials(λ)

$f_{Y_k}(y) = \frac{\lambda^k y^{k-1} e^{-\lambda y}}{(k-1)!}, y \geq 0$

$\mathbb{E}[Y_k] = \frac{k}{\lambda}$

$Var(Y_k) = \frac{k}{\lambda^2}$

Merged process

$\text{Poisson}(\lambda_1) + \text{Poisson}(\lambda_2) = \text{Poisson}(\lambda_1 + \lambda_2)$

$P(k\text{-th arrival comes from first process}) = \frac{\lambda_1}{\lambda_1 + \lambda_2}$

Independence for different arrivals

Splitting process

Split arrivals into two streams, using independent coin flips of a coin with bias q :

Resulting streams are Poisson, rates λq and $\lambda(1-q)$.

Resulting streams are independent!

Source: Cheatsheet @r2cp
Page 4