

Modeling Toronto Apartment Rental Prices

Austin Hart, Olivia Jiang, Woobin Kim, Raina Tian

2026-07-25

Introduction

This section should describe the background and objectives of the study.

Data Description


We obtained the data from [this](#) Kaggle dataset. The amount of rows in the original dataset before any data cleaning is 1,124. The data is made up of rental prices and contains data on apartment rental units in Toronto collected in 2018. It contains the apartment's geographic information (address, latitude, longitude), information on the rooms in each unit (the number of bedrooms, bathrooms, and dens), and the unit's monthly rental cost in CAD.

Exploratory Data Analysis (EDA)

The total number of rows in the original dataset is 1,124. During data cleaning, we removed 308 duplicates. We also removed 5 rows with unusual values for `price` (\$65/month, \$99/month, \$99/month, \$36,900/month, \$535,000/month). These outliers may skew our results so we decided to remove them. The sample size is now $n = 811$. There are no missing values in any columns.


Before modeling, we want to find out more about the data. The variables we're most interested in are `price`, `bedroom`, `bathroom`, and `den`.

This histogram of `price` shows that the distribution is slightly right-skewed and unimodal around the mean of \$2255. As well, its standard deviation is \$1021, the minimum is \$150, and the maximum is \$9750.




Next, we view the frequencies of each discrete value of `bedroom`, `bathroom`, and `den`. TODO: comment on them!

Frequencies of Room Counts by Room Type



Below are boxplots of `price` split by `bedroom`, `bathroom`, and `den` counts in order to see how `price`'s distribution varies by each of these 3 apartment characteristics. TODO: comment on them!



We're also about the correlations between the variables so we plot a correlation matrix. TODO: comment on this!

```
##          bedroom  bathroom      den      price
## bedroom  1.0000000  0.50147207 -0.36728231  0.37884260
## bathroom 0.5014721  1.00000000 -0.09791753  0.46537487
## den      -0.3672823 -0.09791753  1.00000000 -0.01986721
## price     0.3788426  0.46537487 -0.01986721  1.00000000
```

Models

This section should include at least two statistical models for the same dataset, justifications of the choices of the models, model diagnostics, parameter estimates and their standard errors and p-values, and interpretation of the results.

```
##
## Call:
## lm(formula = price ~ ., data = data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2661.7  -517.0   -45.2    441.3   6515.9
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    677.50     160.03   4.233 2.99e-05 ***
## bedroom         360.51      89.79   4.015 7.36e-05 ***
## bathroom       734.23     114.15   6.432 4.43e-10 ***
## den             277.02     139.37   1.988  0.0477 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 885.9 on 329 degrees of freedom
## Multiple R-squared:  0.2538, Adjusted R-squared:  0.247
## F-statistic: 37.3 on 3 and 329 DF, p-value: < 2.2e-16
```

Conclusion

This section should provide overall conclusions, any limitations in the datasets and analyses, and validity of the conclusions.

References

Any references can go here.

Appendix

```
library(tidyverse)
```