

# Kähler Geometry

Ritwik Chakraborty

Indian Institute of Technology, Kanpur

*ritwikc@iitk.ac.in*

December 12, 2019

# Overview

- 1 Complex Manifolds
- 2 Goal
- 3 Algebraic Dimension
  - Meromorphic functions
  - Moishezon Manifolds
- 4 Vector Bundles, Connection and Curvature
  - Vector Bundles and Almost Complex Structures
  - Hermitian Structures and Connections
- 5 Kähler Manifolds
  - Answers to Some Questions
  - Calabi Conjecture and Uniformization Theorem
- 6 Some more Questions

# Complex Manifolds

**How does one endow a manifold with complex structure?**

# Complex Manifolds

**How does one endow a manifold with complex structure?**

A complex manifold  $X$  of dimension  $n$  is a Hausdorff, second countable space

# Complex Manifolds

## How does one endow a manifold with complex structure?

A complex manifold  $X$  of dimension  $n$  is a Hausdorff, second countable space equipped with a subsheaf of the sheaf of continuous functions on  $X$ , denoted  $\mathcal{O}_X$ ,

# Complex Manifolds

## How does one endow a manifold with complex structure?

A complex manifold  $X$  of dimension  $n$  is a Hausdorff, second countable space equipped with a subsheaf of the sheaf of continuous functions on  $X$ , denoted  $\mathcal{O}_X$ , such that every point  $x \in X$  has a neighbourhood  $U$  and a homeomorphism  $\varphi$  of  $U$  with an open subset  $V$  of  $\mathbb{C}^n$

# Complex Manifolds

## How does one endow a manifold with complex structure?

A complex manifold  $X$  of dimension  $n$  is a Hausdorff, second countable space equipped with a subsheaf of the sheaf of continuous functions on  $X$ , denoted  $\mathcal{O}_X$ , such that every point  $x \in X$  has a neighbourhood  $U$  and a homeomorphism  $\varphi$  of  $U$  with an open subset  $V$  of  $\mathbb{C}^n$  and

$$\mathcal{O}_X|_U = \varphi^*(\mathcal{O}_{\mathbb{C}^n}|_V)$$

where  $\mathcal{O}_{\mathbb{C}^n}$  is the sheaf of holomorphic functions on  $\mathbb{C}^n$ .

# Complex Manifolds

This is equivalent to equipping a topological  $2n$ -manifold  $X$  with a **holomorphic atlas**  $\{U_i, \varphi_i : U_i \rightarrow \varphi_i(U_i) \subset \mathbb{C}^n\}$  such that the transition maps  $\varphi_{ij} := \varphi_i \circ \varphi_j^{-1} \big|_{\varphi_j(U_i \cap U_j)} : \varphi_j(U_i \cap U_j) \rightarrow \varphi_i(U_i \cap U_j)$  are holomorphic for all  $i, j$ .



# Complex Manifolds

This is equivalent to equipping a topological  $2n$ -manifold  $X$  with a **holomorphic atlas**  $\{U_i, \varphi_i : U_i \rightarrow \varphi_i(U_i) \subset \mathbb{C}^n\}$  such that the transition maps  $\varphi_{ij} := \varphi_i \circ \varphi_j^{-1}|_{\varphi_j(U_i \cap U_j)} : \varphi_j(U_i \cap U_j) \rightarrow \varphi_i(U_i \cap U_j)$  are holomorphic for all  $i, j$ .

**Examples:**  $\mathbb{CP}^n$  (Projective space),  $\mathbb{C}^n/\Gamma$  (Complex torii),  
Quotients of the form  $X/G$  (properly discontinuous action)

# Goal

- ① Doing geometry on complex manifolds
- ② Realize that Kähler manifolds are natural sites for seeking relationships between **topological**, **complex** and **geometric** structures on a space.
- ③ Actually see instances of 2

## A General theme

Real manifolds versus complex manifolds

# Meromorphic functions

## Theorem

*For any compact, connected complex manifold  $X$ ,  $\Gamma(X, \mathcal{O}_X) = \mathbb{C}$ .*

# Meromorphic functions

## Theorem

*For any compact, connected complex manifold  $X$ ,  $\Gamma(X, \mathcal{O}_X) = \mathbb{C}$ .*

Partitions of unity on real manifolds make things much simpler there.

# Meromorphic functions

## Theorem

*For any compact, connected complex manifold  $X$ ,  $\Gamma(X, \mathcal{O}_X) = \mathbb{C}$ .*

Partitions of unity on real manifolds make things much simpler there.

A meromorphic function on  $U \subset_{\text{open}} X$  is a map

$$f : U \rightarrow \coprod_{x \in U} \mathbb{C}(\mathcal{O}_{X,x})$$

Sheaf of meromorphic functions:  $\mathcal{K}_X$

Space of global sections:  $K(X)$

# Algebraic dimension

# Algebraic dimension

- $X$  is connected  $\Rightarrow K(X)$  is a field  
We call this the **function field** of  $X$ .

# Algebraic dimension

- $X$  is connected  $\Rightarrow K(X)$  is a field  
We call this the **function field** of  $X$ .
- $\mathcal{O}_X \subset \mathcal{K}_X$



# Algebraic dimension

- $X$  is connected  $\Rightarrow K(X)$  is a field  
We call this the **function field** of  $X$ .
- $\mathcal{O}_X \subset \mathcal{K}_X$
- **Algebraic dimension** of  $X$ ,  $a(X) = \text{trdeg}_{\mathbb{C}}(K(X))$

# Algebraic dimension

- $X$  is connected  $\Rightarrow K(X)$  is a field

We call this the **function field** of  $X$ .

- $\mathcal{O}_X \subset \mathcal{K}_X$

- **Algebraic dimension** of  $X$ ,  $a(X) = \text{trdeg}_{\mathbb{C}}(K(X))$

**Question: Is  $a(X)$  finite? When is it equal to the complex dimension of  $X$  and how “often”?**

# Algebraic dimension

- $X$  is connected  $\Rightarrow K(X)$  is a field

We call this the **function field** of  $X$ .

- $\mathcal{O}_X \subset \mathcal{K}_X$

- **Algebraic dimension** of  $X$ ,  $a(X) = \text{trdeg}_{\mathbb{C}}(K(X))$

**Question:** Is  $a(X)$  finite? When is it equal to the complex dimension of  $X$  and how “often”?

## Theorem

**(Siegel)** *Let  $X$  be a compact, connected complex  $n$ -manifold. Then,*

$$a(X) \leq n$$

Complex Manifolds

Goal

**Algebraic Dimension**

Vector Bundles, Connection and Curvature

Kähler Manifolds

Some more Questions

Meromorphic functions

**Moishezon Manifolds**

# Moishezon Manifolds

# Moishezon Manifolds

## Moishezon Manifolds

A connected, compact complex manifold  $X$  is said to be Moishezon if  $a(X) = \dim_{\mathbb{C}}(X)$

**Example:** Any projective manifold is Moishezon.

# Moishezon Manifolds

## Moishezon Manifolds

A connected, compact complex manifold  $X$  is said to be Moishezon if  $a(X) = \dim_{\mathbb{C}}(X)$

**Example:** Any projective manifold is Moishezon.

**Question:** Is the converse true?

# Complex and Holomorphic Vector Bundles

# Complex and Holomorphic Vector Bundles

A **complex (holomorphic) vector bundle of rank  $r$** :  
 $(E, X, \pi : E \rightarrow X)$  such that



# Complex and Holomorphic Vector Bundles

A **complex (holomorphic) vector bundle of rank  $r$** :

$(E, X, \pi : E \rightarrow X)$  such that:

- $\forall x \in X, \pi^{-1}(x)$  is a  $\mathbb{C}$ -vector space of dimension  $r$

# Complex and Holomorphic Vector Bundles

A **complex (holomorphic) vector bundle of rank  $r$** :

$(E, X, \pi : E \rightarrow X)$  such that:

- $\forall x \in X, \pi^{-1}(x)$  is a  $\mathbb{C}$ -vector space of dimension  $r$
- $X = \bigcup U_i$  and there are diffeomorphisms (biholomorphic maps)  $\psi_i : \pi^{-1}(U_i) \cong U_i \times \mathbb{C}^r$  such that

# Complex and Holomorphic Vector Bundles

A **complex (holomorphic) vector bundle of rank  $r$ :**

$(E, X, \pi : E \rightarrow X)$  such that:

- $\forall x \in X, \pi^{-1}(x)$  is a  $\mathbb{C}$ -vector space of dimension  $r$
- $X = \bigcup U_i$  and there are diffeomorphisms (biholomorphic maps)  $\psi_i : \pi^{-1}(U_i) \cong U_i \times \mathbb{C}^r$  such that:

$$\begin{array}{ccc} U \times \mathbb{C}^r & \xrightarrow{\psi_i^{-1}} & \pi^{-1}(U) \\ & \searrow \pi_U & \swarrow \pi \\ & U & \end{array}$$

and the induced map  $\pi^{-1}(x) \cong \mathbb{C}^r$  is  $\mathbb{C}$ -linear.

# Some Vector Bundles

A key observation:

# Some Vector Bundles

A key observation:

$\{\text{Holomorphic rank } r \text{ vector bundle } \pi : E \rightarrow X\} \leftrightarrow \{\text{holomorphic cocycle } \{(U_i, \psi_{ij} : U_i \cap U_j \rightarrow GL(r, \mathbb{C}))\}\}$

# Some Vector Bundles

A key observation:

$\{\text{Holomorphic rank } r \text{ vector bundle } \pi : E \rightarrow X\} \leftrightarrow \{\text{holomorphic cocycle } \{(U_i, \psi_{ij} : U_i \cap U_j \rightarrow \text{Gl}(r, \mathbb{C}))\}\}$

- Natural operations on vector bundles  $E, F$  on a complex manifold  $X$ :

$$E \oplus F, E \otimes F, E^*, \bigwedge^i E, \det(E) \dots$$

# Some Vector Bundles

A key observation:

$\{\text{Holomorphic rank } r \text{ vector bundle } \pi : E \rightarrow X\} \leftrightarrow \{\text{holomorphic cocycle } \{(U_i, \psi_{ij} : U_i \cap U_j \rightarrow \text{Gl}(r, \mathbb{C}))\}\}$

- Natural operations on vector bundles  $E, F$  on a complex manifold  $X$ :

$$E \oplus F, E \otimes F, E^*, \bigwedge^i E, \det(E) \dots$$

- $\{\{\text{Isomorphism classes of line bundles}\}, \otimes, (-)^*, \mathcal{O}(0)\}$  is an abelian group, called the **Picard group**,  $\text{Pic}(X)$

# Some Vector Bundles

A key observation:

$\{\text{Holomorphic rank } r \text{ vector bundle } \pi : E \rightarrow X\} \leftrightarrow \{\text{holomorphic cocycle } \{(U_i, \psi_{ij} : U_i \cap U_j \rightarrow \text{Gl}(r, \mathbb{C}))\}\}$

- Natural operations on vector bundles  $E, F$  on a complex manifold  $X$ :

$$E \oplus F, E \otimes F, E^*, \bigwedge^i E, \det(E) \dots$$

- $\{\{\text{Isomorphism classes of line bundles}\}, \otimes, (-)^*, \mathcal{O}(0)\}$  is an abelian group, called the **Picard group**,  $\text{Pic}(X)$
- $\text{Pic}(X) \cong H^1(X, \mathcal{O}_X^*)$



# Some Vector Bundles

A key observation:

$\{\text{Holomorphic rank } r \text{ vector bundle } \pi : E \rightarrow X\} \leftrightarrow \{\text{holomorphic cocycle } \{(U_i, \psi_{ij} : U_i \cap U_j \rightarrow GL(r, \mathbb{C}))\}\}$

- Natural operations on vector bundles  $E, F$  on a complex manifold  $X$ :

$$E \oplus F, E \otimes F, E^*, \bigwedge^i E, \det(E) \dots$$

- $\{\{\text{Isomorphism classes of line bundles}\}, \otimes, (-)^*, \mathcal{O}(0)\}$  is an abelian group, called the **Picard group**,  $\text{Pic}(X)$
- $\text{Pic}(X) \cong H^1(X, \mathcal{O}_X^*)$

Follows from two natural isomorphisms:

$$\text{Pic}(X) \cong \check{H}^1(X, \mathcal{O}_X^*) \text{ and } \check{H}^1(X, \mathcal{O}_X^*) \cong H^1(X, \mathcal{O}_X^*)$$

Complex Manifolds

Goal

Algebraic Dimension

Vector Bundles, Connection and Curvature

Kähler Manifolds

Some more Questions

Vector Bundles and Almost Complex Structures

Hermitian Structures and Connections

# Almost Complex Structures

# Almost Complex Structures

**Almost complex manifold:** a smooth manifold  $X$  equipped with a vector bundle endomorphism  $I : TX \rightarrow TX$  such that  $I \circ I = -id$ .  $I$  is called an almost complex structure on  $X$ .

# Almost Complex Structures

**Almost complex manifold:** a smooth manifold  $X$  equipped with a vector bundle endomorphism  $I : TX \rightarrow TX$  such that  $I \circ I = -id$ .  $I$  is called an almost complex structure on  $X$ .

- 1 Any complex manifold admits a natural almost complex structure:

$$I : T_x U \rightarrow T_x U \quad \partial/\partial x_i \mapsto \partial/\partial y_i \quad \partial/\partial y_i \mapsto -\partial/\partial x_i$$

# Almost Complex Structures

**Almost complex manifold:** a smooth manifold  $X$  equipped with a vector bundle endomorphism  $I : TX \rightarrow TX$  such that  $I \circ I = -id$ .  $I$  is called an almost complex structure on  $X$ .

- 1 Any complex manifold admits a natural almost complex structure:

$$I : T_x U \rightarrow T_x U \quad \partial/\partial x_i \mapsto \partial/\partial y_i \quad \partial/\partial y_i \mapsto -\partial/\partial x_i$$

- 2  $T_{\mathbb{C}}X = T^{1,0}X \oplus T^{0,1}X$

# Almost Complex Structures

**Almost complex manifold:** a smooth manifold  $X$  equipped with a vector bundle endomorphism  $I : TX \rightarrow TX$  such that  $I \circ I = -id$ .  $I$  is called an almost complex structure on  $X$ .

- Any complex manifold admits a natural almost complex structure:

$$I : T_x U \rightarrow T_x U \quad \partial/\partial x_i \mapsto \partial/\partial y_i \quad \partial/\partial y_i \mapsto -\partial/\partial x_i$$

- $T_{\mathbb{C}}X = T^{1,0}X \oplus T^{0,1}X$

- $I|_{T^{1,0}X} \stackrel{\xi}{\cong} i \cdot id$  and  $I|_{T^{0,1}X} \stackrel{\eta}{\cong} -i \cdot id$

# Almost Complex Structures

**Almost complex manifold:** a smooth manifold  $X$  equipped with a vector bundle endomorphism  $I : TX \rightarrow TX$  such that  $I \circ I = -id$ .  $I$  is called an almost complex structure on  $X$ .

- Any complex manifold admits a natural almost complex structure:

$$I : T_x U \rightarrow T_x U \quad \partial/\partial x_i \mapsto \partial/\partial y_i \quad \partial/\partial y_i \mapsto -\partial/\partial x_i$$

- $T_{\mathbb{C}}X = T^{1,0}X \oplus T^{0,1}X$

- $I|_{T^{1,0}X} \stackrel{\xi}{\cong} i \cdot id$  and  $I|_{T^{0,1}X} \stackrel{\eta}{\cong} -i \cdot id$

- $\mathcal{A}_X^{p,q}(E)$ , the sheaf  $U \mapsto \mathcal{A}_X^{p,q}(E)(U) := \Gamma(U, \wedge^{p,q} X \otimes E)$

Is the converse to 1 true?

# Hermitian Structures and Hermitan Vector Bundles

A Riemannian metric  $g$  on  $(X, I)$  is an **hermitian structure** on  $X$  if for any point  $x \in X$  the scalar product  $g_x$  on  $T_x X$  is compatible with the almost complex structure  $I_x$ . The induced real  $(1,1)$  form  $\omega := g(I(-), (-))$  is called the **fundamental form**. Locally,

$$\omega = (i/2) \sum_{i,j=1}^n h_{ij} dz^i \wedge d\bar{z}^j$$



# Hermitian Structures and Hermitian Vector Bundles

A Riemannian metric  $g$  on  $(X, I)$  is an **hermitian structure** on  $X$  if for any point  $x \in X$  the scalar product  $g_x$  on  $T_x X$  is compatible with the almost complex structure  $I_x$ . The induced real  $(1,1)$  form  $\omega := g(I(-), (-))$  is called the **fundamental form**. Locally,

$$\omega = (i/2) \sum_{i,j=1}^n h_{ij} dz^i \wedge d\bar{z}^j$$

Suitably generalizing this from  $TX$  to arbitrary complex vector bundles:

**Hermitian vector bundle  $(E, h)$ :** A complex vector bundle  $E$  on a smooth manifold  $M$ ,  $E_x$  is endowed with a hermitian inner product  $h_x$ , smooth in  $x$ .

# Connections and Curvature

A **connection** on a complex vector bundle  $E$  is a  $\mathbb{C}$ -linear sheaf homomorphism  $\nabla : \mathcal{A}^0(E) \rightarrow \mathcal{A}^1(E)$  which satisfies the Leibniz rule:

$$\nabla(f \cdot s) = d(f) \otimes s + f \cdot \nabla(s)$$

# Connections and Curvature

A **connection** on a complex vector bundle  $E$  is a  $\mathbb{C}$ -linear sheaf homomorphism  $\nabla : \mathcal{A}^0(E) \rightarrow \mathcal{A}^1(E)$  which satisfies the Leibniz rule:

$$\nabla(f \cdot s) = d(f) \otimes s + f \cdot \nabla(s)$$

This naturally extends to another Leibniz map  $\nabla : \mathcal{A}^k(E) \rightarrow \mathcal{A}^{k+1}(E)$  as:

$$\nabla(\alpha \otimes s) = d(\alpha) \otimes s + (-1)^k \alpha \wedge \nabla(s)$$

# Connections and Curvature

A **connection** on a complex vector bundle  $E$  is a  $\mathbb{C}$ -linear sheaf homomorphism  $\nabla : \mathcal{A}^0(E) \rightarrow \mathcal{A}^1(E)$  which satisfies the Leibniz rule:

$$\nabla(f \cdot s) = d(f) \otimes s + f \cdot \nabla(s)$$

This naturally extends to another Leibniz map

$\nabla : \mathcal{A}^k(E) \rightarrow \mathcal{A}^{k+1}(E)$  as:

$$\nabla(\alpha \otimes s) = d(\alpha) \otimes s + (-1)^k \alpha \wedge \nabla(s)$$

The **curvature**  $F_\nabla$  of a connection  $\nabla$  on a vector bundle  $E$  is defined as the composition:

$$F_\nabla := \nabla \circ \nabla : \mathcal{A}^0(E) \rightarrow \mathcal{A}^1(E) \rightarrow \mathcal{A}^2(E)$$

# Chern Connection

# Chern Connection

**What happens when  $E$  is equipped with a Hermitian structure  $h$ ?**

# Chern Connection

**What happens when  $E$  is equipped with a Hermitian structure  $h$ ?**

A connection  $\nabla$  on  $E$  is said to be a **hermitian connection** with respect to  $h$  if:

$$d(h(s_1, s_2)) = h(\nabla(s_1), s_2) + h(s_1, \nabla s_2).$$

# Chern Connection

**What happens when  $E$  is equipped with a Hermitian structure  $h$ ?**

A connection  $\nabla$  on  $E$  is said to be a **hermitian connection** with respect to  $h$  if:

$$d(h(s_1, s_2)) = h(\nabla(s_1), s_2) + h(s_1, \nabla s_2).$$

Another upgrade: **What if  $E$  is a holomorphic vector bundle?**



# Chern Connection

**What happens when  $E$  is equipped with a Hermitian structure  $h$ ?**

A connection  $\nabla$  on  $E$  is said to be a **hermitian connection** with respect to  $h$  if:

$$d(h(s_1, s_2)) = h(\nabla(s_1), s_2) + h(s_1, \nabla s_2).$$

Another upgrade: **What if  $E$  is a holomorphic vector bundle?**  
There is a canonical choice!

# Chern Connection

**What happens when  $E$  is equipped with a Hermitian structure  $h$ ?**

A connection  $\nabla$  on  $E$  is said to be a **hermitian connection** with respect to  $h$  if:

$$d(h(s_1, s_2)) = h(\nabla(s_1), s_2) + h(s_1, \nabla s_2).$$

Another upgrade: **What if  $E$  is a holomorphic vector bundle?**  
There is a canonical choice!

## Theorem

*Let  $(E, h)$  be a holomorphic vector bundle. Then there exists a **unique hermitian connection**  $\nabla$  that is **compatible with the holomorphic structure**, i.e  $\nabla^{1,0} = \bar{\partial}$ . This connection is called the **Chern connection** on  $(E, h)$ .*

Complex Manifolds

Goal

Algebraic Dimension

Vector Bundles, Connection and Curvature

**Kähler Manifolds**

Some more Questions

Answers to Some Questions

Calabi Conjecture and Uniformization Theorem

# Riemannian vs Complex Geometry

A **Kähler structure** is a hermitian structure  $g$  for which the fundamental form  $\omega$  is closed, i.e.  $d\omega = 0$ . In this case,  $\omega$  is called a Kähler form on  $X$ .

**Examples:**  $\mathbb{CP}^n$  with the Fubini-Study metric, (in the standard chart  $U_i$ )  $\omega_{FS} = \frac{i}{2\pi} \partial \bar{\partial} \log \left( \sum_{l=0}^n \left| \frac{z_l}{z_i} \right|^2 \right)$ , Complex torii,  $\mathbb{C}^n$ ,

# Riemannian vs Complex Geometry

A **Kähler structure** is a hermitian structure  $g$  for which the fundamental form  $\omega$  is closed, i.e.  $d\omega = 0$ . In this case,  $\omega$  is called a Kähler form on  $X$ .

**Examples:**  $\mathbb{CP}^n$  with the Fubini-Study metric, (in the standard chart  $U_i$ )  $\omega_{FS} = \frac{i}{2\pi} \partial \bar{\partial} \log(\sum_{l=0}^n |\frac{z_l}{z_i}|^2)$ , Complex torii,  $\mathbb{C}^n$ ,

## Theorem

*Let  $(X, g)$  be a Kähler manifold. Then under the isomorphism  $\xi : TX \cong T^{1,0}X$ , the Chern connection  $\nabla$  on the holomorphic tangent space  $T^{1,0}X$  corresponds to the Levi-Civita connection  $D$ .*

Complex Manifolds

Goal

Algebraic Dimension

Vector Bundles, Connection and Curvature

**Kähler Manifolds**

Some more Questions

Answers to Some Questions

Calabi Conjecture and Uniformization Theorem

# Answers to Some Questions Raised so Far

## Theorem

**[Moishe]** *A Moishezon space is a projective algebraic variety if and only if it admits a Kähler metric.*

## Theorem

**[Siu],[Demailly]** *Let  $X$  be a compact manifold carrying an almost positive line bundle. Then  $X$  is a Moishezon space.*

# Answers to Some Questions Raised so Far

## Theorem

**[Moishe]** *A Moishezon space is a projective algebraic variety if and only if it admits a Kähler metric.*

## Theorem

**[Siu],[Demailly]** *Let  $X$  be a compact manifold carrying an almost positive line bundle. Then  $X$  is a Moishezon space.*

A real  $(1,1)$ -form  $\alpha$  is called (semi)-positive if for all non-zero holomorphic tangent vector  $v \in T^{1,0}X$ , one has  $-i\alpha(v, \bar{v}) > 0$  (resp.  $\geq 0$ ).



- The pair  $(I, \omega)$  on  $X$  gives us a natural Riemannian metric  $g$  on  $X$ , defined as  $g(u, v) = \omega(u, I(v))$ .

- The pair  $(I, \omega)$  on  $X$  gives us a natural Riemannian metric  $g$  on  $X$ , defined as  $g(u, v) = \omega(u, I(v))$ .
- The Riemannian Ricci curvature is then given by  $Rc_g(u, v) = \text{Ric}(\omega)(u, I(v))$ , where  $\text{Ric}(\omega) = -i\partial\bar{\partial}\log(\omega^n)$  is the Ricci form.

- The pair  $(I, \omega)$  on  $X$  gives us a natural Riemannian metric  $g$  on  $X$ , defined as  $g(u, v) = \omega(u, I(v))$ .
- The Riemannian Ricci curvature is then given by  $Rc_g(u, v) = \text{Ric}(\omega)(u, I(v))$ , where  $\text{Ric}(\omega) = -i\partial\bar{\partial}\log(\omega^n)$  is the Ricci form.
- The **first Chern class**  $c_1(L)$  of a holomorphic line bundle  $L \in \text{Pic}(X)$  is its image under the boundary map:

$$c_1 : H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \mathbb{Z})$$

- The pair  $(I, \omega)$  on  $X$  gives us a natural Riemannian metric  $g$  on  $X$ , defined as  $g(u, v) = \omega(u, I(v))$ .
- The Riemannian Ricci curvature is then given by  $Rc_g(u, v) = \text{Ric}(\omega)(u, I(v))$ , where  $\text{Ric}(\omega) = -i\partial\bar{\partial}\log(\omega^n)$  is the Ricci form.
- The **first Chern class**  $c_1(L)$  of a holomorphic line bundle  $L \in \text{Pic}(X)$  is its image under the boundary map:

$$c_1 : H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \mathbb{Z})$$

## Conjecture Calabi

Every real  $(1,1)$ -form in  $c_1(X)$  is the Ricci for some Kähler form.

Complex Manifolds

Goal

Algebraic Dimension

Vector Bundles, Connection and Curvature

**Kähler Manifolds**

Some more Questions

Answers to Some Questions

Calabi Conjecture and Uniformization Theorem

# Uniformization theorem

# Uniformization theorem

## Theorem

**Uniformization theorem** *Given any compact, oriented Riemannian surface  $(S, g)$  there exists a metric  $\tilde{g}$  in the conformal class  $[g] = \{e^f g \mid f \in C^\infty(S, \mathbb{R})\}$  having constant curvature  $\text{sgn}(\chi(S))$*

# Uniformization theorem

## Theorem

**Uniformization theorem** *Given any compact, oriented Riemannian surface  $(S, g)$  there exists a metric  $\tilde{g}$  in the conformal class  $[g] = \{e^f g \mid f \in C^\infty(S, \mathbb{R})\}$  having constant curvature  $\text{sgn}(\chi(S))$*

- Equivalent to the equation  $\text{Ric}(\omega_g) = \text{sgn}(\chi(S)) \cdot \omega_g$

# Uniformization theorem

## Theorem

**Uniformization theorem** *Given any compact, oriented Riemannian surface  $(S, g)$  there exists a metric  $\tilde{g}$  in the conformal class  $[g] = \{e^f g \mid f \in C^\infty(S, \mathbb{R})\}$  having constant curvature  $\text{sgn}(\chi(S))$*

- Equivalent to the equation  $\text{Ric}(\omega_g) = \text{sgn}(\chi(S)) \cdot \omega_g$
- Generalizing to Kähler manifolds, a Kähler metric  $\omega$  on  $X$  is said to be **Kähler-Einstein** is  $\text{Ric}(\omega) = \lambda \omega$ .



## Theorem

**(Aubin, Yau)** If  $c_1(X) = 0$  (resp.  $< 0$ ) then for any  $\alpha \in \mathcal{C}_X$ , there exists an  $\omega \in \alpha$  such that  $\text{Ric}(\omega) = 0$  (resp.  $< 0$ ).

# Some More Questions

- Using the homomorphism  $Aut(X) \rightarrow Aut(Pic(X))$  to study action of elements of  $Aut(X)$  on geometric objects on  $X$ .
- Deformation of vector bundles and limiting geometries

# The End