

A guide for checking the paper “Pattern-avoiding binary arrays and a Stirling-type table”

This note accompanies the 39-page version of 10 October 2026. Page numbers refer to that version. The repository is <https://github.com/r5x8ksva-del/a207123-explicit-formulas>.

1. What the paper claims

- **Structure for fixed m** (Theorems A and B; Theorems 3.1 and 4.6): the minimal order of a linear recurrence in k for $U_k(m)$ is exactly $3m + 1$, and $U_k(m)$ has a positive double-sum formula in Stirling and r -Stirling numbers.
- **All recurrences** (Theorem C; Theorems 5.1, 5.4, B.1, Proposition 5.6, Corollary 5.7): the generating function is not D-finite; the operators annihilating U on a quadrant form the left ideal $\mathcal{OL}_1$, and those annihilating the triangle N form $\mathcal{OL}_N$, with exact dimension counts; U is not Stirling-like in the sense of Kauers.
- **Real roots** (Theorem D; Theorem 8.1, Corollaries 8.2 and 8.13): the polynomials h_k have only real, simple zeros, $\lfloor k/3 \rfloor$ of them greater than 1; equivalently the chain polynomials of the posets Λ_k are real-rooted.
- **Further results**: obstructions to specific single-sum formulas (Theorem E, Section 6) and the near-diagonal of the triangle (Theorem F, Section 7).

The paragraph *What is new* on p. 5 states which tools are standard and which steps are specific to these arrays.

2. What is machine-checked

The results in Tables 5 and 6 (pp. 30–31) are proved in Lean 4 with Mathlib from the definitions of $a_k(n)$ and $U_k(m)$: the reduction, the basic recurrence, the triangle N , the exact orders (Theorem 3.1), Theorem 3.2 (the roots, the expansion over them and the asymptotics) with Remark 3.3 (its numerical values by interval arithmetic), Corollary 3.4 with Remark 3.5 (the OEIS recurrences for the columns and the rows, and the minimality of the row orders), the block decomposition (Theorem 4.2), the generating function in x and y and the coefficient formula (Theorem 4.4, Lemma 4.5, including the set-partition meaning of $H(m, s, j)$; Proposition 4.7 by counting) and the explicit formula (Theorem 4.6), through a refined recurrence, the non-D-finiteness (Theorem 5.1), the ${}_1F_1$ form of $\sum_m t^m / P_m$ (Remark 5.2), the descriptions $\text{Rel}(U) = \mathcal{OL}_1$ and $\text{Rel}(N) = \mathcal{OL}_N$ with the dimension formulas (Theorems 5.4 and B.1), Corollary 5.5, Proposition 5.6, Corollary 5.7, Section 6.1 except the last two sentences of Remark 6.4 (no single sum $\sum_s A(s) \binom{k+c-2s}{m+s+d}$ for $U_k(m)$, nor for the sequences ending with an ascent; of the shapes $\binom{k+c-\alpha s}{\beta s+d}$ with $\alpha, \beta \geq 0$ only $\alpha + \beta = 1$ occurs; the formal proofs use coordinates with respect to $1, y, \dots, y^{\alpha+\beta-1}$ and Vieta’s formulas instead of residues, poles and norms), and the near-diagonal theorem (Theorem 7.1, Proposition 7.2), Lemmas 8.4–8.9, Propositions 8.10 and 8.11 and Theorem 8.1 (the zeros of every n_k are real and, except -1 , simple; those of h_k are real and simple), Corollary 8.2 (each row of N is positive, strictly log-concave and unimodal; proved without Newton’s inequalities), Lemma 8.12 and Corollary 8.13 (where the zeros of h_k lie and how often its coefficients change sign, with Descartes’ rule from Mathlib), Corollary 8.14 (the central limit theorem; the formal proof estimates characteristic functions directly instead of citing the Lindeberg–Feller theorem), Corollary 8.15 and Remark 8.16 ($N(k, q)$ counts chains in $\tilde{\Lambda}_k$, h_k is the h -polynomial of the order complex and has a negative coefficient for $k \geq 3$). The axiom check reports only `propext`, `Classical.choice` and `Quot.sound`. Lean checks the proofs, not whether the formal statements say what the paper says: **Appendix A (pp. 32–34) transcribes the basic definitions that carry the meaning**, and comparing them with the paper is a short and useful check; the definitions added for later entries of Tables 5 and 6 (blocks, c_m , τ_m , values of rational functions, interlacing, the law of X_k , chains in $\tilde{\Lambda}_k$, the binomial convention of Theorem 6.1, the sequences ending with an ascent in Theorem 6.2, the sums in Section 6.1) are in the Lean files and have been compared only by AI.

3. Where a human check is most needed

The proofs were written with substantial assistance from an AI model and cross-checked by further AI runs (see the statement at the end of the paper). The parts below are not formalized, so an independent reading by an expert is the main remaining safeguard. In order of priority:

1. **Section 6, obstructions (pp. 16–22)**. Theorem 6.5 (two single sums) is computer-assisted: a sieve modulo 5040 and Skolem’s p -adic method (Propositions 6.8–6.10, Lemma 6.11); the scripts are in the repository. The

written proofs of Theorems 6.1–6.3 use residues or poles on fibres and norms in number fields; the theorems are formalized by another route.

2. **Section 4, block decomposition (pp. 10–12):** the explicit bijection of Proposition 4.7 (the formal version only compares the two cardinalities).
3. **Section 5:** Remark 5.8, the step from Kauers’ definition of Stirling-like sequences to the formalized Corollary 5.7, and the description of $\mathcal{O}(k, m)L_1$ by normal forms that the formal Corollary 5.5 uses instead of constructing $\mathcal{O}(k, m)$.
4. **Section 8, real-rootedness (pp. 24–29):** formalized except the statements of Remark 8.16 quoted from the literature (the Boolean lattice, Cohen–Macaulayness). Some written proofs differ from the formal ones, which replace the upper half-plane by $g = cf + \sum_a w_a f/(z - a)$ (Lemma 8.5 itself is formalized separately) and the Lindeberg–Feller theorem by a direct estimate of characteristic functions.

Appendix B (pp. 34–37) is formalized, but its written proof is new; comments on its readability are welcome.

4. Questions on novelty

Searches in arXiv, OpenAlex, Crossref and Google Scholar found no previous work on these arrays and no precedent for the two points below; MathSciNet, zbMATH and Web of Science have not been searched yet. An expert’s knowledge would help most here: (a) Is real-rootedness known for chain polynomials of a class of posets containing Λ_k , or for chain polynomials with zeros outside $[-1, 0]$? (b) Has the annihilator ideal of a bivariate sequence been determined for a generator that is not of Stirling-like shape, as in Theorems 5.4 and B.1?

5. Reproducing the computations

From the repository root (Python 3; only `verify_all.py` needs `numpy`):

- `python3 verify_all.py` — 414 automated checks of the results and numbers, about 11 minutes;
- `python3 code/main_extra/check_paper_numbers.py paper/main.tex` — recomputes the numbers printed in the paper (32 checks);
- `python3 code/main_extra/check_appendix_lean.py paper/main.tex` — compares Appendix A with the Lean sources (35 declarations);
- `python3 code/main_extra/check_appendix_reln.py` — exact checks of the operator identities of Lemma B.2 (14 checks);
- `python3 code/main_extra/check_chains_negzeros.py` — Corollary 8.15 and Remark 8.16 (6 checks).

Lean: `toolchain leanprover/lean4:v4.34.1` with `Mathlib v4.34.1` (see `lean/lean-toolchain` and `lean/lake-manifest.json`). The usual `Mathlib` workflow in `lean/` (`lake exe cache get`, then `lake build`) should apply, but the author has built the development only on Windows, with `lean/run_lean_checks_direct.sh`; a single module needs up to about 8 GB of memory. `lake env lean Axioms.lean` prints the axioms used.