

# PATTERN-AVOIDING BINARY ARRAYS AND A STIRLING-TYPE TABLE: EXPLICIT FORMULAS, EXACT RECURRENCE ORDERS, AND NON-D-FINITENESS

YUNKUN SONG

**ABSTRACT.** Let  $a_k(n)$  be the number of  $n \times k$  binary arrays in which no row contains 001 or 010 and no column contains 001 or 011 as three consecutive entries (OEIS A207123). Splitting the rows by parity gives  $a_k(n) = U_k(\lceil n/2 \rceil) U_k(\lfloor n/2 \rfloor)$ , where  $U_k(m)$  counts the sequences in  $\{0, \dots, m\}^k$  in which every three consecutive entries  $(a, b, c)$  satisfy  $b = c$  or  $a \geq \max(b, c)$ ; equivalently,  $U_k(m)$  counts the multichains of length  $m$  in a subposet  $\Lambda_k$  of the Boolean lattice. A unique factorization into blocks gives a positive double-sum formula for  $U_k(m)$  in terms of Stirling and  $r$ -Stirling numbers, and for each  $m$  the minimal order of a linear recurrence in  $k$  is exactly  $3m + 1$ . The bivariate generating function is not D-finite, and every polynomial-coefficient recurrence that holds on a quadrant is a left multiple of the defining four-term recurrence; in particular  $U$  is not Stirling-like in the sense of Kauers. We also determine all binomial shapes with nonnegative parameters in which  $U_k(m)$  is a single sum; the shape of the explicit formula is not among them. The numerators  $h_k(t)$  of  $\sum_m U_k(m)t^m = h_k(t)/(1-t)^{k+1}$ , analogues of the Eulerian polynomials, have only real and simple zeros; equivalently, the chain polynomials of the posets  $\Lambda_k$  are real-rooted, and the rows of the associated Stirling-type triangle are log-concave. The near-diagonal entries  $N(k, k-d)$  of this triangle are polynomials in  $k$  exactly from  $k = 2d + 2$  on. The reduction, the exact orders, the explicit formula, the non-D-finiteness, the description of the quadrant recurrences and the near-diagonal theorem are formalized in Lean 4 with Mathlib.

## 1. INTRODUCTION

**1.1. The arrays.** Let  $a_k(n)$  be the number of  $n \times k$  matrices over  $\{0, 1\}$  such that, reading each row from left to right, no three consecutive entries form 001 or 010, and, reading each column from top to bottom, no three consecutive entries form 001 or 011. The table  $T(n, k) = a_k(n)$  is OEIS A207123, contributed by R. H. Hardin, and its columns  $k = 3, \dots, 7$  are A207118–A207122 [23]. The OEIS records empirical linear recurrences for these columns and for the first rows; as of October 2026 they are marked as empirical or conjectural.

The column condition says exactly that a 0 is never followed, two places further down, by a 1. Hence the odd-numbered rows and the even-numbered rows are, column by column, weakly decreasing, and the two halves do not interact. Encoding each column of each half by its number of ones (its *height*) turns the row condition into a condition on three consecutive heights. This gives (Proposition 2.2)

$$(1) \quad a_k(n) = U_k(\lceil n/2 \rceil) U_k(\lfloor n/2 \rfloor),$$

where  $U_k(m)$  is the number of sequences  $(h_1, \dots, h_k) \in \{0, \dots, m\}^k$  such that every triple  $(a, b, c) = (h_j, h_{j+1}, h_{j+2})$  is *good*:

$$b = c \quad \text{or} \quad a \geq \max(b, c).$$

Equivalently,  $U_k(m)$  counts the words of length  $k$  over the ordered alphabet  $\{0, \dots, m\}$  that avoid the six consecutive patterns 112, 121, 123, 132, 213 and 231 (for patterns in words see [15]). The

---

2020 *Mathematics Subject Classification.* Primary 05A15; Secondary 05A20, 06A07, 11B73, 26C10, 33F10, 68V20.

*Key words and phrases.* pattern-avoiding arrays, Stirling numbers,  $r$ -Stirling numbers, D-finite power series, Ore algebras, real-rooted polynomials, chain polynomials, interlacing, formal verification.

The author is an undergraduate student in the College of Arts and Sciences at Boston University.

column half of this argument is the same elementary observation that C. Krause used in June 2026 to prove formulas for the related arrays A202093–A202099, whose rows and columns both avoid 001 and 011 [18]; there the row condition also decouples and the count is a product of binomial coefficients. For our row condition it does not, and the whole difficulty is in the table  $U_k(m)$ .

Table 1 shows a few values. The numbers  $U_k(1)$  count the binary words of length  $k$  that avoid 001 and 010. In OEIS terms,  $U_k(1) = A038718(k+2)$ ,  $U_3(m) = A084990(m+1)$  and  $U_4(m) = A326247(m+2)$ . The last identity holds because  $U_4(m) = \binom{m+2}{2}^2 - 4\binom{m+2}{4}$ , the number of ordered pairs of 2-element subsets of an  $(m+2)$ -element set that neither cross nor nest, which is the quantity counted by A326247; we know no structural reason for this coincidence.

| $k \backslash m$ | 0 | 1  | 2   | 3    | 4    | 5     |
|------------------|---|----|-----|------|------|-------|
| 0                | 1 | 1  | 1   | 1    | 1    | 1     |
| 1                | 1 | 2  | 3   | 4    | 5    | 6     |
| 2                | 1 | 4  | 9   | 16   | 25   | 36    |
| 3                | 1 | 6  | 17  | 36   | 65   | 106   |
| 4                | 1 | 9  | 32  | 80   | 165  | 301   |
| 5                | 1 | 14 | 64  | 192  | 457  | 938   |
| 6                | 1 | 21 | 119 | 419  | 1136 | 2604  |
| 7                | 1 | 31 | 214 | 873  | 2669 | 6778  |
| 8                | 1 | 46 | 388 | 1837 | 6334 | 17802 |

TABLE 1. The numbers  $U_k(m)$  for  $0 \leq k \leq 8$  and  $0 \leq m \leq 5$ .

**1.2. Main results.** Let  $\Lambda_k \subseteq \{0,1\}^k$  be the set of binary words of length  $k$  that avoid 001 and 010, ordered componentwise. Then  $U_k(m)$  is the number of multichains of length  $m$  in  $\Lambda_k$  (Proposition 2.4), and the triangle  $N(k, q)$  and the polynomials  $h_k$  introduced below count chains in  $\Lambda_k$  and are  $h$ -polynomials of order complexes; for the whole Boolean lattice  $\{0,1\}^k$  the same constructions give  $(m+1)^k$ , the numbers  $q!\binom{k}{q}$  and the Eulerian polynomials (Remark 8.16). Accordingly, several of our results parallel classical facts: the annihilator of the Stirling numbers  $\{k^n\}$  is generated by their triangular recurrence [13, Example 5], the near-diagonals  $\{k_{k-d}^k\}$  are polynomials in  $k$  [11], and the Eulerian polynomials are real-rooted [3]; compare Theorems D(3), E and F. Theorem D(4) and the zeros of  $h_k$  greater than 1 (Theorem F) show where the analogy ends.

Throughout,  $b_i(x) = 1 - x - ix^3$ ,

$$P_m = \prod_{i=0}^m b_i, \quad W_m = 1 + x^2 \sum_{j=1}^m j P_{j-1} \quad (P_{-1} = 1), \quad G_m(x) = \sum_{k \geq 0} U_k(m) x^k.$$

The starting point is the four-term recurrence (Lemma 2.5)

$$(2) \quad U_k(m) = U_k(m-1) + U_{k-1}(m) + m U_{k-3}(m) \quad (k \geq 1, m \geq 0)$$

with suitable boundary values.

**Theorem A** (Exact recurrence order; Theorem 3.1). *For every  $m \geq 0$ ,  $G_m = W_m/P_m$  and  $\gcd(W_m, P_m) = 1$ . Consequently the minimal order of a linear recurrence with constant coefficients satisfied by  $(U_k(m))_{k \geq 0}$  is exactly  $3m+1$ , and its characteristic polynomial is  $(y-1) \prod_{i=1}^m (y^3 - y^2 - i)$ .*

Theorem 3.2 gives the first two terms of the asymptotics: for  $m \geq 1$ ,  $U_k(m) = c_m \rho_m^k - c_{m-1} \rho_{m-1}^{k+3} + O(\tau_m^k)$  as  $k \rightarrow \infty$ , where  $\rho_m$  is the real root of  $y^3 = y^2 + m$ , the constants  $c_m > 0$  are explicit, and  $\tau_m < \rho_{m-1}$ .

**Theorem B** (Positive explicit formula; Theorem 4.6). *With Stirling numbers of the second kind  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  and  $H(m, s, j) = h_s(j, j+1, \dots, m)$ , which for  $1 \leq j \leq m$  is the  $r$ -Stirling number  $\left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\}_j$  of Broder [5],*

$$U_k(m) = \sum_{s \geq 0} \left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\} \binom{k+m-2s}{k-3s} + \sum_{j=1}^m j \sum_{s \geq 0} H(m, s, j) \binom{k-2+m-j-2s}{k-2-3s}.$$

*Every term is a nonnegative integer with a combinatorial meaning, and the formula refines according to the number of ascents.*

**Theorem C** (Obstructions to specific summation formulas; Theorems 6.1, 6.2, 6.3 and 6.5).

- (1) *Let  $m \geq 1$  and let  $\alpha, \beta \geq 0$  be integers, not both 0. There are integers  $c, d, s_0$  and  $k_0 \geq 0$  and complex numbers  $A(s)$  independent of  $k$  such that*

$$U_k(m) = \sum_{s \geq s_0} A(s) \binom{k+c-\alpha s}{\beta s+d} \quad \text{for all } k \geq k_0$$

*if and only if  $\alpha + \beta = 1$ . In particular there is no such single sum of the shape  $(\alpha, \beta) = (2, 1)$  of the explicit formula.*

- (2) *Let  $m \geq 2$ . For large  $k$ ,  $U_k(m)$  is not the sum of two single sums of the shape  $(2, 1)$ ; three of them always suffice.*  
 (3) *For the number of sequences that end with an ascent, single sums of the shape  $(2, 1)$  are excluded for  $m \geq 2$ , and sums of two of them for  $m \geq 3$ .*

Part (1) shows that statements of this kind must specify the shape: the shapes with  $\alpha + \beta = 1$ , for instance Newton's forward difference series  $\sum_s A(s) \binom{k}{s}$ , are available for every sequence. None of the parts excludes every shorter formula.

**Theorem D** (Non-D-finiteness and all quadrant recurrences; Theorems 5.1, 5.4 and Corollary 5.7).

- (1)  *$F(x, t) = \sum_{k,m} U_k(m) x^k t^m$  satisfies no nonzero linear differential equation in  $x$  with polynomial coefficients; hence  $F$  is not  $D$ -finite.*  
 (2) *There is no nontrivial recurrence*

$$\sum_{a \leq A, b \leq B} p_{ab}(k) U_{k-a}(m-b) = 0$$

*with polynomial coefficients depending only on  $k$  that holds on a quadrant  $k \geq k_0, m \geq m_0$ .*

- (3) *In the Ore algebra  $\mathbb{C}[k, m] \langle X, E^{-1} \rangle$  of backward shifts, the operators annihilating  $U$  on some quadrant form exactly the left ideal generated by  $L_1 = 1 - E^{-1} - X - mX^3$ . The space of such operators with support in  $[0, A] \times [0, B]$  and coefficients of total degree at most  $D$  has dimension  $(A-2)BD(D+1)/2$  when  $A \geq 3, B \geq 1, D \geq 1$ , and 0 otherwise.*  
 (4) *Every nonzero such operator has three monomials whose exponent vectors span a lattice triangle of normalized area at least 3. In particular  $U$  is not Stirling-like in the sense of Kauers [13].*

The triangle  $N(k, q)$ , the number of good words of length  $k$  that use exactly the letters  $1, \dots, q$ , plays for  $U$  the role that the numbers  $q! \left\{ \begin{smallmatrix} k \\ q \end{smallmatrix} \right\}$  of surjections play for powers (Remark 8.16 makes this precise):

$$U_k(m) = \sum_q N(k, q) \binom{m+1}{q}, \quad \text{just as} \quad m^k = \sum_q q! \left\{ \begin{smallmatrix} k \\ q \end{smallmatrix} \right\} \binom{m}{q}.$$

**Theorem E** (Near-diagonal; Theorem 7.1). *For every  $d \geq 0$  there is a polynomial  $p_d$  of degree  $2d$  with leading coefficient  $2/(2^d d!)$  such that  $N(k, k-d) = p_d(k)$  for all  $k \geq 2d+2$ , and  $N(2d+1, d+1) \neq p_d(2d+1)$ . For example  $N(k, k-1) = k^2 - k - 4$  for  $k \geq 4$ .*

**Theorem F** (Real roots; Theorem 8.1, Corollaries 8.2 and 8.13). *For  $k \geq 0$  define the polynomial  $h_k$  by  $\sum_{m \geq 0} U_k(m)t^m = h_k(t)/(1-t)^{k+1}$ . All zeros of  $h_k$  are real and simple; exactly  $\lfloor k/3 \rfloor$  of them exceed 1, and the others are negative. Consequently  $N(k, q) > 0$  for  $1 \leq q \leq k$ , and every row  $N(k, 1), \dots, N(k, k)$  of the triangle is strictly log-concave, hence unimodal.*

Since  $N(k, q)$  is the number of chains with  $q - 1$  elements in  $\Lambda_k$  without its least and greatest element (Remark 8.16), Theorem F is a real-rootedness result for chain polynomials, a subject studied for several classes of posets in [2, 1, 4]; unlike in the results there, for  $k \geq 3$  the zeros are not confined to  $[-1, 0]$  (Remark 8.16).

Consecutive polynomials  $h_k$  do not have interlacing zeros, so the usual inductive argument does not apply to them directly. We work instead with the row polynomials  $\sum_q N(k, q)z^{q-1}$ , which are related to  $h_k$  by the substitution  $t = z/(1+z)$ , and show that four interlacing relations among them propagate along the recurrence of  $N$ .

**1.3. Related work.** Lipshitz [20] and Stanley [25] developed the theory of D-finite power series (see Kauers' book [14] for a modern account); Zeilberger [30] and Wilf–Zeilberger [29] the holonomic approach to special function identities. Kauers [13] introduced Stirling-like sequences: bivariate sequences whose annihilator is generated by a three-term operator whose two shift vectors form a basis of  $\mathbb{Z}^2$ . His Example 5 shows that the annihilator of the Stirling numbers of the second kind is exactly the left ideal generated by the triangular recurrence. The argument has two steps: there is no recurrence in one variable only, and every annihilating operator reduces modulo the generator to one in one variable. Our Theorem 5.4 follows the same outline for the four-term operator  $L_1$ , which is not of Stirling-like shape (Corollary 5.7). We work with polynomial coefficients and annihilation on a quadrant, show that every annihilating operator with rational coefficients is a left multiple of  $L_1$  as well (Corollary 5.5), and count the recurrences of bounded size. Chyzak, Kauers and Salvy [6] show that the Stirling numbers are not  $\partial$ -finite; Klazar [17] studies Bell numbers and their relatives from the point of view of algebraic differential equations. The  $r$ -Stirling numbers are due to Broder [5], and polynomiality of diagonals of Stirling-type triangles goes back to Gessel and Stanley [11]. Real-rootedness and interlacing are standard tools for proving log-concavity [26, 3, 10]; Harper [12] used the real-rootedness of the row polynomials of the Stirling numbers to prove their asymptotic normality. The finite computation in Section 6.2 combines a sieve modulo primes with Skolem's  $p$ -adic method for exponential Diophantine equations (see [24]).

Kitaev, Mansour and Vella [16] studied pattern avoidance in binary matrices for two-dimensional patterns (right-angled patterns with three cells and  $2 \times 2$  patterns); here the forbidden patterns are instead consecutive patterns of length three inside the rows and inside the columns. For arrays avoiding finite sets of horizontal and vertical patterns, linear recurrences in one dimension for a fixed size of the other exist by the transfer-matrix method [28, Section 4.7]; see also Theorem 2 of Dougherty-Bliss, Koutschan, Ter-Saakov and Zeilberger [8]. Dougherty-Bliss and Kauers [7] proved closed forms and D-finiteness results for a different family of Hardin's arrays (king-move arrays), and Dougherty-Bliss and Spahn [9] showed by the transfer-matrix method that, for a fixed parameter and a fixed number of columns, the number of these arrays is eventually a polynomial in the number of rows. We are not aware of previous work on the table  $U_k(m)$  itself.

**1.4. Formal verification.** The results listed in Section 9 are formalized in Lean 4 [19] with Mathlib [22], starting from the definitions of  $a_k(n)$  and  $U_k(m)$ . The formal proofs use only the three standard axioms. Lean checks proofs, not the faithfulness of formal statements to the statements in this paper; Appendix A transcribes the definitions that a reader has to compare. The Lean development and all verification scripts are in the public repository given under *Data and code* at the end of the paper. Formal verification has precedent in this area: Mahboubi and Sibut-Pinote [21] formalized Apéry's proof of the irrationality of  $\zeta(3)$  in Coq, verifying a posteriori the recurrences produced by computer algebra.

**1.5. Organization.** Section 2 proves the reduction, the basic recurrence and the basic facts about the triangle  $N$ . Section 3 treats the generating functions in  $k$ , the exact recurrence orders and the asymptotics. Section 4 gives the block decomposition and the explicit formula. Section 5 contains the non-D-finiteness and the description of all quadrant recurrences. Section 6 proves the obstructions of Theorem C. Section 7 treats the near-diagonal, and Section 8 proves Theorem F. Section 9 describes the formalization, and Section 10 lists open problems.

## 2. THE REDUCTION AND THE BASIC RECURRENCE

**Definition 2.1.** A triple  $(a, b, c)$  of integers is *good* if  $b = c$  or  $a \geq \max(b, c)$ . For  $k, m \geq 0$  let  $\mathcal{H}_k(m)$  be the set of  $h \in \{0, \dots, m\}^k$  such that  $(h_j, h_{j+1}, h_{j+2})$  is good for  $1 \leq j \leq k-2$ , and let  $U_k(m) = |\mathcal{H}_k(m)|$ . The elements of  $\mathcal{H}_k(m)$  are called *good sequences*; an index  $i$  with  $h_i < h_{i+1}$  is an *ascent*.

Thus  $U_k(0) = 1$ ,  $U_0(m) = 1$ ,  $U_1(m) = m + 1$  and  $U_2(m) = (m + 1)^2$ .

*Convention.* In all summation formulas,  $\binom{a}{b} = 0$  unless  $0 \leq b \leq a$  (also when  $a < 0$ ), and sums over  $s$  or  $j$  run over the finitely many indices with a nonzero summand. This convention is needed: with the generalized binomial coefficients for  $a < 0$  the formulas below are false. When the upper index is a polynomial variable, as in  $\binom{m+1}{q}$  regarded as a polynomial in  $m$ , the polynomial binomial coefficient is meant.

**Proposition 2.2** (Reduction). *For all  $k, n \geq 0$ ,  $a_k(n) = U_k(\lceil n/2 \rceil) U_k(\lfloor n/2 \rfloor)$ .*

*Proof.* A column  $(e_1, \dots, e_n)$  contains 001 or 011 in three consecutive places if and only if  $e_i = 0$  and  $e_{i+2} = 1$  for some  $i$ . So the column condition says that  $e_i \geq e_{i+2}$  for all  $i$ . It couples only rows of the same parity: the rows  $1, 3, 5, \dots$  form, column by column, a weakly decreasing 0/1 matrix with  $\lceil n/2 \rceil$  rows, and so do the rows  $2, 4, \dots$  with  $\lfloor n/2 \rfloor$  rows. The row condition concerns each row separately.

A column-wise weakly decreasing 0/1 matrix with  $M$  rows is determined by its height vector  $h \in \{0, \dots, M\}^k$ , where  $h_j$  is the number of ones in column  $j$ . Row  $r$  then reads  $([r \leq h_1], \dots, [r \leq h_k])$ . For three consecutive columns with heights  $(a, b, c)$ , some row reads 001 if and only if  $c > \max(a, b)$  (take  $r = c$ ), and some row reads 010 if and only if  $b > \max(a, c)$  (take  $r = b$ ). Neither happens if and only if  $c \leq \max(a, b)$  and  $b \leq \max(a, c)$ . If  $b < c$  the first inequality gives  $c \leq a$ ; if  $b > c$  the second gives  $b \leq a$ . Hence neither happens if and only if  $(a, b, c)$  is good. The two halves are independent, which gives the product.  $\square$

**Example 2.3.** The  $5 \times 4$  matrix

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \end{pmatrix}$$

is counted by  $a_4(5)$ . Its rows 1, 3, 5 have column heights  $(3, 1, 3, 3)$  and its rows 2, 4 have column heights  $(1, 2, 2, 0)$ ; both height vectors are good, so they lie in  $\mathcal{H}_4(3)$  and  $\mathcal{H}_4(2)$ . Conversely every pair in  $\mathcal{H}_4(3) \times \mathcal{H}_4(2)$  comes from exactly one matrix of this kind, and  $a_4(5) = U_4(3) U_4(2) = 80 \cdot 32 = 2560$ .

Reversing all rows (or all columns) is a bijection between our arrays and the arrays defined with the opposite reading direction, so the reading convention does not affect the count.

Let  $\Lambda_k$  be the set of words in  $\{0, 1\}^k$  avoiding 001 and 010, ordered componentwise.

**Proposition 2.4.**  $U_k(m)$  is the number of multichains  $x_1 \leq x_2 \leq \dots \leq x_m$  in  $\Lambda_k$ , that is, the value  $Z(\Lambda_k, m + 1)$  of the zeta polynomial of  $\Lambda_k$  in the normalization of [28, Section 3.12]. In particular  $U_k(1) = |\Lambda_k|$ .

*Proof.* By the proof of Proposition 2.2, a column-wise weakly decreasing 0/1 matrix with  $m$  rows and  $k$  columns all of whose rows lie in  $\Lambda_k$  is the same as an element of  $\mathcal{H}_k(m)$ , namely its height vector. Reading the rows of such a matrix from the bottom up gives a multichain  $x_1 \leq \cdots \leq x_m$  in  $\Lambda_k$ , and every multichain arises in this way exactly once.  $\square$

**Lemma 2.5** (Basic recurrence). *For  $k \geq 1$  and  $m \geq 0$ ,*

$$U_k(m) = U_k(m-1) + U_{k-1}(m) + mU_{k-3}(m),$$

*with the conventions  $U_0 \equiv 1$ ,  $U_{-1} \equiv 1$ ,  $U_{-2} \equiv 0$  and  $U_k(-1) = 0$  for  $k \geq 1$ .*

*Proof.* Classify  $h \in \mathcal{H}_k(m)$  by the first position  $i$  at which the value  $m$  occurs. If  $m$  does not occur,  $h \in \mathcal{H}_k(m-1)$ . If  $i = 1$ , every triple starting at position 1 is good because  $a = m \geq \max(b, c)$ , so  $h_2, \dots, h_k$  is an arbitrary element of  $\mathcal{H}_{k-1}(m)$ . If  $i = 2$ , then  $h_1 < m$  ( $m$  choices). For  $k = 2$  this gives  $m = mU_{-1}(m)$  sequences. For  $k \geq 3$  the triple  $(h_1, m, h_3)$  has  $a < b$ , so it is good only if  $h_3 = m$ ; after that the triples  $(m, m, \cdot)$  and  $(m, \cdot, \cdot)$  are automatically good, and  $h_4, \dots, h_k$  is an arbitrary element of  $\mathcal{H}_{k-3}(m)$ . For  $k = 1$  the case does not occur ( $mU_{-2} = 0$ ). If  $i \geq 3$ , the triple  $(h_{i-2}, h_{i-1}, m)$  has  $b < c$  and  $a < c$ , which is not good. The four cases are disjoint and exhaustive.  $\square$

**2.1. The triangle  $N(k, q)$ .** Goodness is defined using only  $=$  and  $\geq$ , so it is preserved by strictly increasing maps of the alphabet. Hence the number of good words of length  $k$  whose set of values is a given  $q$ -element set of integers depends only on  $k$  and  $q$ . We denote it by  $N(k, q)$  and use the alphabet  $\{1, \dots, q\}$ . Thus  $N(0, 0) = 1$ ,  $N(k, 0) = 0$  for  $k \geq 1$ , and  $N(k, q) = 0$  unless  $0 \leq q \leq k$ . Table 2 shows the first rows. Remark 8.16 interprets  $N(k, q)$  as a number of chains in  $\Lambda_k$ .

**Proposition 2.6.** *For  $k, m \geq 0$ ,*

$$(3) \quad U_k(m) = \sum_{q=0}^k N(k, q) \binom{m+1}{q}, \quad N(k, q) = \sum_{i=0}^q (-1)^{q-i} \binom{q}{i} U_k(i-1),$$

*where in the second formula  $U_0(-1) = 1$  and  $U_k(-1) = 0$  for  $k \geq 1$ .*

*Proof.* Sorting the words in  $\mathcal{H}_k(m)$  by their set of values  $V \subseteq \{0, \dots, m\}$  gives the first formula: there are  $\binom{m+1}{q}$  sets with  $|V| = q$ , and each contributes  $N(k, q)$  words. The first formula also holds for  $m = -1$  with the stated values of  $U_k(-1)$ , because  $\sum_q N(k, q) \binom{0}{q} = N(k, 0)$ . Binomial inversion of  $U_k(i-1) = \sum_q N(k, q) \binom{i}{q}$  ( $i \geq 0$ ) gives the second formula.  $\square$

**Proposition 2.7** (Triangular recurrence). *For  $k \geq 3$  and  $q \geq 1$ , with  $N$  taken to be 0 outside  $0 \leq q \leq k$ ,*

$$N(k, q) = N(k-1, q-1) + N(k-1, q) + (q-1)[N(k-3, q-2) + 2N(k-3, q-1) + N(k-3, q)].$$

*For  $k = 2$  the recurrence holds for  $q = 1$  but not for  $q = 2$ ; the initial values are  $N(0, 0) = 1$ ,  $N(1, 1) = 1$ ,  $N(2, 1) = 1$ ,  $N(2, 2) = 2$ .*

*Proof.* Let  $w$  be a good word of length  $k \geq 3$  with set of values  $\{1, \dots, q\}$ , and let  $i$  be the first position of the letter  $q$ . As in the proof of Lemma 2.5,  $i \in \{1, 2\}$ , and if  $i = 2$  then  $w_3 = q$ .

If  $i = 1$ , then  $w = qw'$ , where  $w'$  is a good word whose set of values is  $\{1, \dots, q\}$  or  $\{1, \dots, q-1\}$ . Conversely, for every such  $w'$  the word  $qw'$  is good, because its only new triple starts with  $q$ . This gives  $N(k-1, q) + N(k-1, q-1)$  words.

If  $i = 2$ , then  $w = aqqw''$  with  $1 \leq a \leq q-1$  and a good word  $w''$  of length  $k-3$ . Conversely every such word is good, since its new triples are  $(a, q, q)$ ,  $(q, q, w''_1)$  and  $(q, w''_1, w''_2)$ . The word uses all letters if and only if the set of values  $V$  of  $w''$  is one of  $\{1, \dots, q\}$ ,  $\{1, \dots, q-1\}$ ,  $\{1, \dots, q\} \setminus \{a\}$  and  $\{1, \dots, q-1\} \setminus \{a\}$ . By invariance under increasing maps, for each of the  $q-1$  values of  $a$

these give  $N(k-3, q)$ ,  $N(k-3, q-1)$ ,  $N(k-3, q-1)$  and  $N(k-3, q-2)$  words  $w''$ . Adding the two cases gives the formula.

For  $k=2$  the word  $(1, 2)$  has  $i=2$  but no third letter; it is not counted by the right-hand side, which gives  $N(2, 2) = 1$  instead of 2.  $\square$

**Corollary 2.8.**  $N(k, k) = 2$  for  $k \geq 2$ ,  $N(k, k-1) = k^2 - k - 4$  for  $k \geq 4$ , and  $N(3, 2) = 4$ .

*Proof.* For  $q = k \geq 3$  the recurrence reduces to  $N(k, k) = N(k-1, k-1)$ , and  $N(2, 2) = 2$ . (Directly: a good word with  $k$  distinct letters satisfies  $w_1 > w_2 > \dots > w_{k-1}$  and  $w_k < w_{k-2}$ , so the letter  $w_k$  is 1 or 2.) For  $q = k-1 \geq 2$  it reads  $N(k, k-1) = N(k-1, k-2) + N(k-1, k-1) + (k-2)N(k-3, k-3)$ . This gives  $N(3, 2) = 1 + 2 + 1 = 4$  and  $N(4, 3) = 4 + 2 + 2 = 8$ , and for  $k \geq 5$  the increment is  $2 + 2(k-2) = 2k-2$ , which is the increment of  $k^2 - k - 4$ .  $\square$

**Corollary 2.9.** For every  $k \geq 0$  there is a unique polynomial  $u_k \in \mathbb{Q}[m]$  with  $u_k(m) = U_k(m)$  for all  $m \geq 0$ . It has degree  $k$ , leading coefficient  $2/k!$  for  $k \geq 2$  (and 1 for  $k \leq 1$ ), and  $u_k(-1) = 0$  for  $k \geq 1$ .

*Proof.* Take  $u_k(m) = \sum_q N(k, q) \binom{m+1}{q}$  and use  $N(k, k) = 2$  for  $k \geq 2$ ,  $N(1, 1) = N(0, 0) = 1$  and  $N(k, 0) = 0$  for  $k \geq 1$ .  $\square$

| $k \backslash q$ | 0 | 1 | 2  | 3   | 4   | 5   | 6   | 7  | 8 |
|------------------|---|---|----|-----|-----|-----|-----|----|---|
| 0                | 1 |   |    |     |     |     |     |    |   |
| 1                | 0 | 1 |    |     |     |     |     |    |   |
| 2                | 0 | 1 | 2  |     |     |     |     |    |   |
| 3                | 0 | 1 | 4  | 2   |     |     |     |    |   |
| 4                | 0 | 1 | 7  | 8   | 2   |     |     |    |   |
| 5                | 0 | 1 | 12 | 25  | 16  | 2   |     |    |   |
| 6                | 0 | 1 | 19 | 59  | 65  | 26  | 2   |    |   |
| 7                | 0 | 1 | 29 | 124 | 199 | 139 | 38  | 2  |   |
| 8                | 0 | 1 | 44 | 253 | 557 | 574 | 277 | 52 | 2 |

TABLE 2. The triangle  $N(k, q)$  for  $0 \leq k \leq 8$ .

### 3. GENERATING FUNCTIONS IN $k$ , EXACT RECURRENCE ORDERS AND ASYMPTOTICS

**Theorem 3.1.** Let  $m \geq 0$ .

- (1)  $(1 - x - mx^3)G_m = G_{m-1} + mx^2$  with  $G_{-1} = 1$ ; hence  $G_m = W_m/P_m$ .
- (2)  $\gcd(W_m, P_m) = 1$  in  $\mathbb{Q}[x]$ . Since  $\deg P_m = 3m + 1$ ,  $\deg W_m = 3m$  and  $P_m(0) = 1$ , the minimal order of a linear recurrence with constant coefficients satisfied by  $U_k(m)$  for all large  $k$  is exactly  $3m + 1$ , with characteristic polynomial  $(y-1) \prod_{i=1}^m (y^3 - y^2 - i)$ .
- (3)  $P_m$  is squarefree. In particular all poles of  $G_m$  are simple, and  $G_m$  has a pole with nonzero residue at every root of  $P_m$ .

*Proof.* (1) Multiply Lemma 2.5 by  $x^k$  and sum over  $k \geq 1$ . Since  $U_{-2}(m) = 0$  and  $U_{-1}(m) = 1$ , the last term gives  $\sum_{k \geq 1} mU_{k-3}(m)x^k = mx^2 + mx^3G_m$ ; the other terms give  $G_m - 1$ ,  $G_{m-1} - 1$  and  $xG_m$  (for  $m=0$  use  $U_k(-1) = 0$  for  $k \geq 1$  and  $G_{-1} = 1$ ). This is the first identity. The closed form follows by induction on  $m$ , since  $W_m = W_{m-1} + mx^2P_{m-1}$  and  $P_m = b_mP_{m-1}$ .

(2) For  $j > i$  the polynomial  $b_i$  divides  $P_{j-1}$ , so  $W_m \equiv W_i \pmod{b_i}$  for  $i \leq m$ . Let  $d \in \mathbb{Z}[x]$  be a primitive irreducible common factor of  $W_m$  and  $P_m$ . Then  $d$  divides some  $b_i$  and hence  $W_i$ . By Gauss's lemma  $d$  divides  $W_i$  in  $\mathbb{Z}[x]$ , so  $d(1)$  divides  $W_i(1) = 1$ ; here  $P_{j-1}(1) = 0$  for  $j \geq 1$  because

$b_0 = 1 - x$ . This excludes  $d = b_0$ . For  $i \geq 1$ ,  $b_i$  has a rational root if and only if  $i = y^2(y - 1)$  for an integer  $y \geq 2$  (rational root theorem applied to  $y^3 - y^2 - i$ ), and then

$$b_i = (1 - yx)(1 + (y - 1)x + y(y - 1)x^2),$$

where the quadratic factor has discriminant  $-(y - 1)(3y + 1) < 0$ . The values at  $x = 1$  of the possible factors are  $b_i(1) = -i$ ,  $1 - y$  and  $y^2$ . The condition  $d(1) = \pm 1$  leaves only  $d = b_1$  and  $d = 1 - 2x$  (from  $b_4$ ). But  $W_1 - b_1 = x + x^2$  is coprime to  $b_1$ , and  $W_4(1/2) = 645/512 \neq 0$ .

The statement about recurrences is the standard fact that the minimal order of a linear recurrence satisfied by the coefficients of a rational function  $W/P$  in lowest terms with  $P(0) \neq 0$  and  $\deg W < \deg P$  is  $\deg P$ ; the characteristic polynomial is the reciprocal of  $P_m$ .

(3) A common root of  $b_i$  and  $b_j$  ( $i \neq j$ ) would satisfy  $(i - j)x^3 = 0$ , which is impossible since  $b_i(0) = 1$ . For  $i \geq 1$  the reciprocal  $z^3 - z^2 - i$  of  $b_i$  has discriminant  $-i(4 + 27i) \neq 0$ , and  $b_0$  is linear. The last statement follows from (2).  $\square$

**Theorem 3.2.** *For  $m \geq 1$  let  $\rho_m$  be a real root of  $y^3 - y^2 - m$ , and put  $\rho_0 := 1$ .*

- (1)  $\rho_m$  is the only real root,  $1 = \rho_0 < \rho_1 < \rho_2 < \dots$ , and the two non-real roots  $\sigma, \bar{\sigma}$  of  $y^3 - y^2 - m$  satisfy  $|\sigma|^2 = \rho_m(\rho_m - 1)$ , which increases with  $m$ , and  $|\sigma|^2 < \rho_{m-1}^2$ .
- (2) For every  $m \geq 0$  there are nonzero numbers  $\alpha_m(\sigma)$  such that  $U_k(m) = \sum_{\sigma} \alpha_m(\sigma) \sigma^k$  for all  $k \geq 0$ , where  $\sigma$  runs over the  $3m + 1$  roots of  $(y - 1) \prod_{i=1}^m (y^3 - y^2 - i)$ . If  $\sigma$  is a root of  $y^3 - y^2 - j$  with  $1 \leq j < m$ , or  $\sigma = 1$  and  $j = 0$ , then  $\alpha_m(\sigma) = \alpha_j(\sigma) (-\sigma^3)^{m-j} / (m - j)!$ .
- (3) Let  $c_0 = 1$  and, for  $m \geq 1$ ,  $x_m = 1/\rho_m$  and

$$c_m = \alpha_m(\rho_m) = \frac{G_{m-1}(x_m) + mx_m^2}{x_m(1 + 3mx_m^2)} = \frac{\rho_m^{3m+3}}{m!(\rho_m^2 + 3m)} \left(1 + \sum_{j=1}^m j m^j \rho_m^{-3j-2}\right) > 0.$$

Then for  $m \geq 1$

$$U_k(m) = c_m \rho_m^k - c_{m-1} \rho_{m-1}^{k+3} + O(\tau_m^k) \quad (k \rightarrow \infty),$$

where  $\tau_1 = \sqrt{\rho_1(\rho_1 - 1)} \approx 0.826$  and  $\tau_m = \max(\rho_{m-2}, \sqrt{\rho_m(\rho_m - 1)}) < \rho_{m-1}$  for  $m \geq 2$ .

Here  $m^j = m(m - 1) \cdots (m - j + 1)$ .

*Proof.* (1) The function  $f(y) = y^3 - y^2 - m$  is increasing on  $(-\infty, 0]$ , decreasing on  $[0, 2/3]$  and increasing on  $[2/3, \infty)$ , and  $f(0) = f(1) = -m < 0$ . So  $f$  has exactly one real root, and it exceeds 1. Since  $y^3 - y^2$  is increasing on  $[1, \infty)$ ,  $\rho_m$  increases with  $m$ . The product of the roots is  $m$ , so  $|\sigma|^2 = m/\rho_m = \rho_m(\rho_m - 1)$ , which increases with  $m$ . Finally  $\rho_m \rho_{m-1}^2 > \rho_{m-1}^3 = \rho_{m-1}^2 + m - 1 \geq m$ , which is  $|\sigma|^2 < \rho_{m-1}^2$ .

(2) By Theorem 3.1,  $G_m$  is a proper rational function ( $\deg W_m < \deg P_m$ ) whose poles are the reciprocals of the  $3m + 1$  distinct roots listed, all simple and with nonzero residues. Hence  $G_m = \sum_{\sigma} \alpha_m(\sigma)/(1 - \sigma x)$  with  $\alpha_m(\sigma) \neq 0$ , which is the first claim. If  $1/\sigma$  is a root of  $b_j$  with  $j < m$ , then  $b_m(1/\sigma) = b_j(1/\sigma) - (m - j)\sigma^{-3} = -(m - j)\sigma^{-3} \neq 0$ , and  $G_m = (G_{m-1} + mx^2)/b_m$  gives  $\alpha_m(\sigma) = \alpha_{m-1}(\sigma)/b_m(1/\sigma)$ . Iterating gives the formula.

(3) At  $x_m$  we have  $G_m = \tilde{G}_m/b_m$ , where  $\tilde{G}_m := G_{m-1} + mx^2 = W_m/P_{m-1}$  is holomorphic at  $x_m$ , and  $b'_m(x_m) = -(1 + 3mx_m^2)$ . Comparing with  $\alpha/(1 - \rho_m x) = \alpha x_m/(x_m - x)$  gives the first expression for  $c_m$ . Since  $1 - x_m = mx_m^3$ , we have  $b_v(x_m) = (m - v)x_m^3$ , hence  $P_{j-1}(x_m) = m^j x_m^{3j}$  and  $P_{m-1}(x_m) = m! x_m^{3m}$ , and  $W_m(x_m) = 1 + \sum_j j m^j x_m^{3j+2}$ ; this gives the second expression, which is positive. By (1),  $\rho_m$  is the root of largest modulus; the second largest modulus is attained only at  $\rho_{m-1}$ , because the non-real roots of  $y^3 - y^2 - i$  for  $i \leq m$  have modulus at most  $\sqrt{\rho_m(\rho_m - 1)} < \rho_{m-1}$ ; and all other roots have modulus at most  $\tau_m$ . Finally, by (2),  $\alpha_m(\rho_{m-1}) = \alpha_{m-1}(\rho_{m-1}) (-\rho_{m-1}^3) = -c_{m-1} \rho_{m-1}^3$ ; for  $m = 1$  this is  $\alpha_1(1) = -1 = -c_0 \rho_0^3$ .  $\square$

**Remark 3.3.** Consequently  $U_k(m)/(c_m \rho_m^k) - 1 = -\kappa_m \theta_m^k (1 + o(1))$  with  $\theta_m = \rho_{m-1}/\rho_m$  and  $\kappa_m = c_{m-1} \rho_{m-1}^3 / c_m > 0$ , and the rate  $\theta_m$  cannot be improved. Numerically  $\rho_1 \approx 1.46557$ ,  $c_1 \approx 2.20961$ ,  $c_2 \approx 7.84112$  and  $c_3 \approx 28.97686$ ; in fact  $c_1 = (10 + 15\rho_1 + 17\rho_1^2)/31$ . When  $\rho_m$  is an integer, that is  $m = y^2(y-1)$  ( $m = 4, 18, 48, \dots$ ), the constant  $c_m$  is rational; for instance  $\rho_4 = 2$ ,  $c_4 = 215/2$ , and  $U_k(4) = \frac{215}{2} 2^k - c_3 \rho_3^{k+3} + O(\rho_2^k)$ .

All of this concerns fixed  $m$ ; Theorem 3.2 gives no error bound that is uniform in  $m$ . By the mean value theorem  $1 - \theta_m \sim 1/(3m)$  as  $m \rightarrow \infty$ , and in the computed range  $1 \leq m \leq 24$  the constant  $\kappa_m$  increases with  $m$  ( $\kappa_1 \approx 0.453$ ,  $\kappa_4 \approx 1.745$ ,  $\kappa_{24} \approx 9.869$ ). The relative error  $U_k(m)/(c_m \rho_m^k) - 1$  at  $k = 3m$  is about  $-0.36$  for  $m = 2$  and  $-0.996$  for  $m = 24$ , and at  $k = 10m$  about  $-0.05$  and  $-0.41$ . So the leading term alone is a good approximation only when  $k$  is considerably larger than  $3m$ ; these figures are numerical observations, not a uniform estimate.

**Corollary 3.4** (Columns of the original table). *For fixed  $k \geq 1$  there are polynomials  $\lambda_k, \mu_k \in \mathbb{Q}[n]$  with  $\deg \lambda_k = 2k$  and  $\deg \mu_k = 2k - 2$  such that  $a_k(n) = \lambda_k(n) + (-1)^n \mu_k(n)$  for all  $n \geq 0$ . The generating function  $\sum_n a_k(n) x^n$  has reduced denominator  $(1-x)^{2k+1}(1+x)^{2k-1}$ ; in particular, the minimal order of a linear recurrence with constant coefficients for the  $k$ -th column of A207123 is  $4k$ .*

*Proof.* Let  $u = u_k$  be the polynomial of Corollary 2.9, of degree  $k$  with leading coefficient  $\ell \neq 0$ . Put  $a^+(n) = u(n/2)^2$  and  $a^-(n) = u((n+1)/2)u((n-1)/2)$ , polynomials in  $n$  of degree  $2k$ . By Proposition 2.2,  $a_k(n) = a^+(n)$  for even  $n$  and  $a_k(n) = a^-(n)$  for odd  $n$ , so  $a_k(n) = \lambda_k(n) + (-1)^n \mu_k(n)$  with  $\lambda_k = (a^+ + a^-)/2$  and  $\mu_k = (a^+ - a^-)/2$ . The leading coefficient of  $\lambda_k$  is  $\ell^2/4^k$ . With  $y = n/2$  and  $h = 1/2$ , Taylor expansion gives

$$u(y+h)u(y-h) = u(y)^2 + h^2(uu'' - u'^2)(y) + \sum_{r \geq 2} h^{2r} g_r(y), \quad \deg g_r \leq 2k - 2r,$$

and  $uu'' - u'^2$  has leading term  $\ell^2(k(k-1) - k^2)y^{2k-2} = -k\ell^2 y^{2k-2} \neq 0$ . Hence  $\deg \mu_k = 2k - 2$ . For a polynomial  $f$  of degree  $e$  with leading coefficient  $c$  one has  $\sum_n f(n)x^n = A(x)/(1-x)^{e+1}$  with  $\deg A \leq e$  and  $A(1) = e!c \neq 0$ . Applying this to  $\lambda_k$ , and to  $\mu_k$  with  $x$  replaced by  $-x$ , the generating function of  $a_k$  is a proper rational function whose numerator over  $(1-x)^{2k+1}(1+x)^{2k-1}$  does not vanish at  $x = \pm 1$ .  $\square$

**Remark 3.5** (The OEIS entries). The existence of some linear recurrence for each column also follows from the general transfer-matrix argument of [8, Theorem 2]; the point of Corollary 3.4 is the exact order for every  $k$ . The empirical recurrences recorded in the OEIS for the columns  $k = 3, \dots, 7$  (A207118–A207122) have characteristic polynomial exactly  $(x-1)^{2k+1}(x+1)^{2k-1}$ , so by Corollary 3.4 they hold for all  $n$ . For a fixed row  $n$ , the sequence  $k \mapsto a_k(n)$  is the product of  $k \mapsto U_k(\lceil n/2 \rceil)$  and  $k \mapsto U_k(\lfloor n/2 \rfloor)$ , which satisfy recurrences of orders  $3\lceil n/2 \rceil + 1$  and  $3\lfloor n/2 \rfloor + 1$  for all  $k \geq 0$  (Theorem 3.2(2)). Hence the product satisfies a recurrence of order at most  $\Delta_n = (3\lceil n/2 \rceil + 1)(3\lfloor n/2 \rfloor + 1)$ , and a proposed recurrence with constant coefficients holds for all large  $k$  as soon as it holds for  $\Delta_n$  consecutive values in that range. In this way we checked, in exact arithmetic, that the empirical recurrences recorded for the rows  $n = 2, \dots, 7$  (A207069, A207070, A207124–A207127) hold for all  $k$ ; by the Berlekamp–Massey algorithm their orders 10, 22, 28, 49, 55, 85 are minimal, and they are the numbers of distinct products of a characteristic root of the first factor with one of the second. The entries A207069 and A207070 are titled with the vertical rule 001/101 instead of 001/011; for two and three rows both rules give the same numbers (trivially for two rows, and by exchanging the first two rows for three), and A207123 lists them as its rows 2 and 3.

## 4. BLOCK DECOMPOSITION AND AN EXPLICIT FORMULA

**Definition 4.1.** For integers  $0 \leq a < v$  define the blocks

$$S_v = [v], \quad T_{v,a} = [a, v, v], \quad E_{v,a} = [a, v],$$

and also  $S_0 = [0]$ . The number  $v$  is the *level* of the block and  $a$  its *valley*.

**Theorem 4.2** (Unique factorization). *Concatenation is a bijection from the set of finite sequences of blocks with weakly decreasing levels in  $\{0, \dots, m\}$ , in which only the last block may be an  $E$ -block, onto  $\bigcup_k \mathcal{H}_k(m)$ . Equivalently, as an identity between languages,*

$$\mathcal{H}(\leq m) = B_m^* \cdot (\mathcal{H}(\leq m-1) \sqcup E_m), \quad B_m = \{S_m\} \cup \{T_{m,a} : a < m\}, \quad E_m = \{E_{m,a} : a < m\},$$

where  $\mathcal{H}(\leq m) = \bigcup_k \mathcal{H}_k(m)$  and  $\mathcal{H}(\leq -1)$  consists of the empty sequence.

*Proof.* Every concatenation is good. Let  $p$  be a position that is not a valley; its value  $v$  is the level of its block. All later values are at most  $v$ : the rest of the block equals  $v$ , later blocks have level at most  $v$ , and valleys are below levels. So every triple starting at  $p$  has  $a \geq \max(b, c)$ . A triple starting at the valley of a  $T$ -block is  $(a, v, v)$ , which is good. The valley of an  $E$ -block is followed by only one entry.

*Existence.* Induct on the length. Let  $M = \max h$ , first attained at position  $i$ . As in Lemma 2.5,  $i \in \{1, 2\}$ . If  $i = 1$ , split off  $S_M$ . If  $i = 2$  and  $k = 2$ , the sequence is  $E_{M, h_1}$ . If  $i = 2$  and  $k \geq 3$ , then  $h_3 = M$  and we split off  $T_{M, h_1}$ . The remainder is good and has maximum at most  $M$ .

*Uniqueness.* The level of the first block equals  $\max h$ . If  $h_1 = \max h$  the first block is  $S_{h_1}$ ; otherwise it is a  $T$ - or an  $E$ -block, and an  $E$ -block can only be the last block. Finally, if  $a < v$  then  $(a, v, c)$  is good only for  $c = v$ , so a pair  $(a, v)$  that is not at the end must be completed to  $T_{v,a}$ .  $\square$

**Example 4.3.** The good sequence  $(3, 1, 3, 3, 2, 0, 2, 2, 1, 0, 1) \in \mathcal{H}_{11}(3)$  factors as

$$S_3 T_{3,1} S_2 T_{2,0} S_1 E_{1,0} = [3] [1, 3, 3] [2] [0, 2, 2] [1] [0, 1].$$

Its three ascents start at the valleys 1, 0 and 0, and it ends with an ascent because its last block is an  $E$ -block.

In general the ascents  $h_i < h_{i+1}$  of a good sequence occur exactly after the valleys, so the number of ascents equals the number of  $T$ - and  $E$ -blocks, and a sequence ends with an ascent if and only if its last block is an  $E$ -block.

**Theorem 4.4.** *Let  $U_k(m, s)$  be the number of  $h \in \mathcal{H}_k(m)$  with  $s$  ascents, write  $b_v = 1 - x - v y x^3$  in this theorem (so that  $b_v$  becomes  $b_v(x)$  at  $y = 1$ ), and let  $G_m(x, y)$  be the generating function  $\sum_{k,s} U_k(m, s) x^k y^s$ . Then  $G_{-1} = 1$ ,*

$$G_m = \frac{G_{m-1} + m y x^2}{1 - x - m y x^3}, \quad \text{equivalently} \quad G_m = \frac{1}{\prod_{v=0}^m b_v} + y x^2 \sum_{j=1}^m \frac{j}{\prod_{v=j}^m b_v}.$$

*Proof.* Translate Theorem 4.2:  $B_m^*$  contributes  $1/(1 - x - m y x^3)$  (one single block, and  $m$  triple blocks with one ascent each),  $E_m$  contributes  $m y x^2$ , and the sequences with maximum at most  $m-1$  contribute  $G_{m-1}$ . Unfolding the recursion gives the second form.  $\square$

For  $y = 1$  this is a combinatorial proof of Theorem 3.1(1). The first term of the second form counts the sequences that do not end with an ascent, and the term with index  $j$  counts those that end with an  $E$ -block of level  $j$ .

**Lemma 4.5.** *Let  $H(m, s, j) = h_s(j, j+1, \dots, m)$  be the complete homogeneous symmetric polynomial of degree  $s$  evaluated at  $j, \dots, m$ , and  $H(m, -1, j) := 0$ . Then, for  $0 \leq j \leq m$ ,*

$$[x^n y^s] \prod_{v=j}^m \frac{1}{1 - x - v y x^3} = H(m, s, j) \binom{n + m - j - 2s}{n - 3s}.$$

Moreover  $H(m, s, 0) = H(m, s, 1) = \left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\}$ , and for  $1 \leq j \leq m$ ,  $H(m, s, j) = \left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\}_j$  is the number of partitions of  $\{1, \dots, m+s\}$  into  $m$  blocks such that  $1, \dots, j$  lie in distinct blocks.

*Proof.* With  $u = x^3/(1-x)$  we have  $1-x-vyx^3 = (1-x)(1-vyu)$ , hence

$$\prod_{v=j}^m \frac{1}{1-x-vyx^3} = \sum_{s \geq 0} H(m, s, j) y^s \frac{x^{3s}}{(1-x)^{m-j+1+s}},$$

and  $[x^n] x^{3s}(1-x)^{-M} = \binom{n-3s+M-1}{n-3s}$ . The variable 0 does not contribute to  $h_s$ , and  $\sum_n \left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\} z^n = z^m / \prod_{i=1}^m (1-iz)$  gives  $H(m, s, 1) = \left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\}$ . The identification with  $r$ -Stirling numbers is Broder's generating function  $\sum_n \left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\}_j z^n = z^m / \prod_{i=j}^m (1-iz)$  [5].  $\square$

**Theorem 4.6** (Explicit formula). *With the binomial convention of Section 2, for all  $k, m \geq 0$  (and all  $s \geq 0$  in the second formula),*

$$U_k(m) = \sum_{s \geq 0} \left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\} \binom{k+m-2s}{k-3s} + \sum_{j=1}^m j \sum_{s \geq 0} H(m, s, j) \binom{k-2+m-j-2s}{k-2-3s},$$

$$U_k(m, s) = \left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\} \binom{k+m-2s}{k-3s} + \sum_{j=1}^m j H(m, s-1, j) \binom{k+m-j-2s}{k+1-3s}.$$

*Proof.* By Theorem 4.4 and Lemma 4.5 with  $j=0$  and  $n=k$ , the coefficient of  $x^k y^s$  in  $1/\prod_{v=0}^m b_v$  is  $\left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\} \binom{k+m-2s}{k-3s}$ . By Lemma 4.5 with  $n=k-2$  and  $s-1$  in place of  $s$ , the coefficient of  $x^k y^s$  in  $y x^2 j / \prod_{v=j}^m b_v$  is  $j H(m, s-1, j) \binom{k+m-j-2s}{k+1-3s}$ . This is the second formula. Summing over  $s$  and replacing  $s$  by  $s+1$  in the second part gives the first.  $\square$

The first summand counts the sequences that do not end with an ascent; for each  $s$  it is a single product, and the following bijection explains it. The factor  $j$  in the second summand is the choice of the valley of the final  $E$ -block of level  $j$ , the factor  $H(m, s, j)$  accounts for the  $T$ -blocks before it, whose levels lie in  $[j, m]$ , and the binomial coefficient counts the positions of the single blocks.

**Proposition 4.7.** *Let  $k, m, s \geq 0$ . The good sequences of length  $k$  with values in  $\{0, \dots, m\}$  that have exactly  $s$  ascents and do not end with an ascent are in explicit bijection with the pairs consisting of a  $(k-3s)$ -element subset of  $\{1, \dots, k+m-2s\}$  and a partition of  $\{1, \dots, m+s\}$  into  $m$  blocks.*

*Proof.* By Theorem 4.2 such a sequence is a sequence of  $k-3s$  single blocks and  $s$  triple blocks with weakly decreasing levels in  $\{0, \dots, m\}$ . Encode it as a word in the letters  $S, T$  and  $|$ : read the blocks from left to right, starting at level  $m$ ; write one bar for each unit by which the level drops (also before the first block, if its level is below  $m$ ), and add bars at the end until level 0 is reached. The word has  $k-3s$  letters  $S$ ,  $s$  letters  $T$  and  $m$  bars, and the level of each block is the number of bars to its right; conversely the word determines the levels. The positions of the letters  $S$  form the subset. Now number the  $m+s$  letters  $T$  and  $|$  from right to left by  $1, \dots, m+s$ . Every bar opens a new block of a set partition, and a letter  $T$  whose block has valley  $a$  joins the block opened by the  $(a+1)$ -st bar from the right; this bar lies to the right of the letter because  $a$  is smaller than its level. This gives a partition of  $\{1, \dots, m+s\}$  into  $m$  blocks. Conversely, given a partition into  $m$  blocks, the block minima are the bars, and every other element is a letter  $T$  whose valley is one less than the rank of its block when the blocks are ordered by their minima; this valley is smaller than the number of bars to its right.  $\square$

**Example 4.8.** For  $m=1$  we have  $\left\{ \begin{smallmatrix} 1+s \\ 1 \end{smallmatrix} \right\} = 1$  and  $H(1, s, 1) = 1$ , so Theorem 4.6 gives

$$U_k(1) = \sum_{s \geq 0} \binom{k+1-2s}{s+1} + \sum_{s \geq 0} \binom{k-2-2s}{s},$$

for instance  $U_5(1) = (6 + 6) + (1 + 1) = 14$ , the number of binary words of length 5 that avoid 001 and 010.

**Remark 4.9** (Other forms). Using the telescoping identity  $jx^3Q_j = (1-x)Q_j - Q_{j+1}$  for  $Q_j = \prod_{v=j}^m b_v^{-1}$  (with  $y = 1$ ) one obtains a single sum minus a double sum without the weights  $j$ ,

$$U_k(m) = \sum_{s \geq 0} \left\{ \begin{matrix} m+s \\ m \end{matrix} \right\} \binom{k+1+m-2s}{m+s} - \sum_{j=1}^m \sum_{s \geq 0} H(m, s, j) \binom{k+m-j-2s}{m-j+s}.$$

Partial fractions over the nodes  $0, 1, \dots, m$  give a formula with a single division: with  $\psi_i(n) = [x^n] 1/b_i = \sum_{0 \leq l \leq n/3} \binom{n-2l}{l} i^l$  and  $\psi_i(n) = 0$  for  $n < 0$ ,

$$m! U_k(m) = \sum_{i=0}^m (-1)^{m-i} \binom{m}{i} \left[ \psi_i(k+3m) + \sum_{j=1}^i j i^j \psi_i(k+3m-3j-2) \right].$$

If the Stirling-type numbers are expanded, all of these become triple sums. The last formula needs only the sequences  $\psi_0, \dots, \psi_m$ , which satisfy  $\psi_i(0) = 1$  and  $\psi_i(n) = \psi_i(n-1) + i \psi_i(n-3)$  for  $n \geq 1$ .

## 5. NON-D-FINITENESS AND ALL QUADRANT RECURRENCES

Let  $F(x, t) = \sum_{k, m \geq 0} U_k(m) x^k t^m = \sum_m G_m(x) t^m$ . Recall that  $f \in K[[x, t]]$  is *D-finite* if all partial derivatives  $\partial_x^i \partial_t^j f$  span a finite-dimensional vector space over  $K(x, t)$  [20]. If  $f$  is D-finite then, for each variable,  $f$  satisfies a nonzero linear differential equation in that variable with polynomial coefficients.

### Theorem 5.1.

- (1) *There are no  $p_0, \dots, p_r \in \mathbb{C}[x, t]$ , not all zero, with  $\sum_j p_j \partial_x^j F = 0$ . In particular  $F$  is not D-finite (over any subfield of  $\mathbb{C}$ ).*
- (2) *There is no recurrence  $\sum_{a \leq A} \sum_{b \leq B} p_{ab}(k) U_{k-a}(m-b) = 0$  with  $p_{ab} \in \mathbb{C}[k]$  not all zero that holds for all  $k \geq k_0$ ,  $m \geq m_0$  (here  $k_0 \geq A$  and  $m_0 \geq B$ , so that all indices are nonnegative).*

*Proof.* (1) Suppose such an equation exists. Since  $\mathbb{C}[[x, t]]$  is a domain we may divide by a power of  $t$  and assume that the polynomials  $p_{j0}(x) := p_j(x, 0)$  are not all zero. Writing  $p_j = \sum_l p_{jl}(x) t^l$ , the coefficient of  $t^m$  gives the identity of rational functions

$$\sum_j \sum_l p_{jl}(x) G_{m-l}^{(j)}(x) = 0,$$

where here  $G_n := 0$  for  $n < 0$ , because  $F = \sum_{m \geq 0} G_m t^m$  (this is not the convention  $G_{-1} = 1$  of Theorem 3.1). Let  $j^* = \max\{j : p_{j0} \neq 0\}$ . The points  $x_m = 1/\rho_m$  are pairwise distinct (Theorem 3.2), so we can choose  $m$  with  $p_{j^*0}(x_m) \neq 0$ . At  $x_m$  we have  $b_v(x_m) = (m-v)x_m^3 \neq 0$  for  $v \neq m$ , so  $G_{m-l}$  is holomorphic at  $x_m$  for  $l \geq 1$ , while  $G_m$  has a simple pole with nonzero residue because  $W_m(x_m) = 1 + \sum_{j=1}^m j m^j x_m^{3j+2} > 0$ . Hence  $G_m^{(j^*)}$  has a pole of order exactly  $j^* + 1$  at  $x_m$ , and all other terms have poles of order at most  $j^*$ . This is a contradiction.

If  $F$  were D-finite, then  $F, \partial_x F, \dots, \partial_x^d F$  would be linearly dependent over  $\mathbb{C}(x, t)$  for some  $d$ , and clearing denominators would give an equation excluded by (1).

(2) Multiply the recurrence by  $x^k t^m$  and sum over all  $k, m \geq 0$ , with  $U$  extended by 0 to negative indices. With  $\vartheta = x \partial_x$  the left-hand side becomes  $\mathcal{L}F$  for the nonzero operator  $\mathcal{L} = \sum_{a,b} p_{ab}(\vartheta) x^a t^b$ . The coefficients of  $\mathcal{L}F$  vanish on the quadrant. For each fixed  $m < m_0$  the corresponding coefficient series in  $x$  is rational, because each  $G_{m'}$  is rational; for each fixed  $k < k_0$  the coefficient series in  $t$  is rational, because  $U_{k'}(m)$  is a polynomial in  $m$  (Corollary 2.9). So  $\mathcal{L}F = R$  is a rational function. If  $R = 0$  we contradict (1). Otherwise  $(R \partial_x - R_x) \mathcal{L}$  is a nonzero operator (the ring

of differential operators over  $\mathbb{C}(x, t)$  is a domain) annihilating  $F$ , and clearing denominators again contradicts (1).  $\square$

The basic recurrence (2) has coefficients depending only on  $m$ ; it corresponds to the differential equation  $(1 - x - t)F - x^3 t \partial_t F = 1 + x^2 t / (1 - t)^2$  in  $t$ . D-finiteness would require such an equation in every direction; part (1) shows that there is none in  $x$ , and part (2) is the corresponding statement for recurrences in  $k$ . The same statements hold for the triangle. By (3),  $\sum_m U_k(m) t^m = \sum_{q \geq 1} N(k, q) t^{q-1} (1 - t)^{-q-1}$  for  $k \geq 1$ , so  $\mathcal{N}(x, y) = \sum_{k, q} N(k, q) x^k y^q$  satisfies  $1 + tF(x, t) = \mathcal{N}(x, t/(1 - t)) / (1 - t)$ . The substitution  $y = t/(1 - t)$  therefore turns a differential equation in  $x$  for  $\mathcal{N}$  into an inhomogeneous one for  $F$  with polynomial right-hand side, which is excluded as in the proof of part (2).

**Remark 5.2.** The part that does not end with an ascent is hypergeometric in  $t$ . With  $\lambda = (1 - x)/x^3$  we have  $i - \lambda = -b_i/x^3$ , so  $(1 - \lambda)_n = (-1)^n x^{-3n} \prod_{i=1}^n b_i$ , and therefore, in  $\mathbb{Q}(x)[[t]]$ ,

$$\sum_{m \geq 0} \frac{t^m}{P_m} = \frac{1}{1 - x} {}_1F_1\left(1; 1 - \lambda; -\frac{t}{x^3}\right).$$

By Theorem 4.4, the coefficient of  $x^k t^m$  is the number of sequences in  $\mathcal{H}_k(m)$  that do not end with an ascent.

**5.1. The ideal of quadrant recurrences.** Let  $\mathcal{A} = \{f : \mathbb{N}^2 \rightarrow \mathbb{C}\}$ . For  $p \in \mathbb{C}[k, m]$  let  $p$  act by multiplication,  $(pf)(k, m) = p(k, m)f(k, m)$ , and let

$$(Xf)(k, m) = f(k - 1, m), \quad (E^{-1}f)(k, m) = f(k, m - 1),$$

with  $f = 0$  at negative indices. Let  $\mathcal{O} \subset \text{End}_{\mathbb{C}}(\mathcal{A})$  be the algebra generated by  $k$ ,  $m$ ,  $X$  and  $E^{-1}$ . We have  $Xk = (k - 1)X$ ,  $E^{-1}m = (m - 1)E^{-1}$ , and  $X$  commutes with  $m$  and with  $E^{-1}$ . Every element of  $\mathcal{O}$  has a unique normal form  $\sum_{(a, b)} r_{ab}(k, m) X^a E^{-b}$ , so  $\mathcal{O}$  is the Ore algebra  $\mathbb{C}[k, m]\langle X, E^{-1} \rangle$  acting faithfully. Define the *support*  $\text{Supp}(R) = \{(a, b) : r_{ab} \neq 0\}$  and

$$\text{Rel}(U) = \{R \in \mathcal{O} : (RU)(k, m) = 0 \text{ for all } k \geq k_0, m \geq m_0, \text{ for some } k_0, m_0\}.$$

Lemma 2.5 says that  $L_1 = 1 - E^{-1} - X - mX^3 \in \text{Rel}(U)$ : indeed  $(L_1 U)(k, m) = 0$  for  $k \geq 3$ ,  $m \geq 1$ . A recurrence involving forward shifts becomes one of this form after left multiplication by a suitable monomial  $X^a E^{-b}$ , that is, after evaluating it at  $(k - a, m - b)$ .

**Lemma 5.3.** *Let  $R = \sum_{a=0}^A r_a(k, m) X^a$  with  $r_a \in \mathbb{C}[k, m]$ . If  $R \in \text{Rel}(U)$ , then  $R = 0$ .*

*Proof.* Suppose  $R \neq 0$  and  $RU$  vanishes for  $k \geq k_0$ ,  $m \geq m_0$ . Let  $D = \max_a \deg_k r_a$  and write  $r_a(k, m) = \ell_a(m) k^D + (\text{lower terms in } k)$ ; the  $\ell_a \in \mathbb{C}[m]$  are not all zero. Choose  $m \geq m_0$  with  $3m + 1 > A$  and  $(\ell_0(m), \dots, \ell_A(m)) \neq 0$ , and put  $C_m(z) = \sum_a \ell_a(m) z^a$ , a nonzero polynomial of degree at most  $A$ . The series  $S_m(x) = \sum_{k \geq 0} (RU)(k, m) x^k$  is a polynomial, and, with  $\vartheta = x d/dx$ ,

$$S_m(x) = \sum_a r_a(\vartheta, m) (x^a G_m(x)).$$

Let  $z$  be a root of  $P_m$ . By Theorem 3.1,  $G_m$  has a simple pole at  $z$  with a nonzero residue  $\varrho$ , and  $\vartheta^j(c/(x - z)) = (-1)^j j! z^j c (x - z)^{-j-1} + O((x - z)^{-j})$ . So the coefficient of  $(x - z)^{-D-1}$  in  $S_m$  is  $(-1)^D D! z^D \varrho C_m(z)$ , which must vanish. Hence  $C_m$  vanishes at the  $3m + 1$  distinct roots of  $P_m$ , which is impossible.  $\square$

**Theorem 5.4.**

$$(1) \text{ Rel}(U) = \mathcal{O} \cdot L_1.$$

- (2) For  $A, B, D \geq 0$  let  $V(A, B, D)$  be the space of  $R \in \text{Rel}(U)$  with  $\text{Supp}(R) \subseteq [0, A] \times [0, B]$  and all coefficients of total degree at most  $D$ . Then

$$\dim V(A, B, D) = \begin{cases} (A-2)B \frac{D(D+1)}{2} & \text{if } A \geq 3, B \geq 1, D \geq 1, \\ 0 & \text{otherwise.} \end{cases}$$

If instead the degrees in  $k$  and  $m$  are bounded by  $D_k$  and  $D_m$ , the dimension is  $(A-2)B(D_k+1)D_m$  for  $A \geq 3, B \geq 1, D_m \geq 1$ , and 0 otherwise.

*Proof.* (1) If  $Q \in \mathcal{O}$  has support in  $[0, A'] \times [0, B']$ , then  $(QL_1U)(k, m)$  is a combination of the values  $(L_1U)(k-a, m-b)$  with  $a \leq A', b \leq B'$ , so it vanishes for  $k \geq A' + 3, m \geq B' + 1$ . Hence  $\mathcal{O}L_1 \subseteq \text{Rel}(U)$ .

Conversely, every  $R \in \mathcal{O}$  can be written as  $R = QL_1 + R_0$  with  $Q \in \mathcal{O}$  and  $R_0 = \sum_a r_a(k, m)X^a$  free of  $E^{-1}$ . Indeed, for a monomial with  $b \geq 1$ , the identity  $E^{-1} = 1 - X - mX^3 - L_1$  and the commutation rules give

$$c(k, m)X^aE^{-b} = c(k, m)[X^a - X^{a+1} - (m-b+1)X^{a+3}]E^{-(b-1)} - c(k, m)X^aE^{-(b-1)}L_1,$$

which lowers the power of  $E^{-1}$  by one; induction on the highest power of  $E^{-1}$  gives the claim. If  $R \in \text{Rel}(U)$ , then  $R_0 = R - QL_1 \in \text{Rel}(U)$ , so  $R_0 = 0$  by Lemma 5.3, and  $R = QL_1$ .

- (2) For  $Q = \sum q(a, b)X^aE^{-b}$  the normal form of  $QL_1$  is

$$(4) \quad r(a, b) = q(a, b) - q(a, b-1) - q(a-1, b) - (m-b)q(a-3, b),$$

with  $q = 0$  at negative indices. Let  $Q \neq 0$ . If  $\beta^*$  is the largest  $b$  with  $q(\cdot, b) \neq 0$  and  $q(\alpha', \beta^*) \neq 0$ , then  $r(\alpha', \beta^* + 1) = -q(\alpha', \beta^*) \neq 0$ . If  $\alpha^*$  is the largest  $a$  with  $q(a, \cdot) \neq 0$  and  $q(\alpha^*, \beta') \neq 0$ , then  $r(\alpha^* + 3, \beta') = -(m - \beta')q(\alpha^*, \beta') \neq 0$ . If  $e$  is the largest total degree of the coefficients of  $Q$  and  $[q(a, b)]_e \neq 0$  denotes a nonzero homogeneous part of degree  $e$ , then the homogeneous part of degree  $e+1$  of  $r(a+3, b)$  is  $-m[q(a, b)]_e \neq 0$ . In particular right multiplication by  $L_1$  is injective, and  $QL_1 \in V(A, B, D)$  forces  $\text{Supp}(Q) \subseteq [0, A-3] \times [0, B-1]$  and total degree at most  $D-1$ ; conversely every such  $Q$  gives an element of  $V(A, B, D)$ . Counting the monomials  $k^i m^j X^a E^{-b}$  with  $a \leq A-3, b \leq B-1$  and  $i+j \leq D-1$  gives the formula. For separate degree bounds, the coefficient of the top power of  $k$  in  $QL_1$  is  $\hat{Q}L_1$ , where  $\hat{Q} \neq 0$  collects the top  $k$ -coefficients of  $Q$ , so the  $k$ -degree does not change; the  $m$ -degree goes up by exactly one by the argument just given with  $m$ -degrees in place of total degrees.  $\square$

### Corollary 5.5.

- (1) Let  $\mathcal{O}(k, m) = \mathbb{C}(k, m)\langle X, E^{-1} \rangle$  be the Ore algebra with rational coefficients and the same commutation rules; it contains  $\mathcal{O}$  through the normal forms, but it does not act on  $\mathcal{A}$ . Let  $T = \sum_{a,b} t_{ab}(k, m)X^aE^{-b} \in \mathcal{O}(k, m)$ , and suppose that  $\sum_{a,b} t_{ab}(k, m)U_{k-a}(m-b) = 0$  at all points  $(k, m)$  of some quadrant at which all  $t_{ab}$  are defined. Then  $T \in \mathcal{O}(k, m)L_1$ .
- (2)  $\mathcal{O}(k, m)L_1 \cap \mathcal{O} = \mathcal{O}L_1$ .

*Proof.* (1) Shrink the quadrant so that all indices are nonnegative, let  $\Delta \in \mathbb{C}[k, m]$  be a common denominator of the  $t_{ab}$ , and put  $R = \Delta^2 T \in \mathcal{O}$ . At a point of the quadrant where  $\Delta \neq 0$ , all  $t_{ab}$  are defined and  $(RU)(k, m) = \Delta(k, m)^2 \sum_{a,b} t_{ab}(k, m)U_{k-a}(m-b) = 0$ . Where  $\Delta = 0$ , we have  $(RU)(k, m) = \Delta(k, m)((\Delta T)U)(k, m) = 0$ , because  $\Delta T \in \mathcal{O}$ . So  $R \in \text{Rel}(U)$ , hence  $R = QL_1$  with  $Q \in \mathcal{O}$  by Theorem 5.4, and  $T = \Delta^{-2}QL_1$ .

(2) The reduction in the proof of Theorem 5.4(1) works over  $\mathbb{C}(k, m)$  as well, so every element of  $\mathcal{O}(k, m)$  is of the form  $QL_1 + R_0$  with  $R_0$  free of  $E^{-1}$ . This decomposition is unique: if  $QL_1$  is free of  $E^{-1}$ , then  $Q = 0$ , since for  $Q \neq 0$  the normal form of  $QL_1$  has the nonzero term at  $(\alpha', \beta^* + 1)$  found in the proof of Theorem 5.4(2). If  $R \in \mathcal{O}$  lies in  $\mathcal{O}(k, m)L_1$ , its decomposition in  $\mathcal{O}$  is also one in  $\mathcal{O}(k, m)$ , so its remainder vanishes and  $R \in \mathcal{O}L_1$ .  $\square$

Part (1) is the analogue, for  $U$  and annihilation on a quadrant, of the description of the annihilator of  $\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \}$  with rational coefficients in [13, Example 5]. Part (2) only uses that the coefficient of  $E^{-1}$  in  $L_1$  is a unit; for operators whose leading coefficient is not a unit the contraction of the rational left ideal can be strictly larger, see Zhang [31] for the univariate case.

Theorem 5.4 was also confirmed by exact linear algebra (ranks modulo large primes and exact kernels over  $\mathbb{Q}$ ) for many boxes, using the values  $U_k(m)$  in the window  $0 \leq k, m \leq 40$ . For the triangle we only state the analogous result:  $\text{Rel}(N) = \mathcal{O} \cdot L_N$  with  $L_N = 1 - X - XY - (q - 1)X^3(1 + Y)^2$ , where  $Y$  shifts  $q$ , and the dimension in a box is  $(A - 2)(B - 1)D(D + 1)/2$  for  $A \geq 3$ ,  $B \geq 2$ ,  $D \geq 1$ . We omit the proof. It transports the problem to  $U$  through the binomial transform  $V(k, m) = \sum_q \binom{m}{q} N(k, q)$ , which satisfies  $V(k, m + 1) = U_k(m)$  by (3), and needs an additional saturation lemma for the left factors  $Y$  and  $1 + Y$ ; a complete proof is part of the Lean development (`RelN_eq`, `finrank_relN_tot`; see Section 9).

**5.2.  $U$  is not Stirling-like.** For lattice points  $p, p', p''$  write  $\det(p, p', p'') = \det(p' - p, p'' - p)$ ; its absolute value is twice the area of the triangle.

**Proposition 5.6.** *Let  $R \in \text{Rel}(U)$ ,  $R \neq 0$ . Then  $\text{Supp}(R)$  contains three points  $p_0, p_1, p_2$  with  $\det(p_0, p_1, p_2) \geq 3$ . In particular  $|\text{Supp}(R)| \geq 3$ , and if  $\text{Supp}(R) \subseteq \{q_0, q_1, q_2\}$  for some lattice points  $q_0, q_1, q_2$ , then  $|\det(q_0, q_1, q_2)| \geq 3$ .*

*Proof.* By Theorem 5.4,  $R = QL_1$  with  $Q \neq 0$ ; let  $q$  be the normal form of  $Q$ , so that the normal form  $r$  of  $R$  is given by (4). Let  $\beta^-$  and  $\beta^+$  be the smallest and the largest second coordinate in  $\text{Supp}(Q)$ , let  $\alpha^-$  and  $\alpha^+$  be the smallest and the largest first coordinate in row  $\beta^-$ , and choose  $(\alpha', \beta^+) \in \text{Supp}(Q)$ . Then

$$r(\alpha^-, \beta^-) = q(\alpha^-, \beta^-), \quad r(\alpha^+ + 3, \beta^-) = -(m - \beta^-)q(\alpha^+, \beta^-), \quad r(\alpha', \beta^+ + 1) = -q(\alpha', \beta^+),$$

all nonzero because  $\mathbb{C}[k, m]$  is a domain. These three points give  $\det = (\alpha^+ - \alpha^- + 3)(\beta^+ - \beta^- + 1) \geq 3$ . If  $\text{Supp}(R) \subseteq \{q_0, q_1, q_2\}$ , these three points are a permutation of  $q_0, q_1, q_2$  (their determinant is nonzero, so they are distinct), and a permutation changes the determinant at most by a sign.  $\square$

The three points are the vertices of a translate of the Newton triangle of  $L_1$ , with vertices  $(0, 0), (3, 0), (0, 1)$ , stretched in both directions. This is the combinatorial content of the fact that Newton polygons are additive under multiplication.

**Corollary 5.7.** *Let  $\mathbf{v}_i = (s_i, t_i) \in \mathbb{Z}^2$ ,  $i = 0, 1, 2$ , be pairwise distinct with  $|\det(\mathbf{v}_0, \mathbf{v}_1, \mathbf{v}_2)| \leq 2$ , let  $c_0, c_1, c_2$  be rational functions of  $(k, m)$ , not all zero, and let  $T = \sum_{i=0}^2 c_i(k, m) S^{\mathbf{v}_i}$ , where  $(S^{(s,t)}f)(k, m) = f(k + s, m + t)$ . Then  $T$  does not annihilate  $U$  at all points of any quadrant where its coefficients are defined. In particular  $U$  satisfies no recurrence with at most two terms, and  $U$  is not Stirling-like in the sense of [13, Definition 3].*

*Proof.* Suppose  $T$  annihilates  $U$  on a quadrant, away from the poles of the  $c_i$ . Choose integers  $c \geq \max_i s_i$  and  $d \geq \max_i t_i$  and evaluate the relation at  $(k - c, m - d)$ . Write  $c_i(k - c, m - d) = n_i(k, m)/\Delta(k, m)$  with polynomials  $n_i$  and a common denominator  $\Delta$ . Then  $R' = \sum_i n_i X^{c-s_i} E^{-(d-t_i)}$  satisfies  $(R'U)(k, m) = 0$  on a quadrant wherever  $\Delta(k, m) \neq 0$ , and  $R = \Delta \cdot R'$  satisfies it everywhere on that quadrant. So  $R \in \text{Rel}(U)$ . The points  $(c - s_i, d - t_i)$  are pairwise distinct, so the normal form of  $R$  has the coefficients  $\Delta n_i$ , not all zero; hence  $R \neq 0$ , and  $\text{Supp}(R)$  is contained in the image of  $\{(s_i, t_i)\}$  under  $(s, t) \mapsto (c - s, d - t)$ , which preserves  $|\det|$ . This contradicts Proposition 5.6. A Stirling-like sequence is annihilated by its three-term generator  $u + v S^{(v_1, v_2)} - w S^{(w_1, w_2)}$ , whose shift points  $(0, 0), (v_1, v_2), (w_1, w_2)$  are distinct with  $|\det| = 1$  because  $(v_1, v_2), (w_1, w_2)$  form a basis of  $\mathbb{Z}^2$ .  $\square$

In particular, the results of [13] on Stirling-like sequences do not apply to  $U$  directly.

**Remark 5.8.** In [13, Section 2.2] sequences are functions on all of  $\mathbb{Z}^2$ , operators have rational coefficients, and the annihilator of  $f$  consists of the operators  $Q$  with  $Q \cdot f \equiv 0$ ; Kauers extends  $\{n \choose k\}$  and the other classical triangles to  $\mathbb{Z}^2$  so that their recurrences hold everywhere. If some extension of  $U$  to  $\mathbb{Z}^2$  were Stirling-like, its three-term generator would annihilate that extension at all points where its coefficients are defined. Write the shifts of the generator in the coordinates  $(k, m)$  as  $(0, 0)$ ,  $(s_1, t_1)$  and  $(s_2, t_2)$ . On the quadrant  $k \geq \max(0, -s_1, -s_2)$ ,  $m \geq \max(0, -t_1, -t_2)$  the generator reads only values at points of  $\mathbb{N}^2$ , where the extension agrees with  $U$ , so it would annihilate  $U$  at all points of this quadrant where its coefficients are defined, which Corollary 5.7 excludes. In this subsection only Proposition 5.6 is formalized. The passage from [13, Definition 3] to it, that is, the reduction in the proof of Corollary 5.7 together with the argument above, is not.

## 6. OBSTRUCTIONS TO SPECIFIC SUMMATION FORMULAS

The first summand of Theorem 4.6 is a single sum of products of a Stirling number and a binomial coefficient, and the second is a double sum. In this section we show that  $U_k(m)$  itself has no single-sum expression of this kind, and determine all binomial shapes with nonnegative parameters that do occur (Section 6.1); and that it is not a sum of two such single sums for  $m \geq 2$  (Section 6.2). These results exclude specific classes of formulas only: single sums of the shapes with  $\alpha + \beta = 1$  exist for every sequence (Theorem 6.3), and whether the double sum of Theorem 4.6 is shortest in some wider sense is open (Section 10).

### 6.1. Single-family sums.

**Theorem 6.1.** *Let  $m \geq 1$ . There are no integers  $c, d$  and  $k_0 \geq 0$  and complex numbers  $A(s)$  ( $s \in \mathbb{Z}$ ), independent of  $k$ , such that*

$$U_k(m) = \sum_s A(s) \binom{k+c-2s}{m+s+d} \quad \text{for all } k \geq k_0.$$

*Proof. Step 1: shape of the generating function.* We use the binomial convention of Section 2; in particular only  $s \geq -m-d$  contribute, and the sum over  $s$  is finite for each  $k$ . Let  $u = x^3/(1-x)$ . For  $n \geq 0$ ,

$$\sum_k \binom{k+c-2s}{n} x^k = \frac{x^{n-c+2s}}{(1-x)^{n+1}},$$

and for  $n = m+s+d$  this equals  $x^g u^{s+m+d+1}$  with  $g = -2(m+d) - c - 3$ , independent of  $s$ . The right-hand side therefore sums, in  $\mathbb{C}((x))$ , to  $x^g \tilde{A}(u)$  with  $\tilde{A}(u) = \sum_{s \geq -m-d} A(s) u^{s+m+d+1} \in \mathbb{C}[[u]]$ . It differs from  $G_m$  only in finitely many coefficients, so

$$(5) \quad G_m = x^g \tilde{A}(u) + L(x)$$

for a Laurent polynomial  $L$ .

*Step 2:  $\tilde{A}$  is rational.* The series  $x$  is a root of  $T^3 + uT - u$ , which is Eisenstein at the prime  $u$  of  $\mathbb{C}[[u]]$ ; hence  $[\mathbb{C}((u))(x) : \mathbb{C}((u))] = 3$  and  $1, x, x^2$  are linearly independent over  $\mathbb{C}((u))$ . Now  $\tilde{A}(u) = x^{-g}(G_m - L)$  lies in  $\mathbb{C}(x) = \mathbb{C}(u)[x]$ , so  $\tilde{A} = \alpha + \beta x + \gamma x^2$  with  $\alpha, \beta, \gamma \in \mathbb{C}(u)$ . Comparing coordinates over  $\mathbb{C}((u))$  gives  $\beta = \gamma = 0$ , so  $\tilde{A} \in \mathbb{C}(u)$ .

*Step 3: residues on the fibre  $u = 1$ .* The equation  $u(x) = 1$  is equivalent to  $b_1(x) = 1 - x - x^3 = 0$ , so the fibre over  $u = 1$  consists of the three roots  $\xi_1, \xi_2, \xi_3$  of  $b_1$ , which is irreducible over  $\mathbb{Q}$ . Let  $\xi$  be one of them. Then  $1 - \xi = \xi^3$ , hence  $b_w(\xi) = (1 - w)\xi^3$ ,  $\xi b'_1(\xi) = -\xi - 3\xi^3 = 2\xi - 3$ , and  $u'(\xi) = \xi^2(3 - 2\xi)/(1 - \xi)^2 = (3 - 2\xi)/\xi^4 \neq 0$ . Moreover  $P_{j-1}(\xi) = 0$  for  $j \geq 2$ , so  $W_m(\xi) = W_1(\xi) = 1 + \xi^2(1 - \xi) = \xi(1 + \xi) \neq 0$ , and

$$P'_m(\xi) = b'_1(\xi) \prod_{w \neq 1} b_w(\xi) = b'_1(\xi) \xi^{3m} (-1)^{m-1} (m-1)!.$$

So  $G_m$  has a simple pole at  $\xi$  with residue  $W_m(\xi)/P'_m(\xi)$ , and

$$\text{Res}_{x=\xi} G_m \cdot u'(\xi) = \frac{\xi(1+\xi) \cdot \xi \cdot (3-2\xi)}{(2\xi-3)\xi^{3m}(-1)^{m-1}(m-1)!\xi^4} = \frac{(-1)^m(1+\xi)\xi^{-3m-2}}{(m-1)!}.$$

On the other hand  $u-1$  is a local coordinate at  $\xi$ , and  $L$  is holomorphic there. By (5),  $\tilde{A}$  has a simple pole at  $u=1$ , and  $\text{Res}_{x=\xi} G_m = \xi^g \text{Res}_{u=1} \tilde{A}/u'(\xi)$ . Hence  $\theta := (1+\xi)\xi^{-M}$  with  $M = 3m+2+g$  takes the same value  $(-1)^m(m-1)!\text{Res}_{u=1} \tilde{A}$  at the three roots. Since the three roots are conjugate over  $\mathbb{Q}$ , the characteristic polynomial of  $\theta$  over  $\mathbb{Q}$  is  $(X-\theta)^3$ , so  $\theta \in \mathbb{Q}$ .

*Step 4: a norm.* Let  $f = X^3 + X - 1$ , the monic minimal polynomial of  $\xi$ . Then  $\text{Nm}(\xi) = -f(0) = 1$  and  $\text{Nm}(1+\xi) = -f(-1) = 3$ , so  $\theta^3 = \text{Nm}(\theta) = 3$ , which has no rational solution.  $\square$

For  $m=0$  the statement fails ( $U_k(0) = 1$ ). The part of  $U_k(m)$  counting sequences that do not end with an ascent is of the excluded form, with  $A(s) = \left\{ \begin{smallmatrix} m+s \\ m \end{smallmatrix} \right\}$ ; so it is the sequences ending with an ascent that destroy the single-family structure. They cannot be written in this form either:

**Theorem 6.2.** *Let  $U_k^\uparrow(m)$  be the number of  $h \in \mathcal{H}_k(m)$  with  $h_{k-1} < h_k$ . For every  $m \geq 2$  the conclusion of Theorem 6.1 holds with  $U_k(m)$  replaced by  $U_k^\uparrow(m)$ .*

*Proof.* By Theorem 4.4 with  $y=1$ ,  $G_m^\uparrow(x) := \sum_k U_k^\uparrow(m)x^k = (W_m-1)/P_m = x^2 \sum_{j=1}^m j / \prod_{v=j}^m b_v$ . Steps 1 and 2 of the previous proof are unchanged. The fibre  $u=1/i$  is the set of roots of  $b_i$ . Let  $1 \leq i \leq m$  and let  $\eta$  be a root of  $b_i$ . Then  $1-\eta = i\eta^3$ , so

$$b_v(\eta) = (i-v)\eta^3, \quad b'_i(\eta) = \frac{2\eta-3}{\eta}, \quad u'(\eta) = \frac{3-2\eta}{i^2\eta^4}.$$

Only the terms with  $j \leq i$  have a pole at  $\eta$ , and  $\prod_{v=j, v \neq i}^m b_v(\eta) = \eta^{3(m-j)}(i-j)!(-1)^{m-i}(m-i)!$ . Put  $\tilde{W}_i := 1 + \sum_{j=1}^i j i^j x^{3j+2}$ , so that  $\sum_{j=1}^i j \eta^{3j}/(i-j)! = (\tilde{W}_i(\eta) - 1)/(i!\eta^2)$ . Then

$$\text{Res}_{x=\eta} G_m^\uparrow \cdot u'(\eta) = \frac{(-1)^{m-i+1}(\tilde{W}_i(\eta) - 1)\eta^{-3m-3}}{(m-i)!i!\eta^2}.$$

Take  $i=2 \leq m$ . The polynomial  $b_2 = 1 - x - 2x^3$  has no rational root, so it is irreducible over  $\mathbb{Q}$ , and as in Step 3 the quantity  $\theta = (\tilde{W}_2(\eta) - 1)\eta^{-3m-3-g}$  is rational. Now  $\tilde{W}_2 - 1 = 2x^5 + 4x^8$ , and  $2\eta^3 = 1 - \eta$  gives  $\tilde{W}_2(\eta) - 1 = \eta^5(4 - 2\eta)$ , so  $\theta = (4 - 2\eta)\eta^n$  with  $n = 2 - 3m - g$ . The monic minimal polynomial of  $\eta$  is  $X^3 + X/2 - 1/2$ , so in  $\mathbb{Q}(\eta)$  we have  $\text{Nm}(\eta) = 1/2$  and  $\text{Nm}(4 - 2\eta) = 8(2^3 + 2/2 - 1/2) = 68$ , hence  $\text{Nm}(\theta) = 17 \cdot 2^{2-n}$ . But  $\text{Nm}(\theta) = \theta^3$  for rational  $\theta$ , and the 17-adic valuation of a rational cube is divisible by 3.  $\square$

For  $m=1$ ,  $G_1^\uparrow = x^2/b_1 = x^{-1}u/(1-u)$  is of the excluded form, so the bound  $m \geq 2$  is sharp.

The fibre argument also determines all binomial shapes with nonnegative parameters. Here  $\xi \in (0,1)$  is the real root of  $b_1$  (so  $\xi \approx 0.6823$ ). We use that its minimal polynomial over  $\mathbb{Q}$  is  $X^3 + X - 1$ , that  $\text{Nm}(\xi) = 1$ , that  $\mathbb{Q}(\xi)$  is not a normal extension of  $\mathbb{Q}$  (the other two roots of  $b_1$  are not real), and that  $\xi^N \notin \mathbb{Q}$  for every integer  $N \neq 0$ : if  $\xi^N = r \in \mathbb{Q}$ , then  $r^3 = \text{Nm}(\xi)^N = 1$ , so  $\xi^N = 1$ , which is impossible.

**Theorem 6.3.** *Let  $m \geq 1$  and let  $\alpha, \beta \geq 0$  be integers, not both 0. There are integers  $c, d, s_0$  and  $k_0 \geq 0$  and complex numbers  $A(s)$  ( $s \geq s_0$ ), independent of  $k$ , such that*

$$(6) \quad U_k(m) = \sum_{s \geq s_0} A(s) \binom{k+c-\alpha s}{\beta s+d} \quad \text{for all } k \geq k_0$$

*if and only if  $\alpha + \beta = 1$ .*

*Proof.* For  $(\alpha, \beta) = (0, 1)$ , Newton's forward difference formula  $U_k(m) = \sum_{s=0}^k \Delta^s U_\bullet(m) \big|_0 \binom{k}{s}$  has the form (6), and for  $(1, 0)$  so has  $U_k(m) = \sum_{s=0}^k (U_s(m) - U_{s-1}(m)) \binom{k-s}{0}$ , with  $U_{-1}(m) = 0$ . Now let  $\alpha + \beta \geq 2$ . The shape  $(2, 1)$  is Theorem 6.1: there the lower index is written  $m + s + d$ , and terms with a negative lower index vanish, so the lower limit  $s_0$  makes no difference. So let  $(\alpha, \beta) \neq (2, 1)$ , and suppose that (6) holds.

*Step 1: generating functions.* Terms with  $\beta s + d < 0$  vanish, so we may assume  $A(s) = 0$  for them; if  $\beta = 0$  and  $d < 0$ , then all terms vanish, which is impossible. For  $\beta s + d \geq 0$ ,

$$\sum_k \binom{k + c - \alpha s}{\beta s + d} x^k = x^{d-c} (1-x)^{-d-1} v^s, \quad v = \frac{x^{\alpha+\beta}}{(1-x)^\beta}.$$

Since  $v$  has  $x$ -adic valuation  $\alpha + \beta$ , the right-hand side of (6) sums in  $\mathbb{C}((x))$  to  $x^{d-c} (1-x)^{-d-1} \tilde{A}(v)$  with  $\tilde{A} = \sum_s A(s) v^s \in \mathbb{C}((v))$ , and  $G_m = x^{d-c} (1-x)^{-d-1} \tilde{A}(v) + L(x)$  for a Laurent polynomial  $L$ .

*Step 2:  $\tilde{A}$  is rational.* Let  $y = x$  if  $\alpha \geq 1$ , and  $y = x/(1-x)$  if  $\alpha = 0$ ; in the second case  $v = y^\beta$ . Then  $\mathbb{C}(x) = \mathbb{C}(y)$ , the series  $y$  has  $x$ -adic valuation 1, and  $y$  is a root of  $T^{\alpha+\beta} - v(1-T)^\beta$ , respectively of  $T^\beta - v$ ; this polynomial is monic of degree  $\alpha + \beta$  and Eisenstein at  $v$ . Hence  $\mathbb{C}(x) = \bigoplus_{i < \alpha+\beta} \mathbb{C}(v) y^i$ , and  $1, y, \dots, y^{\alpha+\beta-1}$  are linearly independent over  $\mathbb{C}((v))$ , because their valuations are distinct modulo  $\alpha + \beta$ . As  $\tilde{A}(v) = x^{c-d} (1-x)^{d+1} (G_m - L)$  lies in  $\mathbb{C}(x)$ , comparing coordinates gives  $\tilde{A} \in \mathbb{C}(v)$ .

*Step 3: the fibre over  $v(\xi)$ .* The function  $G_m$  has a pole at  $\xi$  (Theorem 3.1), and  $x^{d-c} (1-x)^{-d-1}$  and  $L$  are holomorphic at  $\xi$ , so the rational function  $\tilde{A}$  has a pole at  $v_0 = v(\xi)$ . Let  $\eta \in \mathbb{C} \setminus \{0, 1\}$  with  $v(\eta) = v_0$ . Since  $v$  is holomorphic and not constant near  $\eta$ ,  $\tilde{A}(v(x)) \rightarrow \infty$  as  $x \rightarrow \eta$ , so  $G_m$  has a pole at  $\eta$ , and  $b_w(\eta) = 0$  for some  $w$  with  $1 \leq w \leq m$  ( $w = 0$  is excluded because  $\eta \neq 1$ ). Since  $1 - \xi = \xi^3$ , we have  $v_0 = \xi^e$  with  $e = \alpha - 2\beta$ . We now find a point of the fibre  $\{v = v_0\}$  that violates this, according to the shape.

*Case  $\alpha, \beta \geq 1$  and  $e \neq 0$ .* The fibre is the zero set of the monic polynomial  $\varphi(x) = x^{\alpha+\beta} - \xi^e(1-x)^\beta$ , and  $\varphi(0) = -\xi^e$ ,  $\varphi(1) = 1$ . Let  $\eta_1, \dots, \eta_{\alpha+\beta}$  be the zeros of  $\varphi$ , with multiplicity. Each of them is a zero of some  $b_{w_i}$  with  $w_i \geq 1$ , that is,  $w_i = (1 - \eta_i)/\eta_i^3$ , and therefore

$$\prod_i w_i = \frac{\varphi(1)}{(\prod_i \eta_i)^3} = ((-1)^{\alpha+\beta+1} \xi^e)^{-3} = \pm \xi^{-3e}.$$

The left-hand side is a positive integer, in particular rational, but  $\xi^{-3e} \notin \mathbb{Q}$ .

*Case  $\beta = 0$  and  $\alpha \geq 2$ .* Here  $v = x^\alpha$ , and  $\eta = \xi\zeta$  with  $\zeta = e^{2\pi i/\alpha}$  lies in the fibre. If  $b_w(\xi\zeta) = 0$ , then, using  $\xi^3 = 1 - \xi$ , we get  $1 - w\zeta^3 = \xi(\zeta - w\zeta^3)$ . If  $\zeta \neq w\zeta^3$ , this puts  $\xi$  in the abelian field  $\mathbb{Q}(\zeta)$ , all of whose subfields are normal over  $\mathbb{Q}$ , whereas  $\mathbb{Q}(\xi)$  is not. If  $\zeta = w\zeta^3$ , then  $w\zeta^2 = 1$ , so  $w = 1$  and  $\zeta = -1$ , and the equation reads  $2 = 0$ .

*Case  $\alpha = 2\beta$  and  $\beta \geq 2$ .* Here  $v = u^\beta$  with  $u = x^3/(1-x)$ , and  $v_0 = 1$ . Let  $\zeta = e^{2\pi i/\beta}$  and let  $\eta$  be a zero of  $x^3 - \zeta(1-x)$ . Then  $\eta \notin \{0, 1\}$  and  $u(\eta) = \zeta$ , so  $\eta$  lies in the fibre. But  $b_w(\eta) = 0$  means  $u(\eta) = 1/w$ , which is impossible, since  $\zeta \neq 1$  has absolute value 1.

*Case  $\alpha = 0$  and  $\beta \geq 2$ .* With  $y = x/(1-x)$  we have  $v_0 = y(\xi)^\beta = \xi^{-2\beta}$ , so the fibre consists of the points  $x_\zeta = \zeta/(\xi^2 + \zeta)$  with  $\zeta^\beta = 1$ , none of which is 0 or 1. If  $\beta$  is even, then  $x_{-1} = 1/(1 - \xi^2) > 1$ , but  $b_w(x) < 0$  for  $x > 1$ . Let  $\beta$  be odd. Each  $x_\zeta$  is a zero of some  $b_{w_\zeta}$ , so  $u(x_\zeta) = 1/w_\zeta$ ; as  $1 - x_\zeta = \xi^2/(\xi^2 + \zeta)$ , this reads  $w_\zeta \zeta^3 = \xi^2(\xi^2 + \zeta)^2$ . Taking the product over all  $\zeta$  and using  $\prod_\zeta \zeta = 1$  and  $\prod_\zeta (\xi^2 + \zeta) = \xi^{2\beta} + 1$ , we obtain  $\prod_\zeta w_\zeta = \chi^2$  with  $\chi = \xi^{3\beta} + \xi^\beta = (1 - \xi)^\beta + \xi^\beta$ . So  $\chi^2 \in \mathbb{Q}$ , and since the cubic field  $\mathbb{Q}(\xi)$  has no quadratic subfield,  $\chi \in \mathbb{Q}$ . Then the polynomial  $\chi(X) = (1 - X)^\beta + X^\beta$  takes the same value at all roots of  $b_1$ . But  $0 < \chi(\xi) < 2\xi^\beta \leq 2\xi^3 < 0.64$ , whereas a non-real root  $\xi_2$  of  $b_1$  satisfies  $|\xi_2|^2 = 1/\xi$  and  $1 - \xi_2 = \xi_2^3$ , so that  $|\chi(\xi_2)| = |\xi_2|^\beta |\xi_2^{2\beta} + 1| \geq \xi^{-\beta/2} (\xi^{-\beta} - 1) \geq \xi^{-3/2} (\xi^{-3} - 1) > 3.8$ .  $\square$

**Remark 6.4.** Shapes with  $\alpha < 0$  are not covered by Theorem 6.3. If  $\alpha + \beta = 1$ , for instance  $(\alpha, \beta) = (-1, 2)$ , then  $v$  has  $x$ -adic valuation 1, and every sequence again has a representation (6). If  $\alpha < 0$  and  $\alpha + \beta \geq 2$ , then  $v$  vanishes also at  $x = \infty$ , so  $\mathbb{C}(x)$  has degree  $\beta > \alpha + \beta$  over  $\mathbb{C}(v)$  and the comparison of coordinates in Step 2 breaks down. We do not treat this case here.

**6.2. Sums of two families.** Let  $u = x^3/(1-x)$ . By Step 1 of the proof of Theorem 6.1, a single sum of the shape  $(2, 1)$  that equals  $U_k(m)$  for all large  $k$  amounts to an identity  $G_m = x^g \tilde{A}(u) + L(x)$  with  $\tilde{A} \in \mathbb{C}((u))$  and a Laurent polynomial  $L$ . Conversely,  $x^g u^s = x^{g+3s}(1-x)^{-s}$  is a Laurent polynomial for  $s \leq 0$  and equals  $\sum_k \binom{k-g-2s-1}{s-1} x^k$  for  $s \geq 1$ , so every series  $x^g A(u)$  with  $A \in \mathbb{C}((u))$  is, up to a Laurent polynomial, the generating function of a single sum of the shape  $(2, 1)$ . We call  $x^g A(u)$  a *family*. For example, the formula  $U_k(1) = \sum_{s \geq 0} \binom{k+1-2s}{s+1} + \sum_{s \geq 0} \binom{k-2-2s}{s}$  of Example 4.8 is a sum of two families.

**Theorem 6.5.** *Let  $m \geq 2$ . There are no integers  $g_1, g_2$ , series  $A_1, A_2 \in \mathbb{C}((u))$  and Laurent polynomial  $L$  such that*

$$G_m = x^{g_1} A_1(u) + x^{g_2} A_2(u) + L(x).$$

*For  $m \geq 3$  the same holds with  $G_m$  replaced by  $G_m^\uparrow$ .*

By the remarks above, the first statement says equivalently that there are no integers  $c_1, d_1, c_2, d_2$  and  $k_0 \geq 0$  and complex numbers  $A_1(s), A_2(s)$  such that

$$U_k(m) = \sum_s A_1(s) \binom{k+c_1-2s}{m+s+d_1} + \sum_s A_2(s) \binom{k+c_2-2s}{m+s+d_2} \quad \text{for all } k \geq k_0.$$

Taking  $A_2 = 0$ , the theorem contains Theorem 6.1 for  $m \geq 2$  and Theorem 6.2 for  $m \geq 3$ , with a different proof.

**Corollary 6.6.** *The least number of families whose sum differs from  $G_m$  by a Laurent polynomial is 1 for  $m = 0$ , 2 for  $m = 1$  and 3 for  $m \geq 2$ . For  $G_m^\uparrow$  it is 1 for  $m = 1$ , 2 for  $m = 2$  and 3 for  $m \geq 3$ .*

The proof of Theorem 6.5 has two parts: a necessary condition on each fibre  $u = 1/i$  that does not involve  $m$ , and a finite computation showing that the conditions for  $i = 1, 2$  (and for  $i = 1, 2, 3$  in the case of  $G^\uparrow$ ) are incompatible.

**Lemma 6.7.** *Every  $f \in \mathbb{C}(x)$  can be written uniquely as  $f = a_0 + a_1 x + a_2 x^2$  with  $a_r \in \mathbb{C}(u)$ , and  $1, x, x^2$  are linearly independent over  $\mathbb{C}((u))$ ; we call  $(a_0, a_1, a_2)$  the coordinates of  $f$ . The coordinates  $e_g$  of  $x^g$  lie in  $\mathbb{C}[u, u^{-1}]^3$  for every  $g \in \mathbb{Z}$ , and the coordinates of  $G_m$  and of  $G_m^\uparrow$  have the form  $c(u)/\prod_{v=1}^m (1-vu)$  with  $c \in \mathbb{C}[u, u^{-1}]^3$ .*

*Proof.* The first statement is Step 2 of the proof of Theorem 6.1. The relations  $x^3 = u - ux$  and  $x^{-1} = (x^2 + u)/u$  give the second. Finally  $b_v = (1-x)(1-vu)$  and  $(1-x)^{-1} = ux^{-3}$ , so  $1/P_m = u^{m+1}x^{-3m-3}/\prod_{v=1}^m (1-vu)$ , while  $W_m$  and  $W_m - 1$  are polynomials in  $x$ .  $\square$

Fix  $i \geq 1$ . The fibre  $u = 1/i$  is the set of zeros  $\eta_1, \eta_2, \eta_3$  of  $b_i$ ; they are distinct, and  $u'(\eta_j) \neq 0$ , because  $u' = x^2(3-2x)/(1-x)^2$  and  $0, 1, 3/2$  are not zeros of  $b_i$ . Let  $V$  be the Vandermonde matrix with rows  $(1, \eta_j, \eta_j^2)$ . If the coordinates  $a$  of  $f \in \mathbb{C}(x)$  are regular at  $u = 1/i$ , then  $(f(\eta_j))_j = V a(1/i)$ ; if they have at most a simple pole there, with residue vector  $R$ , then  $(u'(\eta_j) \text{Res}_{x=\eta_j} f)_j = VR$ , since  $u - 1/i$  is a local coordinate at each  $\eta_j$ .

**Proposition 6.8.** *Suppose that  $G_m = x^{g_1} A_1(u) + x^{g_2} A_2(u) + L(x)$  with integers  $g_1, g_2$ , series  $A_1, A_2 \in \mathbb{C}((u))$  and a Laurent polynomial  $L$ , and let  $1 \leq i \leq m$ . Then*

$$\det \left( \eta_j^{g_1}, \eta_j^{g_2}, u'(\eta_j) \text{Res}_{x=\eta_j} G_m \right)_{j=1,2,3} = 0.$$

*The same holds with  $G_m$  replaced by  $G_m^\uparrow$ .*

*Proof.* Let  $a$  and  $\ell$  be the coordinates of  $G_m$  and of  $L$ . By the uniqueness of coordinates over  $\mathbb{C}((u))$ ,  $a = A_1 e_{g_1} + A_2 e_{g_2} + \ell$ . The cross product  $\nu = e_{g_1} \times e_{g_2}$  lies in  $\mathbb{C}[u, u^{-1}]^3$  and is orthogonal to  $e_{g_1}$  and  $e_{g_2}$ , so  $\nu \cdot a = \nu \cdot \ell$ , an element of  $\mathbb{C}[u, u^{-1}]$ . By Lemma 6.7,  $a$  has at most a simple pole at  $u = 1/i$ ; if  $R$  is its residue vector, it follows that  $\nu(1/i) \cdot R = 0$ , that is,  $\det(e_{g_1}(1/i), e_{g_2}(1/i), R) = 0$ . Multiplying by  $V$  gives the claim.  $\square$

For  $1 \leq i \leq m$  and a zero  $\eta$  of  $b_i$ , the computation in the proof of Theorem 6.2, together with  $W_m(\eta) = \tilde{W}_i(\eta)$ , gives

$$(7) \quad u'(\eta) \operatorname{Res}_{x=\eta} G_m = \frac{(-1)^{m-i+1} \tilde{W}_i(\eta) \eta^{-3m-3}}{(m-i)! i! i^2},$$

and the formula there for  $G_m^\dagger$  is the same with  $\tilde{W}_i$  replaced by  $\tilde{W}_i - 1$ .

**Corollary 6.9.** *In the situation of Proposition 6.8, let  $b = g_2 - g_1$  and  $n = -3m - 3 - g_1$ , and let  $W = \tilde{W}_i$  for  $G_m$ , respectively  $W = \tilde{W}_i - 1$  for  $G_m^\dagger$ . If  $b_i$  is irreducible over  $\mathbb{Q}$  and  $\eta$  is a zero of  $b_i$ , then  $1, \eta^b$  and  $W(\eta)\eta^n$  are linearly dependent over  $\mathbb{Q}$ .*

*Proof.* Divide the  $j$ -th row of the determinant by  $\eta_j^{g_1}$ , and the third column by the rational factor in (7); then the determinant of the rows  $(1, \eta_j^b, W(\eta_j)\eta_j^n)$  vanishes. Since the  $\eta_j$  are the conjugates of  $\eta$ , this matrix is  $V$  times the matrix of coordinates of  $1, \eta^b, W(\eta)\eta^n$  with respect to the basis  $1, \eta, \eta^2$  of  $\mathbb{Q}(\eta)$ , and  $V$  is invertible.  $\square$

The condition no longer involves  $m$ , and  $b$  and  $n$  are the same for all fibres, because the exponent  $-3m - 3$  in (7) does not depend on  $i$ . For  $i \in \{1, 2, 3\}$  the cubic  $b_i$  has no rational zero, so it is irreducible. In  $\mathbb{Q}(\eta)$  we have  $\eta^3 = (1 - \eta)/i$  and  $\eta^{-1} = 1 + i\eta^2$ , so the coordinates of  $\eta^b$  and of  $W(\eta)\eta^n$  lie in  $\mathbb{Z}[1/i]$ . Let  $J_i(n, b) \in \mathbb{Z}[1/i]$  be the determinant formed from their last two coordinates, for  $W = \tilde{W}_i$ , and  $J_i^\dagger(n, b)$  the same for  $W = \tilde{W}_i - 1$ . Since the coordinates of  $1$  are  $(1, 0, 0)$ , the condition of Corollary 6.9 is  $J_i(n, b) = 0$ , respectively  $J_i^\dagger(n, b) = 0$ .

**Proposition 6.10.** *There is no  $(n, b) \in \mathbb{Z}^2$  with  $b \neq 0$  and  $J_1(n, b) = J_2(n, b) = 0$ , and none with  $b \neq 0$  and  $J_1^\dagger(n, b) = J_2^\dagger(n, b) = J_3^\dagger(n, b) = 0$ .*

Let  $\ell$  be an odd prime that does not divide  $i$ . Then  $\mathbb{Z}_\ell[\eta] = \mathbb{Z}_\ell[x]/(b_i)$  is a free  $\mathbb{Z}_\ell$ -module with basis  $1, \eta, \eta^2$ , and  $\eta$  is a unit in it. Let  $P \geq 1$  be such that  $\eta^P \equiv 1 \pmod{\ell \mathbb{Z}_\ell[\eta]}$ , and write  $\eta^P = 1 + \ell\mu$  with  $\mu \in \mathbb{Z}_\ell[\eta]$ .

**Lemma 6.11.** (1)  $J_i(n, b) \pmod{\ell}$  depends only on  $n \pmod{P}$  and  $b \pmod{P}$ .

(2) Let  $\Theta(n)$  be the determinant formed from the last two coordinates of  $\mu$  and of  $W(\eta)\eta^n$ . If  $b = Pb'$  with  $b' \neq 0$ , then  $J_i(n, b) = b'\ell(\Theta(n) + \ell\kappa)$  for some  $\kappa \in \mathbb{Z}_\ell$ . In particular  $J_i(n, b) \neq 0$  whenever  $\Theta(n) \not\equiv 0 \pmod{\ell}$ , a condition that depends only on  $n \pmod{P}$ .

The same holds for  $J_i^\dagger$ .

*Proof.* (1) holds because  $\eta^n \pmod{\ell}$  depends only on  $n \pmod{P}$ . For (2), the binomial series gives  $\eta^b = (1 + \ell\mu)^{b'} = \sum_{r \geq 0} \binom{b'}{r} \ell^r \mu^r$ , also for  $b' < 0$ : the series converges in  $\mathbb{Z}_\ell[\eta]$ , and its sum is  $(1 + \ell\mu)^{b'}$  because  $(1 + Y)^a(1 + Y)^c = (1 + Y)^{a+c}$  in  $\mathbb{Z}[[Y]]$  for all integers  $a, c$  and the substitution  $Y = \ell\mu$  is continuous. The determinant is  $\mathbb{Z}_\ell$ -linear and continuous in its first argument and vanishes for the element  $1$ , so  $J_i(n, b) = \sum_{r \geq 1} \binom{b'}{r} \ell^r \Theta_r$ , where  $\Theta_r$  is formed from  $\mu^r$  in place of  $\mu$ ; thus  $\Theta_1 = \Theta(n)$ . For  $r \geq 2$  the number  $\binom{b'}{r} \ell^r / (b'\ell) = \binom{b'-1}{r-1} \ell^{r-1} / r$  has  $\ell$ -adic valuation at least  $r - 1 - v_\ell(r) \geq 1$ , since  $v_\ell(2) = 0$  and  $v_\ell(r) \leq \log_3 r \leq r - 2$  for  $r \geq 3$ .  $\square$

*Proof of Proposition 6.10.* Let  $M = 5040 = 2^4 \cdot 3^2 \cdot 5 \cdot 7$ , and use the primes of Table 3, where  $P$  is the order of  $\eta$  in  $(\mathbb{Z}_\ell[\eta]/\ell)^\times$ . For each certificate prime a computer tabulates, in  $\mathbb{Z}_\ell[\eta]/\ell$ , the last

| fibre   | pairs $(\ell, P)$ , with $P$ the order of $\eta$ modulo $\ell$   |
|---------|------------------------------------------------------------------|
| $i = 1$ | $(3, 8), (11, 60), (13, 168), (29, 840), (2521, 2520)$           |
| $i = 2$ | $(7, 48), (17, 72), (19, 18), (41, 280), (71, 5040), (127, 126)$ |
| $i = 3$ | $(13, 84), (71, 70)$                                             |

TABLE 3. The certificate primes for Proposition 6.10. All orders divide 5040.

two coordinates of  $\eta^b$  and of  $W(\eta)\eta^n$  for  $b, n \in \mathbb{Z}/P\mathbb{Z}$ , and  $\mu \bmod \ell$  from  $\eta^P$  computed modulo  $\ell^2$ ; all of this is integer arithmetic. It then checks the following two finite statements.

- (a) For every  $b_0 \in \{1, \dots, M-1\}$  and every  $n_0 \in \mathbb{Z}/M\mathbb{Z}$ , some prime of the fibres  $i = 1, 2$  satisfies  $J_i(n_0, b_0) \not\equiv 0 \pmod{\ell}$ , and some prime of the fibres  $i = 1, 2, 3$  satisfies  $J_i^\uparrow(n_0, b_0) \not\equiv 0 \pmod{\ell}$ .
- (b) For every  $n_0 \in \mathbb{Z}/M\mathbb{Z}$ , some prime of the fibres  $i = 1, 2$  satisfies  $\Theta(n_0) \not\equiv 0 \pmod{\ell}$  for  $W = \tilde{W}_i$ , and some prime of the fibres  $i = 1, 2, 3$  does so for  $W = \tilde{W}_i - 1$ .

In (a) there are  $M(M-1) = 25,396,560$  classes in each case. In (b), for  $W = \tilde{W}_i$  the primes of the fibre  $i = 1$  suffice: trying them in the order of the table, 3, 11, 13, 29 and 2521 are the first to succeed for 3780, 1176, 72, 6 and 6 classes. For  $W = \tilde{W}_i - 1$  they fail exactly for  $n_0 \in \{2515, 5035\}$ , and there the primes 7 and 17 of the fibre  $i = 2$  succeed, respectively. The failure at  $n_0 = 5035 \equiv -5$  is unavoidable:  $\tilde{W}_1 - 1 = x^5$ , so  $W(\eta)\eta^{-5} = 1$  on the fibre  $i = 1$ , and the condition holds there for  $n = -5$  and every  $b$ .

The script `code/tableB/check_b2.py` of the repository performs these checks in about twenty seconds and is part of `verify_all.py`. The certificate is not unique: separately written programs found complete certificates with the moduli 32760, 55440 and 83160, using other primes.

Now let  $b \neq 0$ , and suppose that the conditions of the proposition hold at  $(n, b)$ . If  $M \nmid b$ , then by (a) and Lemma 6.11(1) one of the determinants is nonzero modulo a certificate prime. If  $M \mid b$ , then  $P \mid b$  for every certificate prime, and by (b) and Lemma 6.11(2) one of the determinants is nonzero. Either way we have a contradiction.  $\square$

*Proof of Theorem 6.5.* If  $g_1 = g_2$ , replace  $A_1$  by  $A_1 + A_2$ , then  $g_2$  by  $g_1 + 1$  and  $A_2$  by 0; so we may assume  $b = g_2 - g_1 \neq 0$ . For  $m \geq 2$ , Corollary 6.9 with  $i = 1, 2$  gives  $J_1(n, b) = J_2(n, b) = 0$ , and for  $G_m^\uparrow$  with  $m \geq 3$  it gives  $J_1^\uparrow(n, b) = J_2^\uparrow(n, b) = J_3^\uparrow(n, b) = 0$ , contradicting Proposition 6.10.  $\square$

*Proof of Corollary 6.6.* Three families always suffice: by Lemma 6.7,  $G_m = a_0 + a_1x + a_2x^2$  with  $a_r \in \mathbb{C}(u)$ , and if  $a_r^+$  denotes the part of the expansion of  $a_r$  at  $u = 0$  with positive powers of  $u$ , then  $G_m - \sum_r x^r a_r^+(u)$  is a Laurent polynomial, because  $x^r u^e = x^{r+3e}(1-x)^{-e}$  for  $e \leq 0$ . The same holds for  $G_m^\uparrow$ . Next,  $G_0 = 1/(1-x) = x^{-3}u$ , and  $G_1$  is a sum of two families by the formula for  $U_k(1)$ , but not of one by Theorem 6.1; for  $m \geq 2$  Theorem 6.5 applies. Finally  $G_1^\uparrow = x^{-1}u/(1-u)$ , and

$$G_2^\uparrow = x^2 \left( \frac{1}{b_1 b_2} + \frac{2}{b_2} \right) = \frac{x^{-4}u^2}{(1-u)(1-2u)} + \frac{2x^{-1}u}{1-2u}$$

is a sum of two families, but not of one by Theorem 6.2; for  $m \geq 3$  Theorem 6.5 applies.  $\square$

## 7. THE NEAR-DIAGONAL OF THE TRIANGLE

For  $d \in \mathbb{Z}$  put  $D(k, d) = N(k, k-d)$ , so that  $D(k, d) = 0$  for  $d < 0$ . Proposition 2.7 with  $q = k-d$  reads

$$(8) \quad D(k, d) = D(k-1, d) + D(k-1, d-1) + (k-d-1)[D(k-3, d-1) + 2D(k-3, d-2) + D(k-3, d-3)]$$

for  $k \geq 3$  and  $k-d \geq 1$ .

**Theorem 7.1.** *For every  $d \geq 0$  there is a unique polynomial  $p_d \in \mathbb{Q}[k]$  with  $D(k, d) = p_d(k)$  for all  $k \geq 2d + 2$ . It has degree  $2d$ , and*

$$p_d(k) = \frac{2}{2^d d!} \left( k^{2d} - (4d^2 - 3d) k^{2d-1} + O(k^{2d-2}) \right);$$

*in particular its leading coefficient is twice that of  $\{k_{-d}\}$ . The defect  $e_d = D(\cdot, d) - p_d$  satisfies*

$$e_d(2d+1) = (-1)^{d+1} (d+1)!, \quad e_d(2d) = (-1)^{d+1} \frac{(d+2)!}{2},$$

*so the threshold  $2d+2$  is sharp for every  $d$ .*

*Proof.* We argue by induction on  $d$ , with  $p_{d'} = 0$  and  $e_{d'} = 0$  for  $d' < 0$ . For  $d = 0$ , (8) reads  $D(k, 0) = D(k-1, 0)$  for  $k \geq 3$ , and  $D(2, 0) = N(2, 2) = 2$ ; so  $p_0 = 2$ ,  $e_0(1) = D(1, 0) - 2 = -1$  and  $e_0(0) = D(0, 0) - 2 = -1$ , as claimed.

Let  $d \geq 1$  and put

$$R_d(k) = p_{d-1}(k-1) + (k-d-1)[p_{d-1}(k-3) + 2p_{d-2}(k-3) + p_{d-3}(k-3)].$$

For  $k \geq 2d+3$  we have  $k-1 \geq 2(d-1)+2$  and  $k-3 \geq 2d \geq 2(d-i)+2$  for  $i = 1, 2, 3$ , so by induction all terms on the right of (8) except  $D(k-1, d)$  are given by the polynomials, and  $D(k, d) - D(k-1, d) = R_d(k)$ . Let  $p_d$  be the polynomial with  $p_d(k) = D(2d+2, d) + \sum_{i=2d+3}^k R_d(i)$  for integers  $k \geq 2d+2$ . Then  $D(k, d) = p_d(k)$  for  $k \geq 2d+2$ , the polynomial is unique, and  $p_d(k) - p_d(k-1) = R_d(k)$  holds identically.

*Leading coefficients.* The polynomial  $R_d$  has degree  $2d-1$ , and its top term comes only from  $(k-d-1)p_{d-1}(k-3)$ ; the other terms have degree at most  $2d-2$ . Hence  $\deg p_d = 2d$  and  $2d \operatorname{lc}(p_d) = \operatorname{lc}(p_{d-1})$ , which gives  $\operatorname{lc}(p_d) = 2/(2^d d!)$ .

*Second coefficient.* Write  $p_d = c_d k^{2d} + c'_d k^{2d-1} + \dots$  and  $r_d = c'_d / c_d$ . The coefficient of  $k^{2d-2}$  in  $p_d(k) - p_d(k-1)$  is  $-\binom{2d}{2} c_d + (2d-1)c'_d$ . In  $R_d(k)$  only  $(k-d-1)p_{d-1}(k-3)$  and  $p_{d-1}(k-1)$  contribute to  $k^{2d-2}$ , namely  $c'_{d-1} - (7d-5)c_{d-1}$  and  $c_{d-1}$ . Using  $c_{d-1} = 2d c_d$  this gives  $(2d-1)r_d = 2d r_{d-1} - 12d^2 + 11d$ , and  $r_d = -(4d^2 - 3d)$  follows by induction from  $r_1 = -1$  (indeed  $p_1 = k^2 - k - 4$ ).

*Defects.* Subtracting  $p_d(k) - p_d(k-1) = R_d(k)$  from (8) gives, wherever (8) applies,

$$e_d(k) - e_d(k-1) = e_{d-1}(k-1) + (k-d-1)[e_{d-1}(k-3) + 2e_{d-2}(k-3) + e_{d-3}(k-3)].$$

At  $k = 2d+2$  we have  $e_d(2d+2) = e_{d-1}(2d+1) = e_{d-2}(2d-1) = e_{d-3}(2d-1) = 0$ , so  $e_d(2d+1) = -(d+1)e_{d-1}(2d-1) = -(d+1)(-1)^d d! = (-1)^{d+1} (d+1)!$ . At  $k = 2d+1$  we have  $e_{d-1}(2d) = e_{d-2}(2d-2) = e_{d-3}(2d-2) = 0$ , so

$$e_d(2d) = e_d(2d+1) - d e_{d-1}(2d-2) = (-1)^{d+1} (d+1)! - d (-1)^d \frac{(d+1)!}{2} = (-1)^{d+1} \frac{(d+2)!}{2}. \quad \square$$

For example  $p_0 = 2$ ,  $p_1 = k^2 - k - 4$  ( $k \geq 4$ ),

$$p_2 = \frac{k^4 - 10k^3 + 43k^2 - 98k + 164}{4} \quad (k \geq 6),$$

$$p_3 = \frac{k^6 - 27k^5 + 331k^4 - 2225k^3 + 8560k^2 - 17392k + 11088}{24} \quad (k \geq 8).$$

Table 4 lists the values below the threshold for  $d \leq 3$ .

**Proposition 7.2.** *For every  $d \geq 0$ , the Newton coefficients  $\Delta^i p_d(2d+2)$ ,  $0 \leq i \leq 2d$ , of  $p_d$  at the threshold are positive integers; the last one is  $2(2d-1)!!$ .*

For example, for  $d = 2$  they are 65, 74, 64, 30, 6, that is,  $p_2(6+n) = 65 + 74\binom{n}{1} + 64\binom{n}{2} + 30\binom{n}{3} + 6\binom{n}{4}$ .

| $d$ | $k$ | $D(k, d)$ | $p_d(k)$ | $e_d(k)$ |
|-----|-----|-----------|----------|----------|
| 0   | 1   | 1         | 2        | -1       |
| 1   | 2   | 1         | -2       | 3        |
| 1   | 3   | 4         | 2        | 2        |
| 2   | 3   | 1         | 17       | -16      |
| 2   | 4   | 7         | 19       | -12      |
| 2   | 5   | 25        | 31       | -6       |
| 3   | 4   | 1         | -114     | 115      |
| 3   | 5   | 12        | -78      | 90       |
| 3   | 6   | 59        | -1       | 60       |
| 3   | 7   | 199       | 175      | 24       |

TABLE 4. The values  $D(k, d) = N(k, k-d)$  below the threshold,  $d+1 \leq k \leq 2d+1$ , compared with  $p_d(k)$ .

*Proof.* They are integers because  $p_d$  takes integer values at integers, and the last one is  $(2d)! \text{lc}(p_d) = 2(2d-1)!!$ . For the positivity we use two facts about the expansion  $f(n) = \sum_l f_l \binom{n}{l}$ . First, if all  $f_l \geq 0$ , then the coefficients of  $f(n+s)$  for  $s \geq 0$  are  $\sum_{i \geq l} f_i \binom{s}{i-l} \geq f_l$ , by  $\binom{n+s}{i} = \sum_l \binom{s}{i-l} \binom{n}{l}$ . Second,  $(\alpha+n)f(n)$  has the coefficients  $(\alpha+l)f_l + lf_{l-1}$ , because  $n \binom{n}{l} = (l+1) \binom{n}{l+1} + l \binom{n}{l}$ . In the polynomial range, (8) gives, with  $k = 2d+3+n$ ,

$$\Delta p_d(2d+2+n) = p_{d-1}(2d+2+n) + (d+2+n)[p_{d-1}(2d+n) + 2p_{d-2}(2d+n) + p_{d-3}(2d+n)].$$

By induction, the coefficients of  $p_{d-1}(2d+n)$  in this basis are positive up to index  $2d-2$ , and those of the other polynomials in the bracket, taken at points at or beyond their thresholds, are nonnegative. By the two facts, the coefficients of the right-hand side are positive up to index  $2d-1$ . These are the numbers  $\Delta^i p_d(2d+2)$ ,  $1 \leq i \leq 2d$ . Finally  $\Delta^0 p_d(2d+2) = D(2d+2, d) > 0$ , since the word  $(d+2, d+1, \dots, 2, 1, 1, \dots, 1)$  of length  $2d+2$  is good and uses all letters  $1, \dots, d+2$ .  $\square$

Two natural strengthenings fail, which shows that positivity in small cases should not be extrapolated. Written in the variable  $k-2d-1$ , the coefficients of  $p_d$  are positive for  $d \leq 46$ , but each  $p_d$  with  $47 \leq d \leq 57$  has a negative coefficient. The Newton coefficients at the point  $2d+1$  are not all positive either:  $p_d(2d+1) = D(2d+1, d) - (-1)^{d+1}(d+1)!$  is negative for every odd  $d$  with  $69 \leq d \leq 101$ . Both statements were obtained by exact computation.

## 8. REAL ROOTS OF THE ROWS OF THE TRIANGLE

For  $k \geq 1$  let  $n_k(z) = \sum_{q=1}^k N(k, q) z^{q-1}$  be the row polynomial of the triangle  $N$ , and define polynomials  $h_k$  ( $k \geq 0$ ) by

$$\sum_{m \geq 0} U_k(m) t^m = \frac{h_k(t)}{(1-t)^{k+1}}$$

(they are not to be confused with the entries  $h_j$  of a sequence). Since  $\sum_{m \geq 0} \binom{m+1}{q} t^m = t^{q-1}/(1-t)^{q+1}$  for  $q \geq 1$  and  $N(k, 0) = 0$  for  $k \geq 1$ , Proposition 2.6 gives, for  $k \geq 1$ ,

$$(9) \quad h_k(t) = \sum_{q=1}^k N(k, q) t^{q-1} (1-t)^{k-q}, \quad n_k(z) = (1+z)^{k-1} h_k\left(\frac{z}{1+z}\right),$$

while  $h_0 = 1$ . The only good words with values in  $\{1\}$  are constant, so  $N(k, 1) = 1$ ; together with Corollary 2.8 this shows that  $n_k$  has degree  $k-1$  and  $n_k(0) = 1$ . For example  $n_2 = 1+2z$ ,  $n_3 = 1+4z+2z^2$ ,  $h_3 = 1+2t-t^2$  and  $h_4 = 1+4t-3t^2$ ; the coefficients of  $h_k$  need not be nonnegative.

**Theorem 8.1.** *For every  $k \geq 1$  all zeros of  $n_k$  are real, and all of them except possibly  $-1$  are simple. Consequently, for every  $k \geq 0$  all zeros of  $h_k$  are real and simple.*

**Corollary 8.2.** *For every  $k \geq 1$  the numbers  $N(k, 1), \dots, N(k, k)$  are positive and satisfy  $N(k, q)^2 > N(k, q-1)N(k, q+1)$  for  $2 \leq q \leq k-1$ . In particular the row  $N(k, 1), \dots, N(k, k)$  is unimodal.*

*Proof.* A real zero of a nonzero polynomial with nonnegative coefficients is  $\leq 0$ . By Theorem 8.1 and  $n_k(0) = 1$ ,  $n_k$  is therefore a positive multiple of a product of factors  $z + r$  with  $r > 0$ , and all its coefficients  $N(k, 1), \dots, N(k, k)$  are positive. By Newton's inequalities (see [26]), the coefficients of a polynomial  $\sum_{j=0}^d a_j z^j$  with only real zeros satisfy  $a_j^2 \geq a_{j-1}a_{j+1}(1 + 1/j)(1 + 1/(d-j))$  for  $1 \leq j \leq d-1$ . With  $a_j = N(k, j+1) > 0$  and  $d = k-1$  this gives the strict inequalities, and a positive log-concave sequence is unimodal.  $\square$

In the variable  $t$  the zeros of consecutive  $h_k$  do not interlace. For example, the zeros of  $h_3$  are  $1 \pm \sqrt{2}$  and those of  $h_4$  are  $(2 \pm \sqrt{7})/3$ , which lie on the real line in the order  $h_3, h_4, h_4, h_3$ . The proof below works with  $n_k$  instead. The substitution  $t = z/(1+z)$  maps  $(-1, 0)$  onto  $(-\infty, 0)$  and  $(-\infty, -1)$  onto  $(1, \infty)$ , and sends  $z = -1$  to  $t = \infty$ ; the zeros  $-1 \pm \sqrt{2}/2$  of  $n_3$  and the zeros  $-1, (-3 \pm \sqrt{7})/2$  of  $n_4 = (1+z)(1+6z+2z^2)$  do interlace.

**8.1. Interlacing.** All polynomials in this subsection are real and nonzero. A polynomial is *real-rooted* if all its zeros are real; nonzero constants are real-rooted.

**Definition 8.3.** Let  $f$  and  $g$  be real-rooted, with zeros  $a_1 \leq \dots \leq a_n$  and  $b_1 \leq \dots \leq b_{n'}$ , listed with multiplicity. We write  $g \ll f$  if either  $n' = n$  and  $b_1 \leq a_1 \leq b_2 \leq a_2 \leq \dots \leq b_n \leq a_n$ , or  $n' = n-1$  and  $a_1 \leq b_1 \leq a_2 \leq \dots \leq b_{n-1} \leq a_n$ .

Thus the zeros alternate, and the largest one is a zero of  $f$ . Equivalently,  $g \ll f$  if and only if for every real  $x$  the number of zeros of  $f$  in  $(x, \infty)$  exceeds the number of zeros of  $g$  in  $(x, \infty)$  by 0 or 1, zeros counted with multiplicity. Indeed, these counts can be read off from the chain of inequalities, and if a link  $p \leq q$  of the chain fails, then  $x = q$  violates the condition. Comparing  $x = \alpha$  with  $x$  slightly smaller than  $\alpha$ , we see that if  $g \ll f$  and  $\alpha$  is a simple zero of  $f$  with  $g(\alpha) \neq 0$ , then  $f$  and  $g$  have equally many zeros larger than  $\alpha$ ; so if moreover both leading coefficients are positive,  $g(\alpha)$  and  $f'(\alpha)$  have the same sign. Let  $H = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ . The following facts are standard (see [10, 3], whose conventions differ slightly from ours); we include proofs to keep the argument self-contained.

**Lemma 8.4.** *If  $p$  is real-rooted, then  $g \ll f$  if and only if  $pg \ll pf$ .*

*Proof.* It suffices to treat  $p = z - r$ . Multiplying  $f$  and  $g$  by  $z - r$  adds 1 to both counts in the characterization above when  $x < r$  and changes neither count when  $x \geq r$ , so the condition holds for  $pf, pg$  if and only if it holds for  $f, g$ .  $\square$

**Lemma 8.5.** *Let  $f$  be real-rooted, and let  $\operatorname{lc}(f) > 0$  and  $\operatorname{lc}(g) > 0$ .*

(1) *If  $\deg g \leq \deg f$ , then  $g \ll f$  if and only if  $\operatorname{Im}(g(z)/f(z)) \leq 0$  for all  $z \in H$ .*

(2) *If  $\deg f \leq \deg g \leq \deg f + 1$ , then  $f \ll g$  if and only if  $\operatorname{Im}(g(z)/f(z)) \geq 0$  for all  $z \in H$ .*

*In both cases the condition on  $H$  implies that  $g$  is real-rooted.*

*Proof.* Dividing  $f$  and  $g$  by their greatest common divisor, normalized to have positive leading coefficient, changes neither  $g/f$  nor, by Lemma 8.4, the relations  $g \ll f$  and  $f \ll g$ ; the divisor is real-rooted because it divides  $f$ . So we may assume that  $f$  and  $g$  are coprime. Let  $n = \deg f$ .

(1) Suppose  $g \ll f$ . A multiple zero of  $f$  would also be a zero of  $g$ , so the zeros  $\alpha_1 < \dots < \alpha_n$  of  $f$  are simple, and by the remark after the definition  $g(\alpha_i)/f'(\alpha_i) > 0$ . Hence

$$(10) \quad \frac{g}{f} = c + \sum_{i=1}^n \frac{c_i}{z - \alpha_i}, \quad c \geq 0, \quad c_i = \frac{g(\alpha_i)}{f'(\alpha_i)} > 0,$$

where  $c = \text{lc}(g)/\text{lc}(f)$  if  $\deg g = n$  and  $c = 0$  otherwise, and  $\text{Im}(g/f) = -\text{Im } z \sum_i c_i |z - \alpha_i|^{-2} \leq 0$  on  $H$ .

Conversely, suppose that  $\text{Im}(g/f) \leq 0$  on  $H$ . Let  $\alpha$  be a zero of  $f$  of multiplicity  $L$ . Since  $g(\alpha) \neq 0$ , near  $\alpha$  we have  $g/f = \gamma(z - \alpha)^{-L}(1 + O(z - \alpha))$  with real  $\gamma \neq 0$ , and for  $z = \alpha + \epsilon e^{i\theta}$  the imaginary part is  $-\gamma\epsilon^{-L} \sin(L\theta) + O(\epsilon^{1-L})$ . If  $L \geq 2$ , then  $\sin(L\theta)$  takes both signs on  $(0, \pi)$ , which is impossible; so  $L = 1$ , and then  $\gamma > 0$ . Hence (10) holds with  $c \geq 0$  and  $c_i > 0$ . On each interval  $(\alpha_i, \alpha_{i+1})$  the function  $g/f$  decreases from  $+\infty$  to  $-\infty$ , so it has a zero there; if  $c > 0$ , it decreases from  $c$  to  $-\infty$  on  $(-\infty, \alpha_1)$  and has one more zero there; and on  $(\alpha_n, \infty)$  it decreases from  $+\infty$  to  $c \geq 0$  and has no zero. This gives  $n - 1 + [c > 0]$  real zeros of  $g$ . Since  $c > 0$  exactly when  $\deg g = n$ , and otherwise  $\deg g \geq n - 1$  by this count, these are all the zeros of  $g$ , and they interlace as required.

(2) If  $f \ll g$ , then (1) with  $f$  and  $g$  exchanged gives  $\text{Im}(f/g) \leq 0$  on  $H$ , and  $\text{Im}(g/f) = -|g/f|^2 \text{Im}(f/g)$ . Conversely, if  $\text{Im}(g/f) \geq 0$  on  $H$ , the argument of (1) shows that the zeros  $\alpha_1 < \dots < \alpha_n$  of  $f$  are simple and that  $g/f = az + b + \sum_i e_i/(z - \alpha_i)$  with  $e_i < 0$ , where  $a = \text{lc}(g)/\text{lc}(f)$  if  $\deg g = n + 1$ , and  $a = 0$  and  $b = \text{lc}(g)/\text{lc}(f) > 0$  if  $\deg g = n$ . Now  $g/f$  increases from  $-\infty$  to  $+\infty$  on each  $(\alpha_i, \alpha_{i+1})$ ; on  $(\alpha_n, \infty)$  it increases from  $-\infty$  to  $+\infty$  if  $a > 0$ , and to  $b > 0$  if  $a = 0$ ; and if  $a > 0$ , it increases from  $-\infty$  to  $+\infty$  on  $(-\infty, \alpha_1)$ . This gives  $\deg g$  real zeros of  $g$ , and they interlace as required.  $\square$

**Lemma 8.6.** *Let  $f, g, h$  have positive leading coefficients, and let  $f$  be real-rooted. If  $g \ll f$  and  $h \ll f$ , then  $g + h \ll f$ . If  $f \ll g$  and  $f \ll h$ , then  $f \ll g + h$ .*

*Proof.* The imaginary parts add, and  $\deg(g + h) = \max(\deg g, \deg h)$  satisfies the degree condition of Lemma 8.5.  $\square$

In the rest of this section all polynomials have nonnegative coefficients, so the zeros of real-rooted ones lie in  $(-\infty, 0]$ .

**Lemma 8.7.** *Let  $f$  and  $g$  be real-rooted.*

- (1)  $f \ll (1 + z)f$ , and  $f' \ll f$  if  $\deg f \geq 1$ .
- (2) If  $g \ll f$ , then  $f \ll zg$ .
- (3) If  $g \ll f$  and  $\deg g \geq 1$ , then  $g' \ll f'$ .

*Proof.* (1) We have  $\text{Im}((1 + z)f/f) = \text{Im } z > 0$  on  $H$ , and  $f'/f = \sum_j \mu_j/(z - \alpha_j)$ , the sum over the distinct zeros  $\alpha_j$  with multiplicities  $\mu_j$ , has negative imaginary part on  $H$ . Apply Lemma 8.5.

(2) After removing the common factor, (10) gives  $g/f = c + \sum_i c_i/(z - \alpha_i)$  with  $c, c_i \geq 0$  and  $\alpha_i \leq 0$ . Since  $zc_i/(z - \alpha_i) = c_i + c_i\alpha_i/(z - \alpha_i)$ , we get  $\text{Im}(zg/f) = \text{Im } z (c - \sum_i c_i\alpha_i|z - \alpha_i|^{-2}) \geq 0$  on  $H$ . As  $\deg zg \in \{\deg f, \deg f + 1\}$ , Lemma 8.5(2) applies.

(3) Let  $w \in H$ . If  $f(z_0) + wg(z_0) = 0$  for some  $z_0 \in H$ , then  $f(z_0) \neq 0$  and  $g(z_0)/f(z_0) = -1/w \in H$ , which contradicts Lemma 8.5(1). As  $w \notin \mathbb{R}$ , the polynomial  $f + wg$  has degree  $\deg f$ , so all its zeros lie in the closed lower half-plane, and by the Gauss–Lucas theorem so do those of  $f' + wg'$ . The polynomial  $f'$  is real-rooted by Rolle's theorem, so  $f'(z_0) \neq 0$  and  $g'(z_0)/f'(z_0) \neq -1/w$  for  $z_0 \in H$ . Since  $-1/w$  runs through all of  $H$ , we get  $\text{Im}(g'/f') \leq 0$  on  $H$ , and Lemma 8.5(1) gives  $g' \ll f'$ .  $\square$

For a polynomial  $p$  put

$$\mathcal{D}p = (1 + z)p' + 2p = \frac{((1 + z)^2 p)'}{1 + z}, \quad \mathcal{T}p = z\mathcal{D}p, \quad \Psi p = (1 + z)\mathcal{T}p.$$

**Lemma 8.8.** *Let  $p, f, g$  be real-rooted.*

- (1)  $\mathcal{D}p, \mathcal{T}p$  and  $\Psi p$  are real-rooted, of degrees  $\deg p, \deg p + 1$  and  $\deg p + 2$ .
- (2) If  $g \ll f$ , then  $\mathcal{T}g \ll \mathcal{T}f$  and  $\Psi g \ll \Psi f$ .
- (3)  $(1 + z)p \ll \mathcal{T}p$ .

$$(4) \quad \mathcal{T}((1+z)p) = (1+z)(\mathcal{T}p + zp).$$

*Proof.* (1) The polynomial  $((1+z)^2p)'$  is real-rooted by Rolle's theorem and divisible by  $1+z$ , so  $\mathcal{D}p$  is real-rooted; the rest is clear. (2) By Lemmas 8.4 and 8.7(3),  $g \ll f$  implies  $(1+z)^2g \ll (1+z)^2f$ , then  $(1+z)\mathcal{D}g \ll (1+z)\mathcal{D}f$ , and then  $\mathcal{D}g \ll \mathcal{D}f$ ; multiplying by  $z$  and by  $z(1+z)$  gives the claim. (3) With  $Q = (1+z)^2p$ , Lemma 8.7(1) gives  $Q' \ll Q$ , that is,  $(1+z)\mathcal{D}p \ll (1+z) \cdot (1+z)p$ ; so  $\mathcal{D}p \ll (1+z)p$  by Lemma 8.4, and  $(1+z)p \ll z\mathcal{D}p$  by Lemma 8.7(2). (4) is a direct computation.  $\square$

## 8.2. The proof.

**Lemma 8.9.** *For  $k \geq 4$ ,  $n_k = (1+z)n_{k-1} + \Psi n_{k-3} = (1+z)(n_{k-1} + \mathcal{T}n_{k-3})$ .*

*Proof.* Multiply the recurrence of Proposition 2.7 by  $z^{q-1}$  and sum over  $q \geq 1$ . Since  $N(k-1, 0) = N(k-3, 0) = 0$ , the first two terms give  $zn_{k-1} + n_{k-1}$ . With  $p = n_{k-3}$  and  $zp' = \sum_q (q-1)N(k-3, q)z^{q-1}$ , the three terms in the bracket give  $z^2(zp' + 2p)$ ,  $2z(zp' + p)$  and  $zp'$ , whose sum is  $z(1+z)^2p' + 2z(1+z)p = \Psi p$ .  $\square$

For  $k = 3$  one has  $n_3 = (1+z)n_2 + z$  instead.

**Proposition 8.10.** *For every  $k \geq 3$ ,*

$$(\alpha_k) \quad n_{k-1} \ll n_k, \quad (\beta_k) \quad n_k \ll \mathcal{T}n_{k-1}, \quad (\delta_k) \quad n_k \ll \Psi n_{k-2}, \quad (\varepsilon_k) \quad \Psi n_{k-2} \ll \mathcal{T}n_k,$$

*and  $(\alpha_2)$  and  $(\beta_2)$  hold as well. In particular every  $n_k$  is real-rooted.*

*Proof.* The polynomials needed for  $k \leq 3$  are  $n_1 = 1$ ,  $n_2 = 1 + 2z$ ,  $n_3 = 1 + 4z + 2z^2$ ,  $\mathcal{T}n_1 = 2z$ ,  $\mathcal{T}n_2 = 2z(2+3z)$ ,  $\Psi n_1 = 2z(1+z)$  and  $\mathcal{T}n_3 = 2z(2z+1)(2z+3)$ , with the zeros  $-1/2$ ;  $-1 \pm \sqrt{2}/2$ ;  $0$ ;  $-2/3$ ;  $0$ ;  $-1$ ;  $0$ ; and  $-3/2$ ,  $-1/2$ ,  $0$ . Since  $-1 - \sqrt{2}/2 < -3/2 < -1 < -2/3 < -1/2 < -1 + \sqrt{2}/2 < 0$ , the relations  $(\alpha_2)$ ,  $(\beta_2)$ ,  $(\alpha_3)$ ,  $(\beta_3)$ ,  $(\delta_3)$  and  $(\varepsilon_3)$  hold.

Let  $k \geq 3$ , and assume  $(\alpha_{k-1})$ ,  $(\beta_{k-1})$ ,  $(\alpha_k)$ ,  $(\beta_k)$ ,  $(\delta_k)$  and  $(\varepsilon_k)$ . Put  $A = n_k$ ,  $B = n_{k-1}$  and  $C = n_{k-2}$ . They are real-rooted, so Lemma 8.8 applies to them, and  $n_{k+1} = (1+z)A + \Psi C = (1+z)(A + \mathcal{T}C)$  by Lemma 8.9. The degrees below always satisfy the conditions of Lemma 8.6.

$(\alpha_{k+1})$   $A \ll (1+z)A$  by Lemma 8.7(1), and  $A \ll \Psi C$  by  $(\delta_k)$ ; add.

$(\beta_{k+1})$   $(1+z)A \ll \mathcal{T}A$  by Lemma 8.8(3), and  $\Psi C \ll \mathcal{T}A$  by  $(\varepsilon_k)$ ; add.

$(\delta_{k+1})$  By Lemma 8.4 it suffices to show  $A + \mathcal{T}C \ll \mathcal{T}B$ . Here  $A \ll \mathcal{T}B$  is  $(\beta_k)$ , and  $\mathcal{T}C \ll \mathcal{T}B$  follows from  $(\alpha_{k-1})$  by Lemma 8.8(2); add.

$(\varepsilon_{k+1})$  By Lemma 8.8(4),

$$\mathcal{T}n_{k+1} = \Psi A + z(1+z)A + (1+z)\mathcal{T}^2C + z(1+z)\mathcal{T}C.$$

First,  $\Psi B \ll \Psi A$  by  $(\alpha_k)$  and Lemma 8.8(2). For the other three terms we cancel the factor  $1+z$  (Lemma 8.4):  $\mathcal{T}B \ll zA$  follows from  $(\beta_k)$  by Lemma 8.7(2);  $\mathcal{T}B \ll \mathcal{T}(\mathcal{T}C)$  follows from  $(\beta_{k-1})$  by Lemma 8.8(2); and  $\mathcal{T}B \ll z\mathcal{T}C$  follows from  $\mathcal{T}C \ll \mathcal{T}B$  by Lemma 8.7(2).

So  $\Psi B$  is related by  $\ll$  to each of the four terms; add.

This proves the four relations for  $k+1$ , and the induction continues.  $\square$

**Proposition 8.11.** *For every  $j \geq 2$  the polynomials  $n_j$  and  $n_{j-1}$  have no common zero other than  $-1$ . Hence every zero of  $n_k$  other than  $-1$  is simple.*

*Proof.* Suppose the first claim fails, and let  $j$  be minimal such that  $n_j$  and  $n_{j-1}$  have a common zero  $\alpha \neq -1$ . Then  $\alpha \neq 0$ , since  $n_i(0) = 1$ . The cases  $j \leq 4$  are excluded directly:  $n_1 = 1$ ;  $n_3(-1/2) = -1/2$ ; and  $n_4 = (1+z)(n_3 + 2z)$  does not vanish at the zeros of  $n_3$ . So  $j \geq 5$ , and Lemma 8.9 for  $n_j$  and for  $n_{j-1}$  gives

$$\mathcal{T}n_{j-3}(\alpha) = 0, \quad n_{j-2}(\alpha) + \mathcal{T}n_{j-4}(\alpha) = 0.$$

Let  $f = \mathcal{T}n_{j-3}$ . Then  $n_{j-2} \ll f$  by  $(\beta_{j-2})$ , and  $\mathcal{T}n_{j-4} \ll f$  by  $(\alpha_{j-3})$  and Lemma 8.8(2).

If  $\alpha$  is a simple zero of  $f$ , the remark after the definition of  $\ll$  gives  $n_{j-2}(\alpha)f'(\alpha) \geq 0$  and  $\mathcal{T}n_{j-4}(\alpha)f'(\alpha) \geq 0$ . The two values add up to 0, so  $n_{j-2}(\alpha) = 0$ , which contradicts the minimality of  $j$ .

If  $\alpha$  is a multiple zero of  $f = z\mathcal{D}n_{j-3}$ , then, as  $\alpha \notin \{0, -1\}$ , it is a multiple zero of  $Q' = (1+z)\mathcal{D}n_{j-3}$ , where  $Q = (1+z)^2n_{j-3}$ . The zeros of  $Q'$  that are not zeros of the real-rooted polynomial  $Q$  are simple, because  $Q'/Q$  is strictly decreasing between consecutive zeros of  $Q$ . So  $\alpha$  is a zero of  $Q$  of multiplicity at least 3, hence a multiple zero of  $n_{j-3}$ , and  $(\alpha_{j-3})$  forces  $n_{j-4}(\alpha) = 0$ , again contradicting the minimality of  $j$ .

Finally, if  $\alpha \neq -1$  were a multiple zero of  $n_k$  with  $k \geq 2$ , then  $(\alpha_k)$  would force  $n_{k-1}(\alpha) = 0$ .  $\square$

*Proof of Theorem 8.1.* For  $k \leq 1$  there is nothing to prove, since  $n_1 = h_0 = h_1 = 1$ . For  $k \geq 2$ , Propositions 8.10 and 8.11 give  $n_k = 2(1+z)^a \prod_{i=1}^r (z - z_i)$  with  $a + r = k - 1$  and distinct real numbers  $z_i \neq -1$ ; the leading coefficient is  $N(k, k) = 2$ . By (9),

$$h_k(t) = (1-t)^{k-1} n_k\left(\frac{t}{1-t}\right) = 2 \prod_{i=1}^r ((1+z_i)t - z_i),$$

whose zeros  $z_i/(1+z_i)$  are real and distinct.  $\square$

For the location of the zeros of  $h_k$  we need the multiplicity of the zero  $-1$  of  $n_k$ .

**Lemma 8.12.** *For  $k \geq 1$  write  $n_k = (1+z)^{a_k} m_k$  with  $m_k(-1) \neq 0$ , and put  $\lambda_k = m_k(-1)$ . Then  $a_k = \lceil k/3 \rceil - 1$ , and  $\lambda_k$  has the sign  $(-1)^{\lfloor k/3 \rfloor}$  if  $k \not\equiv 2 \pmod{3}$  and the sign  $(-1)^{\lfloor k/3 \rfloor + 1}$  if  $k \equiv 2 \pmod{3}$ . Consequently  $\deg h_k = \lfloor 2k/3 \rfloor$ , and the leading coefficient of  $h_k$  has the sign  $(-1)^{\lfloor k/3 \rfloor}$ .*

The statements about  $a_k$ ,  $\deg h_k$  and the sign of the leading coefficient of  $h_k$  are also formalized in Lean, with a different proof (`rootMultiplicity_nrowPoly`, `natDegree_hpoly`, `leadingCoeff_hpoly_sign`).

*Proof.* For  $p = (1+z)^a m$  we have  $\mathcal{T}p = z(1+z)^a((a+2)m + (1+z)m')$ . The claims hold for  $k \leq 3$ , where  $(a_k, \lambda_k) = (0, 1), (0, -1), (0, -1)$ . Let  $k \geq 4$ , assume the claims for smaller indices, and put  $a = a_{k-3}$ . Then  $n_{k-1} + \mathcal{T}n_{k-3} = (1+z)^a B$  with a polynomial  $B$ , and  $n_k = (1+z)^{a+1} B$  by Lemma 8.9. If  $k \not\equiv 1 \pmod{3}$ , then  $a_{k-1} = a + 1$ , so that  $B(-1) = -(a+2)\lambda_{k-3} \neq 0$ . If  $k = 3b + 1$ , then  $a_{k-1} = a$  and  $B(-1) = \lambda_{k-1} - (a+2)\lambda_{k-3}$ , where  $\lambda_{k-1} = \lambda_{3b}$  and  $-\lambda_{k-3} = -\lambda_{3b-2}$  both have the sign  $(-1)^b$ , so again  $B(-1) \neq 0$ . In both cases  $a_k = a + 1$  and  $\lambda_k = B(-1)$ , and the sign of  $\lambda_k$  is as claimed.

Let  $r = k - 1 - a_k = \lfloor 2k/3 \rfloor$ , and write  $m_k = \sum_j \mu_j z^j$ , a polynomial of degree  $r$ . By (9),  $h_k(t) = (1-t)^r m_k(t/(1-t)) = \sum_j \mu_j t^j (1-t)^{r-j}$ , whose coefficient of  $t^r$  is  $\sum_j \mu_j (-1)^{r-j} = (-1)^r \lambda_k \neq 0$ . So  $\deg h_k = r$ , and the three residues of  $k$  modulo 3 give the sign of the leading coefficient.  $\square$

**Corollary 8.13.** *For  $k \geq 1$ , exactly  $\lfloor k/3 \rfloor$  zeros of  $h_k$  lie in  $(1, \infty)$ , the other  $\lfloor (k+1)/3 \rfloor$  zeros are negative, and the coefficient sequence of  $h_k$  has exactly  $\lfloor k/3 \rfloor$  changes of sign.*

*Proof.* By the proof of Theorem 8.1, the zeros of  $h_k$  are the numbers  $z_i/(1+z_i)$ , where  $z_i$  runs over the zeros of  $n_k$  other than  $-1$ . The zeros  $z_i < -1$  give zeros in  $(1, \infty)$ , and the zeros  $z_i \in (-1, 0)$  give negative zeros. Let  $\nu_k$  be the number of zeros of  $n_k$  in  $(-\infty, -1)$ . Since  $\deg n_k = \deg n_{k-1} + 1$ , the relation  $(\alpha_k)$  and the characterization of  $\ll$  show that the number of zeros of  $n_k$  in  $(-\infty, -1)$  exceeds that of  $n_{k-1}$  by 0 or 1, and likewise for  $(-\infty, -1]$ . For  $k \equiv 1 \pmod{3}$  the multiplicity of  $-1$  increases by one (Lemma 8.12), which forces  $\nu_k = \nu_{k-1}$ . The leading coefficient  $2 \prod_i (1+z_i)$  of  $h_k$  has the sign  $(-1)^{\nu_k}$ , so  $\nu_k \equiv \lfloor k/3 \rfloor \pmod{2}$  by Lemma 8.12, which forces  $\nu_k = \nu_{k-1}$  for  $k \equiv 2$  and  $\nu_k = \nu_{k-1} + 1$  for  $k \equiv 0 \pmod{3}$ . With  $\nu_1 = 0$  this gives  $\nu_k = \lfloor k/3 \rfloor$ , and the remaining  $\deg h_k - \nu_k = \lfloor 2k/3 \rfloor - \lfloor k/3 \rfloor = \lfloor (k+1)/3 \rfloor$  zeros are negative. Finally  $h_k(0) = 1$ , and for a polynomial with only real zeros Descartes' rule of signs is exact, so the number of sign changes is the number  $\nu_k$  of positive zeros.  $\square$

**Corollary 8.14.** *Let  $X_k$  be the number of distinct values of a word chosen uniformly at random among the  $n_k(1)$  good words of length  $k$  whose set of values is  $\{1, \dots, q\}$  for some  $q$ . Then  $(X_k - \mathbb{E}X_k)/\sqrt{\text{Var } X_k}$  converges in distribution to the standard normal distribution as  $k \rightarrow \infty$ .*

*Proof.* The probability generating function  $z n_k(z)/n_k(1)$  of  $X_k$  has only real zeros, all of them  $\leq 0$ , so  $X_k - 1$  is a sum of  $k - 1$  independent Bernoulli variables, one for each zero  $-r$  of  $n_k$ , with success probability  $1/(1 + r)$ . The  $a_k = \lceil k/3 \rceil - 1$  variables that belong to the zero  $-1$  have variance  $1/4$  each, so  $\text{Var } X_k \geq (\lceil k/3 \rceil - 1)/4 \rightarrow \infty$ . Since the summands are bounded, the Lindeberg–Feller theorem gives the claim, as in Harper’s argument [12].  $\square$

**Corollary 8.15.** *Let  $k \geq 1$  and let  $u_k$  be the polynomial of Corollary 2.9. Then  $u_k(-j) = 0$  for  $1 \leq j \leq \lceil k/3 \rceil$ , and  $u_k(-\lceil k/3 \rceil - 1) = (-1)^k \text{lc}(h_k) \neq 0$ .*

*Proof.* Let  $s = \deg h_k = \lfloor 2k/3 \rfloor$  (Lemma 8.12), so that  $k - s = \lceil k/3 \rceil$ , and write  $h_k(t) = \sum_{i=0}^s c_i t^i$ . For  $m \geq 0$  we have  $[t^m] t^i (1 - t)^{-k-1} = \binom{m-i+k}{k}$ , where for  $0 \leq m < i$  both sides vanish because  $i \leq k$ . Hence  $u_k(m) = \sum_{i=0}^s c_i \binom{m-i+k}{k}$  as polynomials in  $m$ , with polynomial binomial coefficients. For an integer  $j \geq 1$ ,  $\binom{k-i-j}{k}$  is the product of the  $k$  consecutive integers  $1 - i - j, \dots, k - i - j$  divided by  $k!$ ; it vanishes if  $i + j \leq k$  and equals  $(-1)^k$  if  $i + j = k + 1$ . For  $j \leq k - s$  all terms vanish, and for  $j = k - s + 1$  only the term with  $i = s$  remains, which is  $(-1)^k c_s$ .  $\square$

**Remark 8.16** (Chains and Eulerian polynomials). Let  $k \geq 1$ , let  $\mathbf{0} = (0, \dots, 0)$  and  $\mathbf{1} = (1, \dots, 1)$  be the least and the greatest element of  $\Lambda_k$ , and put  $\bar{\Lambda}_k = \Lambda_k \setminus \{\mathbf{0}, \mathbf{1}\}$ . For  $w \in \{1, \dots, q\}^k$  and  $1 \leq r < q$  let  $y_r = ([w_1 > r], \dots, [w_k > r])$ . By the proof of Proposition 2.2, applied to the heights  $w_j - 1$ , the word  $w$  is good if and only if  $y_1, \dots, y_{q-1} \in \Lambda_k$ ; moreover,  $w$  uses every letter  $1, \dots, q$  if and only if  $\mathbf{1} > y_1 > \dots > y_{q-1} > \mathbf{0}$ , and  $w$  is recovered from the  $y_r$  as  $w_j = 1 + |\{r : (y_r)_j = 1\}|$ . Hence  $N(k, q)$  is the number of chains with  $q - 1$  elements in  $\bar{\Lambda}_k$ , so  $n_k$  is the chain polynomial of  $\bar{\Lambda}_k$ , and by (9) the polynomial  $h_k(t) = (1 - t)^{k-1} n_k(t/(1 - t))$  is the  $h$ -polynomial of the order complex of  $\bar{\Lambda}_k$ , a simplicial complex of dimension  $k - 2$ . For the Boolean lattice  $\{0, 1\}^k$  in place of  $\Lambda_k$  the same constructions give  $(m + 1)^k$ , the numbers  $q! \{q\}^k$  of ordered set partitions, and the Eulerian polynomials  $A_k(t)$ , which satisfy  $\sum_{m \geq 0} (m + 1)^k t^m = A_k(t)/(1 - t)^{k+1}$  [28, Section 1.4]. This is the precise form of the analogy stated in the introduction. In this language, Theorem 8.1 says that the chain polynomial of  $\bar{\Lambda}_k$  is real-rooted, as the Eulerian polynomials are [3]; real-rootedness of chain polynomials has been studied for several other classes of posets [2, 1, 4]. By Corollary 8.13,  $h_k$  has a negative coefficient for  $k \geq 3$ , so the order complex of  $\bar{\Lambda}_k$  is not Cohen–Macaulay for  $k \geq 3$  [27, Chapter II]. If the  $h$ -polynomial  $h$  of an order complex of dimension  $d - 1$  has nonnegative coefficients, as it has for Cohen–Macaulay posets, then the chain polynomial  $f(z) = (1 + z)^d h(z/(1 + z))$  has no zeros in  $(-\infty, -1)$ , since  $h > 0$  on  $(1, \infty)$ ; the general results of [4] on TN-posets and  $P$ -positive posets also place all zeros in  $[-1, 0]$ . By the proof of Corollary 8.13,  $n_k$  has exactly  $\lfloor k/3 \rfloor$  zeros in  $(-\infty, -1)$ , so for  $k \geq 3$  Theorem 8.1 does not follow from results of this kind.

## 9. FORMAL VERIFICATION

The Lean 4 development (Lean v4.34.1, Mathlib for the same version) consists of 27 files that start from the definitions of  $a_k(n)$  (as a number of matrices) and of  $U_k(m)$  (as a number of lists). The main formal statements are listed in Table 5.

The development also formalizes several auxiliary results that are not stated in this paper. The command `#print axioms` reports only `propext`, `Classical.choice` and `Quot.sound`, and an exhaustive scan of all 2675 declarations of the development finds no other axiom.

Not formalized: the block decomposition (Theorem 4.2), the generating function of Theorem 4.4 and the bijection of Proposition 4.7 (the formal proof of Theorem 4.6 goes through the recurrence instead, refined by ascents); in Lemma 4.5, the refinement by  $y$  of the coefficient formula (the

| Result                                             | Lean names                                                                                                                                                                                           |
|----------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Propositions 2.2, 2.4                              | <code>a_eq</code> , <code>U_eq_multichains</code> , <code>card_Lam</code>                                                                                                                            |
| Lemma 2.5                                          | <code>lemma1</code> , <code>lemma1_one</code> , <code>lemma1_two</code> , <code>U_zero_left</code> , <code>U_zero_right</code>                                                                       |
| Propositions 2.6, 2.7                              | <code>U_eq_sum_N</code> , <code>N_inv</code> , <code>N_tri</code> , <code>N_init</code> , <code>N_tri_fails_at_two</code>                                                                            |
| Corollaries 2.8, 2.9                               | <code>N_diag</code> , <code>N_subdiag</code> , <code>existsUnique_upoly</code> , <code>natDegree_upoly</code> ,<br><code>leadingCoeff_upoly_eq</code> , <code>upoly_eval_neg_one</code>              |
| Theorem 3.1                                        | <code>G_rec</code> , <code>P_mul_G</code> , <code>isCoprime_W_P</code> , <code>min_order</code> ,<br><code>order_lower_bound</code> , <code>squarefree_Ppoly</code> , <code>simple_root_Ppoly</code> |
| Theorem 3.2(1), except $ \sigma ^2 < \rho_{m-1}^2$ | <code>existsUnique_rho</code> , <code>one_lt_rho</code> , <code>rho_strictMono</code> ,<br><code>normSq_eq_of_root</code> , <code>norm_lt_rho_of_root</code> , <code>root_P_min</code>               |
| Corollary 3.4                                      | <code>T1_9</code> , <code>parity_min_order</code>                                                                                                                                                    |
| Theorem 4.6, Remark 4.9                            | <code>U_explicit</code> , <code>Us_explicit</code> , <code>U_F3</code> , <code>U_F4</code>                                                                                                           |
| Theorem 5.1                                        | <code>not_isDFinite_FK</code> , <code>no_x_ODE_poly</code> , <code>no_k_only_recurrence</code> ; for<br>$N$ : <code>not_isDFinite_NKser</code> , <code>no_k_only_recurrence_N</code>                 |
| Theorem 5.4                                        | <code>RelU_eq</code> , <code>RelU_iff_mem_span</code> , <code>finrank_relU_tot</code> ,<br><code>finrank_relU_gr</code> ; for $N$ : <code>RelN_eq</code> , <code>finrank_relN_tot</code>             |
| Proposition 5.6                                    | <code>relU_support_det</code> , <code>relU_card_support</code> ,<br><code>not_mem_relU_of_support_subset</code>                                                                                      |
| Theorem 7.1, Proposition 7.2                       | <code>T4_1</code> , <code>T4_1_threshold</code> , <code>T4_2_3</code>                                                                                                                                |

TABLE 5. Formalized results.

case  $y = 1$  is formalized) and the identification of  $H(m, s, j)$  with  $r$ -Stirling numbers for  $j \geq 1$ ; the inequality  $|\sigma|^2 < \rho_{m-1}^2$  in Theorem 3.2(1), Theorem 3.2(2)(3) and Remark 3.3; Remarks 3.5 and 5.2; Corollary 5.5; Corollary 5.7 and Remark 5.8, which lead from Kauers' definition to Proposition 5.6; Section 6; and Section 8, apart from the statements about  $a_k$ ,  $\deg h_k$  and the sign of the leading coefficient of  $h_k$  in Lemma 8.12.

The definitions that carry the meaning of the formal statements are transcribed in Appendix A: good triples and  $U$  (lists of length  $k$  with entries at most  $m$ ); the row and column rules for matrices; D-finiteness (following Lipshitz, with the field  $K(x, t)$  realized inside the fraction field of  $K[[x]][[t]]$ ); the operators  $X$ ,  $E^{-1}$  with the convention that negative indices give 0; and  $\text{Rel}(U)$  as annihilation on some quadrant. The author has compared these definitions with the statements in this paper; the checklist used for the comparison is in the notes folder of the repository.

## 10. OPEN PROBLEMS

- (1) *Shorter formulas.* Find a meaningful class of formulas in which the double sum of Theorem 4.6 is provably shortest.
- (2) *The triangle.* Find a single or short double sum for  $N(k, q)$ .
- (3) *Other zeros.* By Corollary 8.15,  $u_k$  vanishes at  $-1, \dots, -\lceil k/3 \rceil$ ; show that it has no other negative integer zeros. A computer search for  $k \leq 300$  (scripts in the repository) found no further negative integer zeros.
- (4) *Signs of  $h_k$ .* The coefficient sequence of  $h_k$  changes sign exactly  $\lfloor k/3 \rfloor$  times (Corollary 8.13). Describe where the sign changes occur. Computations for  $i \leq 579$  (scripts in the repository) suggest that for each  $i$  the coefficient of  $t^i$  in  $h_k$  is positive for all large  $k$ , and that the least  $k$  from which on it stays positive grows like  $\alpha^* i$  with  $\alpha^* \approx 9.6085$ .

## APPENDIX A. THE LEAN DEFINITIONS

The formal development is in the folder `lean/A207123` of the repository given under *Data and code*. We transcribe the definitions that carry the meaning of the main formal statements of Section 9, from the files `Basic.lean`, `Reduction.lean`, `DFinite.lean`, `NonDFinite.lean`,

`OreRel.lean` and `NotStirlingLike.lean`. Unicode symbols of the source are typeset as mathematical symbols, and comments and proofs are omitted. Auxiliary definitions used in other statements of Table 5 (for instance the Lean versions of  $b_i$ ,  $P_m$ ,  $W_m$  and  $\rho_m$ , and of the spaces in Theorem 5.4(2)) are in the same folder; the script `code/main_extra/lean_statements.py` lists all definitions and statements.

### Good sequences.

```
def Good (a b c : ℕ) : Prop := b = c ∨ (b ≤ a ∧ c ≤ a)

def Legal : List ℕ → Prop
| a :: b :: c :: t => Good a b c ∧ Legal (b :: c :: t)
| _ => True

def seqs (m : ℕ) : ℕ → Finset (List ℕ)
| 0 => {[]}
| k + 1 => (Finset.range (m + 1)).biUnion fun a => (seqs m k).image (List.cons a)

def L (k m : ℕ) : Finset (List ℕ) := (seqs m k).filter Legal

def U (k m : ℕ) : ℕ := (L k m).card

def NW (k q : ℕ) : Finset (List ℕ) := (L k q).filter fun l => l.toFinset = Icc 1 q

def N (k q : ℕ) : ℕ := (NW k q).card
```

The head of a list is  $h_1$ , so  $a$  is the earliest entry of a triple. The lemma `mem_seqs` states that `seqs m k` consists of the lists of length  $k$  with entries at most  $m$ ; `L k m` is the set  $\mathcal{H}_k(m)$ , and `N k q` counts the good lists of length  $k$  whose set of values is  $\{1, \dots, q\}$ .

### The arrays.

```
abbrev Mat (n k : ℕ) := Fin n → Fin k → Bool

def RowRule {n k : ℕ} (M : Mat n k) : Prop :=
  ∀ (i : Fin n) (j : ℕ) (h : j + 2 < k),
    ¬ (M i ⟨j, by omega⟩ = false ∧ M i ⟨j + 1, by omega⟩ = false
      ∧ M i ⟨j + 2, h⟩ = true) ∧
    ¬ (M i ⟨j, by omega⟩ = false ∧ M i ⟨j + 1, by omega⟩ = true
      ∧ M i ⟨j + 2, h⟩ = false)

def ColRule {n k : ℕ} (M : Mat n k) : Prop :=
  ∀ (j : Fin k) (i : ℕ) (h : i + 2 < n),
    ¬ (M ⟨i, by omega⟩ j = false ∧ M ⟨i + 1, by omega⟩ j = false
      ∧ M ⟨i + 2, h⟩ j = true) ∧
    ¬ (M ⟨i, by omega⟩ j = false ∧ M ⟨i + 1, by omega⟩ j = true
      ∧ M ⟨i + 2, h⟩ j = true)

noncomputable def a (k n : ℕ) : ℕ :=
  Fintype.card {M : Mat n k // RowRule M ∧ ColRule M}

theorem a_eq (k n : ℕ) : a k n = U k ((n + 1) / 2) * U k (n / 2)
```

Here `true` stands for 1, `M i j` is the entry in row  $i$  and column  $j$ , and  $(n + 1) / 2$  and  $n / 2$  are  $\lceil n/2 \rceil$  and  $\lfloor n/2 \rfloor$ .

**D-finiteness.**

abbrev PS2 := PowerSeries (PowerSeries K)

noncomputable def dXK (f : PS2 K) : PS2 K :=  
 PowerSeries.mk fun m => PowerSeries.derivative (R := K) (PowerSeries.coeff m f)

noncomputable def dTK (f : PS2 K) : PS2 K :=  
 PowerSeries.derivative (R := PowerSeries K) f

noncomputable def ratFn : Subfield (FractionRing (PS2 K)) :=  
 Subfield.closure  
 (Set.range fun p => algebraMap (PS2 K) (FractionRing (PS2 K)) (polyToPS2 K p))

def IsDFinite (f : PS2 K) : Prop :=  
 FiniteDimensional (ratFn K) (Submodule.span (ratFn K) (Set.range fun ij :  $\mathbb{N} \times \mathbb{N}$  =>  
 algebraMap (PS2 K) (FractionRing (PS2 K)) ((dXK K)^[ij.1] ((dTK K)^[ij.2] f))))

noncomputable def FK : PS2 K :=  
 PowerSeries.mk fun m => PowerSeries.mk fun k => (U k m : K)

theorem not\_isDFinite\_FK {K : Type\*} [Field K] ( $\sigma : K \rightarrow^{++} \mathbb{C}$ ) :  $\neg$  IsDFinite K (FK K)

The outer variable of PS2 K is  $t$  and the inner one is  $x$ ; polyToPS2 is the embedding of  $K[x, t]$ , and the lemmas mem\_ratFn\_iff and isFractionRing\_ratFn identify ratFn K with  $K(x, t)$ . The identification of  $K[[x]][[t]]$  with  $K[[x, t]]$  is standard and not formalized.

**Recurrences whose coefficients depend only on  $k$ .**

theorem no\_k\_only\_recurrence (A B k0 m0 :  $\mathbb{N}$ ) (p :  $\mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{C}[X]$ )  
 (hp :  $\exists a \leq A, \exists b \leq B, p \ a \ b \neq 0$ ) :  
 $\neg \forall k \ m, k0 \leq k \rightarrow m0 \leq m \rightarrow A \leq k \rightarrow B \leq m \rightarrow$   
 $\sum a \in \text{range } (A + 1), \sum b \in \text{range } (B + 1),$   
 $(p \ a \ b).eval \ (k : \mathbb{C}) * (U \ (k - a) \ (m - b) : \mathbb{C}) = 0$

**Quadrant recurrences.**

abbrev Arr :=  $\mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{C}$

abbrev Coef := Polynomial (Polynomial  $\mathbb{C}$ )

theorem mulOp\_apply (p : Coef) (f : Arr) (k m :  $\mathbb{N}$ ) :  
 mulOp p f k m = p.evalEval (k :  $\mathbb{C}$ ) (m :  $\mathbb{C}$ ) \* f k m

theorem opX\_apply (f : Arr) (k m :  $\mathbb{N}$ ) :  
 opX f k m = if k = 0 then 0 else f (k - 1) m

theorem opEinv\_apply (f : Arr) (k m :  $\mathbb{N}$ ) :  
 opEinv f k m = if m = 0 then 0 else f k (m - 1)

noncomputable def OU : Subalgebra  $\mathbb{C}$  (Module.End  $\mathbb{C}$  Arr) :=  
 Algebra.adjoin  $\mathbb{C}$  {mulOp cK, mulOp cM, opX, opEinv}

noncomputable def L1 : Module.End  $\mathbb{C}$  Arr := 1 - opEinv - opX - mulOp cM \* opX ^ 3

noncomputable def Uarr : Arr := fun k m => (U k m :  $\mathbb{C}$ )

def VanishOn (f : Arr) (k1 m1 :  $\mathbb{N}$ ) : Prop :=  $\forall k \ m, k1 \leq k \rightarrow m1 \leq m \rightarrow f \ k \ m = 0$

def RelU : Set (Module.End  $\mathbb{C}$  Arr) :=  
 {R | R  $\in$  OU  $\wedge \exists k0 \ m0 : \mathbb{N}, \text{VanishOn } (R \ Uarr) \ k0 \ m0$ }

`theorem RelU_eq : RelU = {R |  $\exists Q \in \text{OU}, R = Q * L1$ }`

In `Coef` the inner variable is  $k$  and the outer one is  $m$ ; `cK` and `cM` are the coefficients  $k$  and  $m$ . The theorems `exists_normal_form` and `normal_form_unique` show that every element of `OU` has a unique normal form  $\sum r_{ab}(k, m)X^aE^{-b}$ , so that `OU` is the Ore algebra  $\mathcal{O}$  acting faithfully.

### Supports.

```
def det2 (p q r :  $\mathbb{N} \times \mathbb{N}$ ) :  $\mathbb{Z}$  :=
  ((q.1 :  $\mathbb{Z}$ ) - p.1) * ((r.2 :  $\mathbb{Z}$ ) - p.2) - ((q.2 :  $\mathbb{Z}$ ) - p.2) * ((r.1 :  $\mathbb{Z}$ ) - p.1)
```

```
theorem relU_support_det {R : Module.End  $\mathbb{C}$  Arr} (hR : R  $\in$  RelU) (hR0 : R  $\neq$  0)
  {r :  $\mathbb{N} \times \mathbb{N} \rightarrow$  _0 Coef} (hr : R = normOp r) :
   $\exists p0 \in$  r.support,  $\exists p1 \in$  r.support,  $\exists p2 \in$  r.support,  $3 \leq$  det2 p0 p1 p2
```

```
theorem not_mem_relU_of_support_subset {r :  $\mathbb{N} \times \mathbb{N} \rightarrow$  _0 Coef} (hr : r  $\neq$  0)
  {p0 p1 p2 :  $\mathbb{N} \times \mathbb{N}$ } (hs : r.support  $\subseteq$  {p0, p1, p2})
  (hdet : |det2 p0 p1 p2|  $\leq$  2) : normOp r  $\notin$  RelU
```

Here `normOp r` is the operator  $\sum_{(a,b)} r(a,b)X^aE^{-b}$  with finitely supported coefficients.

## STATEMENT ON THE USE OF AI TOOLS

The mathematical exploration, the proofs in this draft, the computer checks and the Lean formalization were produced with substantial assistance from a large language model (Claude, Anthropic) working under the direction of the author. The written proofs were cross-checked by further runs of AI models acting as reviewers, which are not independent of the model that wrote them, and the computational statements by separately written programs, in exact arithmetic except for a minority of checks that use high-precision decimal arithmetic and are labelled as numerical in the output of the verification scripts. The proofs of the results listed in Section 9 are machine-checked; the faithfulness of the formal statements to the statements in this paper was checked by AI models and by the author (the checklist is in the repository). The author takes full responsibility for the content.

## DATA AND CODE

The verification scripts (414 automated checks, mostly exact; the high-precision numerical ones are labelled), a script that recomputes every number printed in this paper from the definitions (apart from the computations quoted in Section 10, which have their own scripts), the Lean development and the notes are available at

<https://github.com/r5x8ksva-del/a207123-explicit-formulas>.

## REFERENCES

- [1] C. A. Athanasiadis, T. Douvropoulos, K. Kalampogia-Evangelinou, Two classes of posets with real-rooted chain polynomials, *Electron. J. Combin.* **31**(4) (2024), #P4.16, <https://doi.org/10.37236/12218>.
- [2] C. A. Athanasiadis, K. Kalampogia-Evangelinou, Chain enumeration, partition lattices and polynomials with only real roots, *Combin. Theory* **3**(1) (2023), <https://doi.org/10.5070/C63160425>.
- [3] P. Brändén, Unimodality, log-concavity, real-rootedness and beyond, in: *Handbook of Enumerative Combinatorics* (M. Bóna, ed.), Chapman and Hall/CRC, Boca Raton, FL, 2015, pp. 461–508, <https://doi.org/10.1201/b18255-13>; arXiv:1410.6601.
- [4] P. Brändén, L. Saud Maia Leite, Totally nonnegative matrices, chain enumeration and zeros of polynomials, *Adv. Math.* **487** (2026), Paper No. 110760, <https://doi.org/10.1016/j.aim.2025.110760>; arXiv:2412.06595.
- [5] A. Z. Broder, The  $r$ -Stirling numbers, *Discrete Math.* **49** (1984), no. 3, 241–259, [https://doi.org/10.1016/0012-365X\(84\)90161-4](https://doi.org/10.1016/0012-365X(84)90161-4).

- [6] F. Chyzak, M. Kauers, B. Salvy, A non-holonomic systems approach to special function identities, in: *Proceedings of the 2009 International Symposium on Symbolic and Algebraic Computation (ISSAC 2009)*, ACM, 2009, pp. 111–118, <https://doi.org/10.1145/1576702.1576720>; arXiv:0904.2761.
- [7] R. Dougherty-Bliss, M. Kauers, Hardinian arrays, *Electron. J. Combin.* **31**(2) (2024), #P2.9, <https://doi.org/10.37236/12358>.
- [8] R. Dougherty-Bliss, C. Koutschan, N. Ter-Saakov, D. Zeilberger, The (symbolic and numeric) computational challenges of counting 0-1 balanced matrices, preprint, arXiv:2410.07435v2, 2025.
- [9] R. Dougherty-Bliss, G. Spahn, Rectangular Hardinian arrays, *ACM Commun. Comput. Algebra* **58** (2024), no. 3, 81–84, <https://doi.org/10.1145/3717582.3717589>.
- [10] S. Fisk, Polynomials, roots, and interlacing, preprint, arXiv:math/0612833, 2006.
- [11] I. Gessel, R. P. Stanley, Stirling polynomials, *J. Combin. Theory Ser. A* **24** (1978), 24–33, [https://doi.org/10.1016/0097-3165\(78\)90042-0](https://doi.org/10.1016/0097-3165(78)90042-0).
- [12] L. H. Harper, Stirling behavior is asymptotically normal, *Ann. Math. Statist.* **38** (1967), 410–414, <https://doi.org/10.1214/aoms/1177698956>.
- [13] M. Kauers, Summation algorithms for Stirling number identities, *J. Symbolic Comput.* **42** (2007), no. 10, 948–970, <https://doi.org/10.1016/j.jsc.2007.08.002>.
- [14] M. Kauers, *D-Finite Functions*, Algorithms and Computation in Mathematics 30, Springer, Cham, 2023, <https://doi.org/10.1007/978-3-031-34652-1>.
- [15] S. Kitaev, *Patterns in Permutations and Words*, Monographs in Theoretical Computer Science, Springer, Heidelberg, 2011, <https://doi.org/10.1007/978-3-642-17333-2>.
- [16] S. Kitaev, T. Mansour, A. Vella, Pattern avoidance in matrices, *J. Integer Seq.* **8** (2005), Article 05.2.2.
- [17] M. Klazar, Bell numbers, their relatives, and algebraic differential equations, *J. Combin. Theory Ser. A* **102** (2003), 63–87, [https://doi.org/10.1016/S0097-3165\(03\)00014-1](https://doi.org/10.1016/S0097-3165(03)00014-1).
- [18] C. Krause, Proof of formula, attachment to OEIS A202093, 26 June 2026, <https://oeis.org/A202093/a202093.txt>, accessed 7 October 2026.
- [19] L. de Moura, S. Ullrich, The Lean 4 theorem prover and programming language, in: *Automated Deduction – CADE 28*, Lecture Notes in Comput. Sci. 12699, Springer, 2021, pp. 625–635, [https://doi.org/10.1007/978-3-030-79876-5\\_37](https://doi.org/10.1007/978-3-030-79876-5_37).
- [20] L. Lipshitz, D-finite power series, *J. Algebra* **122** (1989), 353–373, [https://doi.org/10.1016/0021-8693\(89\)90222-6](https://doi.org/10.1016/0021-8693(89)90222-6).
- [21] A. Mahboubi, T. Sibut-Pinote, A formal proof of the irrationality of  $\zeta(3)$ , *Log. Methods Comput. Sci.* **17** (2021), no. 1, Paper No. 16, [https://doi.org/10.23638/LMCS-17\(1:16\)2021](https://doi.org/10.23638/LMCS-17(1:16)2021).
- [22] The mathlib Community, The Lean mathematical library, in: *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, ACM, 2020, pp. 367–381, <https://doi.org/10.1145/3372885.3373824>.
- [23] OEIS Foundation Inc., The On-Line Encyclopedia of Integer Sequences, <https://oeis.org>; entries A207118–A207123, A207069, A207070, A207124–A207127, A038718, A084990, A326247; accessed October 2026.
- [24] N. P. Smart, *The Algorithmic Resolution of Diophantine Equations*, London Mathematical Society Student Texts, Cambridge University Press, Cambridge, 1998, <https://doi.org/10.1017/CB09781107359994>.
- [25] R. P. Stanley, Differentiably finite power series, *European J. Combin.* **1** (1980), 175–188, [https://doi.org/10.1016/S0195-6698\(80\)80051-5](https://doi.org/10.1016/S0195-6698(80)80051-5).
- [26] R. P. Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry, *Ann. New York Acad. Sci.* **576** (1989), 500–535, <https://doi.org/10.1111/j.1749-6632.1989.tb16434.x>.
- [27] R. P. Stanley, *Combinatorics and Commutative Algebra*, 2nd ed., Progress in Mathematics 41, Birkhäuser, Boston, MA, 1996, <https://doi.org/10.1007/b139094>.
- [28] R. P. Stanley, *Enumerative Combinatorics, Volume 1*, 2nd ed., Cambridge Studies in Advanced Mathematics 49, Cambridge University Press, 2012, <https://doi.org/10.1017/CB09781139058520>.
- [29] H. S. Wilf, D. Zeilberger, An algorithmic proof theory for hypergeometric (ordinary and “ $q$ ”) multisum/integral identities, *Invent. Math.* **108** (1992), 575–633, <https://doi.org/10.1007/BF02100618>.
- [30] D. Zeilberger, A holonomic systems approach to special functions identities, *J. Comput. Appl. Math.* **32** (1990), 321–368, [https://doi.org/10.1016/0377-0427\(90\)90042-X](https://doi.org/10.1016/0377-0427(90)90042-X).
- [31] Y. Zhang, Contraction of Ore ideals with applications, in: *Proceedings of the 2016 ACM International Symposium on Symbolic and Algebraic Computation (ISSAC 2016)*, ACM, 2016, pp. 413–420, <https://doi.org/10.1145/2930889.2930890>; arXiv:1511.07922.

COLLEGE OF ARTS AND SCIENCES, BOSTON UNIVERSITY, BOSTON, MA 02215, USA

Email address: [r5x8ksva@bu.edu](mailto:r5x8ksva@bu.edu)