

DPP 01

Computer Science & Information Technology

Discrete Mathematics

Set Theory and Algebra

Q1 How many of the following statements are true?

- I. $\phi \in \phi$
- II. $\phi \subset \phi$
- III. $\phi \subseteq \phi$
- IV. $\phi \in \{\phi\}$
- V. $\phi \subset \{\phi\}$
- VI. $\phi \subseteq \{\phi\}$

Q2 If a set A has 63 proper subsets, then what is the cardinality of A?

Q3 If a set A has 64 subsets of odd cardinality, then what is $|A|$?

- (A) 6
- (B) 63
- (C) 7
- (D) 128

Q4 How many subset of $\{1, 2, 3, \dots, 11\}$ contains at least one even integer?

Q5 Let $A = \{1, 2, 3, 4, 5, 7, 8, 10, 11, 14, 17, 18\}$. How many subsets of A contains exactly six elements?

Q6 Let $A = \{2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15\}$. How many six-elements subsets of A contains exactly four even integers and two odd integers?

Q7 Let $A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$. How many subsets of A contains only odd integers?

Q8 Suppose A, B, C and D are subsets of U (the universe) with A as a subset of B and C as a subset of D i.e., $A \subseteq B$ and $C \subseteq D$.

Now consider the following statements

- I. $A \cap C \subseteq B \cap D$
- II. $A \cup C \subseteq B \cup D$

Which of the following option is correct?

- (A) Only I is true
- (B) Only II is true
- (C) Neither I nor II is true
- (D) Both I and II are true

Q9 Let $A = \{1, 2, 3, \dots, 15\}$. How many subsets of A contains all of the odd integers in A?

Q10 Let A and B be sets where $A = \{x \mid x^2 - 7x = -12\}$ and $B = \{x \mid x^2 - x = 6\}$. Which of the following is true regarding $A \cup B$ and $A \cap B$?

- (A) $A \cup B = \{5\}$ and $A \cap B = \{-2, 3, 4\}$
- (B) $A \cup B = \{3\}$ and $A \cap B = \{-2, 3, 4\}$
- (C) $A \cup B = \{-2, 3, 4\}$ and $A \cap B = \{3\}$
- (D) $A \cup B = \{2, 3, 4\}$ and $A \cap B = \{5\}$

Q11 Let A be $\{a, b, c\}$. Let the relation R on A and let $R = \{(b, a), (b, c), (c, a), (c, b), (b, b)\}$. Which of the following statements about R is/are true?

- (A) R is neither reflexive nor irreflexive.
- (B) R is not symmetric.
- (C) R is transitive.
- (D) R is not anti-symmetric.

Q12 Let A and B be two sets such that $A \times B = \{(1, 1), (2, 2), (3, 1), (3, 2), (1, 2), (1, 4), (2, 1), (2, 4), (3, 4)\}$. What is the value of $|P(A)| + |P(B)|$?

Q13 The relation R is defined on $Z = \{0, \pm 1, \pm 2, \pm 3, \dots\}$ by xRy if and only if $(x^2 + y^2) \bmod 4 = 0$. Then R is:

- (A) Reflexive, Symmetric
- (B) Transitive, Symmetric
- (C) Symmetric
- (D) Reflexive, Symmetric and Transitive



Android App

iOS App

PW Website

Q14 Let R be the relation on $\mathbb{R}$ (Where $\mathbb{R}$ is set of Natural Number) given by xRy if and only if $x < 2y + 1$, then R is _____.

- (A) Reflexive, but not symmetric and not transitive
- (B) Reflexive, Symmetric, and not transitive
- (C) Not Reflexive, not symmetric and Transitive
- (D) Reflexive, but not symmetric and Transitive

Q15 Suppose that R is the relation on positive real

numbers such that xRy if and only if $xy = 1$. Then R satisfies how many of the properties given below?

- (i) Reflexive
- (ii) Irreflexive
- (iii) Symmetric
- (iv) Anti-symmetric
- (v) Transitive
- (vi) Asymmetric

[Android App](#)| [iOS App](#)| [PW Website](#)

Answer Key

Q1 4~4

Q2 6~6

Q3 (C)

Q4 1984~1984

Q5 924~924

Q6 225~225

Q7 63~63

Q8 (D)

Q9 128~128

Q10 (C)

Q11 (A, B, D)

Q12 16~16

Q13 (B)

Q14 (A)

Q15 1~1

[Android App](#)| [iOS App](#)| [PW Website](#)

Hints & Solutions

Q1 Text Solution:

- I. $\phi \in \phi$ is false because the empty set has no members.
- II. $\phi \subset \phi$ is false because the empty set is not a proper subset of itself.
- III. $\phi \subseteq \phi$ is true because the empty set is a subset of every set and so it is a subset of itself.
- IV. $\phi \in \{\phi\}$ is true because ϕ is a member here.
- V. $\phi \subset \{\phi\}$ is true because the empty set is a proper subset of itself.
- VI. $\phi \subseteq \{\phi\}$ is true because the empty set is a subset of every set.

Q2 Text Solution:

If a set has n elements then the number of subsets will be 2^n and the number of proper subsets will be $2^n - 1$.

A has 63 proper subsets so,

$$2^n - 1 = 63$$

$$\Rightarrow 2^n = 63 + 1$$

$$\Rightarrow 2^n = 64$$

$$\Rightarrow 2^n = 2^6$$

$$\therefore n = 6$$

The cardinality of A is 6.

Q3 Text Solution:

The number of subsets for $\{1, 2, 3, \dots, n\}$ with odd cardinality is 2^{n-1} .

Now given that,

$$2^{n-1} = 64$$

$$\Rightarrow 2^{n-1} = 2^6$$

$$\Rightarrow n - 1 = 6$$

$$\Rightarrow n = 7$$

Q4 Text Solution:

There are 2^{11} subsets for the set $\{1, 2, 3, \dots, 11\}$.

Out of which, there are 2^6 subset for the set $\{1, 3, 5, 7, 9, 11\}$ that contains none of the even integers.

Hence, there are $2^{11} - 2^6 = 1984$ subsets that contains at least one even integer.

Q5 Text Solution:

If we choose 6 elements from a set of 12

elements then we can do it in ${}^{12}C_6 = 924$ ways.

Q6 Text Solution:

The set A can be bifurcated into Even and Odd subsets as follows:

Even integers = $\{2, 4, 6, 8, 10, 12\}$

Odd integers = $\{5, 7, 9, 11, 13, 15\}$

Out of 6 even integers, we can choose 4 even integers in 6C_4 ways.

Similarly out of 6 odd integers, we can choose 2 odd integers in 6C_2 ways.

Therefore, the total number of required subsets =

$${}^6C_4 * {}^6C_2 = \frac{6^3 \times 5 \times 4!}{2 \times 4!} \times \frac{6^3 \times 5 \times 4!}{4! \times 2} =$$

$$15 * 15 = 225$$

Q7 Text Solution:

The set of odd integers: $\{1, 3, 5, 7, 9, 11\}$

Total subsets (including the empty set) = $2^6 = 64$

Subtracting the empty set: $64 - 1 = 63$

Thus, the number of subsets containing only odd integers is **63**.

Q8 Text Solution:

Statement I: $A \cap C \subseteq B \cap D$

- Since $A \subseteq B$, every element in $A \cap C$ is also in $B \cap C$.
- Since $C \subseteq D$, every element in $B \cap C$ is also in $B \cap D$.
- Thus, $A \cap C \subseteq B \cap D$ is **true**.

Statement II: $A \cup C \subseteq B \cup D$

- Since $A \subseteq B$, every element in A is also in B, so $A \cup C \subseteq B \cup C$.
- Since $C \subseteq D$, every element in C is also in D, so $B \cup C \subseteq B \cup D$.
- Thus, $A \cup C \subseteq B \cup D$ is **true**.

Q9 Text Solution:

In the given set $A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15\}$, there are 8 odd integers and 7 even integers.

For all the even integers we have two choices



Android App

iOS App

PW Website

whether to include it or not.

Therefore, total possibilities = $2^7 = 128$

Q10 Text Solution:

$$x^2 - 7x = -12 \Rightarrow x^2 - 7x + 12 = 0 \Rightarrow (x-4)(x-3) = 0 \Rightarrow x = 4, x = 3.$$

$$x^2 - x = 6 \Rightarrow x^2 - x - 6 = 0 \Rightarrow (x-3)(x+2) = 0 \Rightarrow x = 3, x = -2.$$

Consequently, $A \cap B = \{3\}$

and $A \cup B = \{-2, 3, 4\}$.

Q11 Text Solution:

$R = \{(b, a), (b, c), (c, a), (c, b), (b, b)\}$

(a) **True;** as (a, a) and (c, c) is not there but (b, b) is present,

(b) **True;** as (a, b) is not there.

(c) **False;** as $(c, b), (b, c)$ is there but (c, c) is not there.

(d) **True;** due to existence of pair (b, c) & (c, b)

Q12 Text Solution:

$A = \{1, 2, 3\}$ and $B = \{1, 2, 4\}$

$|P(A)| = 2^3 = 8$, and $|P(B)| = 2^3 = 8$.

So, the value of $|P(A)| + |P(B)| = 8 + 8 = 16$.

Q13 Text Solution:

R is not reflexive because $(1, 1) \notin R$.

R is symmetric because if $(x^2 + y^2) \bmod 4 = 0$ then $(y^2 + x^2) \bmod 4 = 0$, as addition is commutative.

R is also transitive because R contains all those pairs which consists of even numbers.

$R = \{(2, 0), (2, 4), (4, 8), (8, 12) \dots\}$

So if $(x^2 + y^2) \bmod 4 = 0$ and $(y^2 + z^2) \bmod 4 = 0$ then $(x^2 + z^2) \bmod 4 = 0$

Q14 Text Solution:

xRy if $(x < 2y + 1)$

Reflexive as

$x < 2x + 1$ {always true}

Symmetric:

example

$(1, 5)$ holds as $1 < 2(5) + 1$ but $(5, 1) \notin R$ (as $5 > 2(1) + 1$)

Therefore, not symmetric.

For Transitivity:

Take example of $(5, 3) \in R$ & $(3, 2) \in R$ but $(5, 2) \notin R$

Therefore, Not transitive.

Q15 Text Solution:

$(1, 1) \in R$, but $(2, 2) \notin R$. So, it is neither reflexive nor irreflexive. Also, it is not asymmetric.

If $xy = 1$ then $yx = 1$. So, it is symmetric.

Since $(2, 1/2) \in R$ and $(1/2, 2) \in R$ but $(2, 2) \notin R$.

So, it is neither transitive nor anti-symmetric.



[Android App](#)

| [iOS App](#)

| [PW Website](#)