

# Permutation groups, lattices and orthogonal block structures

Rosemary A. Bailey and Peter J. Cameron  
University of St Andrews  
with Marina Anagnostopoulou-Merkouri (Bristol)



St Andrews Algebra and Combinatorics Seminar  
17 October 2024

# How it began

In the summer of 2022, Marina had a research internship in the department to work with Peter. Having finished before the money ran out, we looked at a new property of finite permutation groups which we called **pre-primitivity**. The idea was that pre-primitivity and quasiprimitivity were independent but together were equivalent to primitivity.

# How it began

In the summer of 2022, Marina had a research internship in the department to work with Peter. Having finished before the money ran out, we looked at a new property of finite permutation groups which we called **pre-primitivity**. The idea was that pre-primitivity and quasiprimitivity were independent but together were equivalent to primitivity. In the next academic year, Marina had a STARIS internship, and we looked more generally at properties of transitive imprimitive permutation groups, which led to the work we describe here.

# Combinatorial considerations about permutation groups

This was the title of a lecture course by Donald Higman in Oxford in 1969–1970. If  $G$  is a permutation group on  $\Omega$  which is primitive but not doubly transitive, then the **orbital digraphs** (whose edge sets are non-diagonal orbits of  $G$  on  $\Omega^2$ ) are connected, and together they form what Higman called a **coherent configuration**, whose adjacency matrices span an associative algebra.

# Combinatorial considerations about permutation groups

This was the title of a lecture course by Donald Higman in Oxford in 1969–1970. If  $G$  is a permutation group on  $\Omega$  which is primitive but not doubly transitive, then the **orbital digraphs** (whose edge sets are non-diagonal orbits of  $G$  on  $\Omega^2$ ) are connected, and together they form what Higman called a **coherent configuration**, whose adjacency matrices span an associative algebra.

Similar ideas were being developed by Boris Weisfeiler for the graph isomorphism problem, and by R. C. Bose and his students for design and analysis of experiments.

# Combinatorial considerations about permutation groups

This was the title of a lecture course by Donald Higman in Oxford in 1969–1970. If  $G$  is a permutation group on  $\Omega$  which is primitive but not doubly transitive, then the **orbital digraphs** (whose edge sets are non-diagonal orbits of  $G$  on  $\Omega^2$ ) are connected, and together they form what Higman called a **coherent configuration**, whose adjacency matrices span an associative algebra.

Similar ideas were being developed by Boris Weisfeiler for the graph isomorphism problem, and by R. C. Bose and his students for design and analysis of experiments.

Our aim was to do something similar for transitive but imprimitive groups.

# What is a Latin square?

## Definition

Let  $n$  be a positive integer.

A **Latin square** of order  $n$  is an  $n \times n$  array of cells in which  $n$  symbols are placed, one per cell, in such a way that each symbol occurs once in each row and once in each column.

# What is a Latin square?

## Definition

Let  $n$  be a positive integer.

A **Latin square** of order  $n$  is an  $n \times n$  array of cells in which  $n$  symbols are placed, one per cell, in such a way that each symbol occurs once in each row and once in each column.

The symbols may be letters, numbers, colours, ...



# What is a Latin square?


## Definition

Let  $n$  be a positive integer.

A **Latin square** of order  $n$  is an  $n \times n$  array of cells in which  $n$  symbols are placed, one per cell, in such a way that each symbol occurs once in each row and once in each column.

The symbols may be letters, numbers, colours, ...

A Latin square of order 8



# Partitions

## Definition

A **partition** of a set  $\Omega$  is a set  $\Pi$  of pairwise disjoint non-empty subsets of  $\Omega$ , called **parts**, whose union is  $\Omega$ .

# Partitions

## Definition

A **partition** of a set  $\Omega$  is a set  $\Pi$  of pairwise disjoint non-empty subsets of  $\Omega$ , called **parts**, whose union is  $\Omega$ .

## Definition

A partition  $\Pi$  is **uniform** if all of its parts have the same size, in the sense that, whenever  $\Gamma_1$  and  $\Gamma_2$  are parts of  $\Pi$ , there is a bijection from  $\Gamma_1$  onto  $\Gamma_2$ .

# Partitions

## Definition

A **partition** of a set  $\Omega$  is a set  $\Pi$  of pairwise disjoint non-empty subsets of  $\Omega$ , called **parts**, whose union is  $\Omega$ .

## Definition

A partition  $\Pi$  is **uniform** if all of its parts have the same size, in the sense that, whenever  $\Gamma_1$  and  $\Gamma_2$  are parts of  $\Pi$ , there is a bijection from  $\Gamma_1$  onto  $\Gamma_2$ .

## Example

If  $\Omega$  is the set of cells in a Latin square, then there are five natural uniform partitions of  $\Omega$ :

- $R$  each part is a row;
- $C$  each part is a column;
- $L$  each part consists of the those cells with a given letter;
- $U$  the **universal** partition, with a single part;
- $E$  the **equality** partition, whose parts are singletons.

If  $G$  is transitive but imprimitive on  $\Omega$ , then  $G$  preserves at least one non-trivial equivalence relation on  $\Omega$ , by definition.

If  $G$  is transitive but imprimitive on  $\Omega$ , then  $G$  preserves at least one non-trivial equivalence relation on  $\Omega$ , by definition.

According to the **Equivalence Relation Theorem**, an equivalence relation does the same job as a partition of  $\Omega$ . We mostly phrase our results in terms of partitions.

If  $G$  is transitive but imprimitive on  $\Omega$ , then  $G$  preserves at least one non-trivial equivalence relation on  $\Omega$ , by definition.

According to the **Equivalence Relation Theorem**, an equivalence relation does the same job as a partition of  $\Omega$ . We mostly phrase our results in terms of partitions.

There is a **partial order** on partitions, defined as follows:

$\Pi_1 \preceq \Pi_2$  if every part of  $\Pi_1$  is contained in a part of  $\Pi_2$ .

If  $G$  is transitive but imprimitive on  $\Omega$ , then  $G$  preserves at least one non-trivial equivalence relation on  $\Omega$ , by definition.

According to the **Equivalence Relation Theorem**, an equivalence relation does the same job as a partition of  $\Omega$ . We mostly phrase our results in terms of partitions.

There is a **partial order** on partitions, defined as follows:

$\Pi_1 \preceq \Pi_2$  if every part of  $\Pi_1$  is contained in a part of  $\Pi_2$ .

This can be read “ $\Pi_1$  refines  $\Pi_2$ ” or “ $\Pi_2$  is coarser than  $\Pi_1$ ”.



# Hasse diagrams

Given a collection  $\mathcal{P}$  of partitions of a set  $\Omega$ , we can show them on a Hasse diagram.


- ▶ Draw a dot for each partition in  $\mathcal{P}$ .
- ▶ If  $\Pi_1 \prec \Pi_2$  then put  $\Pi_2$  higher than  $\Pi_1$  in the diagram.
- ▶ If  $\Pi_1 \prec \Pi_2$  but there is no  $\Pi_3$  in  $\mathcal{P}$  with  $\Pi_1 \prec \Pi_3 \prec \Pi_2$  then draw a line from  $\Pi_1$  to  $\Pi_2$ .

# Hasse diagrams

Given a collection  $\mathcal{P}$  of partitions of a set  $\Omega$ , we can show them on a Hasse diagram.

- ▶ Draw a dot for each partition in  $\mathcal{P}$ .
- ▶ If  $\Pi_1 \prec \Pi_2$  then put  $\Pi_2$  higher than  $\Pi_1$  in the diagram.
- ▶ If  $\Pi_1 \prec \Pi_2$  but there is no  $\Pi_3$  in  $\mathcal{P}$  with  $\Pi_1 \prec \Pi_3 \prec \Pi_2$  then draw a line from  $\Pi_1$  to  $\Pi_2$ .

Here is the Hasse diagram for a Latin square.



The partitions of  $\Omega$ , with this order, form a lattice.

The partitions of  $\Omega$ , with this order, form a lattice.

The **meet** or **infimum**  $\Pi_1 \wedge \Pi_2$  is the partition whose parts are all non-empty intersections of parts of  $\Pi_1$  and  $\Pi_2$ .

The partitions of  $\Omega$ , with this order, form a lattice.

The **meet** or **infimum**  $\Pi_1 \wedge \Pi_2$  is the partition whose parts are all non-empty intersections of parts of  $\Pi_1$  and  $\Pi_2$ .

The **join** or **supremum**  $\Pi_1 \vee \Pi_2$  is the partition defined as follows: form a graph where two points are joined if they lie in the same part of either  $\Pi_1$  or  $\Pi_2$ ; the join is the partition into connected components of this graph.

The partitions of  $\Omega$ , with this order, form a lattice.

The **meet** or **infimum**  $\Pi_1 \wedge \Pi_2$  is the partition whose parts are all non-empty intersections of parts of  $\Pi_1$  and  $\Pi_2$ .

The **join** or **supremum**  $\Pi_1 \vee \Pi_2$  is the partition defined as follows: form a graph where two points are joined if they lie in the same part of either  $\Pi_1$  or  $\Pi_2$ ; the join is the partition into connected components of this graph.

As noted, partitions can also be regarded as equivalence relations. The **composition** of two relations  $R_1$  and  $R_2$  is the relation  $R_1 \circ R_2$  consisting of all pairs  $(\alpha, \beta)$  for which there exists  $\gamma$  such that  $(\alpha, \gamma) \in R_1$  and  $(\gamma, \beta) \in R_2$ . Two relations  $R_1$  and  $R_2$  **commute** if  $R_1 \circ R_2 = R_2 \circ R_1$ .

The partitions of  $\Omega$ , with this order, form a lattice.

The **meet** or **infimum**  $\Pi_1 \wedge \Pi_2$  is the partition whose parts are all non-empty intersections of parts of  $\Pi_1$  and  $\Pi_2$ .

The **join** or **supremum**  $\Pi_1 \vee \Pi_2$  is the partition defined as follows: form a graph where two points are joined if they lie in the same part of either  $\Pi_1$  or  $\Pi_2$ ; the join is the partition into connected components of this graph.

As noted, partitions can also be regarded as equivalence relations. The **composition** of two relations  $R_1$  and  $R_2$  is the relation  $R_1 \circ R_2$  consisting of all pairs  $(\alpha, \beta)$  for which there exists  $\gamma$  such that  $(\alpha, \gamma) \in R_1$  and  $(\gamma, \beta) \in R_2$ . Two relations  $R_1$  and  $R_2$  **commute** if  $R_1 \circ R_2 = R_2 \circ R_1$ .

Here is an alternative definition of Latin square.

## Definition

A **Latin square** is a set  $\{R, C, L\}$  of pairwise commuting uniform partitions of a set  $\Omega$  which satisfy

$$R \wedge C = R \wedge L = C \wedge L = E \text{ and } R \vee C = R \vee L = C \vee L = U.$$



Here is an alternative definition of Latin square.

## Definition

A **Latin square** is a set  $\{R, C, L\}$  of pairwise commuting uniform partitions of a set  $\Omega$  which satisfy  $R \wedge C = R \wedge L = C \wedge L = E$  and  $R \vee C = R \vee L = C \vee L = U$ .

## Definition

An **orthogonal block structure** or **OBS** is a sublattice of the partition lattice consisting of commuting uniform partitions.

Here is an alternative definition of Latin square.

## Definition

A **Latin square** is a set  $\{R, C, L\}$  of pairwise commuting uniform partitions of a set  $\Omega$  which satisfy  $R \wedge C = R \wedge L = C \wedge L = E$  and  $R \vee C = R \vee L = C \vee L = U$ .

## Definition

An **orthogonal block structure** or **OBS** is a sublattice of the partition lattice consisting of commuting uniform partitions.

So Latin squares are OBS.

## Proposition

- ▶ *Let  $\Pi_1$  and  $\Pi_2$  be equivalence relations. Then  $\Pi_1 \vee \Pi_2 = \Pi_1 \circ \Pi_2$  if and only if  $\Pi_1$  and  $\Pi_2$  commute.*

## Proposition

- ▶ *Let  $\Pi_1$  and  $\Pi_2$  be equivalence relations. Then  $\Pi_1 \vee \Pi_2 = \Pi_1 \circ \Pi_2$  if and only if  $\Pi_1$  and  $\Pi_2$  commute.*
- ▶ *A lattice of pairwise commuting partitions is modular.*

## Proposition

- ▶ *Let  $\Pi_1$  and  $\Pi_2$  be equivalence relations. Then  $\Pi_1 \vee \Pi_2 = \Pi_1 \circ \Pi_2$  if and only if  $\Pi_1$  and  $\Pi_2$  commute.*
- ▶ *A lattice of pairwise commuting partitions is modular.*

A lattice is **modular** if  $a \preccurlyeq c$  implies

$$a \vee (b \wedge c) = (a \vee b) \wedge c.$$

## Proposition

- ▶ *Let  $\Pi_1$  and  $\Pi_2$  be equivalence relations. Then  $\Pi_1 \vee \Pi_2 = \Pi_1 \circ \Pi_2$  if and only if  $\Pi_1$  and  $\Pi_2$  commute.*
- ▶ *A lattice of pairwise commuting partitions is modular.*

A lattice is **modular** if  $a \preccurlyeq c$  implies

$$a \vee (b \wedge c) = (a \vee b) \wedge c.$$

The modular law is implied by the **distributive laws**

$$a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c), \quad a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c),$$

which are equivalent to one another.

## Proposition

- ▶ Let  $\Pi_1$  and  $\Pi_2$  be equivalence relations. Then  $\Pi_1 \vee \Pi_2 = \Pi_1 \circ \Pi_2$  if and only if  $\Pi_1$  and  $\Pi_2$  commute.
- ▶ A lattice of pairwise commuting partitions is modular.

A lattice is **modular** if  $a \preccurlyeq c$  implies

$$a \vee (b \wedge c) = (a \vee b) \wedge c.$$

The modular law is implied by the **distributive laws**

$$a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c), \quad a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c),$$

which are equivalent to one another.

An OBS which is a distributive lattice is called a **poset block structure** or **PBS**. More on these later!

# Statisticians at Rothamsted

Here are some of the statisticians who have worked at the agricultural research station at Rothamsted.

Ronald Fisher	1919–1933	then UCL, then Cambridge
Frank Yates	1931–1968	
Oscar Kempthorne	1941–1946	then Ames, Iowa
Desmond Patterson	1947–1967	then Edinburgh
John Nelder	1968–1984	previously National Vegetable Research Station
Rosemary Bailey	1981–1990	
Robin Thompson	1997–now	previously Edinburgh, now emeritus






Trivial OBS (only  $U$  and  $E$ ).

Blocks containing plots.

A rectangle with one plot in each  
Row-Column intersection.



Many more OBS, including

- ▶ blocks containing plots containing subplots
- ▶ several rectangles
- ▶ a rectangle with subplots
- ▶ several rectangles with subplots.

# John Nelder: Crossing and Nesting



If  $\Pi_1$  is a partition of  $\Omega_1$  and  $\Pi_2$  is a partition of  $\Omega_2$  then  $\Pi_1 \times \Pi_2$  is the partition of  $\Omega_1 \times \Omega_2$  whose parts are intersections of a part of  $\Pi_1$  with a part of  $\Pi_2$ .

# John Nelder: Crossing and Nesting



If  $\Pi_1$  is a partition of  $\Omega_1$  and  $\Pi_2$  is a partition of  $\Omega_2$  then  $\Pi_1 \times \Pi_2$  is the partition of  $\Omega_1 \times \Omega_2$  whose parts are intersections of a part of  $\Pi_1$  with a part of  $\Pi_2$ .

Put  $\mathcal{P}_i = (\Omega_i, \mathcal{B}_i)$ , where  $\mathcal{B}_i$  is a collection of partitions of  $\Omega_i$ .



If  $\Pi_1$  is a partition of  $\Omega_1$  and  $\Pi_2$  is a partition of  $\Omega_2$  then  $\Pi_1 \times \Pi_2$  is the partition of  $\Omega_1 \times \Omega_2$  whose parts are intersections of a part of  $\Pi_1$  with a part of  $\Pi_2$ .

Put  $\mathcal{P}_i = (\Omega_i, \mathcal{B}_i)$ , where  $\mathcal{B}_i$  is a collection of partitions of  $\Omega_i$ .

**Crossing**  $\mathcal{P}_1$  with  $\mathcal{P}_2$  gives the set  $\mathcal{P}_1 \times \mathcal{P}_2$  of partitions

$$\{\Pi_1 \times \Pi_2 : \Pi_1 \in \mathcal{B}_1, \Pi_2 \in \mathcal{B}_2\}.$$



If  $\Pi_1$  is a partition of  $\Omega_1$  and  $\Pi_2$  is a partition of  $\Omega_2$  then  $\Pi_1 \times \Pi_2$  is the partition of  $\Omega_1 \times \Omega_2$  whose parts are intersections of a part of  $\Pi_1$  with a part of  $\Pi_2$ .

Put  $\mathcal{P}_i = (\Omega_i, \mathcal{B}_i)$ , where  $\mathcal{B}_i$  is a collection of partitions of  $\Omega_i$ .

**Crossing**  $\mathcal{P}_1$  with  $\mathcal{P}_2$  gives the set  $\mathcal{P}_1 \times \mathcal{P}_2$  of partitions

$$\{\Pi_1 \times \Pi_2 : \Pi_1 \in \mathcal{B}_1, \Pi_2 \in \mathcal{B}_2\}.$$

**Nesting**  $\mathcal{P}_2$  within  $\mathcal{P}_1$  gives the set  $\mathcal{P}_1 / \mathcal{P}_2$  of partitions  $\{\Pi_1 \times U_2 : \Pi_1 \in \mathcal{B}_1\} \cup \{E_1 \times \Pi_2 : \Pi_2 \in \mathcal{B}_2\}$ .



If  $\Pi_1$  is a partition of  $\Omega_1$  and  $\Pi_2$  is a partition of  $\Omega_2$  then  $\Pi_1 \times \Pi_2$  is the partition of  $\Omega_1 \times \Omega_2$  whose parts are intersections of a part of  $\Pi_1$  with a part of  $\Pi_2$ .

Put  $\mathcal{P}_i = (\Omega_i, \mathcal{B}_i)$ , where  $\mathcal{B}_i$  is a collection of partitions of  $\Omega_i$ .

**Crossing**  $\mathcal{P}_1$  with  $\mathcal{P}_2$  gives the set  $\mathcal{P}_1 \times \mathcal{P}_2$  of partitions

$$\{\Pi_1 \times \Pi_2 : \Pi_1 \in \mathcal{B}_1, \Pi_2 \in \mathcal{B}_2\}.$$

**Nesting**  $\mathcal{P}_2$  within  $\mathcal{P}_1$  gives the set  $\mathcal{P}_1 / \mathcal{P}_2$  of partitions  $\{\Pi_1 \times U_2 : \Pi_1 \in \mathcal{B}_1\} \cup \{E_1 \times \Pi_2 : \Pi_2 \in \mathcal{B}_2\}$ .

Iterated crossing and nesting gives **simple orthogonal block structures**.

# Desmond Patterson





In 1976–1978 I was employed as a post-doctoral research fellow in the Statistics Department at Edinburgh University. The aim was to apply ideas from combinatorics and group theory to design of experiments.

In 1976–1978 I was employed as a post-doctoral research fellow in the Statistics Department at Edinburgh University.

The aim was to apply ideas from combinatorics and group theory to design of experiments.

At the start, Desmond Patterson gave me copies of John Nelder's two 1965 papers on orthogonal block structure, and told me to read them.

In 1976–1978 I was employed as a post-doctoral research fellow in the Statistics Department at Edinburgh University.

The aim was to apply ideas from combinatorics and group theory to design of experiments.

At the start, Desmond Patterson gave me copies of John Nelder's two 1965 papers on orthogonal block structure, and told me to read them.

After three months, I said "OK, I understand them now."

In 1976–1978 I was employed as a post-doctoral research fellow in the Statistics Department at Edinburgh University.

The aim was to apply ideas from combinatorics and group theory to design of experiments.

At the start, Desmond Patterson gave me copies of John Nelder's two 1965 papers on orthogonal block structure, and told me to read them.

After three months, I said "OK, I understand them now."

Desmond responded "Hmph! That's good. No one else does."

In 1976–1978 I was employed as a post-doctoral research fellow in the Statistics Department at Edinburgh University.

The aim was to apply ideas from combinatorics and group theory to design of experiments.


At the start, Desmond Patterson gave me copies of John Nelder's two 1965 papers on orthogonal block structure, and told me to read them.

After three months, I said "OK, I understand them now."

Desmond responded "Hmph! That's good. No one else does."


I did not believe him then, but, looking back, I can see that his approach did not incorporate Nelder's ideas until much later.

# Oscar Kempthorne's papers



Then my colleague Robin Thompson gave me a 1961 technical report (long, but in typescript) by Oscar Kempthorne and his colleagues in Ames. This developed essentially the same ideas as Nelder's: lattices of partitions using some of the partitions in a Cartesian lattice (not necessarily with all coordinates having the same number of values, for example, the rows and columns of a rectangle).

# Oscar Kempthorne's papers



Then my colleague Robin Thompson gave me a 1961 technical report (long, but in typescript) by Oscar Kempthorne and his colleagues in Ames. This developed essentially the same ideas as Nelder's: lattices of partitions using some of the partitions in a Cartesian lattice (not necessarily with all coordinates having the same number of values, for example, the rows and columns of a rectangle).

Later I learnt that Kempthorne was furious that Nelder had "stolen" his ideas. I believe that they simply developed them independently, building on the work of Fisher and Yates. In those days, it took much longer for ideas to circulate widely.

# Putting the bits together

One morning, I came into work after drinking too much in the pub the previous evening. I realised that my brain was not capable of serious work, so I gave it the apparently simple task of matching Nelder's block structures with those of Kempthorne. Slowly, I worked through dimensions 1, 2 and 3.



# Putting the bits together

One morning, I came into work after drinking too much in the pub the previous evening. I realised that my brain was not capable of serious work, so I gave it the apparently simple task of matching Nelder's block structures with those of Kempthorne. Slowly, I worked through dimensions 1, 2 and 3. At the end of the day, I hit a problem. For dimension 4, Nelder's approach gave 15 possibilities, but Kempthorne's gave 16. I gave up and went home.

# Putting the bits together

One morning, I came into work after drinking too much in the pub the previous evening. I realised that my brain was not capable of serious work, so I gave it the apparently simple task of matching Nelder's block structures with those of Kempthorne. Slowly, I worked through dimensions 1, 2 and 3. At the end of the day, I hit a problem. For dimension 4, Nelder's approach gave 15 possibilities, but Kempthorne's gave 16. I gave up and went home. The next day, with a clear head, I realised that Kempthorne's approach always gives more possibilities than Nelder's in dimensions at least 4.

# SOBS $\Rightarrow$ poset, but not vice versa

crossing



nesting



not from SOBS



# SOBS $\Rightarrow$ poset, but not vice versa

crossing



nesting



not from SOBS



Kemphthorne's method gives all posets.

# SOBS $\Rightarrow$ poset, but not vice versa

crossing



nesting



not from SOBS



Kemphorne's method gives all posets.

Crossing and nesting give a similar formula in the statistical software R for use in analysis of variance.

“(Fields/Plots)  $\times$  Year” becomes “(Fields/Plots) \* Year”.

# SOBS $\Rightarrow$ poset, but not vice versa

crossing



nesting



not from SOBS



Kemphthorne's method gives all posets.


Crossing and nesting give a similar formula in the statistical software R for use in analysis of variance.

“(Fields/Plots)  $\times$  Year” becomes “(Fields/Plots) \* Year”.

When Terry Speed and RAB combined the two approaches in 1982, we called the structures **poset block structures**.

# How do we define PBS?

(Fields/Plots)  $\times$  Year




Same Field, same Plot  $\prec$  Same Field

$$\{F, P\} \supset \{F\}$$

# How do we define PBS?

(Fields/Plots)  $\times$  Year



Same Field, same Plot  $\prec$  Same Field

$$\{F, P\} \supset \{F\}$$


These partial orders correspond, but they are the opposite way round.

For years I have struggled with the problem of how to show these consistently on Hasse diagrams.



# How do we define PBS?

(Fields/Plots)  $\times$  Year



Same Field, same Plot  $\prec$  Same Field

$$\{F, P\} \supset \{F\}$$

These partial orders correspond, but they are the opposite way round.

For years I have struggled with the problem of how to show these consistently on Hasse diagrams.

Fortunately, my co-authors came up with a clever solution.

Not necessarily same Year  $\prec$  Not necessarily same Plot or Year

$$\{Y\} \subset \{P, Y\}$$

# Definition of Poset Block Structure

Let  $(M, \sqsubseteq)$  be a partially ordered set.

## Definition

A **down-set** in  $M$  is a subset  $D$  of  $M$  with the property that, if  $m \in D$  and  $m' \sqsubseteq m$ , then  $m' \in D$ .

The down-sets form a lattice under the operations of intersection and union.

# Definition of Poset Block Structure

Let  $(M, \sqsubseteq)$  be a partially ordered set.

## Definition

A **down-set** in  $M$  is a subset  $D$  of  $M$  with the property that, if  $m \in D$  and  $m' \sqsubseteq m$ , then  $m' \in D$ .

The down-sets form a lattice under the operations of intersection and union.

Let  $N = |M|$ .

For each element  $m_i$  of  $M$ , let  $\Omega_i$  be a set of size  $n_i > 1$ .

Let  $\Omega$  be the Cartesian product of the sets  $\Omega_i$  for all  $m_i$  in  $M$ .

# Definition of Poset Block Structure

Let  $(M, \sqsubseteq)$  be a partially ordered set.

## Definition

A **down-set** in  $M$  is a subset  $D$  of  $M$  with the property that, if  $m \in D$  and  $m' \sqsubseteq m$ , then  $m' \in D$ .

The down-sets form a lattice under the operations of intersection and union.

Let  $N = |M|$ .

For each element  $m_i$  of  $M$ , let  $\Omega_i$  be a set of size  $n_i > 1$ .

Let  $\Omega$  be the Cartesian product of the sets  $\Omega_i$  for all  $m_i$  in  $M$ .

Now we define a partition  $\Pi_D$  for each down-set  $D$  of  $M$ . This is done as follows.

## Definition

Elements  $(\alpha_1, \dots, \alpha_N)$  and  $(\beta_1, \dots, \beta_N)$  are in the same part of  $\Pi_D$  if and only if  $\alpha_i = \beta_i$  for all  $i$  with  $m_i \notin D$ .

# Easy consequences

$$E = \Pi_{\emptyset} \text{ and } U = \Pi_M.$$

# Easy consequences

$$E = \Pi_{\emptyset} \text{ and } U = \Pi_M.$$

If  $D_1$  and  $D_2$  are down-sets of  $M$  then

$$\Pi_{D_1} \wedge \Pi_{D_2} = \Pi_{D_1 \cap D_2} \text{ and } \Pi_{D_1} \vee \Pi_{D_2} = \Pi_{D_1 \cup D_2}.$$

So the partitions  $\Pi_D$  form a lattice isomorphic to the lattice of down-sets of  $M$ .

# Easy consequences

$$E = \Pi_{\emptyset} \text{ and } U = \Pi_M.$$

If  $D_1$  and  $D_2$  are down-sets of  $M$  then

$$\Pi_{D_1} \wedge \Pi_{D_2} = \Pi_{D_1 \cap D_2} \text{ and } \Pi_{D_1} \vee \Pi_{D_2} = \Pi_{D_1 \cup D_2}.$$

So the partitions  $\Pi_D$  form a lattice isomorphic to the lattice of down-sets of  $M$ .

Now we have two posets:

$(M, \sqsubseteq)$  and  $(\{\Pi_D : D \text{ is a downset of } M\}, \preceq)$ .

# Easy consequences

$$E = \Pi_{\emptyset} \text{ and } U = \Pi_M.$$

If  $D_1$  and  $D_2$  are down-sets of  $M$  then

$$\Pi_{D_1} \wedge \Pi_{D_2} = \Pi_{D_1 \cap D_2} \text{ and } \Pi_{D_1} \vee \Pi_{D_2} = \Pi_{D_1 \cup D_2}.$$

So the partitions  $\Pi_D$  form a lattice isomorphic to the lattice of down-sets of  $M$ .

Now we have two posets:

$(M, \sqsubseteq)$  and  $(\{\Pi_D : D \text{ is a downset of } M\}, \preceq)$ .

Do not confuse these with each other!



# Easy consequences

$$E = \Pi_{\emptyset} \text{ and } U = \Pi_M.$$

If  $D_1$  and  $D_2$  are down-sets of  $M$  then

$$\Pi_{D_1} \wedge \Pi_{D_2} = \Pi_{D_1 \cap D_2} \text{ and } \Pi_{D_1} \vee \Pi_{D_2} = \Pi_{D_1 \cup D_2}.$$

So the partitions  $\Pi_D$  form a lattice isomorphic to the lattice of down-sets of  $M$ .

Now we have two posets:

$(M, \sqsubseteq)$  and  $(\{\Pi_D : D \text{ is a downset of } M\}, \preceq)$ .

Do not confuse these with each other!

Kemphorne and his colleagues did make this mistake.

They tried to draw a single Hasse diagram showing both of these posets at once.

# Easy consequences

$$E = \Pi_{\emptyset} \text{ and } U = \Pi_M.$$

If  $D_1$  and  $D_2$  are down-sets of  $M$  then

$$\Pi_{D_1} \wedge \Pi_{D_2} = \Pi_{D_1 \cap D_2} \text{ and } \Pi_{D_1} \vee \Pi_{D_2} = \Pi_{D_1 \cup D_2}.$$

So the partitions  $\Pi_D$  form a lattice isomorphic to the lattice of down-sets of  $M$ .

Now we have two posets:

$(M, \sqsubseteq)$  and  $(\{\Pi_D : D \text{ is a downset of } M\}, \preceq)$ .

Do not confuse these with each other!

Kemphorne and his colleagues did make this mistake.

They tried to draw a single Hasse diagram showing both of these posets at once.

In June 1988 I took time out from a 2-week conference in Minneapolis to visit Kemphorne. He was very friendly, and said that he much appreciated my work on PBS.

## Definition

For each  $i$  in  $\{1, \dots, N\}$ , let  $A(i)$  be the set  $\{j \in \{1, \dots, N\} : m_i \sqsubset m_j\}$  (these are the **ancestors** of  $i$ .)

# Generalised wreath product

## Definition

For each  $i$  in  $\{1, \dots, N\}$ , let  $A(i)$  be the set  $\{j \in \{1, \dots, N\} : m_i \sqsubset m_j\}$  (these are the **ancestors** of  $i$ .)

Let  $\Omega^i$  be the Cartesian product  $\prod_{j \in A(i)} \Omega_j$ ,

# Generalised wreath product

## Definition

For each  $i$  in  $\{1, \dots, N\}$ , let  $A(i)$  be the set  $\{j \in \{1, \dots, N\} : m_i \sqsubset m_j\}$  (these are the **ancestors** of  $i$ .)  
Let  $\Omega^i$  be the Cartesian product  $\prod_{j \in A(i)} \Omega_j$ ,  
and let  $\pi^i$  be the natural projection from  $\Omega$  onto  $\Omega^i$ .

# Generalised wreath product

## Definition

For each  $i$  in  $\{1, \dots, N\}$ , let  $A(i)$  be the set  $\{j \in \{1, \dots, N\} : m_i \sqsubset m_j\}$  (these are the **ancestors** of  $i$ .)

Let  $\Omega^i$  be the Cartesian product  $\prod_{j \in A(i)} \Omega_j$ ,

and let  $\pi^i$  be the natural projection from  $\Omega$  onto  $\Omega^i$ .

Let  $G(m_i)$  be a permutation group on  $\Omega_i$ ,

# Generalised wreath product

## Definition

For each  $i$  in  $\{1, \dots, N\}$ , let  $A(i)$  be the set  $\{j \in \{1, \dots, N\} : m_i \sqsubset m_j\}$  (these are the **ancestors** of  $i$ .)

Let  $\Omega^i$  be the Cartesian product  $\prod_{j \in A(i)} \Omega_j$ ,

and let  $\pi^i$  be the natural projection from  $\Omega$  onto  $\Omega^i$ .

Let  $G(m_i)$  be a permutation group on  $\Omega_i$ ,

and let  $F_i$  be the set of all functions from  $\Omega^i$  into  $G(m_i)$ .

# Generalised wreath product

## Definition

For each  $i$  in  $\{1, \dots, N\}$ , let  $A(i)$  be the set

$\{j \in \{1, \dots, N\} : m_i \sqsubset m_j\}$  (these are the **ancestors** of  $i$ .)

Let  $\Omega^i$  be the Cartesian product  $\prod_{j \in A(i)} \Omega_j$ ,

and let  $\pi^i$  be the natural projection from  $\Omega$  onto  $\Omega^i$ .

Let  $G(m_i)$  be a permutation group on  $\Omega_i$ ,

and let  $F_i$  be the set of all functions from  $\Omega^i$  into  $G(m_i)$ .

Each  $f_i \in F_i$  allocates a permutation in  $G(m_i)$  to each tuple in  $\Omega^i$ .



# Generalised wreath product

## Definition

For each  $i$  in  $\{1, \dots, N\}$ , let  $A(i)$  be the set

$\{j \in \{1, \dots, N\} : m_i \sqsubset m_j\}$  (these are the **ancestors** of  $i$ .)

Let  $\Omega^i$  be the Cartesian product  $\prod_{j \in A(i)} \Omega_j$ ,

and let  $\pi^i$  be the natural projection from  $\Omega$  onto  $\Omega^i$ .

Let  $G(m_i)$  be a permutation group on  $\Omega_i$ ,

and let  $F_i$  be the set of all functions from  $\Omega^i$  into  $G(m_i)$ .

Each  $f_i \in F_i$  allocates a permutation in  $G(m_i)$  to each tuple in  $\Omega^i$ .

The **generalized wreath product**  $G$  of the groups  $G(m_1), \dots,$

$G(m_N)$  over the poset  $M$  is the group  $\prod_{i=1}^N F_i$ , acting on  $\Omega$  as follows:

# Generalised wreath product

## Definition

For each  $i$  in  $\{1, \dots, N\}$ , let  $A(i)$  be the set

$\{j \in \{1, \dots, N\} : m_i \sqsubset m_j\}$  (these are the **ancestors** of  $i$ .)

Let  $\Omega^i$  be the Cartesian product  $\prod_{j \in A(i)} \Omega_j$ ,

and let  $\pi^i$  be the natural projection from  $\Omega$  onto  $\Omega^i$ .

Let  $G(m_i)$  be a permutation group on  $\Omega_i$ ,

and let  $F_i$  be the set of all functions from  $\Omega^i$  into  $G(m_i)$ .

Each  $f_i \in F_i$  allocates a permutation in  $G(m_i)$  to each tuple in  $\Omega^i$ .

The **generalized wreath product**  $G$  of the groups  $G(m_1), \dots,$

$G(m_N)$  over the poset  $M$  is the group  $\prod_{i=1}^N F_i$ , acting on  $\Omega$  as

follows: if  $\omega = (\omega_1, \dots, \omega_N) \in \Omega$  and  $f = \prod_{i=1}^N f_i \in G$ , then

$$(\omega f)_i = \omega_i(\omega \pi^i f_i) \quad \text{for } i = 1, \dots, N.$$

## Theorem

*The automorphism group of the PBS is the generalized wreath product of symmetric groups  $S_{n_i}$  over the poset  $(M, \sqsubseteq)$ .*

## Theorem

*The automorphism group of the PBS is the generalized wreath product of symmetric groups  $S_{n_i}$  over the poset  $(M, \sqsubseteq)$ .*

By contrast, an orthogonal block structure may have only the trivial group of automorphisms.

## Theorem

*The automorphism group of the PBS is the generalized wreath product of symmetric groups  $S_{n_i}$  over the poset  $(M, \sqsubseteq)$ .*

By contrast, an orthogonal block structure may have only the trivial group of automorphisms.

We saw earlier that Latin squares give rise to OBSs (consisting of the two partitions  $E$  and  $U$  and the row, column, and letter partitions). It is known that almost all Latin squares have trivial automorphism group.

# Some comments on generalised wreath products

1. If  $N = 2$  and  $(M, \sqsubseteq)$  is an antichain then the GWP is  $G(m_1) \times G(m_2)$ .

# Some comments on generalised wreath products

1. If  $N = 2$  and  $(M, \sqsubseteq)$  is an antichain then the GWP is  $G(m_1) \times G(m_2)$ .
2. If  $N = 2$  and  $(M, \sqsubseteq)$  is a chain with  $m_1 \sqsubset m_2$  then the GWP is  $G(m_1) \wr G(m_2)$ .

# Some comments on generalised wreath products

1. If  $N = 2$  and  $(M, \sqsubseteq)$  is an antichain then the GWP is  $G(m_1) \times G(m_2)$ .
2. If  $N = 2$  and  $(M, \sqsubseteq)$  is a chain with  $m_1 \sqsubset m_2$  then the GWP is  $G(m_1) \wr G(m_2)$ .
3. If the poset  $(M, \sqsubseteq)$  can be made by iterated crossing and nesting (as in a simple orthogonal block structure) then the GWP can be made by iterating the corresponding direct and wreath products.



## Some comments on generalised wreath products

1. If  $N = 2$  and  $(M, \sqsubseteq)$  is an antichain then the GWP is  $G(m_1) \times G(m_2)$ .
2. If  $N = 2$  and  $(M, \sqsubseteq)$  is a chain with  $m_1 \sqsubset m_2$  then the GWP is  $G(m_1) \wr G(m_2)$ .
3. If the poset  $(M, \sqsubseteq)$  can be made by iterated crossing and nesting (as in a simple orthogonal block structure) then the GWP can be made by iterating the corresponding direct and wreath products.
4. If the poset  $(M, \sqsubseteq)$  cannot be made in this way, then neither can the GWP.

## Proposition

*The invariant partitions for a transitive permutation group form a lattice of uniform partitions.*

## Proposition

*The invariant partitions for a transitive permutation group form a lattice of uniform partitions.*

Thus if the partitions commute pairwise, they form an orthogonal block structure. This seems to happen remarkably often for small transitive groups: for example, it holds for 1886 of the 1954 transitive groups of degree 16.

## Proposition

*The invariant partitions for a transitive permutation group form a lattice of uniform partitions.*

Thus if the partitions commute pairwise, they form an orthogonal block structure. This seems to happen remarkably often for small transitive groups: for example, it holds for 1886 of the 1954 transitive groups of degree 16.

If the invariant partitions for  $G$  form a chain, then they commute pairwise. We do not know a weaker lattice property that forces the partitions to commute. Even requiring the lattice to be **Boolean** (isomorphic to the lattice of subsets of a set) does not suffice for this.

## Proposition

*The invariant partitions for a transitive permutation group form a lattice of uniform partitions.*

Thus if the partitions commute pairwise, they form an orthogonal block structure. This seems to happen remarkably often for small transitive groups: for example, it holds for 1886 of the 1954 transitive groups of degree 16.

If the invariant partitions for  $G$  form a chain, then they commute pairwise. We do not know a weaker lattice property that forces the partitions to commute. Even requiring the lattice to be **Boolean** (isomorphic to the lattice of subsets of a set) does not suffice for this.

We say that the transitive group  $G$  has the **OB property** if the invariant partitions commute (and so form an orthogonal block structure). It has the **PB property** if the lattice is distributive (and so is a poset block structure).

## Commuting subgroups

Let  $H$  and  $K$  be subgroups of a group  $G$ , with  $H \leq K$ . The corresponding interval in the subgroup lattice consists of all subgroups  $X$  for which  $H \leq X \leq K$ ; it is a lattice, with  $X \wedge Y = X \cap Y$  and  $X \vee Y = \langle X, Y \rangle$ .

# Commuting subgroups

Let  $H$  and  $K$  be subgroups of a group  $G$ , with  $H \leq K$ . The corresponding interval in the subgroup lattice consists of all subgroups  $X$  for which  $H \leq X \leq K$ ; it is a lattice, with  $X \wedge Y = X \cap Y$  and  $X \vee Y = \langle X, Y \rangle$ .

Let  $G$  be transitive on  $\Omega$ , and  $G_\alpha$  the stabiliser of  $\alpha \in \Omega$ . There is a natural isomorphism between the lattice of  $G$ -invariant partitions of  $\Omega$  and the interval from  $G_\alpha$  to  $G$ : the part of  $\Pi$  containing  $\alpha$  is the orbit containing  $\alpha$  of the corresponding subgroup. Moreover, partitions corresponding to  $X$  and  $Y$  commute if and only if  $XY = YX$ : we say that  $X$  and  $Y$  **commute** (to avoid confusion with permutations). Thus:

# Commuting subgroups

Let  $H$  and  $K$  be subgroups of a group  $G$ , with  $H \leq K$ . The corresponding interval in the subgroup lattice consists of all subgroups  $X$  for which  $H \leq X \leq K$ ; it is a lattice, with  $X \wedge Y = X \cap Y$  and  $X \vee Y = \langle X, Y \rangle$ .

Let  $G$  be transitive on  $\Omega$ , and  $G_\alpha$  the stabiliser of  $\alpha \in \Omega$ . There is a natural isomorphism between the lattice of  $G$ -invariant partitions of  $\Omega$  and the interval from  $G_\alpha$  to  $G$ : the part of  $\Pi$  containing  $\alpha$  is the orbit containing  $\alpha$  of the corresponding subgroup. Moreover, partitions corresponding to  $X$  and  $Y$  commute if and only if  $XY = YX$ : we say that  $X$  and  $Y$  **commute** (to avoid confusion with permutations). Thus:

## Proposition

*$G$  has the OB property if and only if the subgroups containing  $G_\alpha$  commute pairwise.*



# Commuting subgroups

Let  $H$  and  $K$  be subgroups of a group  $G$ , with  $H \leq K$ . The corresponding interval in the subgroup lattice consists of all subgroups  $X$  for which  $H \leq X \leq K$ ; it is a lattice, with  $X \wedge Y = X \cap Y$  and  $X \vee Y = \langle X, Y \rangle$ .

Let  $G$  be transitive on  $\Omega$ , and  $G_\alpha$  the stabiliser of  $\alpha \in \Omega$ . There is a natural isomorphism between the lattice of  $G$ -invariant partitions of  $\Omega$  and the interval from  $G_\alpha$  to  $G$ : the part of  $\Pi$  containing  $\alpha$  is the orbit containing  $\alpha$  of the corresponding subgroup. Moreover, partitions corresponding to  $X$  and  $Y$  commute if and only if  $XY = YX$ : we say that  $X$  and  $Y$  **commute** (to avoid confusion with permutations). Thus:

## Proposition

*$G$  has the OB property if and only if the subgroups containing  $G_\alpha$  commute pairwise.*

In particular, a regular permutation group has the OB property if and only if all its subgroups commute pairwise. These groups were determined by Iwasawa.

My earlier paper with Marina studied the following property. A transitive permutation group  $G$  is **pre-primitive** if every  $G$ -invariant partition is the orbit partition of a subgroup of  $G$ . We can take this subgroup to be the full stabiliser of the partition, and so it is a normal subgroup of  $G$ . Since normal subgroups commute, we see:

My earlier paper with Marina studied the following property. A transitive permutation group  $G$  is **pre-primitive** if every  $G$ -invariant partition is the orbit partition of a subgroup of  $G$ . We can take this subgroup to be the full stabiliser of the partition, and so it is a normal subgroup of  $G$ . Since normal subgroups commute, we see:

## Proposition

*A pre-primitive group has the OB property.*

My earlier paper with Marina studied the following property. A transitive permutation group  $G$  is **pre-primitive** if every  $G$ -invariant partition is the orbit partition of a subgroup of  $G$ . We can take this subgroup to be the full stabiliser of the partition, and so it is a normal subgroup of  $G$ . Since normal subgroups commute, we see:

## Proposition

*A pre-primitive group has the OB property.*

For the record: A permutation group is **quasiprimitive** if every non-trivial normal subgroup is transitive; thus, as mentioned earlier, a group is primitive if and only if it is pre-primitive and quasiprimitive.

Our first main theorem is the following:

## Theorem

*A generalised wreath product of primitive permutation groups  $(G_m : m \in M)$  is pre-primitive and has the OB property; it has the PB property if and only if there do not exist incomparable elements  $m_1, m_2 \in M$  such that  $G_{m_1}$  and  $G_{m_2}$  are cyclic of the same prime order.*

Our first main theorem is the following:

## Theorem

*A generalised wreath product of primitive permutation groups  $(G_m : m \in M)$  is pre-primitive and has the OB property; it has the PB property if and only if there do not exist incomparable elements  $m_1, m_2 \in M$  such that  $G_{m_1}$  and  $G_{m_2}$  are cyclic of the same prime order.*

The reason is that, if  $G$  and  $H$  are primitive, then the only non-trivial invariant partitions of  $G \times H$  correspond to orbits of  $G$  and of  $H$ , unless  $G = H = C_p$  for some prime  $p$ , in which case there are  $p + 1$  invariant partitions. (Recall that  $G \times H$  is the g.w.p. of  $G$  and  $H$  over the poset consisting of two incomparable elements.)

# The Krasner–Kaloujnine theorem

This is the following well-known result:

## Theorem

*Let  $G$  be a transitive imprimitive permutation group, with non-trivial invariant partition  $\Pi$ . Then  $G$  is naturally embeddable in the wreath product  $H \wr K$ , where  $H$  is the permutation group induced on a part of  $\Pi$  by its setwise stabiliser, and  $K$  the permutation group induced on the set of parts of  $\Pi$  by  $G$ .*

# The Krasner–Kaloujnine theorem

This is the following well-known result:

## Theorem

*Let  $G$  be a transitive imprimitive permutation group, with non-trivial invariant partition  $\Pi$ . Then  $G$  is naturally embeddable in the wreath product  $H \wr K$ , where  $H$  is the permutation group induced on a part of  $\Pi$  by its setwise stabiliser, and  $K$  the permutation group induced on the set of parts of  $\Pi$  by  $G$ .*

Is there an extension to our situation? The answer is yes ...



Let  $G$  be a transitive group with the PB property, and let  $\Lambda(G)$  be the lattice of  $G$ -invariant partitions. Then  $\Lambda$  is isomorphic to the lattice of down-sets in the poset  $M$ , whose elements can be recovered as the non- $E$  join-indecomposable (JI) elements of  $\Lambda$ . If  $\Pi$  is the partition corresponding to  $m \in M$ , then there is a unique maximal partition  $\Pi^-$  below  $\Pi$ , and we could define  $G_m$  to be the stabiliser of a part of  $\Pi$  acting on the set of parts of  $\Pi^-$  below it.

Let  $G$  be a transitive group with the PB property, and let  $\Lambda(G)$  be the lattice of  $G$ -invariant partitions. Then  $\Lambda$  is isomorphic to the lattice of down-sets in the poset  $M$ , whose elements can be recovered as the non- $E$  join-indecomposable (JI) elements of  $\Lambda$ . If  $\Pi$  is the partition corresponding to  $m \in M$ , then there is a unique maximal partition  $\Pi^-$  below  $\Pi$ , and we could define  $G_m$  to be the stabiliser of a part of  $\Pi$  acting on the set of parts of  $\Pi^-$  below it.

Unfortunately this does not work. The symmetric group  $S_6$  has an outer automorphism, so acts in different ways on two sets of size 6. Let  $\Omega$  be their Cartesian product. The only non-trivial partitions for  $G$  on  $\Omega$  are given by the coordinate projections, and the stabiliser of a part acts on it as  $\text{PGL}(2, 5)$ . But  $S_6$  is not a subgroup of  $\text{PGL}(2, 5) \times \text{PGL}(2, 5)$ .

## Second attempt

There is a way round this. Show that there is a unique maximal  $G$ -invariant partition  $\Psi$  such that  $\Psi \wedge \Pi = \Pi^-$ , and that  $\Psi$  is also JI. Then let  $G_m^*$  be the group induced by the stabiliser of a part of  $\Psi$  on the parts of  $\Psi^-$  it contains. Now  $G_m^* \geq G_m$ , and we have:

There is a way round this. Show that there is a unique maximal  $G$ -invariant partition  $\Psi$  such that  $\Psi \wedge \Pi = \Pi^-$ , and that  $\Psi$  is also JI. Then let  $G_m^*$  be the group induced by the stabiliser of a part of  $\Psi$  on the parts of  $\Psi^-$  it contains. Now  $G_m^* \geq G_m$ , and we have:

### Theorem

*Let  $G$  be a permutation group on  $\Omega$  with the PB property, and let  $M$  be the corresponding poset and  $G_m^*$  the group defined above for  $m \in M$ . Then  $G$  is naturally embedded in the generalised wreath product of the groups  $G_m^*$  over the poset  $M$ .*

The direct product  $G_1 \times G_2$  of transitive groups is naturally embedded in the wreath product in either order; indeed, it is their intersection. Can we generalise this?

## Varying the poset

The direct product  $G_1 \times G_2$  of transitive groups is naturally embedded in the wreath product in either order; indeed, it is their intersection. Can we generalise this?

Given two partial orders on a set  $M$ , their intersection is just the intersection of the sets of ordered pairs forming the two relations.

# Varying the poset

The direct product  $G_1 \times G_2$  of transitive groups is naturally embedded in the wreath product in either order; indeed, it is their intersection. Can we generalise this?

Given two partial orders on a set  $M$ , their intersection is just the intersection of the sets of ordered pairs forming the two relations.

## Theorem

*Let  $\sqsubseteq_1$  and  $\sqsubseteq_2$  be two partial orders on a set  $M$ , and*

*$\mathcal{F} = (G_m : m \in M)$  a family of transitive groups indexed by  $M$ .*

# Varying the poset

The direct product  $G_1 \times G_2$  of transitive groups is naturally embedded in the wreath product in either order; indeed, it is their intersection. Can we generalise this?

Given two partial orders on a set  $M$ , their intersection is just the intersection of the sets of ordered pairs forming the two relations.

## Theorem

*Let  $\sqsubseteq_1$  and  $\sqsubseteq_2$  be two partial orders on a set  $M$ , and*

*$\mathcal{F} = (G_m : m \in M)$  a family of transitive groups indexed by  $M$ .*

- ▶ *The intersection of the g.w.p.s of  $\mathcal{F}$  over  $\sqsubseteq_1$  and  $\sqsubseteq_2$  is the g.w.p. over their intersection.*



# Varying the poset

The direct product  $G_1 \times G_2$  of transitive groups is naturally embedded in the wreath product in either order; indeed, it is their intersection. Can we generalise this?

Given two partial orders on a set  $M$ , their intersection is just the intersection of the sets of ordered pairs forming the two relations.

## Theorem

*Let  $\sqsubseteq_1$  and  $\sqsubseteq_2$  be two partial orders on a set  $M$ , and*

*$\mathcal{F} = (G_m : m \in M)$  a family of transitive groups indexed by  $M$ .*

- ▶ *The intersection of the g.w.p.s of  $\mathcal{F}$  over  $\sqsubseteq_1$  and  $\sqsubseteq_2$  is the g.w.p. over their intersection.*
- ▶ *If  $\sqsubseteq_1$  is contained in  $\sqsubseteq_2$ , then the g.w.p. over  $\sqsubseteq_1$  is embedded in the g.w.p. over  $\sqsubseteq_2$ .*

# Linear extensions

Let  $M$  be a totally ordered set, say  $\{1 < 2 < \cdots < r\}$ , and let  $G_i$  be a transitive permutation group for each  $i \in M$ . The generalised wreath product over this poset is simply the **iterated wreath product**

$$G_1 \wr G_2 \wr \cdots \wr G_r.$$

# Linear extensions

Let  $M$  be a totally ordered set, say  $\{1 < 2 < \cdots < r\}$ , and let  $G_i$  be a transitive permutation group for each  $i \in M$ . The generalised wreath product over this poset is simply the **iterated wreath product**

$$G_1 \wr G_2 \wr \cdots \wr G_r.$$

Since the wreath product is associative, we do not need to bracket this expression.

# Linear extensions

Let  $M$  be a totally ordered set, say  $\{1 < 2 < \cdots < r\}$ , and let  $G_i$  be a transitive permutation group for each  $i \in M$ . The generalised wreath product over this poset is simply the **iterated wreath product**

$$G_1 \wr G_2 \wr \cdots \wr G_r.$$

Since the wreath product is associative, we do not need to bracket this expression.

A **linear extension** of a poset is a linear order containing the given poset. It is well known that any finite poset is the intersection of its linear extensions.

# Linear extensions

Let  $M$  be a totally ordered set, say  $\{1 < 2 < \cdots < r\}$ , and let  $G_i$  be a transitive permutation group for each  $i \in M$ . The generalised wreath product over this poset is simply the **iterated wreath product**

$$G_1 \wr G_2 \wr \cdots \wr G_r.$$

Since the wreath product is associative, we do not need to bracket this expression.

A **linear extension** of a poset is a linear order containing the given poset. It is well known that any finite poset is the intersection of its linear extensions.

## Theorem

*The generalised wreath product of a family  $(G_m : m \in M)$  of transitive permutation groups over a poset  $(M, \sqsubseteq)$  is the intersection of the iterated wreath products over all linear extensions of  $(M, \sqsubseteq)$ .*

This is immediate from the preceding theorem.