

Detecting changes in dispersion in COVID-19 case counts using a negative binomial model

JSM 2024

Rachael Aber, Yanming Di, Ben Dalziel

August 2, 2024

Table of Contents

Why study variability?

Negative binomial model

Results

Concluding remarks

References

Table of Contents

Why study variability?


Negative binomial model

Results

Concluding remarks


References

Why study variability?




- ▶ Highly variable case count time series suggest transmission heterogeneity or changes in R

Why study variability?



- ▶ Highly variable case count time series suggest transmission heterogeneity or changes in R
- ▶ Metrics of variability are overlooked: "How is variability related to different phases of an epidemic?" [3]

Why study variability?



- ▶ Highly variable case count time series suggest transmission heterogeneity or changes in R
- ▶ Metrics of variability are overlooked: "How is variability related to different phases of an epidemic?" [3]
- ▶ Adam et al.[1] found that COVID-19 transmission heterogeneity decreased over time

Why study variability?

- ▶ Dispersion of a case count time series may be a useful metric—part of a framework that models variance flexibly

Why study variability?

- ▶ Dispersion of a case count time series may be a useful metric—part of a framework that models variance flexibly
- ▶ A 'mean crowding' parameter [5] was proposed: mean number per individual of other individuals in the same quadrat

Why study variability?

- ▶ Dispersion of a case count time series may be a useful metric—part of a framework that models variance flexibly
- ▶ A 'mean crowding' parameter [5] was proposed: mean number per individual of other individuals in the same quadrat
- ▶ Useful way to think about dispersion in case count time series, degree of dispersion is degree of clustering/crowding of cases (from the perspective)

Why study variability?

- ▶ Dispersion of a case count time series may be a useful metric—part of a framework that models variance flexibly
- ▶ A 'mean crowding' parameter [5] was proposed: mean number per individual of other individuals in the same quadrat
- ▶ Useful way to think about dispersion in case count time series, degree of dispersion is degree of clustering/crowding of cases (from the perspective)




Table of Contents

Why study variability?

Negative binomial model

Results

Concluding remarks

References

Introduction to the method

- ▶ Let λ_t be epidemic intensity, I_t be incidence at time t , and θ_t be dispersion at time t

Introduction to the method

- ▶ Let λ_t be epidemic intensity, I_t be incidence at time t , and θ_t be dispersion at time t



$$I_t = \text{NB}(\mu = \lambda_t, \theta_t = I_{t-1})[4] \quad (1)$$

Introduction to the method

- ▶ Let λ_t be epidemic intensity, I_t be incidence at time t , and θ_t be dispersion at time t



$$I_t = \text{NB}(\mu = \lambda_t, \theta_t = I_{t-1})[4] \quad (1)$$

- ▶ Adjusted for population size using an offset in the model

Introduction to the method

- ▶ Let λ_t be epidemic intensity, I_t be incidence at time t , and θ_t be dispersion at time t



$$I_t = \text{NB}(\mu = \lambda_t, \theta_t = I_{t-1})[4] \quad (1)$$

- ▶ Adjusted for population size using an offset in the model
- ▶ Linear predictor includes a natural spline in time to account for autocorrelation in case counts (ns are)

$$\log(E[Y_i]/n_i) = \beta_1 h_1(t_i) + \beta_2 h_2(t_i) + \beta_3 h_3(t_i) \quad (2)$$

$$\log(E[Y_i]) - \log(n_i) = \beta_1 h_1(t_i) + \beta_2 h_2(t_i) + \beta_3 h_3(t_i) \quad (3)$$

$$\log(E[Y_i]) = \beta_1 h_1(t_i) + \beta_2 h_2(t_i) + \beta_3 h_3(t_i) + \log(n_i) \quad (4)$$

Introduction to the method

- ▶ Let λ_t be epidemic intensity, I_t be incidence at time t , and θ_t be dispersion at time t



$$I_t = \text{NB}(\mu = \lambda_t, \theta_t = I_{t-1})[4] \quad (1)$$

- ▶ Adjusted for population size using an offset in the model
- ▶ Linear predictor includes a natural spline in time to account for autocorrelation in case counts (ns are)

$$\log(E[Y_i]/n_i) = \beta_1 h_1(t_i) + \beta_2 h_2(t_i) + \beta_3 h_3(t_i) \quad (2)$$

$$\log(E[Y_i]) - \log(n_i) = \beta_1 h_1(t_i) + \beta_2 h_2(t_i) + \beta_3 h_3(t_i) \quad (3)$$

$$\log(E[Y_i]) = \beta_1 h_1(t_i) + \beta_2 h_2(t_i) + \beta_3 h_3(t_i) + \log(n_i) \quad (4)$$

- ▶ Model fit on a rolling basis to each time series (one estimate)



Negative binomial model



$$f_t(l) = \binom{l + \theta - 1}{l} \left(\frac{\mu}{\mu + \theta} \right)^l \left(\frac{\theta}{\mu + \theta} \right)^\theta \quad (5)$$

Negative binomial model



$$f_t(I) = \binom{I + \theta - 1}{I} \left(\frac{\mu}{\mu + \theta} \right)^I \left(\frac{\theta}{\mu + \theta} \right)^\theta \quad (5)$$



$$E(I) = \mu \quad (6)$$

$$Var(I) = \mu + \frac{\mu^2}{\theta} \quad (7)$$

Negative binomial model



$$f_t(I) = \binom{I + \theta - 1}{I} \left(\frac{\mu}{\mu + \theta} \right)^I \left(\frac{\theta}{\mu + \theta} \right)^\theta \quad (5)$$




$$E(I) = \mu \quad (6)$$

$$Var(I) = \mu + \frac{\mu^2}{\theta} \quad (7)$$


- ▶ LRT framework: at each time step along a time series, fit null and full(θ change) model, conduct LRT to produce p-value at each time point

Simulated time series data set




- ▶ Validity/power simulations: Gaussian and uniform epidemic curves with an attack rate of 0.1

Simulated time series data set



- ▶ Validity/power simulations: Gaussian and uniform epidemic curves with an attack rate of 0.1
- ▶ Varying magnitude of θ change, location of the change, underlying population size, and epidemic curve shape (allowed)

Simulated time series data set



- ▶ Validity/power simulations: Gaussian and uniform epidemic curves with an attack rate of 0.1
- ▶ Varying magnitude of θ change, location of the change, underlying population size, and epidemic curve shape (allowed)
- ▶ Epidemic curves over 60 time steps each were produced, and the likelihood-ratio test (LRT) procedure was applied to each

Application to empirical data

- ▶ Weekly case counts from US counties between 2020-01-04 and 2023-03-18

Application to empirical data

- ▶ Weekly case counts from US counties between 2020-01-04 and 2023-03-18
- ▶ Estimated θ_t for 154 time steps for 144 US counties (IRLS procedure implemented via the NBPSeq package[7] and from Di et al.[6])

Application to empirical data

- ▶ Weekly case counts from US counties between 2020-01-04 and 2023-03-18
- ▶ Estimated θ_t for 154 time steps for 144 US counties (IRLS procedure implemented via the NBPSeq package[7] and from Di et al.[6])
- ▶ We investigated large counties (largest three counties in each state)

Table of Contents

Why study variability?

Negative binomial model

Results

Concluding remarks

References

Results: simulated data

- ▶ Negative binomial/LRT method is robust to differences in population size (for population sizes examined)

Results: simulated data

- ▶ Negative binomial/LRT method is robust to differences in population size (for population sizes examined)
- ▶ Illustrated that an increase in θ is associated with decreased variability in simulated incidence time series (same relationship is observable in the empirical time series), with an increase in θ corresponding to a decrease in variability around the trend in incidence

Results: simulated data




Figure: Detecting dispersion changes in incidence time series in populations of different sizes. A/B: Simulated incidence when dispersion is constant/changes.

C/D: Constant/changing dispersion used in generation of above. E: Performance of LRT with simulated data that has different absolute differences in theta (horizontal axis of each pane) illustrates p-value distribution across




Figure: Method applied to case counts between 2020-01-04 and 2023-03-18 for Jefferson County, AL. A: Case counts . B: LRT statistic C: Log dispersion parameter.

Results: empirical data

- ▶ Highly overdispersed incidence observed more frequently later in time series (consistent) (Most dispersed category in Fig reaches)

Results: empirical data

- ▶ Highly overdispersed incidence observed more frequently later in time series (consistent) (Most dispersed category in Fig reaches)
- ▶ Evidence for a change in θ was observed across many counties (evidenced by concentration of low p-values around peak incidence)

Results: empirical data

- ▶ Highly overdispersed incidence observed more frequently later in time series (consistent) (Most dispersed category in Fig reaches)
- ▶ Evidence for a change in θ was observed across many counties (evidenced by concentration of low p-values around peak incidence)
- ▶ High dispersion may indicate less diffuse epidemics that are potentially more subject to climate forcing[2], or increased locally experienced mean density [5]

Results: empirical data

- ▶ Highly overdispersed incidence observed more frequently later in time series (consistent) (Most dispersed category in Fig reaches)
- ▶ Evidence for a change in θ was observed across many counties (evidenced by concentration of low p-values around peak incidence)
- ▶ High dispersion may indicate less diffuse epidemics that are potentially more subject to climate forcing[2], or increased locally experienced mean density [5]
- ▶ Raising variance relative to mean implies spatiotemporal "crowding" of cases (i.e. localized surges) which may necessitate more surge capacity in hospitals and testing centers

Results: empirical data




Figure: Incidence and dispersion in large counties in the US. A: Binned log of the dispersion parameter. B: Log of the dispersion parameter for each of the large counties (y-axis). C: Log incidence for each of the large counties (y-axis). D: LRT p-values for each of the large counties (y-axis).

Table of Contents

Why study variability?

Negative binomial model

Results

Concluding remarks

References

Concluding remarks

- ▶ LRT procedure performs well for relevant range of population sizes

Concluding remarks

- ▶ LRT procedure performs well for relevant range of population sizes
- ▶ Dispersion parameter and LRT statistic don't simply reflect changes in process mean

Concluding remarks

- ▶ LRT procedure performs well for relevant range of population sizes
- ▶ Dispersion parameter and LRT statistic don't simply reflect changes in process mean
- ▶ Dispersion is high at unexpected times (near peak incidence)(changes)

Concluding remarks

- ▶ LRT procedure performs well for relevant range of population sizes
- ▶ Dispersion parameter and LRT statistic don't simply reflect changes in process mean
- ▶ Dispersion is high at unexpected times (near peak incidence)(changes)
- ▶ Methods that use time series are crucial (due to); timing/allocation of public health resources can be achieved (with, pop less effective)

Table of Contents

Why study variability?

Negative binomial model

Results

Concluding remarks

References

- [1] Dillon Adam et al. *Time-varying transmission heterogeneity of SARS and COVID-19 in Hong Kong*. Tech. rep. ISSN: 2693-5015 Type: article. Mar. 2022. DOI: 10.21203/rs.3.rs-1407962/v1. URL: <https://www.researchsquare.com/article/rs-1407962/v1> (visited on 03/28/2022).
- [2] Benjamin D. Dalziel et al. "Urbanization and humidity shape the intensity of influenza epidemics in U.S. cities". In: *Science* 362.6410 (Oct. 2018). Publisher: American Association for the Advancement of Science, pp. 75–79. DOI: 10.1126/science.aat6030. URL: <https://www.science.org/doi/10.1126/science.aat6030> (visited on 10/05/2022).
- [3] Matthew Graham et al. "Measles and the canonical path to elimination". en. In: *Science* 364.6440 (May 2019), pp. 584–587. ISSN: 0036-8075, 1095-9203. DOI: 10.1126/science.aau6299. URL: <https://www.sciencemag.org/lookup/doi/10.1126/science.aau6299> (visited on 08/02/2021).

Thanks!