

## Two antiretroviral drugs likely to be confused

Two antiretroviral drugs likely to be confused

1728 Jill T. Johnson,  
Eddie B. Dunn, and  
Jonathan J. Wolfe

**M**edications for the treatment of human immunodeficiency virus (HIV) infection continue to be developed and placed on the market at a rapid pace. Many of these products receive FDA approval under an accelerated process and therefore may arrive at practice sites before staff members can familiarize themselves with brand and generic names. This lack of product knowledge can easily translate into medication errors.

The names of many antiretroviral agents look alike when written and sound alike when spoken.<sup>1</sup> Contributing to the confusion are caregivers who refer to

these medications by abbreviations instead of actual drug names.<sup>2</sup>

Viramune (nevirapine, Roxane Laboratories) and Viracept (nelfinavir mesylate, Agouron Pharmaceuticals) are used to treat HIV infection. Although their names are similar, these medications differ in important ways. Nevirapine is a nonnucleoside reverse-transcriptase inhibitor that binds directly to reverse transcriptase, inhibiting transcription of viral RNA into DNA. Nelfinavir is a protease inhibitor that prevents immature HIV from budding from the cell membrane and producing mature viral particles.<sup>3</sup>

The U.S. panel of the International AIDS Society recommends using one of two regimens to minimize the development of resistant virions during initial treatment of patients with HIV infection. One regimen combines two nucleoside reverse-transcriptase inhibitors with a protease inhibitor; the other regimen uses two nucleoside reverse-transcriptase inhibitors and a nonnucleoside reverse-transcriptase inhibitor.<sup>4</sup> If nelfinavir were the protease inhibitor prescribed in the first regimen, and nevirapine were inadvertently dispensed on refill, the patient would inappropriately stop protease inhibitor therapy. This could lead to the development of resistance to other HIV protease inhibitors, for cross-resistance among these agents is known to occur.<sup>5</sup> Correcting the error and resuming a protease inhibitor is unlikely to yield sustained suppression of viral replication.

On the other hand, if nevirapine were erroneously dispensed during the ini-

### Orthographic Similarities between Two Antiretroviral Drugs

Drug Names	Bigram Similarity	Trigram Similarity	Levenshtein Distance
Viramune and Viracept	0.75	0.50	5
Nevirapine and nelfinavir	0.80	0.14	6

*The Letters column is a forum for rapid exchange of ideas among readers of AJHP. Liberal criteria are applied in the review of submissions to encourage contributions to this column.*

*The Letters column includes the following types of contributions: (1) comments, addenda, and minor updates on previously published work, (2) alerts on potential problems in practice, (3) observations or comments on trends in drug use, (4) opinions on apparent trends or controversies in drug therapy or clinical research, (5) opinions on public health issues of interest to pharmacists in health systems, (6) comments on ASHP activities, and (7) human interest items about life as a pharmacist. Reports of adverse drug reactions must present a reasonably clear description of causality.*

*Short papers on practice innovations and other original work are*

*included in the Notes section rather than in Letters.*

*Letters need not be submitted with AJHP's manuscript checklist. The following conditions, however, must be adhered to: (1) the body of the letter must be no longer than two typewritten pages, (2) the use of references and tables should be minimized, (3) the number of authors should be no more than three, (4) the authors' names, affiliations, and mailing addresses must be typed at the end of the letter in the format used by AJHP, and (5) the entire letter (including references, tables, and authors' names) must be typed double-spaced. After acceptance of a letter, the authors are required to sign an exclusive publication statement and a copyright transfer form. All letters are subject to revision by the editors. Authors do not receive proofs of edited letters.*

*Letters may be sent via the Internet to [ajhp@ashp.org](mailto:ajhp@ashp.org).*

tial fill and the error went undetected during subsequent refills, the benefits of early intervention with a protease inhibitor would be lost.

The orthographic similarity between the generic names nevirapine and nelfinavir and their brand names Viramune and Viracept may be analyzed by bigram and trigram similarities and Levenshtein distance. These methods are useful in determining the potential for errors with medications whose names look alike and sound alike. Bigram and trigram pair scores between 0.5 and 0.99 indicate an increasing likelihood that names will be confused for one another. The bigram and trigram values in the table predict a strong likelihood of product mix-up. The Levenshtein values also indicate a high probability of error. A Levenshtein score of 6 is associated with a high likelihood of confusion between names.<sup>6</sup>

The National Coordinating Council for Medication Error Reporting and Prevention recognizes circumstances likely to produce an error as its category A for error classification. The facts about Viramune and Viracept fit this category well. In this instance, recognition of the potential for error can guide practitioners toward

appropriate safeguards in ordering and dispensing these medications.<sup>7</sup>

1. Errors with antivirals increase risk to HIV patients. *Qual Rev.* 1997. No. 59.
2. Davis NM. Antiretroviral drug name abbreviations—do your part to discourage their use. *Hosp Pharm.* 1997; 32:1186. Editorial.
3. Sachs MK. Antiretroviral chemotherapy of human immunodeficiency virus infections other than with azidothymidine. *Arch Intern Med.* 1992; 152:485-501.
4. Carpenter CCJ, Fischl MA, Hammer SM et al. Antiretroviral therapy for HIV infection in 1997: updated recommendations of the International AIDS Society—USA panel. *JAMA.* 1997; 277:1962-9.
5. Deeks SG, Smith M, Holodniy M et al. HIV-1 protease inhibitors: a review for clinicians. *JAMA.* 1997; 277:145-53.
6. Lambert BL. Predicting look-alike and sound-alike medication errors. *Am J Health-Syst Pharm.* 1997; 54:1161-71.
7. Raffalli J, Nowakowski J, Wormser GP. "Vira something": a taste of the wrong medicine. *Lancet.* 1997; 350:887. Letter.

**Jill T. Johnson, Pharm.D., BCPS**, Assistant Professor  
Department of Pharmacy Practice

**Eddie B. Dunn, Pharm.D.**, Assistant Professor  
Department of Pharmaceutics

**Jonathan J. Wolfe, Ph.D.**, Associate Professor  
Department of Pharmacy Practice

College of Pharmacy  
University of Arkansas for Medical Sciences  
4301 West Markham Street,  
Slot 522  
Little Rock, AR 72205-7122