

# Social buffering of pesticides in bumblebees: agent-based modeling of the effects of colony size and neonicotinoid exposure on behavior within nests

James D. Crall<sup>2</sup>, Benjamin L. De Bivort<sup>2</sup>, Biswadip Dey<sup>3</sup>, Ashlee N. Ford Versypt<sup>1\*</sup>

<sup>1</sup>Interdisciplinary Toxicology Program, Oklahoma State University, United States,

<sup>2</sup>Department of Organismic and Evolutionary Biology, Harvard University, United States,

<sup>3</sup>Department of Mechanical and Aerospace Engineering, Princeton University, United States

*Submitted to Journal:*

Frontiers in Ecology and Evolution

*Specialty Section:*

Behavioral and Evolutionary Ecology

*ISSN:*

2296-701X

*Article type:*

Brief Research Report Article

*Received on:*

16 Apr 2018

*Accepted on:*

12 Feb 2019

*Provisional PDF published on:*

12 Feb 2019

*Frontiers website link:*

[www.frontiersin.org](http://www.frontiersin.org)

*Citation:*

Crall JD, De\_bivort BL, Dey B and Ford\_versypt AN(2018) Social buffering of pesticides in bumblebees: agent-based modeling of the effects of colony size and neonicotinoid exposure on behavior within nests. *Front. Ecol. Evol.* 7:51. doi:10.3389/fevo.2019.00051

*Copyright statement:*

© 2019 Crall, De\_bivort, Dey and Ford\_versypt. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

Provisional

**Title:** Social buffering of pesticides in bumblebees: agent-based modeling of the effects of colony size and neonicotinoid exposure on behavior within nests

## Authors:

James D. Crall\*, Department of Organismic and Evolutionary Biology, Harvard University, [jcrall@oeb.harvard.edu](mailto:jcrall@oeb.harvard.edu)

Benjamin L. de Bivort, Department of Organismic and Evolutionary Biology, Harvard University, [debivort@oeb.harvard.edu](mailto:debivort@oeb.harvard.edu)

Biswadip Dey\*, Department of Mechanical & Aerospace Engineering, Princeton University, [biswadip@princeton.edu](mailto:biswadip@princeton.edu)

Ashlee N. Ford Versypt<sup>\*,†</sup>, School of Chemical Engineering, Interdisciplinary Toxicology Program, Oklahoma State University, [ashleefv@okstate.edu](mailto:ashleefv@okstate.edu)

\*Authors contributed equally to this work.

<sup>+</sup>Author for correspondence

## Authors listed alphabetically

## Author Contributions

JDC, BD, and ANFV contributed equally to this work. JDC, BD, and ANFV designed the study and wrote the computational code. JDC shared relevant experimental data from a previous study and conducted the in silico experiments. JDC, BdB, BD, and ANFV analyzed results of the in silico experiments and wrote the manuscript.

## Abstract

Neonicotinoids are a globally prevalent class of pesticides that can negatively impact bees and the pollination services they provide. While there is evidence suggesting that colony size may play an important role in mitigating neonicotinoid exposure in bees, mechanisms underlying these effects are not well understood. Here, a recently developed, agent-based computational model is used to investigate how the effects of sub-lethal neonicotinoid exposure on intranest behavior of bumblebees (*Bombus impatiens*) are modulated by colony size. Simulations from the model, parameterized using empirical data on bumblebee workers exposed to imidacloprid (a common neonicotinoid pesticide), suggest that colony size has significant effects on behavior neonicotinoid-sensitivity within bumblebee nests. Specifically, differences are reduced between treated and untreated workers in larger colonies for several key aspects of behavior within nests. Our results suggest that changes in both number of workers and nest architecture may contribute to making larger colonies less sensitive to pesticide exposure.

**Keywords:** Pollinator, Neonicotinoids, Bees, Collective behavior, Social insects, Pesticides, Agent-based modeling

47     **1. Introduction**

48         Agricultural use of neonicotinoid pesticides may negatively affect bees  
49         (Cresswell, 2011; Rundlöf et al., 2015; Woodcock et al., 2016; 2017) and the ecosystem  
50         services they provide (Stanley et al., 2015a) by altering worker behavior (Gill et al.,  
51         2012; Kessler et al., 2015; Whitehorn et al., 2012). Neonicotinoids disrupt learning and  
52         navigation of foraging bees (Palmer et al., 2013; Stanley et al., 2015b; Tan et al., 2015)  
53         and may impair colony growth (Laycock et al., 2012; Whitehorn et al., 2017) through  
54         reductions in resource intake (Feltham et al., 2014; Gill et al., 2012).

55         Workers within bee nests are likely exposed to neonicotinoids (Mitchell et al.,  
56         2017; Rundlöf et al., 2015), which can affect their locomotion and attraction to light (Tosi  
57         and Nieh, 2017), thermoregulation (Crall et al., 2018b; Potts et al., 2018; Tosi et al.,  
58         2016), and hygienic behaviors (Tsvetkov et al., 2017; Wu-Smart and Spivak, 2016). ,  
59         The neonicotinoid imidacloprid reduces activity and alters spatial dynamics and social  
60         interactions within bumblebee (*Bombus impatiens*) colonies (Crall et al., 2018b).

61         Mechanistic models of behavior within bee colonies can consider complex  
62         impacts of pesticides and other stressors under different scenarios (Bryden et al., 2013;  
63         Cresswell, 2017; Henry et al., 2017; Laycock and Cresswell, 2013; Sponsler and  
64         Johnson, 2016). Models have been developed for populations of bumblebees (Becher  
65         et al., 2018; Betti et al., 2017; Henry et al., 2017; Sponsler and Johnson, 2016; Thorbek  
66         et al., 2016) with and without exposure to neonicotinoids.

67         Colony size is a factor in sensitivity to environmental stressors as it plays a role  
68         in the resilience and conservation of social insects (Fisher, 2018; Straub et al., 2015).  
69         Colony size may affect susceptibility to neonicotinoid pesticides. One study (Rundlöf et  
70         al., 2015) found no significant effects of neonicotinoid exposure on honeybees (large  
71         colony sizes, >10,000), intermediate effects on bumblebees (moderate colony size, <  
72         500 workers), and very strong effects on solitary bees, under the same exposure  
73         conditions. Some negative effects of imidacloprid on honeybees were shown to be  
74         stronger in smaller colonies (Wu-Smart and Spivak, 2016).

75         The effects of colony size on worker behavior within nests in response to  
76         pesticide exposure are not well understood. Colony size may improve resilience to  
77         pesticides as a direct effect of larger population size by decreasing the chances of

78 colony failure when a small number of workers are lost to disease or death or behavioral  
79 impairment (Straub et al., 2015). Alternatively, larger colonies could mitigate the effects  
80 of exposure on the behavior of individuals. Changes in interaction rate or nest  
81 architecture associated with larger colony size could modulate the effects of pesticide  
82 exposure on individuals and colony performance through the division of labor,  
83 behavioral development, or social interactions associated with large social insect  
84 colonies.

85 Our goal is to elucidate the mechanisms underlying the effects of neonicotinoid  
86 exposure on worker behavior within bumblebee nests and how these effects are  
87 modulated by colony size. We use our spatially-explicit agent-based model called  
88 BeeNestABM (Crall et al., 2018b; Ford Versypt et al., 2018) to test the hypothesis that  
89 the effects of exposure on worker behavior within bumblebee nests are reduced in  
90 larger colonies. Bumblebee colonies range from a single individual during colony  
91 founding by a solitary queen up to several hundred workers (Michener and Laberge,  
92 1954). Inside colonies, interactions with nestmates and nest architecture can have  
93 important effects on worker behavior. Interactions with nestmates directly affect activity  
94 and task switching in colonies (Crall et al., 2018b; Renner and Nieh, 2008) and could  
95 modulate behavioral impacts of pesticides. The physical structure of the nest (e.g., food  
96 pots and developing brood) can modulate worker behavior by carrying information on  
97 larval development and hunger (Boer and Duchateau, 2006; Dornhaus, 2005; Heinrich,  
98 1974). Because neonicotinoids directly reduce activity, which indirectly alters spatial and  
99 social dynamics (Crall et al., 2018b), increased interactions with nestmates or nest  
100 structures in larger colonies could mitigate the impacts of neonicotinoid exposure.

101

## 102 **2. Model description**

103 BeeNestABM was parameterized using empirical data (Crall et al., 2018b) and  
104 simulates bumblebee activity and interactions within a nest. The code and  
105 documentation are available (Ford Versypt et al., 2018). Here, we briefly describe the  
106 model.

107 BeeNestABM simulates movements of bees within a nest on short time scales  
108 allowing for interactions with nestmates and nest structures. The state variables at each

109 time step and for each bee are the x- and y-coordinates, heading angle, speed, and  
110 activity state. The simulation is initialized with a user-specified number of bees. Bees  
111 are initialized with random positions inside a nest arena (0.2 x 0.2 m for all simulations  
112 here). The model is iterated in steps of 0.5 s. Nest structures are placed in 1-cm apart in  
113 a square grid covering a centered, user-specified fraction of the arena.

114 The model consists of three processes that occur at each time step. First,  
115 distances between bees and the nest structures (Fig 1A) are calculated. Next, the  
116 activity state for each bee (Fig 1C-G) is updated. Finally, the coordinates for each active  
117 bee are updated using the movement rules (Fig 1B).

118

## 119 **2.1. Activity state updates**

120 Activity-switching probabilities depend on location on or off the nest structures  
121 (Fig 1D), proximity to nestmates (Fig 1E), and treatment group (Fig 1C-G). Workers are  
122 assigned probabilities calculated previously (Crall et al., 2018b Supplement) from the  
123 means of empirical data of automatically tracked (Crall et al., 2015) bumblebees. These  
124 probabilities represent the probabilities of switching between active (moving) and  
125 inactive (stationary) between time steps, calculated for different combinations of social  
126 interaction state and location (Fig 1). Probabilities for treated bees are set equivalent to  
127 controls for parameters that did not differ significantly between control and treated  
128 workers (Fig 1F-G). Treated bees have activity-switching probabilities that make bees  
129 less likely to initiate movement either spontaneously or after physical contact with a  
130 nestmate and more likely to become immobilized, with stronger effects when treated  
131 bees are located off the nest structures (Fig 1F-G).

132 In the dataset (Crall et al., 2018b), workers were exposed orally to an acute dose  
133 of either 0 ng (control) or 1.0 ng imidacloprid at 87 ppb (treated) in a sucrose solution  
134 and allowed to behave freely in a nest arena. 1.0 ng was approximately equal to the  
135 cumulative imidacloprid consumed per worker in a single day of chronic feeding on  
136 nectar with environmentally realistic imidacloprid concentrations (6 ppb). While that  
137 study found qualitatively similar effects on behavior in workers exposed to imidacloprid  
138 either acutely or chronically, the behavioral effects during chronic exposure varied with

139 time of day. To limit model complexity, we did not incorporate circadian effects but  
140 focused on acute exposure data collected at the same time of day.

141

## 142 **2.2 Movement rules**

143 Previously inactive bees are set at a speed sampled from the empirical speed  
144 distribution. Bees previously moving are updated to the sampled speed if it is within  
145 specified limits of the current speed; otherwise, there is a 10% probability of switching to  
146 the sampled value from the current speed. The heading angle for each bee is the  
147 weighted angular mean of a random walk angle fluctuating within 90° of the current  
148 angle and the angle toward the environmental stimuli determined by the net  
149 displacement between the bee and nest structures (Fig 1B). The environmental stimuli  
150 represent attractions to the nest structures with weights estimated using empirical data  
151 ( $\mu = 0.067$  and 0.052 for control and treated bees, respectively). After location updates,  
152 movement is truncated for each bee that would move outside the perimeter.

153

## 154 **3. Results and Discussion**

155 We used BeeNestABM to examine the effects of colony size and exposure  
156 treatment intensities (the proportion of workers within the colony exposed to  
157 imidacloprid) on worker behavior. We ran simulations where two aspects of colony size  
158 (numbers of nest structures and of workers) were varied simultaneously from 4 to 441,  
159 spanning the ecologically relevant colony sizes for bumblebees, to maintain a 1:1  
160 worker:nest structure ratio across a range of treatment intensities (Fig 2). To examine  
161 the relative importance of nest architecture and number of workers, we ran simulations  
162 varying these parameters independently (Figs S1-S2). For each combination of colony  
163 size and exposure intensity, we estimated mean values across all simulations and  
164 individuals for four behavioral metrics (Figs 2A-D, S1, S2).

165 Exposure intensity and colony size significantly affected worker behavior within  
166 nests. Larger colonies had higher activity levels (Fig 2A), mean distance to the nest  
167 center (Fig 2B), and proportions of time spent on the nest structure (Fig 2C), with  
168 interaction rates (Fig 2D) decreasing at intermediate colony sizes. Higher treatment  
169 intensity was associated with substantial declines in activity and interaction rate (Figs

170 2A,D), a weak increase in  $D_{NC}$  (Fig 2B), and a weak decrease in  $P_{nest}$  (Fig 2C).  
171 Simulations isolating the effects of variation in nest size and number of workers  
172 separately suggest that the effects of colony size are driven primarily by changes in the  
173 size and number of nest structures, especially for  $D_{NC}$  and (trivially)  $P_{nest}$  (Fig S1).

174 We found evidence that the relative difference between control and treated  
175 workers varied significantly within colony size (Fig 2E-H); relative differences between  
176 control and treated workers in activity (Fig 2E),  $D_{NC}$  (Fig 2F), and  $P_{nest}$  (Fig 2G) were  
177 reduced in larger colonies, while relative differences in interaction rate were greatest at  
178 intermediate colony sizes (Fig 2H). The reduced differences between control and  
179 treated workers in larger colonies is likely driven primarily by the weaker effects of  
180 imidacloprid on activity parameters when located on the nest structures (Fig 1F-G, Fig  
181 S2). We found qualitatively similar effects for absolute differences between control and  
182 treated workers for most metrics (Fig S3), with the exception of proportion of time spent  
183 on the nest structure, which varied most with colony size (Figs 2, S3).

184

#### 185 **4. Conclusions**

186 Our results support the hypothesis that larger colonies buffer the effects of  
187 neonicotinoid exposure on behavior within bumblebee nests: differences between  
188 control and treated workers in several key emergent behavioral parameters were  
189 reduced in larger colonies (Fig 2A-D). The dominant effects of nest structure size (Fig  
190 S1-S2) suggest that nest architecture, rather than altered rates of interaction with  
191 nestmates, is the primary driver of reduced effects of imidacloprid exposure in large  
192 colonies, although higher worker numbers did have qualitatively similar, but  
193 quantitatively weaker, effects for several metrics (Fig S2). The strong effects of nest  
194 architecture found here result directly from the empirical observation that neonicotinoids  
195 have particularly strong effects on activity when workers are located on the nest  
196 structure (Crall et al., 2018b). The specific mechanisms driving this patterns are not  
197 clear but suggest that some aspects of the nest structure (potentially including chemical  
198 signals derived from the brood (Boer and Duchateau, 2006) or from nestmates  
199 (Richardson and Gorochowski, 2015) laid on the nest structure) play a direct role in

200 regulating worker behavior. Being on the nest architecture stimulates movement (Fig 1),  
201 potentially mitigating the negative effects of neonicotinoids on activity.

202 The mitigating effects of colony size likely translate into significant impacts on  
203 colony performance. Location on the nest structure is a strong proxy for direct brood  
204 care and maintenance behavior within the nest (Crall et al., 2018a). Our results lead to  
205 the testable predictions that (a) similar levels of exposure will lead to stronger effects on  
206 nest behavior in smaller colonies and that (b) this would lead to stronger reductions in  
207 per-worker nursing rates and brood growth in smaller colonies. For bumblebees,  
208 increased sensitivity at small colony sizes is especially relevant because exposure  
209 during colony initiation by a solitary queen or during early developmental stages can  
210 substantially impair colony performance (Baron et al., 2017; Leza et al., 2018).

211 Our model improves understanding of how colony size and pesticide exposure  
212 affect behavior within bumblebee nests. However, the scope of the model is limited to a  
213 simplified set of behaviors. In the future, the model could be expanded to more complex  
214 behaviors of bumblebees (Cameron, 1989; Crall et al., 2018a). Combining this model  
215 with colony dynamic processes at larger spatial and longer time scales, including  
216 foraging, growth, and landscape dynamics (Becher et al., 2018) could help assess how  
217 the impacts of neonicotinoids on behavior translate to long-term colony performance.  
218

## 219 **5. Acknowledgments**

220 The authors acknowledge support from the Statistical and Applied Mathematical  
221 Sciences Institute (SAMSI) and the organizers of the workshop titled Data-Driven  
222 Modeling of Collective Behavior and Emergent Phenomena in Biology that enabled the  
223 collaboration. JDC was supported in part by funding from the Rockefeller Foundation  
224 and the Winslow Foundation. BdB was supported by the NSF (IOS-1557913), the Alfred  
225 P. Sloan Foundation, The Klingenstein-Simons Fellowship, and the Smith Family  
226 Foundation.

## 227 **6. References**

228  
229  
230 Baron, G. L., Jansen, V. A. A., Brown, M. J. F., and Raine, N. E. (2017). Pesticide  
231 reduces bumblebee colony initiation and increases probability of population  
232 extinction. *Nature Ecology & Evolution* 1, 1308–1316. doi:10.1038/s41559-017-


- 233 0260-1.
- 234 Becher, M. A., Twiston-Davies, G., Penny, T. D., Goulson, D., Rotheray, E. L., and  
235 Osborne, J. L. (2018). Bumble-BEEHAVE: A systems model for exploring  
236 multifactorial causes of bumblebee decline at individual, colony, population and  
237 community level. *J Appl Ecol* 351, 14–12. doi:10.1111/1365-2664.13165.
- 238 Betti, M., LeClair, J., Wahl, L., and Zamir, M. (2017). Bee++: An Object-Oriented, Agent-  
239 Based Simulator for Honey Bee Colonies. *Insects* 8, 31–26.  
240 doi:10.3390/insects8010031.
- 241 Boer, S. P. A. D., and Duchateau, M. J. H. M. (2006). A larval hunger signal in the  
242 bumblebee *Bombus terrestris*. *Insect. Soc.* 53, 369–373. doi:10.1007/s00040-006-  
243 0883-8.
- 244 Bryden, J., Gill, R. J., Mitton, R. A. A., Raine, N. E., and Jansen, V. A. A. (2013).  
245 Chronic sublethal stress causes bee colony failure. *Ecol Letters* 16, 1463–1469.  
246 doi:10.1111/ele.12188.
- 247 Cameron, S. A. (1989). Temporal Patterns of Division of Labor among Workers in the  
248 Primitively Eusocial Bumble Bee, *Bombus griseocollis* (Hymenoptera: Apidae)1).  
249 *Ethology* 80, 137–151. doi:10.1111/j.1439-0310.1989.tb00735.x.
- 250 Crall, J. D., Gravish, N., Mountcastle, A. M., and Combes, S. A. (2015). BEEtag: A Low-  
251 Cost, Image-Based Tracking System for the Study of Animal Behavior and  
252 Locomotion. *PLoS ONE* 10, e0136487. doi:10.1371/journal.pone.0136487.
- 253 Crall, J. D., Gravish, N., Mountcastle, A. M., Kocher, S. D., Oppenheimer, R. L., Pierce,  
254 N. E., et al. (2018a). Spatial fidelity of workers predicts collective response to  
255 disturbance in a social insect. *Nat Commun*, 1–13. doi:10.1038/s41467-018-03561-  
256 w.
- 257 Crall, J. D., Switzer, C. M., Oppenheimer, R. L., Ford Versypt, A. N., Dey, B., Brown, A.,  
258 et al. (2018b). Neonicotinoid exposure disrupts bumblebee nest behavior, social  
259 networks, and thermoregulation. *Science* 362, 683–686.  
260 doi:10.1126/science.aat1598.
- 261 Cresswell, J. E. (2011). A meta-analysis of experiments testing the effects of a  
262 neonicotinoid insecticide (imidacloprid) on honey bees. *Ecotoxicology* 20, 149–157.  
263 doi:10.1007/s10646-010-0566-0.
- 264 Cresswell, J. E. (2017). A demographic approach to evaluating the impact of stressors  
265 on bumble bee colonies. *Ecol Ent* 42, 221–229. doi:10.1111/een.12376.
- 266 Dornhaus, A. (2005). Bumble bees (*Bombus terrestris*) store both food and information  
267 in honeypots. *Behavioral Ecology* 16, 661–666. doi:10.1093/beheco/ari040.
- 268 Feltham, H., Park, K., and Goulson, D. (2014). Field realistic doses of pesticide

- 269 imidacloprid reduce bumblebee pollen foraging efficiency. *Ecotoxicology* 23, 317–  
270 323. doi:10.1007/s10646-014-1189-7.
- 271 Fisher, K. (2018). Are societies resilient? Challenges faced by social insects in a  
272 changing world. *Insect. Soc.* 0, 0–0. doi:10.1007/s00040-018-0663-2.
- 273 Ford Versypt, A. N., Crall, J. D., and Dey, B. (2018). BeeNestABM: An open-source  
274 agent-based model of spatiotemporal distribution of bumblebees in nests. *JOSS* 3,  
275 718. doi:10.21105/joss.00718.
- 276 Gill, R. J., Ramos-Rodriguez, O., and Raine, N. E. (2012). Combined pesticide  
277 exposure severely affects individual- and colony-level traits in bees. *Nature* 490,  
278 105–108. doi:10.1038/nature11585.
- 279 Heinrich, B. (1974). Pheromone induced brooding behavior in *Bombus vosnesenskii*  
280 and *B. edwardsii* (Hymenoptera: Bombidae). *Journal of the Kansas Entomological  
281 Society* 47, 396–404.
- 282 Henry, M., Becher, M. A., Osborne, J. L., Kennedy, P. J., Aupinel, P., Bretagnolle, V., et  
283 al. (2017). Predictive systems models can help elucidate bee declines driven by  
284 multiple combined stressors. *Apidologie* 48, 1–12. doi:10.1007/s13592-016-0476-0.
- 285 Kessler, S. C., Tiedeken, E. J., Simcock, K. L., Derveau, S., Mitchell, J., Softley, S., et  
286 al. (2015). Bees prefer foods containing neonicotinoid pesticides. *Nature* 521, 74–  
287 76. doi:10.1038/nature14414.
- 288 Laycock, I., and Cresswell, J. E. (2013). Repression and Recuperation of Brood  
289 Production in *Bombus terrestris* Bumble Bees Exposed to a Pulse of the  
290 Neonicotinoid Pesticide Imidacloprid. *PLoS ONE* 8, e79872–8.  
291 doi:10.1371/journal.pone.0079872.
- 292 Laycock, I., Lenthall, K. M., Barratt, A. T., and Cresswell, J. E. (2012). Effects of  
293 imidacloprid, a neonicotinoid pesticide, on reproduction in worker bumble bees  
294 (*Bombus terrestris*). *Ecotoxicology* 21, 1937–1945. doi:10.1007/s10646-012-0927-y.
- 295 Leza, M., Watrous, K. M., Bratu, J., and Woodard, S. H. (2018). Effects of neonicotinoid  
296 insecticide exposure and monofloral diet on nest-founding bumblebee queens.  
297 *Proceedings of the Royal Society B: Biological Sciences* 285, 20180761–9.  
298 doi:10.1098/rspb.2018.0761.
- 299 Michener, C. D., and Laberge, W. E. (1954). A Large *Bombus* Nest from Mexico.  
300 *Psyche: A Journal of Entomology* 61, 63–67. doi:10.1155/1954/41470.
- 301 Mitchell, E. A. D., Mulhauser, B., Mulot, M., Mutabazi, A., Glauser, G., and Aebi, A.  
302 (2017). A worldwide survey of neonicotinoids in honey. *Science* 358, 109–111.  
303 doi:10.1126/science.aan3684.
- 304 Palmer, M. J., Moffat, C., Saranzewa, N., Harvey, J., Wright, G. A., and Connolly, C. N.

- 305 (2013). Cholinergic pesticides cause mushroom body neuronal inactivation in  
306 honeybees. *Nat Commun* 4, 1634. doi:10.1038/ncomms2648.
- 307 Potts, R., Clarke, R. M., Oldfield, S. E., Wood, L. K., Hempel de Ibarra, N., and  
308 Cresswell, J. E. (2018). The effect of dietary neonicotinoid pesticides on non-flight  
309 thermogenesis in worker bumble bees (*Bombus terrestris*). *J Insect Phys* 104, 33–  
310 39. doi:10.1016/j.jinsphys.2017.11.006.
- 311 Renner, M. A., and Nieh, J. C. (2008). Bumble bee olfactory information flow and  
312 contact-based foraging activation. *Insect. Soc.* 55, 417–424. doi:10.1007/s00040-  
313 008-1021-6.
- 314 Richardson, T. O., and Gorochowski, T. E. (2015). Beyond contact-based transmission  
315 networks: the role of spatial coincidence. *Journal of The Royal Society Interface* 12,  
316 20150705–11. doi:10.1098/rsif.2015.0705.
- 317 Rundlöf, M., Andersson, G. K. S., Bommarco, R., Fries, I., Hederström, V., Herbertsson,  
318 L., et al. (2015). Seed coating with a neonicotinoid insecticide negatively affects wild  
319 bees. *Nature* 521, 77–80. doi:10.1038/nature14420.
- 320 Sponsler, D. B., and Johnson, R. M. (2016). Mechanistic modeling of pesticide  
321 exposure: The missing keystone of honey bee toxicology. *Environ. Toxicol. Chem.*,  
322 1–42. doi:10.1002/etc.3661.
- 323 Stanley, D. A., Garratt, M. P. D., Wickens, J. B., Wickens, V. J., Potts, S. G., and Raine,  
324 N. E. (2015a). Neonicotinoid pesticide exposure impairs crop pollination services  
325 provided by bumblebees. *Nature* 528, 548–550. doi:10.1038/nature16167.
- 326 Stanley, D. A., Smith, K. E., and Raine, N. E. (2015b). Bumblebee learning and memory  
327 is impaired by chronic exposure to a neonicotinoid pesticide. *Sci. Rep.* 5, 16508.  
328 doi:10.1038/srep16508.
- 329 Straub, L., Williams, G. R., Pettis, J., Fries, I., and Neumann, P. (2015). Superorganism  
330 resilience: eusociality and susceptibility of ecosystem service providing insects to  
331 stressors. *Curr Opin Ins Sci* 12, 109–112. doi:10.1016/j.cois.2015.10.010.
- 332 Tan, K., Chen, W., Dong, S., Liu, X., Wang, Y., and Nieh, J. C. (2015). A neonicotinoid  
333 impairs olfactory learning in Asian honey bees (*Apis cerana*) exposed as larvae or  
334 as adults. *Sci. Rep.* 5, 1–8. doi:10.1038/srep10989.
- 335 Thorbek, P., Campbell, P. J., and Thompson, H. M. (2016). Colony impact of pesticide-  
336 induced sublethal effects on honeybee workers: A simulation study using  
337 BEEHAVE. *Environ. Toxicol. Chem.* 36, 831–840. doi:10.1002/etc.3581.
- 338 Tosi, S., and Nieh, J. C. (2017). A common neonicotinoid pesticide, thiamethoxam,  
339 alters honey bee activity, motor functions, and movement to light. *Sci. Rep.* 7,  
340 15132. doi:10.1038/s41598-017-15308-6.

- 341 Tosi, S., Démaries, F. J., Nicolson, S. W., Medrzycki, P., Pirk, C. W. W., and Human, H.  
342 (2016). Effects of a neonicotinoid pesticide on thermoregulation of African honey  
343 bees (*Apis mellifera scutellata*). *J Insect Phys* 93-94, 56–63.  
344 doi:10.1016/j.jinsphys.2016.08.010.
- 345 Tsvetkov, N., Samson-Robert, O., Sood, K., Patel, H. S., Malena, D. A., Gajiwala, P. H.,  
346 et al. (2017). Chronic exposure to neonicotinoids reduces honey bee health near  
347 corn crops. *Science* 356, 1395–1397. doi:10.1126/science.aam7470.
- 348 Whitehorn, P. R., O'Connor, S., Wackers, F. L., and Goulson, D. (2012). Neonicotinoid  
349 pesticide reduces bumble bee colony growth and queen production. *Science* 336,  
350 351–352. doi:10.1126/science.1215025.
- 351 Whitehorn, P. R., Wallace, C., and Vallejo-Marin, M. (2017). Neonicotinoid pesticide  
352 limits improvement in buzz pollination by bumblebees. *Sci. Rep.* 7, 1–8.  
353 doi:10.1038/s41598-017-14660-x.
- 354 Woodcock, B. A., Bullock, J. M., Shore, R. F., Heard, M. S., Pereira, M. G., Redhead,  
355 J., et al. (2017). Country-specific effects of neonicotinoid pesticides on honey bees  
356 and wild bees. *Science* 356, 1393–1395. doi:10.1126/science.aaa1190.
- 357 Woodcock, B. A., Isaac, N. J. B., Bullock, J. M., Roy, D. B., Garthwaite, D. G., Crowe,  
358 A., et al. (2016). Impacts of neonicotinoid use on long-term population changes in  
359 wild bees in England. *Nat Commun* 7, 12459. doi:10.1038/ncomms12459.
- 360 Wu-Smart, J., and Spivak, M. (2016). Sub-lethal effects of dietary neonicotinoid  
361 insecticide exposure on honey bee queen fecundity and colony development. *Sci.  
362 Rep.* 6, 32108. doi:10.1038/srep32108.
- 363
- 364

365 **Figure Captions**




366

367 **Figure 1. Parameter estimation and model description.** (A) Schematic diagram of model simulation,  
 368 including individual workers, nest structure (filled tan circles), nest center (blue circle), and physical  
 369 boundaries. (B) Model for simulated worker movement showing the random walk ( $v_{\text{rw}}$ ), nest attraction  
 370 ( $v_{\text{nest}}$ ), and resultant ( $v_{\text{res}}$ ) movement vectors. (C) Activity switching probabilities ( $p_{A \rightarrow I}$ ,  $p_{I \rightarrow A}$ ), are measured separately depending on whether bees are (D) located on or


372 off the nest structure or (E) in physical contact with a nestmate. (F-G) Changes in probability of switching  
373 from active to inactive ( $p_{A \rightarrow I}$ , F) and switching from inactive to active state ( $p_{I \rightarrow A}$ , G) as a result of  
374 imidacloprid exposure. Probability changes are shown separately for combinations of location (Loc) and  
375 social interaction state (Soc). For each box, upper values show parameter estimates for untreated bees,  
376 middle values (in parentheses) show the percent change in the estimated parameters in bees fed 1.0 ng  
377 imidacloprid (compared to controls), and lower values are p-values. Bold values indicate significant  
378 effects, with box color indicating the direction of change (red for decreasing, green for increasing). Data  
379 from Fig S5 in (Crall et al., 2018b).

380




382  
 383  
 384  
 385  
 386  
 387  
 388  
 389  
 390  
 391  
 392  
 393  
 394  
 395  
 396


**Figure 2. Effects of colony size on emergent behavior in bumblebee nests.** Mean behavioral metrics (left column) and relative differences between treated and control workers (right column) as a function of colony size and treatment intensity (% of workers treated). Data are shown for four metrics: (A, E) activity, (B, F) distance to the nest center ( $D_{\text{NC}}$ ), (C, G) proportion of time spent on the nest structure ( $P_{\text{nest}}$ ), and (D, H) interaction rate ( $R_{\text{int}}$ ). Values are the mean across all workers and replicate simulations at a given parameter set. The number of simulation iterations at each parameter set was varied with colony size (# of iterations =  $10000/N$ , where  $N$  = # of workers). Phase spaces smoothed using local averaging over adjacent values (smoothing window: +/- 10% for treatment intensity,  $(\sqrt{N-2})^2:(\sqrt{N+2})^2$  for  $N_{\text{nest}}/N_{\text{workers}}$ ). For all simulations, the number of nest structure elements ( $N_{\text{nest}}$ ) and the number of workers ( $N_{\text{workers}}$ ) were varied simultaneously to maintain a 1:1 nest structure:worker ratio. Differences between control and treated workers (right column) were calculated as a percentage of mean absolute value,  $\text{abs}([trt - ctrl]/\text{mean}(trt, ctrl)]) * 100$ , where  $trt$  denotes the treated workers and  $ctrl$  denotes the control group.



**Figure S1. Independent effects of nest structure size and worker number on emergent behavior in bumblebee nests.** Mean behavioral metrics when the number of nest structural elements ( $N_{\text{nest}}$ , left column) and number of workers ( $N_{\text{workers}}$ , right column) are independently varied. Data are shown for four metrics: (A, E) activity, (B, F) distance to the nest center ( $D_{\text{NC}}$ ), (C, G) proportion of time spent on the nest structure ( $P_{\text{nest}}$ ), and (D, H) interaction rate ( $R_{\text{int}}$ ). Values are the mean across all workers and replicate simulations at a given parameter set. The number of simulation iterations at each parameter set was varied with colony size (# of iterations =  $10000/N$ , where  $N$  = # of workers). Phase spaces smoothed using local averaging over adjacent values (smoothing window: +/- 10% for treatment intensity,  $(\sqrt{N}-2)^2:(\sqrt{N+2})^2$  for  $N_{\text{nest}}/N_{\text{workers}}$ ). For each metric, identical color scales are used for the left and right columns.




412 **Figure S2. Independent effects of nest structure size and worker number on relative effect size of**  
 413 **imidacloprid.** Relative difference between treated and untreated workers when the number of nest  
 414 structures ( $N_{\text{nest}}$ , left column) and number of workers ( $N_{\text{workers}}$ , right column) are independently varied.  
 415 Data are shown for four metrics: (A, E) activity, (B, F) distance to the nest center ( $D_{\text{NC}}$ ), (C, G) proportion  
 416 of time spent on the nest structure ( $P_{\text{nest}}$ ), and (D, H) interaction rate ( $R_{\text{int}}$ ). Differences calculated as in  
 417 Fig 2. The number of simulation iterations at each parameter set was varied with colony size (# of  
 418 iterations =  $10000/N$ , where  $N$  = # of workers). Phase spaces smoothed using local averaging over  
 419 adjacent values (smoothing window: +/- 10% for treatment intensity,  $(\sqrt{N-2})^2:(\sqrt{N+2})^2$  for  
 420  $N_{\text{nest}}/N_{\text{workers}}$ ). For each metric, identical color scales are used for the left and right columns.  
 421




422  
423 **Figure S3. Absolute differences between treated and control workers across treatment intensity**  
424 **and colony size.** Simulations are the same as in Fig 2. Relative differences for (A) activity,  
425 (B) distance to the nest center ( $D_{\text{NC}}$ ), (C) proportion of time spent on the nest structure ( $P_{\text{nest}}$ ), and (D)  
426 interaction rate ( $R_{\text{int}}$ ) are shown as the absolute difference (rather than % difference in the right column of Fig S2).  
427

# Agent-based model (BeeNestABM)




## Activity state parameter estimation (empirical)

**C**




**D**

Location (Loc)



**E**

Social interaction state (Soc)



**F**

$p_{A \rightarrow I}$

	0.031 (+163%) $<< 0.01$	0.021 (+250%) $<< 0.01$
Off-nest		
On-nest	0.012 (+125%) $<< 0.01$	0.009 (+132%) 0.11

Spont-  
aneous

Socially-  
modulated

**G**

$p_{I \rightarrow A}$

	0.21 (-52%) $<< 0.01$	0.27 (-48%) $<< 0.01$
Off-nest		
On-nest	0.29 (-13%) 0.11	0.34 (-30%) 0.14

Spont-  
aneous

Socially-  
modulated

Figure 02.JPG

## Colony Size ( $N_{\text{workers}} = N_{\text{nest}}$ ) model

