

Article

Honey bees' behavior is impaired by chronic exposure to the neonicotinoid thiacloprid in the field

Lea Tison, Marie-Luise Hahn, Sophie Holtz, Alexander Rößner, Uwe Greggers, Gabriela Bischoff, and Randolph Menzel

Environ. Sci. Technol., **Just Accepted Manuscript** • DOI: 10.1021/acs.est.6b02658 • Publication Date (Web): 07 Jun 2016

Downloaded from <http://pubs.acs.org> on June 9, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

1 **Honey bees' behavior is impaired by chronic exposure to the neonicotinoid thiacloprid**
2 **in the field**

3

4 Léa Tison^{1*}, Marie-Luise Hahn¹, Sophie Holtz¹, Alexander Rößner¹, Uwe Greggers¹, Gabriela
5 Bischoff² and Randolph Menzel¹.

6

7 ¹ *Free University Berlin, Institute for Biology-Neurobiology, D-14195 Berlin, Germany;*

8 ² *Julius Kühn-Institut, Institute for Bee Protection, D-14195 Berlin, Germany.*

9

10

11

12 *** Corresponding author:**

13 Léa Tison

14 Tel: +49 308 385 6284

15 Email: lea.tison@gmail.com

16

Abstract

The decline of pollinators worldwide is of growing concern and has been related to the use of plant protecting chemicals. Most studies have focused on three neonicotinoid insecticides, clothianidin, imidacloprid and thiamethoxam, currently subject to a moratorium in the EU. Here we focus on thiacloprid, a widely used cyano-substituted neonicotinoid thought to be less toxic to honey bees and of which use has increased in the last years. Honey bees (*Apis mellifera carnica*) were exposed chronically to thiacloprid in the field for several weeks at a sublethal concentration. Foraging behavior, homing success, navigation performance, and social communication were impaired, and thiacloprid residue levels increased both in the foragers and the nest mates over time. The effects observed in the field were not due to a repellent taste of the substance. For the first time, we present the necessary data for the risk evaluation of thiacloprid taken up chronically by honey bees in field conditions.

29 **Introduction**

30 Bees, including honey bees, bumble bees and solitary bees represent the most prominent
 31 group of pollinators worldwide and contribute largely to agriculture as 35 % of the food crop
 32 production depends on them¹. The recent loss of pollinator populations can be attributed to multiple
 33 factors such as habitat loss and fragmentation, colony management, bee pests and parasites, and
 34 additional environmental and anthropogenic elements. Doubtlessly the use of pesticides for crop
 35 protection contributes to the loss of pollinator abundance both at the species level and the quantity of
 36 a particular species^{2,3,4}. It has also become evident that neonicotinoids (and other insecticides like
 37 fipronil) play a crucial role as the promoters of pathogen and parasite infections that effectively drive
 38 colony losses^{5,6,7}. Thanks to their systemic properties, neonicotinoids are present in the pollen and
 39 nectar and will thus be continuously collected by pollinators for as long as flowering persists. They
 40 are agonists of nicotinic acetylcholine receptors (nAChR) which are normally activated by the
 41 neurotransmitter acetylcholine⁸. Nicotinic synaptic transmission is a major component of neural
 42 integration in the circuits related to sensory integration and functions related to the mushroom bodies,
 43 mediating multisensory integration, learning, and memory formation^{9,10}. Neonicotinoids negatively
 44 affect the mushroom bodies' physiology¹¹ and function¹² in honey bees. It was already proven that
 45 neonicotinoids compromise olfactory learning¹³ as well as the ability of worker bees to forage and to
 46 communicate^{14,15,16,17}. The toxicity of sublethal doses is also expected to be reinforced over time^{18,19}.
 47 However, a detailed analysis of the chronic exposure to thiacloprid on foraging, navigation, and social
 48 communication is lacking.
 49 The cyano-substituted neonicotinoid thiacloprid is declared less toxic to bees than nitro-substituted
 50 compounds like imidacloprid and thiamethoxam^{20,21,22,23}. The formulations based on thiacloprid are
 51 registered and sold in more than 70 countries worldwide²⁴ and act against sucking and chewing pest
 52 insects of more than 50 crops^{25,26}. The formulations based on thiacloprid are used in the field for
 53 spraying treatment at application rates much higher than for the 3 neonicotinoids suspended in Europe
 54 ^{21,27}. These formulations are allowed to be sprayed during flowering because less damage to
 55 pollinators is expected. Thiacloprid is also used in a maize seed treatment since the withdrawal of
 56 clothianidin and thiamethoxam on maize across Europe in 2013.

57 Toxicity tests performed by the company at the time before releasing thiacloprid on the market
58 evaluated only the short term and lethal effects on worker honeybees. In contrast to acute effects, no
59 standardized protocol exists for measuring chronic effects on individual bees under semi natural
60 conditions²³. The value of tests on single animals has been questioned because a whole colony may be
61 more robust to pesticide exposure²⁹. However, honey bees are acting as single animals during
62 foraging; they need to adjust their behavior to the changing availability of food sources, return to the
63 colony for survival, deliver the collected food and communicate with other foragers. Therefore,
64 testing single foraging honeybees represents best conditions faced by honey bee foragers and other
65 insect pollinators in nature. A few lab studies have shown that chronic exposure to sublethal doses of
66 thiacloprid affects honey bees' sensitivity to the gut pathogen *Nosema cerenae*^{30,31,32} and a field study
67 has shown that navigation is compromised when thiacloprid was given as a single acute dose³³.
68 Chronic and sub-lethal exposure to the substance is the most likely exposure scenario in the field^{26,34}
69 but no field study to our knowledge has yet uncovered any specific behavioral effect of such condition
70 of exposure. In our experiments honey bee foragers were exposed chronically for several weeks in the
71 field to a concentration similar or lower to those used in previous chronic exposure studies with
72 thiacloprid^{30,31,32}. The concentration of thiacloprid in the contaminated sucrose solutions was 5.4 ng/μl
73 whereas the concentration of thiacloprid in the formulation Calypso® directly sprayed on plants and
74 flowers at a distance of 30 to 40 cm is 150 ng/μl.
75 Since most of the collected sucrose solution will be deposited by the forager inside the hive, and only
76 a small proportion will be taken up and metabolized by the bee during its return flight from the feeder
77 to the hive, only a small amount of thiacloprid will reach the brain and interfere with nicotinic
78 synaptic transmission.
79 We found that a chronic exposure to thiacloprid significantly impaired honeybees' foraging
80 behaviour, communication, and navigation. The substance increased in the foragers over time
81 affecting also the animals indirectly exposed in the colony. We found no avoidance of or preference to
82 the substance, supporting the idea that a neural impairment was responsible for affecting the honey
83 bees' abilities to forage, communicate, and navigate rather than a repelling effect.

84

85 **Material and methods**

86 **Preparation of the solutions**

87 Stock solution: 10 mg thiacloprid ([3-[(6-chloro-3-pyridinyl) methyl]-2-thiazolidinylidene]
88 cyanamide, Sigma-Aldrich Pestanal) diluted in 1 mL acetone ($\geq 99.9\%$, Sigma-Aldrich) plus 39 mL
89 distilled water leading to a concentration of 0.25 g/L. Acetone was chosen as the solvent following the
90 EPPO guidelines³⁵. The final concentration of acetone (0.05 %) in the contaminated sucrose solutions
91 was shown to not have an effect on honeybee navigation³³. The thiacloprid sucrose solutions used in
92 the field (0.02 mM, 4.5 ppm) as well as for the taste and choice experiments (0.025 mM, 5 ppm) were
93 freshly made every morning from the stock solution. The concentration of thiacloprid at the treated
94 feeder was always the same regardless of the sucrose solution concentration. The concentration of the
95 solutions used were confirmed by LC-MS/MS (Methods S1).

96

97 **Field experimental design**

98 The experimental area is a highly structured agricultural landscape (trees and bushes, pathways, creek,
99 grass fields, etc) nearby Großseelheim, Germany. Two colonies housed in two observation hives
100 (W.Seip, Bienenzuchtgerätefabrik) were put up on two opposite sides of a cabin at the western border
101 of the experimental area (50°48'51.9"N). Each colony of *Apis mellifera carnica* was equipped with
102 one comb of sealed brood plus newborn bees and one comb of food (Deutsch Normal Mass combs)
103 originating from the same honey bee colony. The queens were kindly provided by the Bieneninstitut
104 Kirchhain, they derived from selected breeder colonies of the carnica breeding population of the
105 institute. They were open mated and aged 1 year old. Sister queens were used in an attempt to keep
106 the genetic difference among the honey bee individuals from each colony at a low level.

107 Training to the feeders

108 Two feeders (F1 and F2) were established 350 meters northeast and 340 southeast respectively and
109 were separated by an angle of 90° as seen from the cabin. The release site (RS) was located 780
110 meters east of the cabin. A group of foragers from each of the two colonies was trained to its
111 respective feeder and marked individually with number tags. The origin of each newly marked bee
112 from the two colonies was controlled at the respective hive entrance. In Experiment 1, one group of

bees (treated group) foraged during 19 days on a sucrose solution containing thiacloprid (4.5 ppm), and the other group (control group) foraged over the same time at a feeder containing only sucrose solution. In Experiment 2, the control hive became the treated hive and the treated hive was removed and replaced by a new control hive. The feeders' locations were exchanged between Experiment 1 and 2 in order to exclude any possible landscape effect related to the feeders' position. In Experiment 2, the two groups of foragers were feeding at their respective feeder during 29 days. Each feeder was placed in a little wooden box to allow counting the entrances and exits of foragers with a retro-reflective sensor (Baumer GmbH). The total number and the identity of bees visiting their feeder throughout each day was known as well as the amount of sucrose solution consumed at both feeders. The concentration of the sucrose solution at each feeder was adjusted during the day in order to regulate the traffic at the feeder (25 - 40 bees) following evaluation by the experimenter of the number of trained foragers visiting the feeder. Dance recruitment was induced 19 times on 19 different days (time: 1500 - 1700 hours) by first halving the sucrose concentration at both feeders for one hour and then increasing it twofold for another hour.

Homing experiment

Colonies were settled in the field for at least a week before the homing experiments started. After a certain number of days foraging at the feeders, single bees were caught on their departure at their respective feeder and transferred into a glass vial after they had freely drunk either a 1 M sucrose solution (control bees) or a 1 M sucrose solution containing 4.5 ppm thiacloprid (treated bees). They were kept in the dark for 45 min while they were transported to the release site. Then a transponder was fixed to thorax and the bee was released (time: 1100 - 1700 hours, temperature: 17-30°C, wind < 15 km/h). No release was made when the sky was evaluated too cloudy or totally overcast, nor when it was raining. Care was taken that the number of control and treated bees released every day were evenly distributed and it was ensured that each bee was released only once. The radar was shut down not before 120 min after the last bee was released if the bee was not yet back to its hive. Since none of the bees that did not return to the hive after being released was seen at the feeder or at the hive entrance on the same or the following days, we assume that they died in the field.

The method used for tracking bees with a harmonic radar system has been described before^{36,37,38}. The transponders were built by ourselves following the procedure from Riley et al. (1996), their attachment and carrying by the bees do not alter honeybees' flight behavior^{39,40}. The flights of the released bees carrying a transponder were monitored using the radar system over a distance of up to 900 meters radius and at a temporal resolution of 1/3 Hertz³⁷.

Electric field recordings

The electric fields emitted by dancing bees⁴¹ consist of low-frequency (movements of the abdomen, 16 Hz on average) and high-frequency (buzzing of the wings, 230 Hz) components synchronization, leading to an average of three to seven electric pulses per waggle. The distance from the hive to a feeding site is encoded in the number of waggle runs and 1 sec is known to represent a distance of about 1 km⁴². The feeders were located 350 meters northeast (F1) and 340 southeast (F2) of the hives and since very few natural food sources existed in the experimental area and none of them were present at the same distance as the feeders, the distinction between dances from trained and untrained foraging bees was possible. Electric field measurements were performed at the same time on both sides of the lower comb in the control and treated hives using 4 copper wires with a silver coating positioned in the dance area (12 cm² covered), connected on each side to a stereo amplifier (USB - Soundbox 7.1, Conrad electronics SE) with a sample rate of 44.1 KHz. Each amplifier was connected to a laptop and the software Presonus Studio One (version 2.4) was used for saving the data as wave files. We recorded in total 340 hours of electric fields on 32 different days (average of 2.67 hours per day).

Thiacloprid residues analysis

Bees were caught at their feeder after foraging for a certain number of days and after they had filled their crop with a 1 M sucrose solution contaminated or not. They were then kept in the dark for 45 minutes before being killed by chilling and put into a -20° C deep-freezer. We also collected unmarked forager bees at the entrance of the treated and control hives when flying out on a foraging trip in order to assess the in-hive contamination of foragers not visiting the feeders but exposed

indirectly to thiacloprid inside the hive via the stored food. See Methods S1 for details about the residue analysis by LC-MS/MS.

Repellent effect

PER experiment

The Proboscis Extension Response (PER) was used to sample hungry bees' sensitivity to varying concentrations of sucrose^{43,44} containing or not thiacloprid (5 ppm). Honeybees were captured at 1400 hours when leaving the hive, immobilized by chilling, and mounted in small brass tubes which restrained body movements but allowed the antennae and the mouthparts to move freely⁴³. One hour later they were tested in the laboratory by touching both antennae with a droplet of ascending concentrations of sugar concentrations (dry sugar diluted in tap water + 0.05 % acetone, 0.1 %, 0.3 %, 1 %, 3 %, 10 %, 30 % and 50 %, w/v). Only the bees which showed a PER for the 50 % sugar concentration were considered as the non-responders (control: 1/74, treated: 3/74) were considered physically unable to extend their proboscis.

Choice experiment

In May, a group of bees was trained to a training/feeding platform located about 30 meters from the hive. The platform was composed of a yellow background and 10 blue squares randomly distributed and containing a mini-feeder from which the bees could freely drink a 1 M sucrose solution. The test platform contained only 6 mini-feeders. During testing of single bees three feeders contained 8 µl of a 1 M control sucrose solution each and the other three 8 µl of a 1 M sucrose solution with thiacloprid (5 ppm) each. The positions of the control and treated mini-feeders were randomly allocated on the platform. The number of feeders drunk and the time a bee took to drink at each of the 6 feeders was recorded. At the end of the test the bee was killed and the same test was repeated with a new naive bee.

Flight tracks and statistical analysis

From the x/y coordinates collected by the radar, the length and duration of the flight from the first to the last signal was calculated. The x/y-coordinates were fitted into a google map scaled in meters

using CorelDraw.X5. The criteria used to categorize the different flight parameters were arbitrarily defined. A “vector flight” was considered as such when fitting into an angle of 45° as seen from the release site ($\pm 22.5^\circ$ each side of the feeder-hive vector direction, F1: 313° , F2: 222°) and had a minimal length of 200 m. The angle of a vector component is the angle between the crossing point of the vector track with the 200 m circle around the release and the direction towards north. The criterion “pass close to F” and “Return to RS” was attributed respectively to bees getting closer than 100 m from their feeder or from the release site during their flight.

The electric field data were transformed to SMR files, preliminary filtered in Spike 2 (version 8.03) and further analyzed using custom-made programs written in Visual Basic 2013 (Microsoft). An amount of 6 ± 2 waggles per run (about 360 ± 120 meters) was used as a criteria to select the dances indicating the location of the feeders. If the number of waggles per run was exceeding this range, the waggle runs were attributed to the “other bees” group.

For the statistical analysis of the data, we used R and Prism 5 and 6. The normality of the data was tested using the D'Agostino-Pearson omnibus test. If the data were normally distributed, we used a paired/unpaired t.test or an analysis of variances with Tukey's post-hoc tests. Otherwise non-parametric tests were performed (Mann-Whitney test, Wilcoxon signed rank test). The Fischer's Exact Test was used to compare proportions. . For the PER data we performed a mixed effects logistic regression in R (lme4 package) with “Bee” and “Date” as random effects to account for the difference between individuals and the date. This was followed by Overall Likelihood Ratio Tests and Tukey's post-hoc tests (multcomb package). The Wheeler-Watson test was used to calculate the angular distribution of the vector components. The survival analysis was conducted using censored Kaplan Meier Log-Rank in R and the influence of multiple variables was investigated using a Cox-regression model. The numbers of bees tested for each experiment and test groups are indicated in the legends of the figures and in the text.

Results

Honey bees' foraging behavior and dance communication are compromised by chronic exposure to thiacloprid.

223 The total foraging span of honey bees foraging at the control feeder was significantly longer than the
224 foraging span of honey bees foraging at the treated feeder (Table 1, Kruskal-Wallis, $P < 0.0001$).
225 Control bees foraged at their feeder on average 0.78 days longer than treated bees (“Total”, Table 1).
226 The significance was different between the groups according to the Experiment (see Table 1).
227 Sucrose consumption at the control and treated feeder was significantly different in both experiments
228 (Paired t-test, $P < 0.0001$). Control bees consumed 1.7 times more sugar solution per day than treated
229 bees (Table S1). The average amount of thiacloprid collected per bee and per day at the treated feeder
230 was estimated at 12118 ± 900 ng in Experiment 1 and 10990 ± 833 ng in Experiment 2 (Table S1).
231 Treated bees performed on average 1.8 times and 1.4 times less foraging trips per day than control
232 bees in Experiment 1 and 2 respectively. On one trip, we estimate that a bee collected on average 216
233 ng of thiacloprid (40 μ l of solution). The total amount of thiacloprid metabolized by a bee per day
234 during the return flights to the hive ranges between 141 and 212 ng (Table S1). This calculation is
235 based on the data related by Rortais et al.⁴⁵ that a bee needs 8 - 12 mg of sugar per hour to fly^{45,46} and
236 on our measurements (treated bees collected on average 1 M sucrose solution and flew on average 2
237 minutes from the feeder to the hive).
238 The reduced sugar consumption is linked to a reduced visitation rates of foragers at the contaminated
239 feeder. Indeed, treated bees visited their feeder significantly less frequently than the control bees and
240 higher sucrose concentrations were needed at the contaminated feeder in order to keep the bees
241 visiting the feeder (Fig. 1 a). The median sucrose concentration used for regular foraging was 0.5 M at
242 the control feeder and 1 M at the treated feeder. Recruitment of foragers via the waggle dance was
243 induced by raising the sucrose concentration at the feeder⁴². First the sucrose concentration at both
244 feeders was reduced to half of the current concentration for one hour, then it was increased twofold
245 for another hour. Sucrose concentrations as high as 2 M during dance induction did not significantly
246 increase the traffic at the treated feeder (ANOVA, $F_{3,72} = 14.01$, $P < 0.0001$), whereas a median
247 concentration of 1 M increased significantly the number of visits at the control feeder ($p < 0.05$, Fig.
248 1b).

Reduced recruitment at the feeder could indicate less waggle dances or compromised dance performance. Therefore, we monitored and estimated the number of waggle runs performed by the dancing bees in both colonies, taking advantage of the fact that waggle dances can be measured by the temporal modulation of the electrostatic field emanating from the dancing bee⁴¹. The number of waggles performed by the bees trained to the control feeder was significantly higher than those of the bees trained to the contaminated feeder (Fig. 2, Wilcoxon signed rank test, $p < 0.0001$) although the sucrose concentration during dance induction was higher at the contaminated feeder (Fig. 1.a). Indeed, honey bees foraging at the control feeder performed on average 3.2 times more waggles per hour than honey bees foraging at the treated feeder. The reduced dance activity of treated bees explains the lower foraging activity at the contaminated feeder.

We also differentiated dances for feeders and dances to unknown natural food sources on the basis of the number of waggle runs as indicators of distance to the respective food source^{41,42}. We found significantly lower dance activity advertising for natural food sources in the treated colony (Fig. S1) indicating that the accumulation of thiacloprid inside the colony also affected bees that did not forage at the contaminated feeder but were on contaminated stored food.

No repellent effect of thiacloprid.

One explanation for lower foraging activity found in treated bees could be an aversive taste of the substance in contaminated sucrose solution. In the laboratory experiment, we tested the proboscis extension response (PER) of hungry foragers to water and 7 different sucrose concentrations (0.1 %, 0.3 %, 1 %, 3 %, 10 %, 30 % and 50 % w/v) containing thiacloprid (5 ppm) or not (Fig. 3). No difference was found in the PER of bees stimulated either with the control sucrose solutions or the contaminated sucrose solutions (logistic regression with random effects “Bee” and “Date”, Sugar concentration x Treatment: $\chi^2_6 = 2.5224$, $P = 0.866$). The results of the Tukey’s post-hoc tests between the control and treated groups for each of the different sucrose concentrations tested can be found in Table S2.

In the free flight experiment, 45 bees had to choose between feeders containing a 1 M sucrose solution with or without thiacloprid (5 ppm). No significant difference was found in the visitation rate

of the bees to the control (64 %) and contaminated (65 %) feeders (n=135 feeders, Fischer Exact test, $P = 0.8989$). The average (\pm s.e.m.) drinking time per bee and feeder was 6.88 ± 0.27 sec at the control feeders, and 7.37 ± 0.36 sec at the contaminated feeders (no significant difference, Mann Whitney, $P = 0.5578$). These results rule out the possibility that thiacloprid has a repellent taste for honeybees.

Thiacloprid residue levels increase in foragers.

The amount of thiacloprid in bees foraging at the contaminated feeders in Experiment 1 and 2 was analyzed by LC-MS/MS (Methods S1). Fig. 4 shows how it accumulated in different body parts over time. The amount of thiacloprid residues found in bees can be seen as the status of intoxication at the moment a bee is released with a transponder after foraging chronically during 2, 3 or 4 days at the contaminated feeder.

The length of exposure of the foragers at the contaminated feeder as well as the amount of thiacloprid collected is related to the amount of residues found in the bees (Fig. 4, Table S3). The more foraging trips honey bees performed to the treated feeder in a certain number of days, the higher was the cumulated amount of contaminated sucrose solution collected and the higher was the amount of thiacloprid residue found in the bees. Only a fraction of the cumulated total amount of thiacloprid collected by the bees at the feeder will be metabolized and most of this uptake will happen during their return flights from the feeder to the hive. This fraction was found very close to the amount of thiacloprid residues found in bees after a defined number of days foraging at the contaminated feeder (Table S3).

In-hive contamination was assessed by collecting unmarked forager bees at the entrance of the treated hive when flying out on foraging trip. Thiacloprid was found in these bees but at much lower amounts than in the foragers trained to the contaminated feeder (Table S3). Indeed, these foragers did not visit the contaminated feeder but they were exposed to thiacloprid inside the hive via the food collected and stored by the foragers visiting the contaminated feeder. Since their waggle dance activity was significantly reduced (Fig. S1) even these low levels of thiacloprid impaired social communication.

Honey bees' homing success and navigation performance are impaired.

Navigation requires the integration of multisensory cues and the retrieval of appropriate memory about the landscape structure. We tested navigation abilities of the bees trained to feeder 1 and 2 during the Experiments 1 and 2. We found that treated bees returned to their hive at a significantly lower proportion than control bees (Fig. 5, homing success: control 91.76 %, treated 76 %, Fischer Exact Test, $P < 0.01$). Based on the crop-emptying measurements by Fournier et al.⁴⁷ we calculated that the foragers released with a transponder could have assimilated in 45 min up to 7 μ l and thus 38 ng thiacloprid in addition of the residues already assimilated over n days foraging at the feeder. This value is a higher estimate because the amount of assimilated sucrose during the 45 minute waiting time may well be much lower depending on the activity of the waiting bee⁴⁸. In any case the partial acute treatment component involved in the navigation experiments adds to the chronic effect. A survival analysis was conducted on the data and a significant influence of thiacloprid on honey bee homing success was found (Kaplan Meier Log Rank test, $\chi^2_1 = 12.9$, $P < 0.001$). For the survival analysis, a flight duration of 120 min was settled for bees that flew out of the radar range and did not come back within the radar range or to the hive during this time. The flight duration of all other bees was the flight time in minutes from the release site to the hive or from the release site to a point inside of the radar range where the signal was lost. The influence of multiple variables was tested in a cox-regression model (Table 2). The variable "Treatment" shows a significant negative effect on honey bee survival. The hazard rate of the treated bees, representing the likelihood of returning to the hive, is almost half the hazard rate of the control bees. The period during which the experiment was performed ("Experiment"), the number of days a bee foraged at its feeder before being released ("Time foraging"), as well as the number of days from the first day of the experiment until a bee was released ("Time exposure") had no significant effect on honey bee homing abilities. The duration of the exposure had no effect possibly because 45 % of the treated bees individually released foraged at the contaminated feeder for less than 3 days. The temperature at the release time did not seem to play a role in the ability of honey bees to come back to their hive. At their release, 76.5 % of the control honey bees and 61 % of the treated honeybees waited for a short time at the release site before starting to fly. This waiting time ("Time before flying") was not different between the control and the treated

bees (mean \pm s.e.m control = 3.17 ± 0.33 min, treated = 4.53 ± 0.69 min, Mann Whitney, $P = 0.5067$) and had no influence on the homing success (Table 2).

During the flight, 9 pauses were recorded in the control group and 24 in the treated group with a maximum of 3 pauses per bee (Table S5). The probability of making a pause during the return flight to the hive was not found significantly different between the control (13 %) and treated groups (24 %, Fischer Exact test, $P = 0.0617$). However, the mean (\pm s.e.m.) pause duration was higher for the treated bees (20.13 ± 5.28 min) than for the control bees (5.29 ± 2.12) but not significantly different between the two groups (Mann Whitney, $P = 0.0974$) possibly because of the limited number of cases and the large variance. The duration of the pause was deleted from the total flight duration in order to calculate an accurate flight speed (Tables S4 and S5). The total flight duration including pauses was however considered for every other analysis. If we take out the duration of the pauses from the total flight duration of the concerned bees and run the survival analysis again, the variable “Treatment” remains significant (Kaplan Meier Log Rank test, $\chi^2_1 = 8.8$, $P < 0.01$; cox regression Model 1: $P = 0.00435$) and none of the other variables tested before become significant.

Among the bees returning to their respective hives, no significant difference was found between the flight duration of control and treated bees (Table S4, median control = 7.8 min, treated = 7.4 min, Mann Whitney, $P = 0.5741$), and no significant difference was found in the distance flown (Median control = 2032 m, treated = 1908 m, Mann Whitney, $P = 0.4778$). However, the treated bees flew significantly slower than the control bees (Table S4, mean \pm s.e.m., speed treated = 4.32 ± 0.13 m/s, control = 4.78 ± 0.15 m/s, Unpaired t-test, $P < 0.05$). In a catch and release situation like in the test performed here, bees usually fly first along a vector they would have taken if they were departing from the feeder in direction to the hive (vector flight)⁴⁹. Then they usually search for some time before flying back to the hive rather straightly. The proportion of vector flights performed did not differ between the control ($n = 55$, 71 %) and treated ($n = 57$, 76 %) bees which returned to their hive (Fischer Exact test = 0.4703). There was a difference in the duration of the vector component between the control bees in Experiment 1 and 2 ($P < 0.05$). Also, control bees from Experiment 2 flew the vector component faster than control bees from Experiment 1 and treated bees from Experiment 2 ($P < 0.01$ and $P < 0.05$ respectively). Since these bees foraged at different feeding locations the effect

indicates a site specific component. Therefore, we compared the parameters of the flights of control and treated bees separately for the two training sites, and found no differences with respect to the duration, length and the spatial distribution of the vector component (Table S5). The homing flight was considered as the flight component from the end of the vector to the hive. No difference was found in the length, duration, or speed of the homing flight between control and treated bees (Table S5). However, we found that more control bees returned less than 100 m from their release site at least once during their search flight (Fisher Exact test, $P < 0.05$) indicating their ability to remember where they were released and use this location to start over the homing flight. Also, significantly more control bees flew less than 100 meters close to their feeder (Fisher Exact test, $P < 0.01$) before heading to the hive indicating the use of known landmarks for a successful homing. Indeed, all the bees which passed close to their feeder flew directly back to the hive from the feeder. The bees which did not return to the hive performed different kinds of flight trajectories before getting lost (Fig. 6). None of the control bees got lost out of the radar range whereas 9 treated bees out of 20 were lost bees in experiment 2 and flew in the opposite direction of the hive, left the radar range and did not return within the range or to the hive. Interestingly, some treated bees (Fig. 6 c) terminated their flights at the end of the vector component. These bees did not initiate search flights or homing flights and did not arrive at the hive.

Discussion

Our study documents important sublethal effects of a low concentration (4.5 ppm) of thiacloprid taken up chronically by foraging bees. We found that higher-order functions like navigation according to a learned landscape memory, motivation to forage and to communicate in a social context were compromised. Honey bees visiting a feeder containing thiacloprid foraged over shorter periods of time probably because they died earlier than the control bees. This result is not surprising, since a 10-day exposure to a sublethal concentration of another neonicotinoid, thiamethoxam, reduced honey bees' life span by 41 %⁵⁰. Exposure to pesticide residues in brood comb was also shown to shorten adult longevity⁵¹. Overexpression of the vitellogenin transcript in the honey bee brains could be one of the molecular

indicators for the alteration in foraging activity and accelerated aging upon neonicotinoid exposure⁶. Previous studies also demonstrated a reduced foraging activity of honey bees on sucrose solutions contaminated with thiacloprid⁵², imidacloprid^{15,53,54}, or clothianidin¹⁴. These effects could be explained by a prolonged stay inside the hive before returning to the feeder¹⁴. We found that if occurring, a prolonged stay inside the hive was not used for dance communication, as dance activity was highly affected by a chronic uptake of thiacloprid, as already shown with imidacloprid¹⁵. We tried to compensate for the reduced foraging activity by increasing the sucrose concentration at the contaminated feeder, but the reduced dance activity could not be totally compensated for even though very high sucrose concentrations were applied during the dance induction periods. Thiacloprid increased the minimum sucrose concentration that honey bee foragers are willing to gather at the feeder as was found for imidacloprid¹⁵. Since increasing sucrose concentration could partially compensate for the reduced foraging activity observed at the contaminated feeder, it is most likely that thiacloprid did not alter the sensory or motor components of foraging but rather the motivation to forage. The results on dance performance point in the same direction. Pollination would be disturbed because of a reduced visitation of the flower by bees²⁸ leading to less flowers pollinated and thus reduced yields for farmers. In addition, honey bee colonies may suffer from a reduced food inflow, making them more susceptible to other disturbances (weather conditions, additional pesticides intoxication, parasites and pathogens).

Several studies reported low toxicity of thiacloprid^{20,55}. Laurino et al.⁵⁵ reported that acute uptake of thiacloprid (144 ppm) appeared to be not dangerous unless the honey bees were starved. It was thus suggested that thiacloprid acts as a repellent leading to reduced uptake and thus to lower toxicity. Here we disprove this hypothesis, documenting that thiacloprid does not have a repellent effect on honey bees. Furthermore, we show drastic effects on honey bee behavior for a concentration 32 times lower than the one used by Laurino et al. The results of our field study, especially the impairment of the foraging behavior and social communication, cannot be related to an avoidance of the substance, corroborating recent findings with other neonicotinoids⁵⁶.

The chronic exposure to thiacloprid lead to an accumulation over time in both the honey bee foraging at the contaminated feeder as well as in bees of the same colony via a contamination of the stored

417 food. The estimated amount of thiacloprid metabolized by a foraging honey bee can be estimated by
418 the energy supply necessary to perform the return trips from the feeder to the hive assuming that all
419 energy for the return flight is taken up from the collected sucrose solution. Applying a concentration
420 of 5.4 ng/μl at the feeder, we calculated that a foraging bee collected on average 216 ng of thiacloprid
421 (40 μl of solution) on one trip (80 times less than the acute oral LD50^(48h) of 17320 ng a.s per
422 bee). Based on the data about metabolic rates in flying bees^{45,46} the bee will metabolize only 0.53 - 0.8
423 μl of the sucrose solution and thus incorporates 2.86 - 4.32 ng thiacloprid while flying back to the
424 hive from the feeder (2 min return flight, 1 M sucrose solution). In natural conditions, foraging bees
425 can be exposed to different concentrations of the substance in nectar. Pohorecka et al.⁵⁷ report data on
426 thiacloprid residues in nectar from flowers, combs and in honey up to 208.8 ng/g. The amount of the
427 substance a bee will metabolize when foraging on nectar sources contaminated with 208.8 ng/g (0.25
428 ng/μl) thiacloprid depends on the distance from the food source to the hive, the flight time during
429 foraging, the motivational state⁴⁶ and the reward rate^{46,47}. If a bee performs a 20 minutes foraging
430 flight and forages on a 50 % nectar concentration, we can estimate that it will metabolize rather
431 similar amounts of thiacloprid (2.6 - 4 ng) as in our study.”

432 Furthermore, we estimated an amount of metabolized thiacloprid between 141 and 212 ng per day and
433 per bee foraging at the contaminated feeder. The lower range of this estimation, which is the most
434 probable, is not far from the daily consumption and thus exposure of 112.1 ± 4.4 ng per bee and per
435 day measured by Vidau et al.³² in his experiment.

436 Homing flight performance has been considered by the EFSA as a relevant criterion for measuring
437 sublethal effects in free-ranging pollinators²¹. Indeed, in order to perform a successful homing flight, a
438 bee has to use its sensory, motor and cognitive functions for successful foraging trips. We showed
439 here that the sensory and motor functions are not compromised but rather specifically their cognitive
440 abilities, such as retrieval of spatial memory about the landscape and motivation to forage and
441 communicate. The homing success of the foragers exposed to thiacloprid was impaired, supporting
442 previous findings on the effects of thiacloprid, imidacloprid, clothianidin³³ and
443 thiamethoxam^{16,29}. Honeybee colonies are behaving like a ‘superorganism’⁵⁸ and a sufficient number
444 of honey bees in each class is needed to perform the various and different tasks in order to keep the

information flow going and to adapt efficiently to changing environmental conditions⁵⁹. High forager death rates can induce a shift in the age that honey bees are starting to forage⁶⁰ and a change in the relative proportions of worker brood versus drone brood production²⁹ which might affect the fitness of the colony⁵⁹.

The radar tracking method applied here allows identification of which components of navigational tasks necessary for successfully return to the hive are compromised. The catch and release test exposes the bee to the condition of localizing itself after being released at an unexpected place within the area around the hive which it had explored during its orientation flights³⁹. Treated bees were more frequently lost than control bees, particularly during the initial part of their homing flight. Treated bees also had a higher probability to start their flight by taking a wrong direction, and they had a tendency to interrupt their flights towards the hive, indicating their inability to recall their memory and locate themselves. Our results also corroborates previous findings³³ that the vector flight of bees acutely treated with thiacloprid was not altered, indicating an uncompromised application of the recently learned vector memory if it is retrieved. Homing, however, requires the activation of a remote memory acquired during exploratory orientation flights and the recognition of landmarks as indicators for the route towards the hive from an unexpected location. The flight trajectories recorded in the Fischer et al. study³³ and here strongly indicate a loss of memory retrieval that differs from the recently learned route flight. Neonicotinoids affect predominantly higher-order cognitive functions of the bee brain that are related to the integrative properties of the mushroom bodies. These structures are known to be essential for across sensory integration, learning, and memory formation^{9,10}, and they require functional nicotinic acetylcholine synaptic transmission both at their input site and their output site. It is thus likely that neonicotinoids at low level doses interfere predominantly with mushroom body functions^{11,12}.

Moreover, thiacloprid is often used together with other pesticides in mixtures⁶¹ and some synergism effect between thiacloprid and ergosterol biosynthesis inhibiting fungicides has already been observed in honey bees, increasing the toxicity by up to 560-fold^{22,48}. For Mullin et al.⁶² “the formulation and not just the dose makes the poison”. Future studies should concentrate their efforts on investigating the effects of neonicotinoids not only as active substances but also as formulations. It should also be

noted that the risk of neonicotinoids to bumble bees or solitary bees is about two to three times as high as for honey bees, due to the different sensitivity among the species⁶³. Dramatic consequences on honey bees and more generally pollinators chronically exposed to very low concentrations of thiacloprid are thus to be expected. Therefore, thiacloprid cannot be considered a less harmful neonicotinoid. Our results also demonstrate how important it is to include field test procedures directed towards chronic exposure to sublethal doses of these pesticides and how essential it is to test a large range of possible behavioral effects of a substance before commercializing it.

Supporting Information Available:

Information about residues analysis by LC-MS/MS can be found in Methods S1. Number of waggle runs performed by bees foraging at food sources other than the feeders (Fig. S1), sucrose consumption at the feeders and estimated amounts of thiacloprid collected and metabolized (Table S1), Tuckey's post-hoc tests of the Proboscis Extension Response experiment (Table S2), pesticide residues analysis of honey bees directly and indirectly exposed to thiacloprid (Table S3), flight data of honey bees returned to the hive (Table S4), detailed flight parameters of honey bees returned to the hive (Table S5). This material is available free of charge via the Internet at <http://pubs.acs.org>.

Acknowledgments:

We thank the farmer Mr. Lemmer for providing us with the access to his grass fields and Dr. R. Büchler for the supply of the honeybee colonies and queens. Special thanks also go to A-C. Frank, A. Duer, B. Paffhausen, S. Balbuena, L. Crastre, H. Zwaka and R. Bartels during the field experiments as well as to V. Pucikova, K. Lehmann and J. Degen for their help with the analysis of the flight tracks. The help of M. Groß in statistics and C. Stemler for the correction of the English text is highly appreciated. We are also very grateful to K. Jaenicke and I. Stachewicz-Voigt from the Julius Kühn-Institute in Berlin for their valuable support in the residue analysis. Finally, we thank the editor and three anonymous reviewers for their constructive comments on the manuscript. The work was financially supported by Olin gGmbH, the Freie Universität Berlin, The Deutsche Forschungsgemeinschaft, and a DAAD Postgraduate Scholarship to L.T.

References

1. Klein, A. M.; Vaissière, B. E. ; Cane, J. H. ; Steffan Dewenter, I.; Cunningham, S. A. ; Kremen, C.; Tscharntke, T. Importance of pollinators in changing landscapes for world crops. *Proc. Biol. Sci.* **2007**, 274 (1608), 303–313.
2. Brittain, C.; Potts, S. G. The potential impacts of insecticides on the life-history traits of bees and the consequences for pollination. *Basic Appl. Ecol.* **2011**, 12 (4), 321–331.
3. Rundlöf, M.; Andersson, G. K. S.; Bommarco, R.; Fries, I.; Hederström, V.; Herbertsson, L.; Jonsson, O.; Klatt, B. K.; Pedersen, T. R.; Yourstone, J.; Smith, H. G. Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature*. **2015**, 521 (7550), 77- 80.
4. Whitehorn, P. R.; O'Connor, S.; Wackers, F. L.; Goulson, D. Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science*. **2012**, 336 (6079), 351–352.
5. Brandt, A.; Gorenflo, A.; Siede, R.; Meixner, M.; Büchler, R. The neonicotinoids thiacloprid, imidacloprid, and clothianidin affect the immunocompetence of honey bees (*Apis mellifera* L.). *J Insect Physiol.* **2016**, 86, 40-7.
6. Christen, V.; Mittner, F.; Fent, K. Molecular Effects of Neonicotinoids in Honey Bees (*Apis mellifera*). *Environ. Sci. Technol.*, **2016**, 50 (7), 4071–408.
7. Sánchez-Bayo, F.; Desneux, N. Neonicotinoids and the prevalence of parasites and disease in bees, *Bee World*. **2015**, 92 (2), 34-40.
8. Liu, Z.; Williamson, M. S.; Lansdell, S. J.; Han, Z.; Denholm, I.; Millar, N. S. A nicotinic acetylcholine receptormutation (Y151S) causes reduced agonist potency to a range of neonicotinoid insecticides. *J. Neurochem.* **2006**, 99 (4), 1273–1281.
9. Heisenberg, M. Mushroom body memoir: from maps to models. *Nat. Rev. Neurosci.* **2003**, 4 (4), 266-275.
10. Menzel, R. The honeybee as a model for understanding the basis of cognition. *Nat. Rev. Neurosci.* **2012**, 13 (11), 758-768.
11. Peng, Y.-C., Yang, E.-C. Sublethal Dosage of Imidacloprid Reduces the Microglomerular Density of Honey Bee Mushroom Bodies. *Sci. Rep.* **2016**, 6, 19298.

- 529 12. Palmer, M. J.; Moffat, C.; Saranzewa, N.; Harvey, J.; Wright, G. A.; Connolly, C. N. Cholinergic
530 pesticides cause mushroom body neuronal inactivation in honeybees. *Nat Commun.* **2013**, 4, 1634.
- 531 13. Williamson, S. M.; Wright, G. A. Exposure to multiple cholinergic pesticides impairs olfactory
532 learning and memory in honeybees. *J Exp Biol.* **2013**, 216 (10), 1799-807.
- 533 14. Desneux, N.; Decourtye, A.; Delpuech, J. M. The sublethal effects of pesticides on beneficial
534 arthropods. *Annu. Rev. Entomol.* **2007**, 52, 81–106.
- 535 15. Eiri, D. M.; Nieh, J. C. A nicotinic acetylcholine receptor agonist affects honey bee sucrose
536 responsiveness and decreases waggle dancing. *J. Exp. Biol.* **2012**, 215 (12), 2022-2029.
- 537 16. Henry, M.; Beguin, M.; Requier, F.; Rollin, O.; Odoux, J. F.; Aupinel, P.; Aptel, J.; Tchamitchian,
538 S.; Decourtye, A. A Common Pesticide Decreases Foraging Success and Survival in Honey Bees.
539 *Science.* **2012**, 336 (6079), 348-350.
- 540 17. Schneider, C. W.; Tautz, J.; Grünewald, B.; Fuchs, S. RFID tracking of sublethal effects of two
541 neonicotinoid insecticides on the foraging behaviour of *Apis mellifera*. *PLoS One.* **2012**, 7 (1),
542 e30023.
- 543 18. Tennekes, H. A. The significance of the Druckrey-Küpfmüller equation for risk assessment--the
544 toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time. *Toxicology.*
545 **2010**, 276 (1), 1-4.
- 546 19. Tennekes, H. A.; Sánchez-Bayo, F. Time-Dependent Toxicity of Neonicotinoids and Other
547 Toxicants: Implications for a New Approach to Risk Assessment. *J. Environ. Anal. Toxicol.* **2011**,
548 S4:001.
- 549 20. EFSA. Scientific Opinion of the Panel on Plant Protection Products and their Residues on the
550 science behind the development of a risk assessment of Plant Protection Products on bees (*Apis*
551 *mellifera*, *Bombus spp.* and solitary bees). *EFSA J.* **2012**, 10:2668.
- 552 21. EFSA. Statement on the findings in recent studies investigating sub-lethal effects in bees of some
553 neonicotinoids in consideration of the uses currently authorised in Europe. *EFSA J.* **2012**, 10, 2752.
- 554 22. Iwasa, T.; Motoyama, N.; Ambrose, J. T.; Roe, R. M. Mechanism for the differential toxicity of
555 neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Prot.* **2004**, 23 (5), 371–378.

23. Pisa, L.; Amaral-Rogers, W. V.; Belzunces, L. P.; Bonmatin, J. M.; Downs, C. A.; Goulson, D.; Kreutzweiser, D. P.; Krupke, C.; Liess, M.; McField, M.; Morrissey, C. A.; Noome, D. A.; Settele, J.; Simon-Delso, N.; Stark, J. D.; Van der Sluijs, J. P.; Van Dyck, H.; Wiemers, M. Effects of neonicotinoids and fipronil on non-target invertebrates. *Environ. Sci. Pollut. R.* **2015**, *22* (1), 68–102.
24. FAO specifications and evaluations for thiacloprid.
http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/JMPR/Evaluation06/Thiacloprid06.pdf (accessed April 21, 2016)
25. Elbert, A.; Haas, M.; Springer, B.; Thielert, W.; Nauen, R. Applied aspects of neonicotinoid uses in crop protection. *Pest Manag. Sci.* **2008**, *64* (11), 1099–1105.
26. Simon-Delso, N.; Amaral-Rogers, V.; Belzunces, L. P.; Bonmatin, J. M.; Chagnon, M.; Downs, C.; Furlan, L.; Gibbons, D. W.; Giorio, C.; Girolami, V.; Goulson, D.; Kreutzweiser, D. P.; Krupke, C. H.; Liess, M.; Long, E.; McField, M.; Mineau, P.; Mitchell, E. A.; Morrissey, C. A.; Noome, D. A.; Pisa, L.; Settele, J.; Stark, J. D.; Tapparo, A.; Van Dyck, H.; Van Praagh, J.; Van der Sluijs, J. P.; Whitehorn, P. R.; Wiemers, M. Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environ. Sci. Pollut. R.* **2014**, *22* (1), 2–34.
27. Poquet, Y.; Bodin, L.; Tchamitchian, M.; Fusellier, M.; Giroud, B.; Lafay, F.; Buleté, A.; Tchamitchian, S.; Cousin, M.; Péliissier, M.; Brunet, J. L.; Belzunces, L. P. A pragmatic approach to assess the exposure of the honey bee (*Apis mellifera*) when subjected to pesticide spray. *PLoS ONE*. **2014**, *9* (11), e113728.
28. Van der Sluijs, J. P.; Simon-Delso, N.; Goulson, D.; Maxim, L.; Bonmatin, J. M.; Belzunces, L. P. Neonicotinoids, bee disorders and the sustainability of pollinator services. *Curr. Opin. Environ. Sustain.* **2013**, *5* (3-4), 293–305.
29. Henry, M.; Cerrutti, N.; Aupinel, P.; Decourtye, A.; Gayrard, M.; Odoux, J. F.; Pissard, A.; Rüger, C.; Bretagnolle, V. Reconciling laboratory and field assessments of neonicotinoid toxicity to honeybees. *Proc. Biol. Sci.* **2015**, *282* (1819), 20152110.

- 582 30. Doublet, V.; Labarussias, M.; de Miranda, J. R.; Moritz, R. F. A.; Paxton, R. J. Bees under stress:
 583 sublethal doses of a neonicotinoid pesticide and pathogens interact to elevate honey bee mortality
 584 across the life cycle. *Environ. Microbiol.* **2014**, 17 (4), 969–983.
- 585 31. Retschnig, G.; Neumann, P.; Williams, G. R. Thiacloprid-*Nosema ceranae* interactions in honey
 586 bees: Host survivorship but not parasite reproduction is dependent on pesticide dose. *J. Invertebr.*
 587 *Pathol.* **2014**, 118, 18-19.
- 588 32. Vidau, C.; Diogon, M.; Aufauvre, J.; Fontbonne, R.; Viguès, B.; Brunet, J. L.; Texier, C.; Biron,
 589 D. G.; Blot, N.; El Alaoui, H.; Belzunces, L. P.; Delbac, F. Exposure to sublethal doses of fipronil
 590 and thiacloprid highly increases mortality of honeybees previously infected by *Nosema ceranae*.
 591 *PLoS One.* **2011**, 6 (6), e21550.
- 592 33. Fischer, J.; Müller, T.; Spatz, A. K.; Greggers, U.; Grünewald, B.; Menzel, R. Neonicotinoids
 593 Interfere with Specific Components of Navigation in Honeybees. *PLoS One.* **2014**, 9 (3), e91364.
- 594 34. Krupke, C. H.; Hunt, G. J.; Eitzer, B. D.; Andino, G.; Given, K. Multiple routes of pesticide
 595 exposure for honey bees living near agricultural fields. *PLoS One.* **2012**, 7 (1), e29268.
- 596 35. European and Mediterranean Plant Protection Organisation (EPPO). Guideline on test methods for
 597 evaluating the side effects of plant protection products on honey bees. *EPPO Bull.* **1992**, 22, 203–
 598 215.
- 599 36. Menzel, R.; Kirbach, A.; Haass, W. D.; Fischer, B.; Fuchs, J.; Koblofsky, M.; Lehmann, K.;
 600 Reiter, L.; Meyer, H.; Nguyen, H.; Jones, S.; Norton, P.; Greggers, U. A common frame of
 601 reference for learned and communicated vectors in honeybee navigation. *Curr. Biol.* **2011**, 21 (8),
 602 645–650.
- 603 37. Riley, J. R.; Smith, A. D.; Reynolds, D. R.; Edwards, A. S.; Osborne, J. L.; Williams, I. H.;
 604 Carreck, N. L.; Poppy, G. M. Tracking bees with harmonic radar. *Nature.* **1996**, 379 (6560), 29–
 605 30.
- 606 38. Scheiner, R.; Abramson, C. I.; Brodschneider, R.; Crailsheim, K.; Farina, W. M.; Fuchs, S.;
 607 Grünewald, B. Hahshold, S.; Karrer, M.; Koeniger, G.; Koeniger, N.; Menzel, R.; Mujagic, S.;
 608 Radspieler, G.; Schmickl, T.; Schneider, C.; Siegel, A. J.; Szopek, M.; Thenius, R. Standard
 609 methods for behavioural studies of *Apis mellifera*. *J. Apicult. Res.* **2013**, 52 (4).

- 610 39. Degen, J.; Kirbach, A.; Reiter, L.; Lehmann, K.; Norton, P.; Storms, M.; Koblofskya, M.;
611 Wintera, S.; Georgievaa, P. B.; Nguyen, H.; Chamkhia, H.; Greggers, U.; Menzel, R. Exploratory
612 behaviour of honeybees during orientation flights. *Anim. Behav.* **2015**, 102, 45-57.
- 613 40. Capaldi, E. A.; Smith, A. D.; Osborne, J. L.; Fahrbach, S. E.; Farris, S. M.; Reynolds, D. R.;
614 Edwards, A. S.; Martin, A.; Robinson, G. E.; Poppy, G. M.; Riley, J. R. Ontogeny of orientation
615 flight in the honeybee revealed by harmonic radar. *Nature.* **2000**, 403 (6769), 537-540.
- 616 41. Greggers, U.; Koch, G.; Schmidt, V.; Dürr, A.; Floriou-Servou, A.; Piepenbrock, D.; Göpfert, M.
617 C.; Menzel, R. Reception and learning of electric fields in bees. *Proc. Biol Sci.* **2013**, 280 (1759),
618 20130528.
- 619 42. von Frisch, K. *The dance language and orientation of bees*. Cambridge: Harvard Univ. Press;
620 **1967**.
- 621 43. Bitterman, M. E.; Menzel, R.; Fietz, A.; Schäfer, S. Classical conditioning proboscis extension in
622 honeybees (*Apis mellifera*). *J. Comp. Psychol.* **1983**, 97 (2), 107-119.
- 623 44. Page, R. E.; Erber, J.; Fondrk, K. M. The effect of genotype on response thresholds to sucrose and
624 foraging behavior of honey bees (*Apis mellifera*). *J. Comp. Physiol.* **1998**, 182 (4), 489-500.
- 625 45. Rortais, A.; Arnold, G.; Halm, M. P.; Touffet-Briens, F. Modes of honeybees exposure to
626 systemic insecticides: estimated amounts of contaminated pollen and nectar consumed by different
627 categories of bees. *Apidologie.* **2005**, 36 (1), 71-83.
- 628 46. Balderrama, N. M.; Almeida, L. O.; Núñez, J. A. Metabolic rate during foraging in the honeybee.
629 *J. Comp. Physiol. B.* **1992**, 162 (5), 440-447.
- 630 47. Fournier, A.; Rollin, O.; Le Féon, V.; Decourtye, A.; Henry, M. Crop-Emptying Rate and the
631 Design of Pesticide Risk Assessment Schemes in the Honey Bee and Wild Bees (Hymenoptera:
632 Apidae). *J. Econ. Entomol.* **2014**, 107 (1), 38-46.
- 633 48. Rothe, U.; Nachtigall, W. Flight of the honey bee IV. Respiratory quotients and metabolic rates
634 during sitting, walking and flying. *J. Comp. Physiol. B.* **1989**, 158 (6), 739-749.
- 635 49. Menzel, R.; Greggers, U.; Smith, A.; Berger, S.; Brandt, R. Brunke, S.; Bundrock, G.; Hülse, S.;
636 Plümpe, T.; Schaupp, F.; Schüttler, E.; Stach, S.; Stindt, J.; Stollhoff, N.; Watzl, S. Honey bees
637 navigate according to a map-like spatial memory. *PNAS.* **2005**, 102 (8), 3040-3045.

- 638 50. Oliveira, R. A.; Roat, T. C.; Carvalho, S. M.; Malaspina, O. Side-effects of thiamethoxam on the
639 brain and midgut of the Africanized honeybee *Apis mellifera* (Hymenoptera: Apidae). *Environ.*
640 *Toxicol.* **2013**, 29 (10), 1122-1133.
- 641 51. Wu, J. Y.; Anelli, C. M.; Sheppard, W. S. Sub-Lethal Effects of Pesticide Residues in Brood
642 Comb on Worker Honey Bee (*Apis mellifera*) Development and Longevity. *PLoS One.* **2011**, 6 (2):
643 e14720.
- 644 52. Schmuck, R.; Stadler, T.; Schmidt, H. W. Field relevance of a synergistic effect observed in the
645 laboratory between an EBI fungicide and a chloronicotinyl insecticide in the honeybee (*Apis*
646 *mellifera* L, Hymenoptera). *Pest Manag. Sci.* **2003**, 59 (3), 279–286.
- 647 53. Colin, M. E.; Bonmatin, J. M.; Moineau, I.; Gaimon, C.; Brun, S.; Vermandere, J. P. A method to
648 quantify and analyze the foraging activity of honey bees: relevance to the sublethal effects induced
649 by systemic insecticides. *Arch. Environ. Contam. Toxicol.* **2004**, 47 (3), 387–395.
- 650 54. Yang, E. C.; Chuang, Y. C.; Chen, Y. L.; Chang, L. H. Abnormal Foraging Behavior Induced by
651 Sublethal Dosage of Imidacloprid in the Honey Bee (Hymenoptera: Apidae). *J. Econ. Entomol.*
652 **2008**, 101 (6), 1743-1748.
- 653 55. Laurino, D.; Porporato, M.; Patteta, A.; Manino, A. Toxicity of neonicotinoid insecticides to
654 honey bees: laboratory tests. *B. Insectol.* **2011**, 64 (1), 107-113.
- 655 56. Kessler, S. C.; Tiedeken, E. J.; Simcock, K. L.; Derveau, S.; Mitchell, J.; Softley, S.; Stout, J. C.;
656 Wright, G. A. Bees prefer foods containing neonicotinoid pesticides. *Nature* **2015**, 521 (7550), 74–
657 76.
- 658 57. Pohorecka, K.; Skubida, P.; Mischczak, A.; Semkiw, P.; Sikorski, P.; Zagibajło, K.; Teper, D.;
659 Kołtowski, Z.; Zdańska, D.; Skubida, M.; Bober, A. Residues of neonicotinoid insecticides in bee
660 collected plant materials from oilseed rape crops and their effect on bee colonies. *J Apic Sci.* **2012**,
661 56 (2), 115-134.
- 662 58. Hölldobler, B.; Wilson, E. O. *The Superorganism: The Beauty, Elegance, and Strangeness of*
663 *Insect Societies*. W. W. Norton & Company; **2008**.
- 664 59. Khoury, D. S.; Myerscough, M. R.; Barron, A. B. A quantitative model of honey bee colony
665 population dynamics. *PLoS One.* **2011**, 6 (4), e18491.

- 666 60. Herb, B. R.; Wolschin, F.; Hansen, K. D.; Aryee, M. J.; Langmead, B.; Irizarry, R.; Amdam,
667 G. V.; Feinberg, A. P. Reversible switching between epigenetic states in honeybee behavioral
668 subcastes. *Nat. Neurosci.* **2012**, 15 (10), 1371–1373.
- 669 61. Mullin, C. A.; Frazier, M. T.; Frazier, J. L.; Ashcraft, S.; Simonds, R.; vanEngelsdorp, D.; Pettis,
670 J. S. High levels of miticides and agrochemicals in North American apiaries: implications for
671 honey bee health. *PLoS One.* **2010**, 5 (3), e9754.
- 672 62. Mullin, C. A.; Chen, J.; Fine, J. D.; Frazier, M. T.; Frazier, J. L. The formulation makes the honey
673 bee poison. *Pestic Biochem Physiol.* **2015**, 120, 27-35.
- 674 63. Sanchez-Bayo, F.; Goka, K. Pesticide residues and bees – A risk assessment. *PLoS One.* **2014**, 9
675 (4), e94482.

676 **Table 1: Foraging span in days of the trained honey bees at the control or treated feeder.**

	Experiment 1	Experiment 2	Total [§]
Control	5.21 ± 0.32 (<i>n</i> = 67) * a	4.19 ± 0.24 (<i>n</i> = 72) a	4.68 ± 0.20 (<i>n</i> = 139)
Treated	4.7 ± 0.22 (<i>n</i> = 79) a	3.34 ± 0.14 (<i>n</i> = 111) * b	3.91 ± 0.13 (<i>n</i> = 190)

677
678 Numbers shown are means (days foraging) ± s.e.m.

679
680 [§] Mann-Whitney, *P* < 0.01

681
682 * The control group in Exp. 1 and the treated group in Exp. 2 correspond to the same colony, as the control colony in Exp. 1
683 became the treated colony in Exp. 2 and continued to forage at the same feeder (F1).

684
685 Different letters indicate significant differences (post-hoc tests with Bonferroni correction): a-b (Exp.2), *P* < 0.05, a-b
686 (Treated), *P* < 0.001, a-b (F1), *P* < 0.001.

687 **Table 2: Summary of the Cox regression model.**

Variables	Model 1				Model 2			
	regression coefficient	exp (coef) *	Z	P	regression coefficient	exp (coef) *	Z	P
Treatment	-0.577213	0.561461	-3.408	0.000656	-0.5866	0.5562	-3.505	0.000456
Experiment	-0.372878	0.688749	-1.563	0.117983	-0.2864	0.7510	-1.745	0.080899
Time foraging ‡	-0.035163	0.965448	-0.674	0.500248				
Time exposure §	-0.013654	0.986439	-0.838	0.402182				
Temperature	-0.007925	0.992106	-0.238	0.811991				
Time before flying	0.017345	1.017496	1.133	0.257266				
	<i>Rsquare: 0.091 (max possible= 0.999), Likelihood Ratio Test: 17.71 on 6 df, P=0.007007</i>				<i>Rsquare: 0.08 (max possible= 0.999), Likelihood Ratio Test: 15.52 on 2 df, P=0.0004268</i>			

688

689 A backward selection on the AIC was performed on Model 1 in order to obtain Model 2

690

690 Values in bold indicate significant differences

691

691 *exp (coef) = Hazard ratio

692

692 ‡ **Time foraging** is the time in days during which a bee is foraging at its feeder before being released

693

693 § **Time exposure** is the time in days from the first day of the experiment until the day the bee is released

Figure 1**Required sucrose concentrations and foraging activity at the control and treated feeders.**

(a) Sucrose concentrations used in order to keep a similar number of foragers coming regularly to the control and treated feeders and to induce dances. Lower sucrose concentrations were required for control bees than for treated bees.. (b) Mean (\pm 95 % confidence limits) number of visits per hour recorded on the same days ($n = 19$) at both feeders during regular foraging (circles) and during dance induction (squares). The foraging behavior of the treated bees (filled marks) as well as their ability to recruit new untrained foragers are significantly reduced (ANOVA, $F_{3,72} = 14.01$, $P < 0.0001$ and Tukey post-hoc tests). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Figure 2**Number of waggles runs performed by the trained bees from the control and treated feeders.**

The number of waggles runs per hour was obtained from electrostatic field recordings performed on the same days in both hives (n days = 32). The mean number of waggles runs per hour is represented with a cross in the box-plots, it was found significantly higher for the bees foraging at the control feeder than for the bees foraging at the contaminated feeder (Wilcoxon signed rank test, $p < 0.0001$).

Figure 3**Proboscis Extension Response (PER) to different sucrose concentrations containing 5 ppm**

thiacloprid (treated) or not (control). N control = 73. N treated = 71. No difference was found between the two groups (logistic regression with random effects, Sugar conc x Treatment: $\chi^2_6 = 2.5224$, $P = 0.866$).

Figure 4

Accumulation of thiacloprid residue in heads, thoraces, abdomens and in the whole body (representing the sum of the measurements) of honey bees foraging at the contaminated feeder

over time. Honey bee foragers were collected at the end of 2, 3 or 4 days of foraging after they had filled their crop at the feeder containing thiacloprid (4.5 ppm). 10 bees per foraging group.


Figure 5

Probability of homing success as a function of time until return. Treated honey bees returned to their hive at a significantly lower proportion than control bees ($n_{\text{treated}} = 100$, 76 % return; $n_{\text{control}} = 85$, 91.76 % return, Fischer Exact Test, $P < 0.01$). The origin of the temporal axis represents the moment of release.

Figure 6

Flight trajectories of the non-returning bees. Map data provided by: Google Earth and GeoBasis - DE BKG. The figures show the flight trajectories of individual bees, each in a different color within a group (**a**, **b**, **c** and **d**). The trained route of the bees released at the release site (RS) is represented with a red line between the hive (H) and the feeders (F1 and F2). In Experiment 1, F1 was the feeder of the control bees and F2 the feeder of the treated bees. In Experiment 2 the situation was reversed (F1: treated bees, F2: control bees). The circle (black dashed line) represents the edge of the radar range (900 m from the radar). Bees leaving the radar range and then returning into it are marked with a black arrow directed to the East (leaving the range) or to the West (returning into the radar range) respectively. A square at the beginning of each flight track marks the first radar signal, and the triangle at the end of the flight marks the last radar signal. See Table S4 for the number of bees lost within each group.


742 Fig. 1:



743


744

Fig. 2:



748 Fig. 3:

749



750

751

Fig. 4:




Fig. 5:




Fig. 6:



763 TOC/Abstract Art:

764



765