

# A unified approach to inference with linked data

Rachel Anderson\*

This Version: September 9, 2019

## Abstract

This paper studies the joint effects of matching algorithms and estimation procedures for linked data on the quality of the estimates that they produce. Specifically, I compare how different combinations of record linkage and estimation methods perform across a variety of data generating processes and research tasks (i.e. linking the dependent variable, independent variable, sample selection, GMM, linear regression) in order to develop a unified approach for doing empirical work with linked data.

## 1 Introduction

In economic history, identifying a common set of individuals appearing in two or more datasets is complicated by the lack of unique identifying variables, so that matching observations across datasets requires linking individuals by characteristics such as name and reported age. For example, Aizer et al. (2016) link children listed on Mothers' Pension program welfare applications from 1911-1935 with Social Security Death Master File records from 1965-2012 using an automated matching procedure based on individuals' names and dates of birth. The authors match 48 percent of children to a unique death record and 4

---

\*Mailing Address: Department of Economics, Julis Romo Rabinowitz Building, Princeton, NJ 08544. Email: rachelsa@Princeton.edu. This project received funding from the NIH (Grant 1 R01 HD077227-01).

percent to multiple possible records, leaving 48 percent unmatched.<sup>1</sup>

To account for the multiple matches, Aizer et al. (2016) use methods from Anderson et al. (2019), which assume that each of an individual’s associated outcomes is equally likely to be the true match; and they use hazard models to account for attrition (i.e. the unmatched individuals). Alternatively, the authors could have generated a “composite match” equal to the average of the linked observations (Bleakley and Ferrie, 2016), or constructed bounds on the parameter of interest using different configurations of matched data (Nix and Qian, 2015). Had they used probabilistic record linkage methods to link the data, they could have used the robust OLS estimators from Lahiri and Larsen (2005), or prior-informed imputation for missing records proposed by Goldstein et al. (2012).

This example shows that inference using matched data requires deciding how to handle observations with unique, multiple, or zero matches, and suggests how the outputs of a matching procedure determine which estimation tools are available. Indeed, there are a number of recent papers that individually compare the performance of different matching algorithms (Bailey et al., 2017; Abramitzky et al., 2018) and estimation methods for linked data (Harron et al., 2014); however, little is known about the *joint* impact of matching and estimation procedures on the quality of inference. This paper seeks to fill this gap by studying how choices in linkage and estimation affect parameter estimates and confidence intervals in standard econometric models, with the hope of guiding researchers on how to choose which combination of methods best suits their projects’ needs.

This paper is most similar in spirit to Scheuren and Winkler (1997), in that the goal is to develop a unified approach toward linking and analyzing matched data. But whereas Scheuren and Winkler (1997) propose a single method for linear regression,<sup>2</sup> this paper studies the impact of multiple methods across a variety of research scenarios. First, I implement


---

<sup>1</sup>The authors estimate that at least 32 percent of individuals in the Mothers’ Pension program data died before 1965, and therefore should have no match in the 1965-2012 data.

<sup>2</sup>The authors advocate iterating between linkage, estimation, and imputation steps before arriving at a final, fitted, regression line.

several of the most commonly used deterministic and probabilistic record linkage procedures to generate multiple matched versions of the data. Then, for each matched version of the data, I estimate different types of econometric models, using different estimation methods, and compare the parameter estimates and confidence intervals that these combinations produce. One iteration of each matching/model/estimation procedure is represented in Figure 1

Figure 1: Research design



At first glance, it's not obvious that the methods studied in this paper should be directly comparable, or that certain combinations of matching and estimation techniques are possible to implement at the same time; however, I will show how, with a few adjustments, both of these tasks are possible.<sup>3</sup> Although the matching algorithms studied in this paper according to whether they are deterministic or probabilistic, I will show that these methods can be made directly comparable by setting thresholds for each method such that the Type I and Type II error rates are equalized.

Similarly, the estimation techniques studied in this paper differ according to whether they require estimating the probability that a record pair refers to a match. When implementing methods that require this extra information in combination with deterministic

<sup>3</sup>maybe even a theoretical result showing equivalence across classes of matching methods and estimators?

matching techniques that *do not* output match probabilities, I will treat the matches as being equally likely. Similarly, if an estimation procedure does not use match probabilities, but a probabilistic matching algorithm outputs them, then the probabilities will be ignored.

Section 2 introduces the general problem that this paper seeks to address and outlines a common framework for analyzing the techniques listed in Table 1. Sections 3 and 4 describe each of the methods in detail. Section 4 describes the implementation of the methods and the data. Section 5 contains the results. Section 6 will be a real empirical application, and Section 7 will conclude.

Table 1: List of methods to be implemented

Matching Methods		Estimation Methods	
Deterministic	Probabilistic	Uses match weights	No match weights
Abramitzky et al. (2012)	Abramitzky et al. (2018)	Lahiri and Larsen (2005)	Anderson et al. (2019)
Aizer et al. (2016)	Feigenbaum (2016)	Goldstein et al. (2012)	Nix and Qian (2015)
		Anderson (2019)	Bleakley and Ferrie (2016)

## 2 General problem

Suppose that the researcher would like to estimate  $\theta_0$  in a parametric model of the form

$$E[m(\mathbf{z}_i; \theta_0)] = 0 \tag{1}$$

where  $m(\cdot)$  is a moment function and  $\mathbf{z}_i = (x_i, y_i)$  is the data associated with an individual  $i$  sampled at random from the population of interest; however, instead of observing  $(x_i, y_i)$  pairs directly, the researcher observes two datasets. The first dataset  $D_1$  contains variables  $x_i$  and identifiers  $w_i$  for individuals  $i = 1, \dots, N_1$ . The second dataset  $D_2$  contains outcomes  $y_j$  and identifiers  $w_j$  for individuals  $j = 1, \dots, N_2$ .

To estimate (1) with standard econometric methods, the researcher must identify which

of the  $x_i$  and  $y_j$  refer to the same individuals. That is, she needs to recover the matching function  $\varphi : \{1, \dots, N_1\} \rightarrow \{1, \dots, N_2\} \cup \emptyset$ , where  $\varphi(i) = j$  if individual  $i$  in dataset  $D_1$  and individual  $j$  in dataset  $D_2$  refer to the same entity; and  $\varphi(i) = \emptyset$  if  $i$  does not have a match in  $D_2$ . If  $w_i$  and  $w_j$  identify individuals uniquely and do not change over time, then  $\varphi(i) = j$  if and only if  $w_i = w_j$ ; otherwise,  $\varphi(i) = \emptyset$ . However, if the identifiers are non-unique or prone to errors introduced by the record-generating process, then  $\varphi$  needs to be estimated, and inference about  $\theta$  needs to be adjusted accordingly.

In statistics, the task of recovering  $\varphi$  is called *record linkage*. A record linkage procedure consists of a set of decisions about (i) selecting and standardizing the identifiers  $w_i$  and  $w_j$ , (ii) choosing which records to consider as potential matches (especially when  $N_1 \times N_2 \times \dim(w_i)$  is large), (iii) defining what patterns of  $(w_i, w_j)$  constitute (partial) agreements, and (iv) designating  $(i, j)$  pairs as matches.<sup>4</sup> Each step of the record linkage process introduces the possibility that a true match is overlooked (Type II error), or that a false match is designated as correct (Type I error), and there is generally a tradeoff between reducing either one of the two (Abramitzky et al., 2019b; Doidge and Harron, 2018).

To fix ideas, suppose that  $\theta_0$  is the long-run impact of cash transfers on the longevity of children raised in poor families. The vector  $x_i$  includes family and child characteristics as observed in welfare program applications; and the outcomes  $y_j$  are constructed by calculating (day of death – day of birth) for all of the observations in a set of death records. For all  $i$  and  $j$ , the identifiers  $w_i$  and  $w_j$  include the individual’s first and last name, and date of birth. Additionally,  $w_i$  includes  $i$ ’s place of birth;  $w_j$  includes  $j$ ’s place of death; and some  $w_i$  and  $w_j$  contain the individual’s middle name or middle initial.

In this setting, an example of a (deterministic) record linkage procedure consists of:

---

<sup>4</sup>Note that this is the author’s own definition. By contrast, Bailey et al. (2017) categorize historical linking algorithms (that match observations using name and age only) according to how they treat candidate pairs in the following four categories: (M1) a perfect, unique match in terms of name and age similarity; (M2) a single, similar match that is slightly different in terms of age, name, or both; (M3) many perfect matches, leading to problems with match disambiguation; (M4) multiple similar matches that are slightly different in terms of age, name or both.

- (i) using a phonetic algorithm to standardize all string variables;
- (ii) considering as potential matches only  $(i, j)$  pairs whose phonetically standardized names begin with the same letter, and whose birth years are within  $\pm 2$  years;
- (iii) measuring agreements among names using Jaro-Winkler string distances, and weighing disagreements in birth year more than differences in birth month (and more than differences in birth day),
- (iv) designating as matches all  $(i, j)$  pairs with scores calculated using the metrics in (iii) exceeding a pre-specified cutoff; and, if a record  $i$  has multiple possible matches  $j$  that exceed the cut-off, then choosing the match with the highest score (or picking a random match if there is a tie).

Another record linkage procedure could be defined using the same rules for steps (i)-(iii), but replace (iv) with a probabilistic matching rule that does not enforce one-to-one matching:

- (iv\*) use the Expectation-Maximization algorithm to compute “match weights” for each record pair; then designate as matches all pairs that exceed thresholds that are set to reflect specific tolerances for Type I and Type II error.

Except in rare cases, the matching function outputted by replacing (iv) with (iv\*) will be different. Whereas the first procedure associates each  $x_i$  with at most one matched  $y_j$ , the second procedure may associate the same  $x_i$  with multiple  $y_j$  (in technical terms, this implies  $\varphi$  is a correspondence). The former case might use a standard GMM model to estimate  $\theta$ ; while the latter requires methods that associates multiple values of  $y_j$  with each  $x_i$  (Anderson et al., 2019). This example shows that not only do the estimates of  $\theta$  likely depend on the estimates of  $\varphi$ , but also the *methods* for estimating  $\theta$  may be also differ.

As observed by Bailey et al. (2017), record linkage procedures differ by the set of assumptions that motivate their use. However, all of the procedures discussed in this paper

will be studied under the following, common set of assumptions (with some departures later on):

1. (De-duplication) Within a given dataset, each observation refers to a distinct entity. That is, if two observations share the same identifier, they represent two different individuals.
2. (No unobserved sample selection) The observed  $x_i$  and  $y_j$  are random samples conditional on  $w_i$  and  $w_j$ , respectively. This means that all individuals with the same identifying information have equal probability of appearing in the sample.
3. There exists a unique  $\theta_0 \in \Theta$  that satisfies the relationship in (1), that can be consistently estimated using standard econometric techniques if  $\varphi_0$  is known.

The next section discusses in detail the exact record linkage techniques that will be studied, and their motivating assumptions.

### 3 Record Linkage Methods

Research on record linkage appears in many fields, such as statistics, computer science, operations research, and epidemiology, under many names, such as data linkage, entity resolution, instance identification, de-duplication, merge/purge processing, and reconciliation. As such, there are several books devoted to its study (Harron et al., 2015; Christen, 2012; Herzog et al., 2007), and dozens of commercial and open source systems software developed for its implementation (Köpcke and Rahm, 2010). Although insights from other fields have been slow to reach economics, recent working papers examine the impact of different record linkage techniques on inference using historic U.S. census data (Abramitzky et al., 2019b; Bailey et al., 2017); and the most recent version of the Handbook of Econometrics includes

a chapter on the “Econometrics of Data Combination” (Ridder and Moffitt, 2007).<sup>5</sup>

Record linkage techniques are broadly categorized as deterministic or probabilistic<sup>6</sup>; however, every deterministic linkage method has an equivalent probabilistic version (Doidge and Harron, 2018). For this reason, I will focus on developing estimation methods that use the probabilistic record linkage framework, but I will show how they can be applied to data that are linked deterministically.

### 3.1 Steps of the record linkage task

Here I will write a little bit more about each of the steps introduced in Section 2.

#### (i) selecting and standardizing the identifiers $w_i$ and $w_j$

phonetic algorithms

#### (ii) choosing which records to consider as potential matches

blocking

#### (iii) defining what patterns of $(w_i, w_j)$ constitute (partial) agreements

Jaro-winkler distances, for example

---

<sup>5</sup>Similar survey papers also exist in fields outside of economics, such as epidemiology and computer science (Doidge and Harron, 2018; Winkler, 1999). In fact, that record linkage is studied by many fields makes writing (and reading!) such surveys difficult, because authors are constantly writing the same things. For example, Goldstein et al. (2012) prove similar results to those published in Hirukawa and Prokhorov (2018).

<sup>6</sup>define them here



### 3.1.1 (iv) designating $(i, j)$ pairs as matches

this is where the differences between deterministic/probabilistic really arise!

## 3.2 Probabilistic Record Linkage

In describing the record linkage techniques implemented in this paper, I use notation from Fellegi and Sunter (1969). As before, consider two datafiles  $D_1$  and  $D_2$  that record information from two overlapping sets of individuals or entities. The files originate from two record-generating processes that may induce errors and missing values, so that identifying which individuals are represented in both  $D_1$  and  $D_2$  is nontrivial. I assume that individuals appear at most once in each datafile, so that the goal of record linkage is to identify which records in files  $D_1$  and  $D_2$  refer to the same entities.

Suppose that files  $D_1$  and  $D_2$  contain  $N_1$  and  $N_2$  records, respectively, and without loss of generality that  $N_2 \geq N_1$ . Denote also the number of entities represented in both files as  $N_M$ , so that  $N_1 \geq N_M \geq 0$ .

We say that the set of ordered record pairs  $D_1 \times D_2$  is the union of two disjoint sets, *matches* ( $M$ ) and *non-matches* ( $U$ ):

$$M = \{(i, j) : i \in D_1, j \in D_2, i = j\}$$

$$U = \{(i, j) : i \in D_1, j \in D_2, i \neq j\}$$

Hence, the formal goal of record linkage is to identify whether an arbitrary record pair  $(i, j) \in D_1 \times D_2$  belongs to  $M$  or  $U$ . Note that this task is identical to

To perform this task, each record pair is evaluated according to  $L$  different comparison criteria, which are the result of comparing data fields for records  $i$  and  $j$ . For example, if a record pair  $(i, j)$  represents two individuals, the pair may be evaluated according to whether

they share a first name or have the same birthday. These comparisons are represented by a *comparison vector*,

$$\boldsymbol{\gamma}_{ij} = (\gamma_{ij}^1, \dots, \gamma_{ij}^\ell, \dots, \gamma_{ij}^L)$$

where each comparison field  $\gamma_{ij}^\ell$  may be binary-valued, as in “ $i$  and  $j$  have the same birthday,” or use levels to account for partial agreement between strings (see ?, for details). The models presented herein use only binary comparison vectors, however they may be extended to allow for partial agreement using the methods from ?.

The probability of observing a particular configuration of  $\boldsymbol{\gamma}_{ij}$  can be modeled as arising from the mixture distribution:

$$P(\boldsymbol{\gamma}_{ij}) = P(\boldsymbol{\gamma}_{ij}|M)p_M + P(\boldsymbol{\gamma}_{ij}|U)p_U \quad (2)$$

where  $P(\boldsymbol{\gamma}_{ij}|M)$  and  $P(\boldsymbol{\gamma}_{ij}|U)$  are the probabilities of observing the pattern  $\boldsymbol{\gamma}_{ij}$  conditional on the record pair  $(i, j)$  belonging to  $M$  or  $U$ , respectively. The proportions  $p_M$  and  $p_U = 1 - p_M$  are the marginal probabilities of observing a matched or unmatched pair. Applying Bayes’ Rule, we obtain the probability of  $(i, j) \in M$  conditional on observing  $\boldsymbol{\gamma}_{ij}$ ,

$$P(M|\boldsymbol{\gamma}_{ij}) = \frac{p_M P(\boldsymbol{\gamma}_{ij}|M)}{P(\boldsymbol{\gamma}_{ij})} \quad (3)$$

so that if we can estimate the variables in (2), we can estimate the probability that any two records refer to the same entity in (3).

As shown by Fellegi and Sunter (1969), it is possible to use the estimated probabilities to construct an “optimal” matching, given any threshold for false positive and false negative match rates. Conversely, the probabilities also allow us to estimate the false positive rate for any configuration of matches (?). Given the usefulness of the quantities in (3), the next sections will introduce two methods for estimating them.

### 3.3 Simplifying assumptions

Let  $\mathbf{\Gamma} \equiv \{\gamma_{ij} : (i, j) \in X_1 \times X_2\}$  denote the set of comparison vectors for all records pairs  $(i, j) \in X_1 \times X_2$ . Note that  $\mathbf{\Gamma}$  contains potentially  $n_1 \times n_2$  elements, so that calculating  $\mathbf{\Gamma}$  may be computationally expensive when  $X_1$  or  $X_2$  is large. In practice, researchers partition  $X_1 \times X_2$  into “blocks,” such that only records belonging to the same block are attempted to be linked, and records belonging to different blocks are assumed to be nonmatches. For example, postal codes and household membership are often used to define blocks when linking census files (?). Importantly, the blocking variables should be recorded without error, and sometimes there are none available. This paper assumes that no blocking is used; or, alternatively, that records are already divided into blocks that can be analyzed independently using the methods outlined below.

#### Conditional independence

In principle, we can model,

$$\gamma_{ij} \mid M \sim \text{Dirichlet}(\boldsymbol{\delta}_M)$$

$$\gamma_{ij} \mid U \sim \text{Dirichlet}(\boldsymbol{\delta}_U)$$

However, there are  $2^L - 1$  possible configurations of each  $\gamma_{ij}$ , so that  $\boldsymbol{\delta}_M$  and  $\boldsymbol{\delta}_U$  may be very high-dimensional if we want to allow weights to vary across different comparison criteria.

A common assumption in the literature is that the comparison fields  $\ell$  are defined so that  $\gamma_{ij}^\ell$  are independent across  $\ell$  conditional on match status. This implies:

$$P(\gamma_{ij}|C) = \prod_{\ell=1}^L P(\gamma_{ij}^\ell|C)^{\gamma_{ij}^\ell} (1 - Pr(\gamma_{ij}^\ell|C))^{1-\gamma_{ij}^\ell} \quad C \in \{M, U\} \quad (4)$$

Hence the number of parameters used to describe each mixture class is reduced to  $L$ .

? have shown how to relax this assumption using log-linear models, but for now I assume conditional independence to ease computation.

### 3.4 Measuring record linkage performance

(could also summarize) They are evaluated according to matching rates, type I and type II error rates; robustness to selection/attrition?

List of record linkage methods Link of estimation methods

## 4 Estimation Methods

### 4.1 Standard assumptions

### 4.2 Relaxed assumptions

Matching variables correlated with variable of interest; true link not included

## 5 Overview of results

## References

- A. Aizer, S. Eli, J. Ferrie, and A. Lleras-Muney, “The long-run impact of cash transfers to poor families,” *American Economic Review*, vol. 106, no. 4, pp. 935–71, April 2016. [Online]. Available: <http://www.aeaweb.org/articles?id=10.1257/aer.20140529>
- R. Anderson, B. Honore, and A. Lleras-Muney, “Estimation and inference using imperfectly matched data,” *Working paper*, August 2019. [Online]. Available: <http://www.github.com/rachelsanderson/ImperfectMatching>

- H. Bleakley and J. Ferrie, "Shocking behavior: Random wealth in antebellum georgia and human capital across generations," *The Quarterly Journal of Economics*, vol. 131, no. 3, pp. 1455–1495, 2016. [Online]. Available: <https://doi.org/10.1093/qje/qjw014>
- E. Nix and N. Qian, "The fluidity of race: Passing in the united states, 1880-1940," National Bureau of Economic Research, Working Paper 20828, January 2015. [Online]. Available: <http://www.nber.org/papers/w20828>
- P. Lahiri and M. D. Larsen, "Regression analysis with linked data," *Journal of the American Statistical Association*, vol. 100, no. 469, pp. 222–230, 2005. [Online]. Available: <http://www.jstor.org/stable/27590532>
- H. Goldstein, K. L. Harron, and A. M. Wade, "The analysis of record-linked data using multiple imputation with data value priors." *Statistics in medicine*, vol. 31 28, pp. 3481–93, 2012.
- M. Bailey, C. Cole, M. Henderson, and C. Massey, "How well do automated linking methods perform? lessons from u.s. historical data," National Bureau of Economic Research, Working Paper 24019, November 2017. [Online]. Available: <http://www.nber.org/papers/w24019>
- R. Abramitzky, R. Mill, and S. Perez, "Linking individuals across historical sources: a fully automated approach," National Bureau of Economic Research, Working Paper 24324, February 2018. [Online]. Available: <http://www.nber.org/papers/w24324>
- K. Harron, A. Wade, R. Gilbert, B. Muller-Pebody, and H. Goldstein, "Evaluating bias due to data linkage error in electronic healthcare records," *BMC medical research methodology*, vol. 14, p. 36, 03 2014.
- F. Scheuren and W. Winkler, "Regression analysis of data files that are computer matched - part ii," *Survey Methodology*, vol. 23, 01 1997.
- J. P. Ferrie, "A new sample of males linked from the public use microdata sample of the 1850 u.s. federal census of population to the 1860 u.s. federal census manuscript schedules," *Historical Methods*, vol. 29, no. 4, pp. 141–156, 1 1996.
- R. Abramitzky, L. P. Boustan, and K. Eriksson, "Europe's tired, poor, huddled masses: Self-selection and economic outcomes in the age of mass migration," *American Economic Review*, vol. 102, no. 5, pp. 1832–56, May 2012. [Online]. Available: <http://www.aeaweb.org/articles?id=10.1257/aer.102.5.1832>
- R. Abramitzky, L. P. Boustan, and K. Eriksson, "To the new world and back again: Return migrants in the age of mass migration," *ILR Review*, vol. 72, no. 2, pp. 300–322, 2019. [Online]. Available: <https://doi.org/10.1177/0019793917726981>

- T. Enamorado, B. Fifield, and K. Imai, “Using a probabilistic model to assist merging of large-scale administrative records,” *American Political Science Review*, vol. 113, no. 2, p. 353–371, 2019.
- J. J. Feigenbaum, “A machine learning approach to census record linking ?” 2016.
- R. Abramitzky, L. Boustan, K. Eriksson, J. Feigenbaum, and S. Perez, “Automated linking of historical data,” *NBER Working Paper*, 2019.
- J. Doidge and K. Harron, “Demystifying probabilistic linkage,” *International Journal for Population Data Science*, vol. 3, 01 2018.
- K. Harron, H. Goldstein, and C. Dibben, *Methodological Developments in Data Linkage*. United States: John Wiley Sons Inc., 2015.
- P. Christen, *Data Matching: Concepts and Techniques for Record Linkage, Entity Resolution, and Duplicate Detection*. Springer Publishing Company, Incorporated, 2012.
- T. N. Herzog, F. J. Scheuren, and W. E. Winkler, *Data Quality and Record Linkage Techniques*, 1st ed. Springer Publishing Company, Incorporated, 2007.
- H. Köpcke and E. Rahm, “Frameworks for entity matching: A comparison,” *Data Knowledge Engineering*, vol. 69, pp. 197–210, 02 2010.
- G. Ridder and R. Moffitt, “The Econometrics of Data Combination,” in *Handbook of Econometrics*, ser. Handbook of Econometrics, J. Heckman and E. Leamer, Eds. Elsevier, January 2007, vol. 6, ch. 75. [Online]. Available: <https://ideas.repec.org/h/eee/ecochp/6b-75.html>
- W. Winkler, “The state of record linkage and current research problems,” *Statist. Med.*, vol. 14, 10 1999.
- M. Hirukawa and A. Prokhorov, “Consistent estimation of linear regression models using matched data,” *Journal of Econometrics*, vol. 203, no. 2, pp. 344 – 358, 2018. [Online]. Available: <http://www.sciencedirect.com/science/article/pii/S0304407617302464>
- I. P. Fellegi and A. B. Sunter, “A theory for record linkage,” *Journal of the American Statistical Association*, vol. 64, pp. 1183–1210, 1969.