

COMPUTER SCIENCE E-20, SPRING 2014
Homework Problems
Induction I, Strong Induction

Due Thursday, February 12, 2015 before 9PM EST. Upload a PDF of your answers at <https://canvas.harvard.edu/courses/1815/assignments/17757>

1. Prove that for all nonnegative integers n

$$\sum_{i=0}^n i^3 = \left(\sum_{i=0}^n i \right)^2$$

Hint: the following identity may be useful

$$\sum_{i=0}^n i = \frac{n(n+1)}{2}$$

Solution:

- Proof: for all nonnegative integers n

$$\sum_{i=0}^n i^3 = \left(\sum_{i=0}^n i \right)^2$$

- The Induction hypothesis $P(n)$, is:

$$0^3 + 1^3 + 2^3 \dots + n^3 = (0 + 1 + 2 + \dots + n)^2$$

$$0^3 + 1^3 + 2^3 \dots + n^3 = (n(n+1)/2)^2$$

- Base Case ($n = 0$) : $0 == 0$
- Base Case ($n = 1$) : $1 == 1$
- Inductive Step : Assume $P(n)$ is true $\forall n \geq 0$, and prove $P(n+1)$

$$0^3 + 1^3 + 2^3 \dots n^3 + (n+1)^3 = (0 + 1 + 2 + \dots + n + n + 1)^2$$

$$0^3 + 1^3 + 2^3 \dots n^3 + (n+1)^3 = ((n+1)(n+2)/2)^2$$

from Inductive Step:

$$0^3 + 1^3 + 2^3 \dots n^3 = (n(n+1)/2)^2$$

Therefore

$$(n(n+1)/2)^2 + (n+1)^3 = ((n+1)(n+2)/2)^2$$

$$\begin{aligned}
(n+1)^2/2^2 (n^2 + 4(n+1)) &= ((n+1)(n+2)/2)^2 \\
(n+1)^2/2^2 (n^2 + 4n + 4) &= ((n+1)(n+2)/2)^2 \\
(n+1)^2/2^2 (n+2)^2 &= ((n+1)(n+2)/2)^2 \\
(n+1)^2(n+2)^2/2^2 &= ((n+1)(n+2)/2)^2 \\
(n+1)(n+2)/2^2 &= ((n+1)(n+2)/2)^2
\end{aligned}$$

This proves $P(n+1)$, completing the proof by induction.

2. Consider the sequence $a_1 = 1, a_2 = 3, \dots, a_n = a_{n-1} + a_{n-2}$. Using strong induction prove that $a_n \leq \left(\frac{7}{4}\right)^n$ for all positive integers n .


Solution:

- Proof: By strong induction on n that for all positive integers n , $a_n \leq \left(\frac{7}{4}\right)^n$.
- The Induction hypothesis $P(n)$, is: $a_n \leq \left(\frac{7}{4}\right)^n \forall n \geq 1$
- Base Case ($n = 1$) : $a_1 \leq \left(\frac{7}{4}\right)^1 = 1 \leq \left(\frac{7}{4}\right) = 1 \leq 1.75$
- Base Case ($n = 2$) : $a_2 \leq \left(\frac{7}{4}\right)^2 = 3 \leq \left(\frac{7}{4}\right)^2 = 3 \leq 1.75 * 1.75 = 3 \leq 3.06$
- Inductive Step : Assume $P(n)$ is true $\forall n \geq 1$, and prove $P(n+1)$: $a_{n+1} \leq \left(\frac{7}{4}\right)^{n+1}$
substituting $(n+1)$ in $a_n = a_{n-1} + a_{n-2}$
 $a_{n+1} = a_{n+1-1} + a_{n+1-2} \leq \left(\frac{7}{4}\right)^{n+1}$
 $= a_n + a_{n-1} \leq \left(\frac{7}{4}\right)^{n+1}$
 $= \left(\frac{7}{4}\right)^n + \left(\frac{7}{4}\right)^{n-1} \leq \left(\frac{7}{4}\right)^{n+1}$
 $= \left(\frac{7}{4}\right)^{n-1} * \left(\frac{7}{4} + 1\right) \leq \left(\frac{7}{4}\right)^{n+1}$
 $= \left(\frac{7}{4}\right)^{n-1} * \left(\frac{7}{4} + 1\right) \leq \left(\frac{7}{4}\right)^n * \left(\frac{7}{4}\right)$
 $= \left(\frac{7}{4}\right)^{n-1} * \left(\frac{7}{4} + 1\right) \leq \left(\frac{7}{4}\right)^{n-1} * \left(\frac{7}{4}\right)^2$
 $= \left(\frac{11}{4}\right) \leq \left(\frac{49}{16}\right)$

This proves $P(n+1)$, completing the proof by induction.

3. Prove using strong induction that any square can be subdivided into n smaller squares, where $n > 5$. For example, the large square below has been subdivided into 6 squares.

Hint: first show that any square subdivided into k squares can easily be subdivided into $k + 3$ squares, then think how many base cases you need show are true (it is not just the case of $n=6$).



4. The Fibonacci numbers are defined by $F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2}$.
Prove using strong induction that for all $n \geq 0$, F_{3n} is even.

Solution:

- Proof: By strong induction on n that for all $n \geq 0$, F_{3n} is even.
- The Induction hypothesis $P(n)$, is: F_{3n} is even $\forall n \geq 0$
- Base Case ($n = 1$) : $F_3 = F_2 + F_1 = F_1 + F_0 + F_1 = 1+0+1 = 2$.
- Base Case ($n = 2$) : $F_6 = F_5 + F_4$
 $F_4 = F_3 + F_2 = 2+1 = 3$
 $F_6 = F_4 + F_3 + F_4 = 3 + 2 + 3 = 8$
- Inductive Step : Assume $P(n)$ is true $\forall n \geq 0$, and prove
 $P(n+1)$: $F_{3(n+1)}$ is even
 substituting $(n+1)$ in $F_n = F_{n-1} + F_{n-2}$

$$F_{3(n+1)} = F_{n+1-1} + F_{n+1-2}$$

$$F_{3(n+1)} = F_n + F_{n-1}$$

From the inductive hypothesis $P(n)$:
 F_n is even. So, $F_n = 2 * a(\text{some integer } a)$
 F_{n-1} is even. So, $F_{n-1} = 2 * b(\text{some integer } b)$

$$F_{3(n+1)} = 2 * a + 2 * b$$

$$F_{3(n+1)} = 2(a + b), \text{ which implies } F_{3(n+1)} \text{ is even.}$$

This proves $P(n+1)$, completing the proof by induction.