

Summary of Halmos' Naive Set Theory

Robin Adams

August 22, 2023

Contents

1	Primitive Terms and Axioms	2
2	The Subset Relation	3
3	Comprehension Notation	4
4	Unordered Pairs	5
5	Unions	6
6	Intersections	7
7	Unordered Triples	9
8	Relative Complements	10
9	Symmetric Difference	13
10	Power Sets	14
11	Ordered Pairs	16
12	Relations	18
13	Functions	20
14	Families	22
15	Inverses and Composites	24

Chapter 1

Primitive Terms and Axioms

Let there be *sets*. We assume that everything is a set.

Let there be a binary relation of *membership*, \in . If $x \in A$ we say that x *belongs to* A , x is an *element* of A , or x is *contained in* A . If this does not hold we write $x \notin A$.

Axiom 1.1 (Axiom of Extensionality). *Two sets are equal if and only if they have the same elements.*

Axiom 1.2 (Axiom of Comprehension, Aussonderungsaxiom). *To every set A and to every condition $S(x)$ there corresponds a set B whose elements are exactly those elements x of A for which $S(x)$ holds.*

Axiom 1.3. *A set exists.*

Axiom 1.4 (Axiom of Pairing). *For any two sets, there exists a set that they both belong to.*

Axiom 1.5 (Union Axiom). *For every set A , there exists a set that contains all the elements that belong to at least one element of A .*

Definition 1.6 (Subset). Let A and B be sets. We say that A is a *subset* of B , or B *includes* A , and write $A \subseteq B$ or $B \supseteq A$, iff every element of A is an element of B .

Axiom 1.7 (Power Set Axiom). *For any set A , there exists a set that contains all the subsets of A .*

Chapter 2

The Subset Relation

Theorem 2.1. *For any set A , we have $A \subseteq A$.*

PROOF: Every element of A is an element of A . \square

Theorem 2.2. *For any sets A , B and C , if $A \subseteq B$ and $B \subseteq C$ then $A \subseteq C$.*

PROOF: If every element of A is an element of B , and every element of B is an element of C , then every element of A is an element of C . \square

Theorem 2.3. *For any sets A and B , if $A \subseteq B$ and $B \subseteq A$ then $A = B$.*

PROOF: If every element of A is an element of B , and every element of B is an element of A , then A and B have the same elements, and therefore are equal by the Axiom of Extensionality. \square

Definition 2.4 (Proper Subset). Let A and B be sets. We say that A is a *proper* subset of B , or B *properly* includes A , and write $A \subsetneq B$ or $B \supsetneq A$, iff $A \subseteq B$ and $A \neq B$.

Chapter 3

Comprehension Notation

Definition 3.1. Given a set A and a condition $S(x)$, we write $\{x \in A : S(x)\}$ for the set whose elements are exactly those elements x of A for which $S(x)$ holds.

PROOF: This exists by the Axiom of Comprehension and is unique by the Axiom of Extensionality. \square

Theorem 3.2. *There is no set that contains every set.*

PROOF:

$\langle 1 \rangle 1.$ LET: A be a set.

PROVE: There exists a set B such that $B \notin A$.

$\langle 1 \rangle 2.$ LET: $B = \{x \in A : x \notin x\}$

$\langle 1 \rangle 3.$ If $B \in A$ then we have $B \in B$ if and only if $B \notin B$.

$\langle 1 \rangle 4.$ $B \notin A$

\square

Chapter 4

Unordered Pairs

Theorem 4.1. *There exists a set with no elements.*

PROOF: Pick a set A by Axiom 1.3. Then the set $\{x \in A : x \neq x\}$ has no elements. \square

Definition 4.2 (Empty Set). The *empty set* \emptyset is the set with no elements.

Theorem 4.3. *For any set A we have $\emptyset \subset A$.*

PROOF: Vacuous. \square

Definition 4.4 ((Unordered) Pair). For any sets a and b , the *(unordered) pair* $\{a, b\}$ is the set whose elements are just a and b .

PROOF: This exists by the Axioms of Pairing and Comprehension, and is unique by the Axiom of Extensionality. \square

Definition 4.5 (Singleton). For any set a , the *singleton* $\{a\}$ is defined to be $\{a, a\}$.

Chapter 5

Unions

Definition 5.1 (Union). For any set \mathcal{C} , the *union* of \mathcal{C} , $\bigcup \mathcal{C}$, is the set whose elements are the elements of the elements of \mathcal{C} .

We write $\bigcup_{X \in \mathcal{A}} t[X]$ for $\bigcup \{t[X] \mid X \in \mathcal{A}\}$.

PROOF: This exists by the Union Axiom and Comprehension Axiom, and is unique by the Axiom of Extensionality. \square

Proposition 5.2.

$$\bigcup \emptyset = \emptyset$$

PROOF: There is no set that is an element of an element of \emptyset . \square

Proposition 5.3. *For any set A , we have $\bigcup \{A\} = A$.*

PROOF: For any x , we have x is an element of an element of $\{A\}$ if and only if x is an element of A . \square

Definition 5.4. We write $A \cup B$ for $\bigcup \{A, B\}$.

Proposition 5.5. *For any set A , we have $A \cup \emptyset = A$.*

PROOF: $x \in A \cup \emptyset$ iff $x \in A$ or $x \in \emptyset$, iff $x \in A$. \square

Proposition 5.6 (Idempotence). *For any set A , we have $A \cup A = A$.*

PROOF: $x \in A$ or $x \in A$ is equivalent to $x \in A$. \square

Proposition 5.7. *For any sets A and B , we have $A \subseteq B$ if and only if $A \cup B = B$.*

PROOF: For any x , the statement "if $x \in A$ then $x \in B$ " is equivalent to " $x \in A$ or $x \in B$ if and only if $x \in B$ ". \square

Proposition 5.8. *For any sets a and b , we have $\{a\} \cup \{b\} = \{a, b\}$.*

PROOF: Immediate from definitions. \square

Chapter 6

Intersections

Definition 6.1 (Intersection). For any sets A and B , the *intersection* $A \cap B$ is defined to be $\{x \in A : x \in B\}$.

Proposition 6.2. For any set A , we have $A \cap \emptyset = \emptyset$.

PROOF: There is no x such that $x \in A$ and $x \in \emptyset$. \square

Proposition 6.3. For any set A , we have

$$A \cap A = A .$$

PROOF: We have $x \in A$ and $x \in A$ if and only if $x \in A$. \square

Proposition 6.4. For any sets A and B , we have $A \subseteq B$ if and only if $A \cap B = A$.

PROOF: For any x , the statement "if $x \in A$ then $x \in B$ " is equivalent to " $x \in A$ and $x \in B$ if and only if $x \in A$ ". \square

Proposition 6.5. For any sets A , B and C , we have $C \subseteq A$ if and only if $(A \cap B) \cup C = A \cap (B \cup C)$.

PROOF: The statement "if $x \in C$ then $x \in A$ " is equivalent to the statement " $((x \in A \wedge x \in B) \vee x \in C) \Leftrightarrow (x \in A \wedge (x \in B \vee x \in C))$ ". \square

Definition 6.6 (Disjoint). Two sets A and B are *disjoint* if and only if $A \cap B = \emptyset$.

Definition 6.7 (Pairwise Disjoint). Let A be a set. We say the elements of A are *pairwise disjoint* if and only if, for all $x, y \in A$, if $x \cap y \neq \emptyset$ then $x = y$.

Definition 6.8 (Intersection). For any nonempty set \mathcal{C} , the *intersection* of \mathcal{C} , $\bigcap \mathcal{C}$, is the set that contains exactly those sets that belong to every element of \mathcal{C} .

We write $\bigcap_{X \in \mathcal{A}} t[X]$ for $\bigcap \{t[X] \mid X \in \mathcal{A}\}$.

PROOF:

$\langle 1 \rangle 1$. LET: \mathcal{C} be a nonempty set.

$\langle 1 \rangle 2$. There exists a set I whose elements are exactly the sets that belong to every element of \mathcal{C} .

PROOF: Pick $A \in \mathcal{C}$, and take $I = \{x \in A : \forall X \in \mathcal{C}. x \in X\}$.

$\langle 1 \rangle 3$. For any sets I, J , if the elements of I and J are exactly the sets that belong to every element of \mathcal{C} then $I = J$.

PROOF: Axiom of Extensionality.

□

Chapter 7

Unordered Triples

Definition 7.1 ((Unordered) Triple). Given sets a_1, \dots, a_n , define the (*unordered*) n -tuple $\{a_1, \dots, a_n\}$ to be

$$\{a_1, \dots, a_n\} := \{a_1\} \cup \dots \cup \{a_n\} \ .$$

Chapter 8

Relative Complements

Definition 8.1 (Relative Complement). For any sets A and B , the *difference* or *relative complement* $A - B$ is defined to be

$$A - B := \{x \in A : x \notin B\} .$$

Proposition 8.2. For any sets A and E , we have $A \subseteq E$ if and only if

$$E - (E - A) = A$$

PROOF:

$\langle 1 \rangle 1$. LET: A and E be sets.

$\langle 1 \rangle 2$. If $A \subseteq E$ then $E - (E - A) = A$

$\langle 2 \rangle 1$. ASSUME: $A \subseteq E$

$\langle 2 \rangle 2$. $E - (E - A) \subseteq A$

PROOF: If $x \in E$ and $x \notin E - A$ then $x \in A$.

$\langle 2 \rangle 3$. $A \subseteq E - (E - A)$

PROOF: If $x \in A$ then $x \in E$ and $x \notin E - A$.

$\langle 1 \rangle 3$. If $E - (E - A) = A$ then $A \subseteq E$.

PROOF: Since $E - (E - A) \subseteq E$.

□

Proposition 8.3. For any set E we have

$$E - \emptyset = E$$

PROOF: $x \in E$ if and only if $x \in E$ and $x \notin \emptyset$. □

Proposition 8.4. For any set E we have

$$E - E = \emptyset .$$

PROOF: There is no x such that $x \in E$ and $x \notin E$. □

Proposition 8.5. For any sets A and E , we have

$$A \cap (E - A) = \emptyset .$$

PROOF: There is no x such that $x \in A$ and $x \in E - A$. \square

Proposition 8.6. *Let A and E be sets. Then $A \subseteq E$ if and only if*

$$A \cup (E - A) = E .$$

PROOF:

$\langle 1 \rangle 1$. LET: A and E be sets.

$\langle 1 \rangle 2$. If $A \subseteq E$ then $A \cup (E - A) = E$.

$\langle 2 \rangle 1$. ASSUME: $A \subseteq E$

$\langle 2 \rangle 2$. $A \cup (E - A) \subseteq E$

PROOF: If $x \in A$ or $x \in E - A$ then $x \in E$.

$\langle 2 \rangle 3$. $E \subseteq A \cup (E - A)$

PROOF: If $x \in E$ then either $x \in A$ or $x \notin A$. In the latter case, $x \in E - A$.

$\langle 1 \rangle 3$. If $A \cup (E - A) = E$ then $A \subseteq E$

PROOF: Since $A \subseteq A \cup (E - A)$.

\square

Proposition 8.7. *Let A , B and E be sets. Then:*

1. *If $A \subseteq B$ then $E - B \subseteq E - A$.*

2. *If $A \subseteq E$ and $E - B \subseteq E - A$ then $A \subseteq B$.*

PROOF:

$\langle 1 \rangle 1$. LET: A , B and E be sets.

$\langle 1 \rangle 2$. If $A \subseteq B$ then $E - B \subseteq E - A$.

PROOF: If $A \subseteq B$, $x \in E$ and $x \notin B$, then we have $x \in E$ and $x \notin A$.

$\langle 1 \rangle 3$. If $A \subseteq E$ and $E - B \subseteq E - A$ then $A \subseteq B$.

$\langle 2 \rangle 1$. ASSUME: $A \subseteq E$

$\langle 2 \rangle 2$. ASSUME: $E - B \subseteq E - A$

$\langle 2 \rangle 3$. LET: $x \in A$

$\langle 2 \rangle 4$. $x \in E$

$\langle 2 \rangle 5$. $x \notin E - A$

$\langle 2 \rangle 6$. $x \notin E - B$

$\langle 2 \rangle 7$. $x \in B$

\square

Example 8.8. We cannot remove the hypothesis $A \subseteq E$ in item 2 above. Let $E = \emptyset$, $A = \{\emptyset\}$ and $B = \emptyset$. Then $E - B = E - A = \emptyset$ but $A \not\subseteq B$.

Proposition 8.9 (De Morgan's Law). *For any sets A , B and E , we have $E - (A \cup B) = (E - A) \cap (E - B)$.*

PROOF: $(x \in E \wedge \neg(x \in A \vee x \in B)) \Leftrightarrow (x \in E \wedge x \notin A \wedge x \in E \wedge x \notin B)$. \square

Proposition 8.10 (De Morgan's Law). *For any sets A , B and E , we have $E - (A \cap B) = (E - A) \cup (E - B)$.*

PROOF: $(x \in E \vee \neg(x \in A \wedge x \in B)) \Leftrightarrow (x \in E \wedge x \notin A) \vee (x \in E \wedge x \notin B)$. \square

Proposition 8.11. *For any sets A , B and E , if $A \subseteq E$ then*

$$A - B = A \cap (E - B) .$$

PROOF: If $A \subseteq E$ then we have $(x \in A \wedge x \notin B) \Leftrightarrow (x \in A \wedge x \in E \wedge x \notin B)$. \square

Proposition 8.12. *For any sets A and B , we have $A \subseteq B$ if and only if $A - B = \emptyset$.*

PROOF: Both are equivalent to the statement that there is no x such that $x \in A$ and $x \notin B$. \square

Proposition 8.13. *For any sets A and B , we have*

$$A - (A - B) = A \cap B .$$

PROOF: $(x \in A \wedge \neg(x \in A \wedge x \notin B)) \Leftrightarrow x \in A \wedge x \in B$. \square

Proposition 8.14. *For any sets A , B and C , we have*

$$A \cap (B - C) = (A \cap B) - (A \cap C) .$$

PROOF: $(x \in A \wedge x \in B \wedge x \notin C) \Leftrightarrow (x \in A \wedge x \in B \wedge \neg(x \in A \wedge x \in C))$. \square

Proposition 8.15. *For any sets A , B , C and E , if $(A \cap B) - C \subseteq E$ then we have*

$$A \cap B \subseteq (A \cap C) \cup (B \cap (E - C)) .$$

PROOF:

$\langle 1 \rangle 1$. LET: $x \in A \cap B$

PROVE: $x \in (A \cap C) \cup (B \cap (E - C))$

$\langle 1 \rangle 2$. CASE: $x \in C$

PROOF: Then $x \in A \cap C$.

$\langle 1 \rangle 3$. CASE: $x \notin C$

PROOF: Then $x \in E$ and so $x \in B \cap (E - C)$.

\square

Proposition 8.16. *For any sets A , B , C and E , we have*

$$(A \cup C) \cap (B \cup (E - C)) \subseteq A \cup B .$$

PROOF: The statement $(x \in A \vee x \in C) \wedge (x \in B \vee (x \in E \wedge x \notin C))$ implies $x \in A \vee x \in B$. \square

Proposition 8.17 (De Morgan's Law). *Let E be a set and \mathcal{C} a nonempty set. Then*

$$E - \bigcup_{X \in \mathcal{C}} X = \bigcap_{X \in \mathcal{C}} (E - X) .$$

PROOF: Easy. \square

Proposition 8.18 (De Morgan's Law). *Let E be a set and \mathcal{C} a nonempty set. Then*

$$E - \bigcap_{X \in \mathcal{C}} X = \bigcup_{X \in \mathcal{C}} (E - X) .$$

PROOF: Easy. \square

Chapter 9

Symmetric Difference

Definition 9.1 (Symmetric Difference). For any sets A and B , the *symmetric difference* $A + B$ is defined to be

$$A + B := (A - B) \cup (B - A) .$$

Proposition 9.2. For any sets A and B , we have

$$A + B = B + A$$

PROOF: From the commutativity of union. \square

Proposition 9.3. For any sets A , B and C , we have

$$A + (B + C) = (A + B) + C .$$

PROOF: Each is the set of all x that belong to either exactly one or all three of A , B and C . \square

Proposition 9.4. For any set A , we have

$$A + \emptyset = A .$$

PROOF:

$$\begin{aligned} A + \emptyset &= (A - \emptyset) \cup (\emptyset - A) \\ &= A \cup \emptyset \\ &= A \end{aligned}$$

\square

Proposition 9.5. For any set A we have

$$A + A = \emptyset .$$

PROOF:

$$\begin{aligned} A + A &= (A - A) \cup (A - A) \\ &= \emptyset \cup \emptyset \\ &= \emptyset \end{aligned}$$

\square

Chapter 10

Power Sets

Definition 10.1 (Power Set). For any set A , the *power set* of A , $\mathcal{P}A$, is the set whose elements are exactly the subsets of A .

PROOF: This exists by the Power Set Axiom and Axiom of Comprehension, and is unique by the Axiom of Extensionality. \square

Proposition 10.2.

$$\mathcal{P}\emptyset = \{\emptyset\}$$

PROOF: The only subset of \emptyset is \emptyset . \square

Proposition 10.3. For any set a , we have

$$\mathcal{P}\{a\} = \{\emptyset, \{a\}\} .$$

PROOF: The only subsets of $\{a\}$ are \emptyset and $\{a\}$. \square

Proposition 10.4. For any sets a and b , we have

$$\mathcal{P}\{a, b\} = \{\emptyset, \{a\}, \{b\}, \{a, b\}\} .$$

PROOF: The only subsets of $\{a, b\}$ are \emptyset , $\{a\}$, $\{b\}$ and $\{a, b\}$. \square

Proposition 10.5. For any nonempty set \mathcal{C} we have

$$\bigcap_{X \in \mathcal{C}} \mathcal{P}X = \mathcal{P}\left(\bigcap \mathcal{C}\right) .$$

PROOF:

$$x \in \bigcup_{X \in \mathcal{C}} \mathcal{P}X \Leftrightarrow \forall X \in \mathcal{C}. x \subseteq X$$

$$\Leftrightarrow \forall X \in \mathcal{C}. \forall y \in x. y \in X$$

$$\Leftrightarrow \forall y \in x. \forall X \in \mathcal{C}. y \in X$$

$$\Leftrightarrow x \subseteq \bigcap \mathcal{C}$$

\square

Proposition 10.6. *For any set \mathcal{C} we have*

$$\bigcup_{X \in \mathcal{C}} \mathcal{P}X \subseteq \mathcal{P} \bigcup \mathcal{C} .$$

PROOF: If there exists $X \in \mathcal{C}$ such that $x \subseteq X$ then $x \subseteq \bigcup \mathcal{C}$. \square

Proposition 10.7. *For any set E , we have*

$$\bigcap \mathcal{P}E = \emptyset .$$

PROOF: Since $\emptyset \in \mathcal{P}E$. \square

Proposition 10.8. *For any sets E and F , if $E \subseteq F$ then $\mathcal{P}E \subseteq \mathcal{P}F$.*

PROOF: If $E \subseteq F$ and $X \subseteq E$ then $X \subseteq F$. \square

Chapter 11

Ordered Pairs

Definition 11.1 (Ordered Pair). For any sets a and b , the *ordered pair* (a, b) is defined by

$$(a, b) := \{\{a\}, \{a, b\}\} .$$

Proposition 11.2. For any sets a, b, x and y , if $(a, b) = (x, y)$ then $a = x$ and $b = y$.

PROOF:

$\langle 1 \rangle 1$. LET: a, b, x and y be sets.

$\langle 1 \rangle 2$. ASSUME: $(a, b) = (x, y)$

$\langle 1 \rangle 3$. $a = x$

PROOF: $\{a\} = \bigcap (a, b) = \bigcap (x, y) = \{x\}$.

$\langle 1 \rangle 4$. $\{a, b\} = \{x, y\}$

$\langle 1 \rangle 5$. CASE: $a = b$

$\langle 2 \rangle 1$. $x = y$

PROOF: Since $\{x, y\} = \{a, b\}$ is a singleton.

$\langle 2 \rangle 2$. $b = y$

PROOF: $b = a = x = y$

$\langle 1 \rangle 6$. CASE: $a \neq b$

$\langle 2 \rangle 1$. $x \neq y$

PROOF: Since $\{x, y\} = \{a, b\}$ is not a singleton.

$\langle 2 \rangle 2$. $b = y$

PROOF: $\{b\} = \{a, b\} - \{a\} = \{x, y\} - \{x\} = \{y\}$.

□

Definition 11.3 (Cartesian Product). For any sets A and B , the *Cartesian product* $A \times B$ is

$$A \times B := \{p \in \mathcal{PP}(A \cup B) : \exists a \in A. \exists b \in B. p = (a, b)\} .$$

Proposition 11.4. For any sets A, B and X , we have

$$(A - B) \times X = (A \times X) - (B \times X) .$$

PROOF: Easy. \square

Proposition 11.5. *For any sets A and B , we have $A \times B = \emptyset$ if and only if $A = \emptyset$ or $B = \emptyset$.*

PROOF: Easy. \square

Proposition 11.6. *For any sets A , B , X and Y , if $A \subseteq X$ and $B \subseteq Y$ then $A \times B \subseteq X \times Y$. The converse holds assuming $A \neq \emptyset$ and $B \neq \emptyset$.*

PROOF: Easy. \square

Chapter 12

Relations

Definition 12.1 (Relation). A *relation* is a set of ordered pairs.

If R is a relation, we write xRy for $(x, y) \in R$.

Given sets X and Y , a relation *between X and Y* is a subset of $X \times Y$.

Given a set X , a relation *on X* is a relation between X and X .

Definition 12.2 (Domain). The *domain* of a relation R is the set

$$\text{dom } R := \{x \in \bigcup \bigcup R : \exists y. (x, y) \in R\} \ .$$

Definition 12.3 (Range). The *range* of a relation R is the set

$$\text{ran } R := \{y \in \bigcup \bigcup R : \exists x. (x, y) \in R\} \ .$$

Definition 12.4 (Reflexive). Let R be a relation on X . Then R is *reflexive* iff, for all $x \in X$, we have xRx .

Definition 12.5 (Symmetric). Let R be a relation on X . Then R is *symmetric* iff, whenever xRy , then yRx .

Definition 12.6 (Transitive). Let R be a relation on X . Then R is *transitive* iff, whenever xRy and yRz , then xRz .

Definition 12.7 (Equivalence Relation). Let R be a relation on X . Then R is an *equivalence relation* iff it is reflexive, symmetric and transitive.

Definition 12.8 (Partition). Let X be a set. A *partition* of X is a pairwise disjoint set of nonempty subsets of X whose union is X .

Definition 12.9 (Equivalence Class). Let R be an equivalence relation on X . Let $x \in X$. The *equivalence class* of x with respect to R is

$$x/R := \{y \in X : xRy\} \ .$$

We write X/R for the set of all equivalence classes with respect to R .

Definition 12.10 (Induced). Let P be a partition of X . The relation *induced* by P is X/P where $x(X/P)y$ iff there exists $X \in P$ such that $x \in X$ and $y \in X$.

Theorem 12.11. *Let R be an equivalence relation on X . Then X/R is a partition of X that induces the relation R .*

PROOF: Easy. \square

Theorem 12.12. *Let P be a partition of X . Then X/P is an equivalence relation on X , and $P = X/(X/P)$.*

PROOF: Easy. \square

Chapter 13

Functions

Definition 13.1 (Function). Let X and Y be sets. A *function*, *map*, *mapping*, *transformation* or *operator* f from X to Y , $f : X \rightarrow Y$, is a relation f between X and Y such that, for all $x \in X$, there exists a unique $f(x) \in Y$, called the *value* of f at the *argument* x , such that $(x, f(x)) \in f$.

Definition 13.2 (Onto). Let $f : X \rightarrow Y$. We say f maps X *onto* Y iff $\text{ran } f = Y$.

Definition 13.3 (Image). Let $f : X \rightarrow Y$ and $A \subseteq X$. The *image* of A under f is

$$f(A) := \{f(x) : x \in A\} .$$

Definition 13.4 (Inclusion Map). Let Y be a set and $X \subseteq Y$. Then the *inclusion map* $i : X \hookrightarrow Y$ is the function defined by $i(x) = x$ for all $x \in X$.

Definition 13.5 (Identity Map). For any set X , the *identity map* on X is the inclusion map of X into X .

Definition 13.6 (Restriction). Let $f : Y \rightarrow Z$ and $X \subseteq Y$. The *restriction* of f to X is the function $f \upharpoonright X : X \rightarrow Z$ defined by

$$(f \upharpoonright X)(x) = f(x) \quad (x \in X) .$$

Given sets X , Y and Z with $X \subseteq Y$, if $f : X \rightarrow Z$ and $g : Y \rightarrow Z$, we say g is an *extension* of f to Y iff $f = g \upharpoonright X$.

Definition 13.7 (Projection). Given sets X and Y , the *projection* maps $\pi_1 : X \times Y \rightarrow X$ and $\pi_2 : X \times Y \rightarrow Y$ are defined by

$$\pi_1(x, y) = x, \quad \pi_2(x, y) = y \quad (x \in X, y \in Y) .$$

Definition 13.8 (Canonical Map). Let X be a set and R an equivalence relation on X . The *canonical map* $\pi : X \rightarrow X/R$ is the map defined by $\pi(x) = x/R$.

Definition 13.9 (One-to-One). A function $f : X \rightarrow Y$ is *one-to-one*, or a *one-to-one correspondence*, iff, for all $x, y \in X$, if $f(x) = f(y)$ then $x = y$.

Definition 13.10.

$$0 = \emptyset$$

Definition 13.11.

$$1 = \{\emptyset\}$$

Definition 13.12.

$$2 = \{\emptyset, \{\emptyset\}\}$$

Definition 13.13 (Characteristic Function). Let X be a set and $A \subseteq X$. The *characteristic function* of A is the function $\chi_A : X \rightarrow 2$ defined by

$$\chi_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

Chapter 14

Families

Definition 14.1 (Family). Let I and X be sets. A *family* of elements of X indexed by I is a function $a : I \rightarrow X$. We write a_i for $a(i)$, and $\{a_i\}_{i \in I}$ for a .

Proposition 14.2 (Generalized Associative Law for Unions). Let $\{I_j\}_{j \in J}$ be a family of sets. Let $K = \bigcup_{j \in J} I_j$. Let $\{A_k\}_{k \in K}$ be a family of sets indexed by K . Then

$$\bigcup_{k \in K} A_k = \bigcup_{j \in J} \bigcup_{i \in I_j} A_i .$$

PROOF: Easy. \square

Proposition 14.3 (Generalized Commutative Law for Unions). Let $\{I_j\}_{j \in J}$ be a family of sets. Let $f : J \rightarrow J$ be a one-to-one correspondence from J onto J . Then

$$\bigcup_{j \in J} I_j = \bigcup_{j \in J} I_{f(j)} .$$

PROOF: Easy. \square

Proposition 14.4 (Generalized Associative Law for Intersections). Let $\{I_j\}_{j \in J}$ be a nonempty family of nonempty sets. Let $K = \bigcup_{j \in J} I_j$. Let $\{A_k\}_{k \in K}$ be a family of sets indexed by K . Then

$$\bigcap_{k \in K} A_k = \bigcap_{j \in J} \bigcap_{i \in I_j} A_i .$$

PROOF: Easy. \square

Proposition 14.5 (Generalized Commutative Law for Intersections). Let $\{I_j\}_{j \in J}$ be a nonempty family of sets. Let $f : J \rightarrow J$ be a one-to-one correspondence from J onto J . Then

$$\bigcap_{j \in J} I_j = \bigcap_{j \in J} I_{f(j)} .$$

PROOF: Easy. \square

Proposition 14.6. *Let B be a set and $\{A_i\}_{i \in I}$ a family of sets. Then*

$$B \cap \bigcup_{i \in I} A_i = \bigcup_{i \in I} (B \cap A_i)$$

PROOF: Easy. \square

Proposition 14.7. *Let B be a set and $\{A_i\}_{i \in I}$ a nonempty family of sets. Then*

$$B \cup \bigcap_{i \in I} A_i = \bigcap_{i \in I} (B \cup A_i)$$

PROOF: Easy. \square

Definition 14.8 (Cartesian Product of a Family of Sets). Let $\{A_i\}_{i \in I}$ be a family of sets. The *Cartesian product* $\times_{i \in I} A_i$ is the set of all families $\{a_i\}_{i \in I}$ such that $\forall i \in I, a_i \in A_i$.

We write A^I for $\times_{i \in I} A_i$.

Theorem 14.9. *Let X be a set. The function $\chi : \mathcal{P}X \rightarrow 2^X$ that maps a subset A of X to χ_A is a one-to-one correspondence.*

PROOF: Easy. \square

Definition 14.10 (Projection). Let $\{A_i\}_{i \in I}$ be a family of sets and $i \in I$. The *projection* function $\pi_i : \times_{i \in I} A_i \rightarrow A_i$ is defined by $\pi_i(a) = a_i$.

Proposition 14.11. *Let $\{A_i\}_{i \in I}$ and $\{B_j\}_{j \in J}$ be families of sets. Then*

$$\left(\bigcup_{i \in I} A_i \right) \times \left(\bigcup_{j \in J} B_j \right) = \bigcup_{i \in I} \bigcup_{j \in J} (A_i \times B_j) .$$

PROOF: Easy. \square

Proposition 14.12. *Let $\{A_i\}_{i \in I}$ and $\{B_j\}_{j \in J}$ be nonempty families of sets. Then*

$$\left(\bigcap_{i \in I} A_i \right) \times \left(\bigcap_{j \in J} B_j \right) = \bigcap_{i \in I} \bigcap_{j \in J} (A_i \times B_j) .$$

PROOF: Easy. \square

Proposition 14.13. *Let $f : X \rightarrow Y$. Let $\{A_i\}_{i \in I}$ be a family of subsets of X . Then*

$$f \left(\bigcup_{i \in I} A_i \right) = \bigcup_{i \in I} f(A_i) .$$

PROOF: Easy. \square

Example 14.14. It is not true in general that, if $f : X \rightarrow Y$ and $\{A_i\}_{i \in I}$ is a nonempty family of subsets of X , then $f(\bigcap_{i \in I} A_i) = \bigcap_{i \in I} f(A_i)$.

Take $X = \{a, b\}$ and $Y = \{c\}$ where $a \neq b$. Take $I = \{i, j\}$ with $i \neq j$. Let $A_i = \{a\}$ and $A_j = \{b\}$. Let f be the unique function $X \rightarrow Y$. Then $f(\bigcap_{i \in I} A_i) = f(\emptyset) = \emptyset$ but $\bigcap_{i \in I} f(A_i) = \{c\}$.

Chapter 15

Inverses and Composites

Definition 15.1 (Inverse). Given a function $f : X \rightarrow Y$, the *inverse* of f is the function $f^{-1} : \mathcal{P}Y \rightarrow \mathcal{P}X$ defined by

$$f^{-1}(B) = \{x \in X : f(x) \in B\} \ .$$

We call $f^{-1}(B)$ the *inverse image* of B under f .

Proposition 15.2. *Let $f : X \rightarrow Y$. Then f maps X onto Y if and only if the inverse image of any nonempty subset of Y is nonempty.*

PROOF: Easy. \square

Proposition 15.3. *Let $f : X \rightarrow Y$. Then f is one-to-one if and only if the inverse image of any singleton subset of Y is a singleton.*

PROOF: Easy. \square

Proposition 15.4. *Let $f : X \rightarrow Y$. Let $B \subseteq Y$. Then*

$$f(f^{-1}(B)) \subseteq B \ .$$

PROOF: Easy. \square

Proposition 15.5. *Let $f : X \rightarrow Y$. Let $A \subseteq X$. Then*

$$A \subseteq f^{-1}(f(A)) \ .$$

Equality holds if f is one-to-one.

PROOF: Easy. \square

Proposition 15.6. *Let $f : X \rightarrow Y$. Let $\{B_i\}_{i \in I}$ be a family of subsets of Y . Then*

$$f^{-1}\left(\bigcup_{i \in I} B_i\right) = \bigcup_{i \in I} f^{-1}(B_i) \ .$$

PROOF: Easy. \square

Proposition 15.7. *Let $f : X \rightarrow Y$. Let $\{B_i\}_{i \in I}$ be a nonempty family of subsets of Y . Then*

$$f^{-1} \left(\bigcap_{i \in I} B_i \right) = \bigcap_{i \in I} f^{-1}(B_i) .$$

PROOF: Easy. \square

Proposition 15.8. *Let $f : X \rightarrow Y$ and $B \subseteq Y$. Then $f^{-1}(Y - B) = X - f^{-1}(B)$.*

PROOF: Easy. \square

Definition 15.9. Let $f : X \rightarrow Y$ be one-to-one. Then we write f^{-1} for the function $f^{-1} : \text{ran } f \rightarrow X$ defined by: $f^{-1}(y)$ is the unique x such that $f(x) = y$.

Definition 15.10 (Composite). Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. The *composite* $g \circ f = gf : X \rightarrow Z$ is the function defined by

$$(g \circ f)(x) = g(f(x)) \quad (x \in X) .$$

Proposition 15.11. *Let $f : X \rightarrow Y$, $g : Y \rightarrow Z$ and $h : Z \rightarrow W$. Then*

$$h(gf) = (hg)f .$$

PROOF: Easy. \square

Example 15.12. Function composition is not commutative in general.

Let $X = \{a, b\}$ where $a \neq b$. Define $f : X \rightarrow Y$ by $f(a) = f(b) = a$, and g by $g(a) = g(b) = b$. Then $gf = g$ but $fg = f$.

Proposition 15.13. *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. Then*

$$(gf)^{-1} = f^{-1}g^{-1} : \mathcal{P}Z \rightarrow \mathcal{P}X .$$

PROOF: Easy. \square

Definition 15.14 (Inverse). Let R be a relation between X and Y . The *inverse* or *converse* R^{-1} is the relation between Y and X defined by

$$yR^{-1}x \Leftrightarrow xRy .$$

Proposition 15.15. *For any relation R , we have*

$$\text{dom } R^{-1} = \text{ran } R .$$

PROOF: Easy. \square

Proposition 15.16. *For any relation R , we have*

$$\text{ran } R^{-1} = \text{dom } R .$$

PROOF: Easy. \square