

# Summary of Halmos' Naive Set Theory

Robin Adams

August 23, 2023

# Contents

1	Primitive Terms and Axioms	2
2	The Subset Relation	3
3	Comprehension Notation	4
4	Unordered Pairs	5
5	Unions	6
6	Intersections	7
7	Unordered Triples	9
8	Relative Complements	10
9	Symmetric Difference	13
10	Power Sets	14
11	Ordered Pairs	16
12	Relations	18
13	Functions	21
14	Families	23
15	Inverses and Composites	25
16	Numbers	27
17	The Peano Axioms	29
18	Arithmetic	33
19	Order	45

# Chapter 1

## Primitive Terms and Axioms

Let there be *sets*. We assume that everything is a set.

Let there be a binary relation of *membership*,  $\in$ . If  $x \in A$  we say that  $x$  *belongs to*  $A$ ,  $x$  is an *element* of  $A$ , or  $x$  is *contained in*  $A$ . If this does not hold we write  $x \notin A$ .

**Axiom 1.1** (Axiom of Extensionality). *Two sets are equal if and only if they have the same elements.*

**Axiom 1.2** (Axiom of Comprehension, Aussonderungsaxiom). *To every set  $A$  and to every condition  $S(x)$  there corresponds a set  $B$  whose elements are exactly those elements  $x$  of  $A$  for which  $S(x)$  holds.*

**Axiom 1.3** (Axiom of Pairing). *For any two sets, there exists a set that they both belong to.*

**Axiom 1.4** (Union Axiom). *For every set  $A$ , there exists a set that contains all the elements that belong to at least one element of  $A$ .*

**Definition 1.5** (Subset). Let  $A$  and  $B$  be sets. We say that  $A$  is a *subset* of  $B$ , or  $B$  *includes*  $A$ , and write  $A \subseteq B$  or  $B \supseteq A$ , iff every element of  $A$  is an element of  $B$ .

**Axiom 1.6** (Power Set Axiom). *For any set  $A$ , there exists a set that contains all the subsets of  $A$ .*

**Axiom 1.7** (Axiom of Infinity). *There exists a set  $I$  such that:*

- *$I$  has an element that has no elements*
- *for all  $x \in I$ , there exists  $y \in I$  such that the elements of  $y$  are exactly  $x$  and the elements of  $x$ .*

## Chapter 2

# The Subset Relation

**Theorem 2.1.** *For any set  $A$ , we have  $A \subseteq A$ .*

PROOF: Every element of  $A$  is an element of  $A$ .  $\square$

**Theorem 2.2.** *For any sets  $A$ ,  $B$  and  $C$ , if  $A \subseteq B$  and  $B \subseteq C$  then  $A \subseteq C$ .*

PROOF: If every element of  $A$  is an element of  $B$ , and every element of  $B$  is an element of  $C$ , then every element of  $A$  is an element of  $C$ .  $\square$

**Theorem 2.3.** *For any sets  $A$  and  $B$ , if  $A \subseteq B$  and  $B \subseteq A$  then  $A = B$ .*

PROOF: If every element of  $A$  is an element of  $B$ , and every element of  $B$  is an element of  $A$ , then  $A$  and  $B$  have the same elements, and therefore are equal by the Axiom of Extensionality.  $\square$

**Definition 2.4** (Proper Subset). Let  $A$  and  $B$  be sets. We say that  $A$  is a *proper* subset of  $B$ , or  $B$  *properly* includes  $A$ , and write  $A \subsetneq B$  or  $B \supsetneq A$ , iff  $A \subseteq B$  and  $A \neq B$ .

## Chapter 3

# Comprehension Notation

**Definition 3.1.** Given a set  $A$  and a condition  $S(x)$ , we write  $\{x \in A : S(x)\}$  for the set whose elements are exactly those elements  $x$  of  $A$  for which  $S(x)$  holds.

PROOF: This exists by the Axiom of Comprehension and is unique by the Axiom of Extensionality.  $\square$

**Theorem 3.2.** *There is no set that contains every set.*

PROOF:

$\langle 1 \rangle 1.$  LET:  $A$  be a set.

PROVE: There exists a set  $B$  such that  $B \notin A$ .

$\langle 1 \rangle 2.$  LET:  $B = \{x \in A : x \notin x\}$

$\langle 1 \rangle 3.$  If  $B \in A$  then we have  $B \in B$  if and only if  $B \notin B$ .

$\langle 1 \rangle 4.$   $B \notin A$

$\square$

## Chapter 4

# Unordered Pairs

**Theorem 4.1.** *There exists a set with no elements.*

PROOF: Immediate from the Axiom of Infinity.  $\square$

**Definition 4.2** (Empty Set). The *empty set*  $\emptyset$  is the set with no elements.

**Theorem 4.3.** *For any set  $A$  we have  $\emptyset \subset A$ .*

PROOF: Vacuous.  $\square$

**Definition 4.4** ((Unordered) Pair). For any sets  $a$  and  $b$ , the *(unordered) pair*  $\{a, b\}$  is the set whose elements are just  $a$  and  $b$ .

PROOF: This exists by the Axioms of Pairing and Comprehension, and is unique by the Axiom of Extensionality.  $\square$

**Definition 4.5** (Singleton). For any set  $a$ , the *singleton*  $\{a\}$  is defined to be  $\{a, a\}$ .

# Chapter 5

## Unions

**Definition 5.1** (Union). For any set  $\mathcal{C}$ , the *union* of  $\mathcal{C}$ ,  $\bigcup \mathcal{C}$ , is the set whose elements are the elements of the elements of  $\mathcal{C}$ .

We write  $\bigcup_{X \in \mathcal{A}} t[X]$  for  $\bigcup \{t[X] \mid X \in \mathcal{A}\}$ .

PROOF: This exists by the Union Axiom and Comprehension Axiom, and is unique by the Axiom of Extensionality.  $\square$

**Proposition 5.2.**

$$\bigcup \emptyset = \emptyset$$

PROOF: There is no set that is an element of an element of  $\emptyset$ .  $\square$

**Proposition 5.3.** *For any set  $A$ , we have  $\bigcup \{A\} = A$ .*

PROOF: For any  $x$ , we have  $x$  is an element of an element of  $\{A\}$  if and only if  $x$  is an element of  $A$ .  $\square$

**Definition 5.4.** We write  $A \cup B$  for  $\bigcup \{A, B\}$ .

**Proposition 5.5.** *For any set  $A$ , we have  $A \cup \emptyset = A$ .*

PROOF:  $x \in A \cup \emptyset$  iff  $x \in A$  or  $x \in \emptyset$ , iff  $x \in A$ .  $\square$

**Proposition 5.6** (Idempotence). *For any set  $A$ , we have  $A \cup A = A$ .*

PROOF:  $x \in A$  or  $x \in A$  is equivalent to  $x \in A$ .  $\square$

**Proposition 5.7.** *For any sets  $A$  and  $B$ , we have  $A \subseteq B$  if and only if  $A \cup B = B$ .*

PROOF: For any  $x$ , the statement "if  $x \in A$  then  $x \in B$ " is equivalent to " $x \in A$  or  $x \in B$  if and only if  $x \in B$ ".  $\square$

**Proposition 5.8.** *For any sets  $a$  and  $b$ , we have  $\{a\} \cup \{b\} = \{a, b\}$ .*

PROOF: Immediate from definitions.  $\square$

## Chapter 6

# Intersections

**Definition 6.1** (Intersection). For any sets  $A$  and  $B$ , the *intersection*  $A \cap B$  is defined to be  $\{x \in A : x \in B\}$ .

**Proposition 6.2.** For any set  $A$ , we have  $A \cap \emptyset = \emptyset$ .

PROOF: There is no  $x$  such that  $x \in A$  and  $x \in \emptyset$ .  $\square$

**Proposition 6.3.** For any set  $A$ , we have

$$A \cap A = A .$$

PROOF: We have  $x \in A$  and  $x \in A$  if and only if  $x \in A$ .  $\square$

**Proposition 6.4.** For any sets  $A$  and  $B$ , we have  $A \subseteq B$  if and only if  $A \cap B = A$ .

PROOF: For any  $x$ , the statement "if  $x \in A$  then  $x \in B$ " is equivalent to " $x \in A$  and  $x \in B$  if and only if  $x \in A$ ".  $\square$

**Proposition 6.5.** For any sets  $A$ ,  $B$  and  $C$ , we have  $C \subseteq A$  if and only if  $(A \cap B) \cup C = A \cap (B \cup C)$ .

PROOF: The statement "if  $x \in C$  then  $x \in A$ " is equivalent to the statement " $((x \in A \wedge x \in B) \vee x \in C) \Leftrightarrow (x \in A \wedge (x \in B \vee x \in C))$ ".  $\square$

**Definition 6.6** (Disjoint). Two sets  $A$  and  $B$  are *disjoint* if and only if  $A \cap B = \emptyset$ .

**Definition 6.7** (Pairwise Disjoint). Let  $A$  be a set. We say the elements of  $A$  are *pairwise disjoint* if and only if, for all  $x, y \in A$ , if  $x \cap y \neq \emptyset$  then  $x = y$ .

**Definition 6.8** (Intersection). For any nonempty set  $\mathcal{C}$ , the *intersection* of  $\mathcal{C}$ ,  $\bigcap \mathcal{C}$ , is the set that contains exactly those sets that belong to every element of  $\mathcal{C}$ .

We write  $\bigcap_{X \in \mathcal{A}} t[X]$  for  $\bigcap \{t[X] \mid X \in \mathcal{A}\}$ .



PROOF:

$\langle 1 \rangle 1$ . LET:  $\mathcal{C}$  be a nonempty set.

$\langle 1 \rangle 2$ . There exists a set  $I$  whose elements are exactly the sets that belong to every element of  $\mathcal{C}$ .

PROOF: Pick  $A \in \mathcal{C}$ , and take  $I = \{x \in A : \forall X \in \mathcal{C}. x \in X\}$ .

$\langle 1 \rangle 3$ . For any sets  $I, J$ , if the elements of  $I$  and  $J$  are exactly the sets that belong to every element of  $\mathcal{C}$  then  $I = J$ .

PROOF: Axiom of Extensionality.

□

## Chapter 7

# Unordered Triples

**Definition 7.1** ((Unordered) Triple). Given sets  $a_1, \dots, a_n$ , define the (*unordered*)  $n$ -tuple  $\{a_1, \dots, a_n\}$  to be

$$\{a_1, \dots, a_n\} := \{a_1\} \cup \dots \cup \{a_n\} \ .$$

## Chapter 8

# Relative Complements

**Definition 8.1** (Relative Complement). For any sets  $A$  and  $B$ , the *difference* or *relative complement*  $A - B$  is defined to be

$$A - B := \{x \in A : x \notin B\} .$$

**Proposition 8.2.** For any sets  $A$  and  $E$ , we have  $A \subseteq E$  if and only if

$$E - (E - A) = A$$

PROOF:

$\langle 1 \rangle 1$ . LET:  $A$  and  $E$  be sets.

$\langle 1 \rangle 2$ . If  $A \subseteq E$  then  $E - (E - A) = A$

$\langle 2 \rangle 1$ . ASSUME:  $A \subseteq E$

$\langle 2 \rangle 2$ .  $E - (E - A) \subseteq A$

PROOF: If  $x \in E$  and  $x \notin E - A$  then  $x \in A$ .

$\langle 2 \rangle 3$ .  $A \subseteq E - (E - A)$

PROOF: If  $x \in A$  then  $x \in E$  and  $x \notin E - A$ .

$\langle 1 \rangle 3$ . If  $E - (E - A) = A$  then  $A \subseteq E$ .

PROOF: Since  $E - (E - A) \subseteq E$ .

□

**Proposition 8.3.** For any set  $E$  we have

$$E - \emptyset = E$$

PROOF:  $x \in E$  if and only if  $x \in E$  and  $x \notin \emptyset$ . □

**Proposition 8.4.** For any set  $E$  we have

$$E - E = \emptyset .$$

PROOF: There is no  $x$  such that  $x \in E$  and  $x \notin E$ . □

**Proposition 8.5.** For any sets  $A$  and  $E$ , we have

$$A \cap (E - A) = \emptyset .$$

PROOF: There is no  $x$  such that  $x \in A$  and  $x \in E - A$ .  $\square$

**Proposition 8.6.** *Let  $A$  and  $E$  be sets. Then  $A \subseteq E$  if and only if*

$$A \cup (E - A) = E .$$

PROOF:

$\langle 1 \rangle 1$ . LET:  $A$  and  $E$  be sets.

$\langle 1 \rangle 2$ . If  $A \subseteq E$  then  $A \cup (E - A) = E$ .

$\langle 2 \rangle 1$ . ASSUME:  $A \subseteq E$

$\langle 2 \rangle 2$ .  $A \cup (E - A) \subseteq E$

PROOF: If  $x \in A$  or  $x \in E - A$  then  $x \in E$ .

$\langle 2 \rangle 3$ .  $E \subseteq A \cup (E - A)$

PROOF: If  $x \in E$  then either  $x \in A$  or  $x \notin A$ . In the latter case,  $x \in E - A$ .

$\langle 1 \rangle 3$ . If  $A \cup (E - A) = E$  then  $A \subseteq E$

PROOF: Since  $A \subseteq A \cup (E - A)$ .

$\square$

**Proposition 8.7.** *Let  $A$ ,  $B$  and  $E$  be sets. Then:*

1. *If  $A \subseteq B$  then  $E - B \subseteq E - A$ .*

2. *If  $A \subseteq E$  and  $E - B \subseteq E - A$  then  $A \subseteq B$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $A$ ,  $B$  and  $E$  be sets.

$\langle 1 \rangle 2$ . If  $A \subseteq B$  then  $E - B \subseteq E - A$ .

PROOF: If  $A \subseteq B$ ,  $x \in E$  and  $x \notin B$ , then we have  $x \in E$  and  $x \notin A$ .

$\langle 1 \rangle 3$ . If  $A \subseteq E$  and  $E - B \subseteq E - A$  then  $A \subseteq B$ .

$\langle 2 \rangle 1$ . ASSUME:  $A \subseteq E$

$\langle 2 \rangle 2$ . ASSUME:  $E - B \subseteq E - A$

$\langle 2 \rangle 3$ . LET:  $x \in A$

$\langle 2 \rangle 4$ .  $x \in E$

$\langle 2 \rangle 5$ .  $x \notin E - A$

$\langle 2 \rangle 6$ .  $x \notin E - B$

$\langle 2 \rangle 7$ .  $x \in B$

$\square$

**Example 8.8.** We cannot remove the hypothesis  $A \subseteq E$  in item 2 above. Let  $E = \emptyset$ ,  $A = \{\emptyset\}$  and  $B = \emptyset$ . Then  $E - B = E - A = \emptyset$  but  $A \not\subseteq B$ .

**Proposition 8.9** (De Morgan's Law). *For any sets  $A$ ,  $B$  and  $E$ , we have  $E - (A \cup B) = (E - A) \cap (E - B)$ .*

PROOF:  $(x \in E \wedge \neg(x \in A \vee x \in B)) \Leftrightarrow (x \in E \wedge x \notin A \wedge x \in E \wedge x \notin B)$ .  $\square$

**Proposition 8.10** (De Morgan's Law). *For any sets  $A$ ,  $B$  and  $E$ , we have  $E - (A \cap B) = (E - A) \cup (E - B)$ .*

PROOF:  $(x \in E \vee \neg(x \in A \wedge x \in B)) \Leftrightarrow (x \in E \wedge x \notin A) \vee (x \in E \wedge x \notin B)$ .  $\square$

**Proposition 8.11.** *For any sets  $A$ ,  $B$  and  $E$ , if  $A \subseteq E$  then*

$$A - B = A \cap (E - B) .$$

PROOF: If  $A \subseteq E$  then we have  $(x \in A \wedge x \notin B) \Leftrightarrow (x \in A \wedge x \in E \wedge x \notin B)$ .  $\square$

**Proposition 8.12.** *For any sets  $A$  and  $B$ , we have  $A \subseteq B$  if and only if  $A - B = \emptyset$ .*

PROOF: Both are equivalent to the statement that there is no  $x$  such that  $x \in A$  and  $x \notin B$ .  $\square$

**Proposition 8.13.** *For any sets  $A$  and  $B$ , we have*

$$A - (A - B) = A \cap B .$$

PROOF:  $(x \in A \wedge \neg(x \in A \wedge x \notin B)) \Leftrightarrow x \in A \wedge x \in B$ .  $\square$

**Proposition 8.14.** *For any sets  $A$ ,  $B$  and  $C$ , we have*

$$A \cap (B - C) = (A \cap B) - (A \cap C) .$$

PROOF:  $(x \in A \wedge x \in B \wedge x \notin C) \Leftrightarrow (x \in A \wedge x \in B \wedge \neg(x \in A \wedge x \in C))$ .  $\square$

**Proposition 8.15.** *For any sets  $A$ ,  $B$ ,  $C$  and  $E$ , if  $(A \cap B) - C \subseteq E$  then we have*

$$A \cap B \subseteq (A \cap C) \cup (B \cap (E - C)) .$$

PROOF:

$\langle 1 \rangle 1$ . LET:  $x \in A \cap B$

PROVE:  $x \in (A \cap C) \cup (B \cap (E - C))$

$\langle 1 \rangle 2$ . CASE:  $x \in C$

PROOF: Then  $x \in A \cap C$ .

$\langle 1 \rangle 3$ . CASE:  $x \notin C$

PROOF: Then  $x \in E$  and so  $x \in B \cap (E - C)$ .

$\square$

**Proposition 8.16.** *For any sets  $A$ ,  $B$ ,  $C$  and  $E$ , we have*

$$(A \cup C) \cap (B \cup (E - C)) \subseteq A \cup B .$$

PROOF: The statement  $(x \in A \vee x \in C) \wedge (x \in B \vee (x \in E \wedge x \notin C))$  implies  $x \in A \vee x \in B$ .  $\square$

**Proposition 8.17** (De Morgan's Law). *Let  $E$  be a set and  $\mathcal{C}$  a nonempty set. Then*

$$E - \bigcup_{X \in \mathcal{C}} X = \bigcap_{X \in \mathcal{C}} (E - X) .$$

PROOF: Easy.  $\square$

**Proposition 8.18** (De Morgan's Law). *Let  $E$  be a set and  $\mathcal{C}$  a nonempty set. Then*

$$E - \bigcap_{X \in \mathcal{C}} X = \bigcup_{X \in \mathcal{C}} (E - X) .$$

PROOF: Easy.  $\square$

## Chapter 9

# Symmetric Difference

**Definition 9.1** (Symmetric Difference). For any sets  $A$  and  $B$ , the *symmetric difference*  $A + B$  is defined to be

$$A + B := (A - B) \cup (B - A) .$$

**Proposition 9.2.** *For any sets  $A$  and  $B$ , we have*

$$A + B = B + A$$

PROOF: From the commutativity of union.  $\square$

**Proposition 9.3.** *For any sets  $A$ ,  $B$  and  $C$ , we have*

$$A + (B + C) = (A + B) + C .$$

PROOF: Each is the set of all  $x$  that belong to either exactly one or all three of  $A$ ,  $B$  and  $C$ .  $\square$

**Proposition 9.4.** *For any set  $A$ , we have*

$$A + \emptyset = A .$$

PROOF:

$$\begin{aligned} A + \emptyset &= (A - \emptyset) \cup (\emptyset - A) \\ &= A \cup \emptyset \\ &= A \end{aligned}$$

$\square$

**Proposition 9.5.** *For any set  $A$  we have*

$$A + A = \emptyset .$$

PROOF:

$$\begin{aligned} A + A &= (A - A) \cup (A - A) \\ &= \emptyset \cup \emptyset \\ &= \emptyset \end{aligned}$$

$\square$

# Chapter 10

## Power Sets

**Definition 10.1** (Power Set). For any set  $A$ , the *power set* of  $A$ ,  $\mathcal{P}A$ , is the set whose elements are exactly the subsets of  $A$ .

PROOF: This exists by the Power Set Axiom and Axiom of Comprehension, and is unique by the Axiom of Extensionality.  $\square$

**Proposition 10.2.**

$$\mathcal{P}\emptyset = \{\emptyset\}$$

PROOF: The only subset of  $\emptyset$  is  $\emptyset$ .  $\square$

**Proposition 10.3.** For any set  $a$ , we have

$$\mathcal{P}\{a\} = \{\emptyset, \{a\}\} .$$

PROOF: The only subsets of  $\{a\}$  are  $\emptyset$  and  $\{a\}$ .  $\square$

**Proposition 10.4.** For any sets  $a$  and  $b$ , we have

$$\mathcal{P}\{a, b\} = \{\emptyset, \{a\}, \{b\}, \{a, b\}\} .$$

PROOF: The only subsets of  $\{a, b\}$  are  $\emptyset$ ,  $\{a\}$ ,  $\{b\}$  and  $\{a, b\}$ .  $\square$

**Proposition 10.5.** For any nonempty set  $\mathcal{C}$  we have

$$\bigcap_{X \in \mathcal{C}} \mathcal{P}X = \mathcal{P}\left(\bigcap \mathcal{C}\right) .$$

PROOF:

$$x \in \bigcup_{X \in \mathcal{C}} \mathcal{P}X \Leftrightarrow \forall X \in \mathcal{C}. x \subseteq X$$

$$\Leftrightarrow \forall X \in \mathcal{C}. \forall y \in x. y \in X$$

$$\Leftrightarrow \forall y \in x. \forall X \in \mathcal{C}. y \in X$$

$$\Leftrightarrow x \subseteq \bigcap \mathcal{C}$$

$\square$

**Proposition 10.6.** *For any set  $\mathcal{C}$  we have*

$$\bigcup_{X \in \mathcal{C}} \mathcal{P}X \subseteq \mathcal{P} \bigcup \mathcal{C} .$$

PROOF: If there exists  $X \in \mathcal{C}$  such that  $x \subseteq X$  then  $x \subseteq \bigcup \mathcal{C}$ .  $\square$

**Proposition 10.7.** *For any set  $E$ , we have*

$$\bigcap \mathcal{P}E = \emptyset .$$

PROOF: Since  $\emptyset \in \mathcal{P}E$ .  $\square$

**Proposition 10.8.** *For any sets  $E$  and  $F$ , if  $E \subseteq F$  then  $\mathcal{P}E \subseteq \mathcal{P}F$ .*

PROOF: If  $E \subseteq F$  and  $X \subseteq E$  then  $X \subseteq F$ .  $\square$



# Chapter 11

## Ordered Pairs

**Definition 11.1** (Ordered Pair). For any sets  $a$  and  $b$ , the *ordered pair*  $(a, b)$  is defined by

$$(a, b) := \{\{a\}, \{a, b\}\} .$$

**Proposition 11.2.** For any sets  $a, b, x$  and  $y$ , if  $(a, b) = (x, y)$  then  $a = x$  and  $b = y$ .

PROOF:

$\langle 1 \rangle 1$ . LET:  $a, b, x$  and  $y$  be sets.

$\langle 1 \rangle 2$ . ASSUME:  $(a, b) = (x, y)$

$\langle 1 \rangle 3$ .  $a = x$

PROOF:  $\{a\} = \bigcap(a, b) = \bigcap(x, y) = \{x\}$ .

$\langle 1 \rangle 4$ .  $\{a, b\} = \{x, y\}$

$\langle 1 \rangle 5$ . CASE:  $a = b$

$\langle 2 \rangle 1$ .  $x = y$

PROOF: Since  $\{x, y\} = \{a, b\}$  is a singleton.

$\langle 2 \rangle 2$ .  $b = y$

PROOF:  $b = a = x = y$

$\langle 1 \rangle 6$ . CASE:  $a \neq b$

$\langle 2 \rangle 1$ .  $x \neq y$

PROOF: Since  $\{x, y\} = \{a, b\}$  is not a singleton.

$\langle 2 \rangle 2$ .  $b = y$

PROOF:  $\{b\} = \{a, b\} - \{a\} = \{x, y\} - \{x\} = \{y\}$ .

□

**Definition 11.3** (Cartesian Product). For any sets  $A$  and  $B$ , the *Cartesian product*  $A \times B$  is

$$A \times B := \{p \in \mathcal{PP}(A \cup B) : \exists a \in A. \exists b \in B. p = (a, b)\} .$$

**Proposition 11.4.** For any sets  $A, B$  and  $X$ , we have

$$(A - B) \times X = (A \times X) - (B \times X) .$$

PROOF: Easy.  $\square$

**Proposition 11.5.** *For any sets  $A$  and  $B$ , we have  $A \times B = \emptyset$  if and only if  $A = \emptyset$  or  $B = \emptyset$ .*

PROOF: Easy.  $\square$

**Proposition 11.6.** *For any sets  $A$ ,  $B$ ,  $X$  and  $Y$ , if  $A \subseteq X$  and  $B \subseteq Y$  then  $A \times B \subseteq X \times Y$ . The converse holds assuming  $A \neq \emptyset$  and  $B \neq \emptyset$ .*

PROOF: Easy.  $\square$

## Chapter 12

# Relations

**Definition 12.1** (Relation). A *relation* is a set of ordered pairs.

If  $R$  is a relation, we write  $xRy$  for  $(x, y) \in R$ .

Given sets  $X$  and  $Y$ , a relation *between  $X$  and  $Y$*  is a subset of  $X \times Y$ .

Given a set  $X$ , a relation *on  $X$*  is a relation between  $X$  and  $X$ .

**Definition 12.2** (Domain). The *domain* of a relation  $R$  is the set

$$\text{dom } R := \{x \in \bigcup \bigcup R : \exists y. (x, y) \in R\} \ .$$

**Definition 12.3** (Range). The *range* of a relation  $R$  is the set

$$\text{ran } R := \{y \in \bigcup \bigcup R : \exists x. (x, y) \in R\} \ .$$

**Definition 12.4** (Reflexive). Let  $R$  be a relation on  $X$ . Then  $R$  is *reflexive* iff, for all  $x \in X$ , we have  $xRx$ .

**Definition 12.5** (Symmetric). Let  $R$  be a relation on  $X$ . Then  $R$  is *symmetric* iff, whenever  $xRy$ , then  $yRx$ .

**Definition 12.6** (Antisymmetric). A relation  $R$  is *antisymmetric* iff, whenever  $xRy$  and  $yRx$ , then  $x = y$ .

**Definition 12.7** (Transitive). Let  $R$  be a relation on  $X$ . Then  $R$  is *transitive* iff, whenever  $xRy$  and  $yRz$ , then  $xRz$ .

**Definition 12.8** (Equivalence Relation). Let  $R$  be a relation on  $X$ . Then  $R$  is an *equivalence relation* iff it is reflexive, symmetric and transitive.

**Definition 12.9** (Partition). Let  $X$  be a set. A *partition* of  $X$  is a pairwise disjoint set of nonempty subsets of  $X$  whose union is  $X$ .

**Definition 12.10** (Equivalence Class). Let  $R$  be an equivalence relation on  $X$ . Let  $x \in X$ . The *equivalence class* of  $x$  with respect to  $R$  is

$$x/R := \{y \in X : xRy\} \ .$$

We write  $X/R$  for the set of all equivalence classes with respect to  $R$ .

**Definition 12.11** (Induced). Let  $P$  be a partition of  $X$ . The relation *induced* by  $P$  is  $X/P$  where  $x(X/P)y$  iff there exists  $X \in P$  such that  $x \in X$  and  $y \in X$ .

**Theorem 12.12.** Let  $R$  be an equivalence relation on  $X$ . Then  $X/R$  is a partition of  $X$  that induces the relation  $R$ .

PROOF: Easy.  $\square$

**Theorem 12.13.** Let  $P$  be a partition of  $X$ . Then  $X/P$  is an equivalence relation on  $X$ , and  $P = X/(X/P)$ .

PROOF: Easy.  $\square$

**Definition 12.14** (Composition). Let  $R$  be a relation between  $X$  and  $Y$ , and  $S$  a relation between  $Y$  and  $Z$ . The *composite* or *relative product*  $S \circ R = SR$  is the relation between  $X$  and  $Z$  defined by

$$x(S \circ R)z \Leftrightarrow \exists y \in Y (xRy \wedge ySz) .$$

**Proposition 12.15.** Let  $R$  be a relation between  $X$  and  $Y$ ,  $S$  a relation between  $Y$  and  $Z$ , and  $T$  a relation between  $Z$  and  $W$ . Then

$$T(SR) = (TS)R .$$

PROOF: Easy.  $\square$

**Example 12.16.** Composition of relations is not commutative in general. Let  $X = \{a, b\}$  where  $a \neq b$ . Let  $R = \{(a, a), (b, a)\}$  and  $S = \{(a, b), (b, b)\}$ . Then  $SR = S$  but  $RS = R \neq S$ .

**Proposition 12.17.** A relation  $R$  is transitive if and only if  $RR \subseteq R$ .

PROOF: Easy.  $\square$

**Definition 12.18** (Inverse). Let  $R$  be a relation between  $X$  and  $Y$ . The *inverse* or *converse*  $R^{-1}$  is the relation between  $Y$  and  $X$  defined by

$$yR^{-1}x \Leftrightarrow xRy .$$

**Proposition 12.19.** For any relation  $R$ , we have

$$\text{dom } R^{-1} = \text{ran } R .$$

PROOF: Easy.  $\square$

**Proposition 12.20.** For any relation  $R$ , we have

$$\text{ran } R^{-1} = \text{dom } R .$$

PROOF: Easy.  $\square$

**Proposition 12.21.** *Let  $R$  be a relation between  $X$  and  $Y$ , and  $S$  a relation between  $Y$  and  $Z$ . Then*

$$(SR)^{-1} = R^{-1}S^{-1} .$$

PROOF: Easy.  $\square$

**Proposition 12.22.** *A relation  $R$  is symmetric if and only if  $R \subseteq R^{-1}$ .*

PROOF: Easy.  $\square$

**Definition 12.23** (Identity Relation). For any set  $X$ , the *identity relation*  $I_X$  on  $X$  is

$$I_X = \{(x, x) : x \in X\} .$$

**Proposition 12.24.** *Let  $R$  be a relation between  $X$  and  $Y$ . Then*

$$I_Y R = R I_X = R .$$

PROOF: Easy.  $\square$

**Proposition 12.25.** *A relation  $R$  on a set  $X$  is reflexive if and only if  $I_X \subseteq R$ .*

PROOF: Easy.  $\square$

# Chapter 13

## Functions

**Definition 13.1** (Function). Let  $X$  and  $Y$  be sets. A *function*, *map*, *mapping*, *transformation* or *operator*  $f$  from  $X$  to  $Y$ ,  $f : X \rightarrow Y$ , is a relation  $f$  between  $X$  and  $Y$  such that, for all  $x \in X$ , there exists a unique  $f(x) \in Y$ , called the *value* of  $f$  at the *argument*  $x$ , such that  $(x, f(x)) \in f$ .

**Definition 13.2** (Onto). Let  $f : X \rightarrow Y$ . We say  $f$  maps  $X$  *onto*  $Y$  iff  $\text{ran } f = Y$ .

**Definition 13.3** (Image). Let  $f : X \rightarrow Y$  and  $A \subseteq X$ . The *image* of  $A$  under  $f$  is

$$f(A) := \{f(x) : x \in A\} .$$

**Definition 13.4** (Inclusion Map). Let  $Y$  be a set and  $X \subseteq Y$ . Then the *inclusion map*  $i : X \hookrightarrow Y$  is the function defined by  $i(x) = x$  for all  $x \in X$ .

**Proposition 13.5.** For any set  $X$ , the identity relation  $I_X$  is a function  $X \rightarrow X$ .

PROOF: Easy.  $\square$

**Definition 13.6** (Restriction). Let  $f : Y \rightarrow Z$  and  $X \subseteq Y$ . The *restriction* of  $f$  to  $X$  is the function  $f \upharpoonright X : X \rightarrow Z$  defined by

$$(f \upharpoonright X)(x) = f(x) \quad (x \in X) .$$

Given sets  $X$ ,  $Y$  and  $Z$  with  $X \subseteq Y$ , if  $f : X \rightarrow Z$  and  $g : Y \rightarrow Z$ , we say  $g$  is an *extension* of  $f$  to  $Y$  iff  $f = g \upharpoonright X$ .

**Definition 13.7** (Projection). Given sets  $X$  and  $Y$ , the *projection* maps  $\pi_1 : X \times Y \rightarrow X$  and  $\pi_2 : X \times Y \rightarrow Y$  are defined by

$$\pi_1(x, y) = x, \quad \pi_2(x, y) = y \quad (x \in X, y \in Y) .$$

**Definition 13.8** (Canonical Map). Let  $X$  be a set and  $R$  an equivalence relation on  $X$ . The *canonical map*  $\pi : X \rightarrow X/R$  is the map defined by  $\pi(x) = x/R$ .

**Definition 13.9** (One-to-One). A function  $f : X \rightarrow Y$  is *one-to-one*, or a *one-to-one correspondence*, iff, for all  $x, y \in X$ , if  $f(x) = f(y)$  then  $x = y$ .

**Proposition 13.10.** Let  $f : X \rightarrow Y$ . Then the following are equivalent:

1.  $f$  is one-to-one.
2. For all  $A, B \subseteq X$ , we have  $f(A \cap B) = f(A) \cap f(B)$ .
3. For all  $A \subseteq X$ , we have  $f(X - A) \subseteq Y - f(A)$ .

PROOF: Easy.  $\square$

**Proposition 13.11.** Let  $f : X \rightarrow Y$ . Then  $f$  maps  $X$  onto  $Y$  if and only if, for all  $A \subseteq X$ , we have  $Y - f(A) \subseteq f(X - A)$ .

PROOF: Easy.  $\square$

# Chapter 14

## Families

**Definition 14.1** (Family). Let  $I$  and  $X$  be sets. A *family* of elements of  $X$  indexed by  $I$  is a function  $a : I \rightarrow X$ . We write  $a_i$  for  $a(i)$ , and  $\{a_i\}_{i \in I}$  for  $a$ .

**Proposition 14.2** (Generalized Associative Law for Unions). Let  $\{I_j\}_{j \in J}$  be a family of sets. Let  $K = \bigcup_{j \in J} I_j$ . Let  $\{A_k\}_{k \in K}$  be a family of sets indexed by  $K$ . Then

$$\bigcup_{k \in K} A_k = \bigcup_{j \in J} \bigcup_{i \in I_j} A_i .$$

PROOF: Easy.  $\square$

**Proposition 14.3** (Generalized Commutative Law for Unions). Let  $\{I_j\}_{j \in J}$  be a family of sets. Let  $f : J \rightarrow J$  be a one-to-one correspondence from  $J$  onto  $J$ . Then

$$\bigcup_{j \in J} I_j = \bigcup_{j \in J} I_{f(j)} .$$

PROOF: Easy.  $\square$

**Proposition 14.4** (Generalized Associative Law for Intersections). Let  $\{I_j\}_{j \in J}$  be a nonempty family of nonempty sets. Let  $K = \bigcup_{j \in J} I_j$ . Let  $\{A_k\}_{k \in K}$  be a family of sets indexed by  $K$ . Then

$$\bigcap_{k \in K} A_k = \bigcap_{j \in J} \bigcap_{i \in I_j} A_i .$$

PROOF: Easy.  $\square$

**Proposition 14.5** (Generalized Commutative Law for Intersections). Let  $\{I_j\}_{j \in J}$  be a nonempty family of sets. Let  $f : J \rightarrow J$  be a one-to-one correspondence from  $J$  onto  $J$ . Then

$$\bigcap_{j \in J} I_j = \bigcap_{j \in J} I_{f(j)} .$$



PROOF: Easy.  $\square$

**Proposition 14.6.** *Let  $B$  be a set and  $\{A_i\}_{i \in I}$  a family of sets. Then*

$$B \cap \bigcup_{i \in I} A_i = \bigcup_{i \in I} (B \cap A_i)$$

PROOF: Easy.  $\square$

**Proposition 14.7.** *Let  $B$  be a set and  $\{A_i\}_{i \in I}$  a nonempty family of sets. Then*

$$B \cup \bigcap_{i \in I} A_i = \bigcap_{i \in I} (B \cup A_i)$$

PROOF: Easy.  $\square$

**Definition 14.8** (Cartesian Product of a Family of Sets). Let  $\{A_i\}_{i \in I}$  be a family of sets. The *Cartesian product*  $\times_{i \in I} A_i$  is the set of all families  $\{a_i\}_{i \in I}$  such that  $\forall i \in I. a_i \in A_i$ .

We write  $A^I$  for  $\times_{i \in I} A$ .

**Definition 14.9** (Projection). Let  $\{A_i\}_{i \in I}$  be a family of sets and  $i \in I$ . The *projection* function  $\pi_i : \times_{i \in I} A_i \rightarrow A_i$  is defined by  $\pi_i(a) = a_i$ .

**Proposition 14.10.** *Let  $\{A_i\}_{i \in I}$  and  $\{B_j\}_{j \in J}$  be families of sets. Then*

$$\left( \bigcup_{i \in I} A_i \right) \times \left( \bigcup_{j \in J} B_j \right) = \bigcup_{i \in I} \bigcup_{j \in J} (A_i \times B_j) .$$

PROOF: Easy.  $\square$

**Proposition 14.11.** *Let  $\{A_i\}_{i \in I}$  and  $\{B_j\}_{j \in J}$  be nonempty families of sets. Then*

$$\left( \bigcap_{i \in I} A_i \right) \times \left( \bigcap_{j \in J} B_j \right) = \bigcap_{i \in I} \bigcap_{j \in J} (A_i \times B_j) .$$

PROOF: Easy.  $\square$

**Proposition 14.12.** *Let  $f : X \rightarrow Y$ . Let  $\{A_i\}_{i \in I}$  be a family of subsets of  $X$ . Then*

$$f \left( \bigcup_{i \in I} A_i \right) = \bigcup_{i \in I} f(A_i) .$$

PROOF: Easy.  $\square$

**Example 14.13.** It is not true in general that, if  $f : X \rightarrow Y$  and  $\{A_i\}_{i \in I}$  is a nonempty family of subsets of  $X$ , then  $f(\bigcap_{i \in I} A_i) = \bigcap_{i \in I} f(A_i)$ .

Take  $X = \{a, b\}$  and  $Y = \{c\}$  where  $a \neq b$ . Take  $I = \{i, j\}$  with  $i \neq j$ . Let  $A_i = \{a\}$  and  $A_j = \{b\}$ . Let  $f$  be the unique function  $X \rightarrow Y$ . Then  $f(\bigcap_{i \in I} A_i) = f(\emptyset) = \emptyset$  but  $\bigcap_{i \in I} f(A_i) = \{c\}$ .

## Chapter 15

# Inverses and Composites

**Definition 15.1** (Inverse). Given a function  $f : X \rightarrow Y$ , the *inverse* of  $f$  is the function  $f^{-1} : \mathcal{P}Y \rightarrow \mathcal{P}X$  defined by

$$f^{-1}(B) = \{x \in X : f(x) \in B\} \ .$$

We call  $f^{-1}(B)$  the *inverse image* of  $B$  under  $f$ .

**Proposition 15.2.** *Let  $f : X \rightarrow Y$ . Then  $f$  maps  $X$  onto  $Y$  if and only if the inverse image of any nonempty subset of  $Y$  is nonempty.*

PROOF: Easy.  $\square$

**Proposition 15.3.** *Let  $f : X \rightarrow Y$ . Then  $f$  is one-to-one if and only if the inverse image of any singleton subset of  $Y$  is a singleton.*

PROOF: Easy.  $\square$

**Proposition 15.4.** *Let  $f : X \rightarrow Y$ . Let  $B \subseteq Y$ . Then*

$$f(f^{-1}(B)) \subseteq B \ .$$

PROOF: Easy.  $\square$

**Proposition 15.5.** *Let  $f : X \rightarrow Y$ . Let  $A \subseteq X$ . Then*

$$A \subseteq f^{-1}(f(A)) \ .$$

*Equality holds if  $f$  is one-to-one.*

PROOF: Easy.  $\square$

**Proposition 15.6.** *Let  $f : X \rightarrow Y$ . Let  $\{B_i\}_{i \in I}$  be a family of subsets of  $Y$ . Then*

$$f^{-1}\left(\bigcup_{i \in I} B_i\right) = \bigcup_{i \in I} f^{-1}(B_i) \ .$$

PROOF: Easy.  $\square$

**Proposition 15.7.** *Let  $f : X \rightarrow Y$ . Let  $\{B_i\}_{i \in I}$  be a nonempty family of subsets of  $Y$ . Then*

$$f^{-1} \left( \bigcap_{i \in I} B_i \right) = \bigcap_{i \in I} f^{-1}(B_i) .$$

PROOF: Easy.  $\square$

**Proposition 15.8.** *Let  $f : X \rightarrow Y$  and  $B \subseteq Y$ . Then  $f^{-1}(Y - B) = X - f^{-1}(B)$ .*

PROOF: Easy.  $\square$

**Proposition 15.9.** *Let  $f : X \rightarrow Y$  be one-to-one. Then the inverse of  $f$  as a relation,  $f^{-1}$ , is a function  $f^{-1} : \text{ran } f \rightarrow X$ , and for all  $y \in \text{ran } f$ , we have  $f^{-1}(y)$  is the unique  $x$  such that  $f(x) = y$ .*

PROOF: Easy.  $\square$

**Proposition 15.10.** *Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ . Then  $gf : X \rightarrow Z$  and, for all  $x \in X$ , we have*

$$(g \circ f)(x) = g(f(x)) .$$

PROOF: Easy.  $\square$

**Example 15.11.** Example 12.15 shows that function composition is not commutative in general.

**Proposition 15.12.** *Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ . Then*

$$(gf)^{-1} = f^{-1}g^{-1} : \mathcal{P}Z \rightarrow \mathcal{P}X .$$

PROOF: Easy.  $\square$

**Proposition 15.13.** *Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$ . If  $gf = I_X$  then  $f$  is one-to-one and  $g$  maps  $Y$  onto  $X$ .*

PROOF: Easy.  $\square$

# Chapter 16

## Numbers

**Definition 16.1** (Successor). The *successor* of a set  $x$ ,  $x^+$ , is defined by

$$x^+ := x \cup \{x\} .$$

**Definition 16.2.** We define

$$0 = \emptyset$$

$$1 = 0^+$$

$$2 = 1^+$$

etc.

**Definition 16.3** (Characteristic Function). Let  $X$  be a set and  $A \subseteq X$ . The *characteristic function* of  $A$  is the function  $\chi_A : X \rightarrow 2$  defined by

$$\chi_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

**Theorem 16.4.** Let  $X$  be a set. The function  $\chi : \mathcal{P}X \rightarrow 2^X$  that maps a subset  $A$  of  $X$  to  $\chi_A$  is a one-to-one correspondence.

PROOF: Easy.  $\square$

**Definition 16.5.** The set  $\omega$  of *natural numbers* is the set such that:

- $0 \in \omega$
- For all  $n \in \omega$  we have  $n^+ \in \omega$
- For any set  $X$ , if  $0 \in X$  and  $\forall n \in X. n^+ \in X$  then  $\omega \subseteq X$

PROOF: To show this exists, pick a set  $A$  such that  $0 \in A$  and  $\forall n \in A. n^+ \in A$  (by the Axiom of Infinity), and let  $\omega = \bigcap \{X \in \mathcal{P}A : 0 \in X \wedge \forall n \in X. n^+ \in X\}$ .  
 $\square$

**Definition 16.6** (Sequence). A *finite sequence* is a family whose index set is a natural number. An *infinite sequence* is a family whose index set is  $\omega$ .

Given a finite sequence of sets  $\{A_i\}_{i \in n^+}$ , we write  $\bigcup_{i=0}^n A_i$  for  $\bigcup_{i \in n^+} A_i$ . Given an infinite sequence of sets  $\{A_i\}_{i \in \omega}$ , we write  $\bigcup_{i=0}^{\infty} A_i$  for  $\bigcup_{i \in \omega} A_i$ .

We make similar definitions for  $\bigcap$  and  $\times$ .

**Proposition 16.7.** *For any natural numbers  $m$  and  $n$ , if  $m \in n$  then  $m^+ \in n^+$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $P(n)$  be the property  $\forall m \in n. m^+ \in n^+$

$\langle 1 \rangle 2$ .  $P(0)$

PROOF: Vacuous.

$\langle 1 \rangle 3$ . For any natural number  $n$ , if  $P(n)$  then  $P(n^+)$ .

$\langle 2 \rangle 1$ . LET:  $n$  be a natural number.

$\langle 2 \rangle 2$ . ASSUME:  $P(n)$

$\langle 2 \rangle 3$ . LET:  $m \in n^+$

$\langle 2 \rangle 4$ .  $m \in n$  or  $m = n$

$\langle 2 \rangle 5$ .  $m^+ \in n^+$  or  $m^+ = n^+$

PROOF:  $\langle 2 \rangle 2$

$\langle 2 \rangle 6$ . CASE:  $m^+ \in n^{++}$

□

## Chapter 17

# The Peano Axioms

**Theorem 17.1** (Principle of Mathematical Induction). *For any subset  $S$  of  $\omega$ , if  $0 \in S$  and  $\forall n \in S. n^+ \in S$ , then  $S = \omega$ .*

PROOF: From the definition of  $\omega$ .  $\square$

**Proposition 17.2.**

$$\forall n \in \omega. \forall x \in n. n \not\subseteq x$$

PROOF:

$\langle 1 \rangle 1.$   $\forall x \in 0. 0 \not\subseteq x$

PROOF: Vacuous.

$\langle 1 \rangle 2.$  For any natural number  $n$ , if  $\forall x \in n. n \not\subseteq x$  then  $\forall x \in n^+. n^+ \not\subseteq x$ .

$\langle 2 \rangle 1.$  LET:  $n$  be a natural number.

$\langle 2 \rangle 2.$  ASSUME:  $\forall x \in n. n \not\subseteq x$

$\langle 2 \rangle 3.$  LET:  $x \in n^+$

$\langle 2 \rangle 4.$  ASSUME: for a contradiction  $n^+ \subseteq x$

$\langle 2 \rangle 5.$   $x \in n$  or  $x = n$

$\langle 2 \rangle 6.$  CASE:  $x \in n$

PROOF: Then we have  $n \subseteq n^+ \subseteq x$  contradicting  $\langle 2 \rangle 2.$

$\langle 2 \rangle 7.$  CASE:  $x = n$

PROOF: Then we have  $n \in n^+ \subseteq x = n$  and  $n \subseteq n$  contradicting  $\langle 2 \rangle 2.$

$\square$

**Corollary 17.2.1.** *For any natural number  $n$  we have  $n \notin n$ .*

**Corollary 17.2.2.** *For any natural number  $n$  we have  $n \neq n^+$ .*

**Definition 17.3** (Transitive Set). A set  $E$  is a *transitive* set iff, whenever  $x \in y \in E$ , then  $x \in E$ .

**Proposition 17.4.** *Every natural number is a transitive set.*

PROOF:

$\langle 1 \rangle 1.$   $0$  is a transitive set.

PROOF: Vacuously, if  $x \in y \in 0$  then  $x \in 0$ .

$\langle 1 \rangle 2$ . For any natural number  $n$ , if  $n$  is a transitive set, then  $n^+$  is a transitive set.

$\langle 2 \rangle 1$ . LET:  $n$  be a natural number.

$\langle 2 \rangle 2$ . ASSUME:  $n$  is a transitive set.

$\langle 2 \rangle 3$ . LET:  $x \in y \in n^+$

$\langle 2 \rangle 4$ .  $y \in n$  or  $y = n$

$\langle 2 \rangle 5$ . CASE:  $y \in n$

$\langle 3 \rangle 1$ .  $x \in n$

PROOF:  $\langle 2 \rangle 2$ ,  $\langle 2 \rangle 3$ ,  $\langle 2 \rangle 5$ .

$\langle 3 \rangle 2$ .  $x \in n^+$

$\langle 2 \rangle 6$ . CASE:  $y = n$

$\langle 3 \rangle 1$ .  $x \in n$

PROOF:  $\langle 2 \rangle 3$ ,  $\langle 2 \rangle 6$

$\langle 3 \rangle 2$ .  $x \in n^+$

□

**Proposition 17.5.** *For any natural numbers  $m$  and  $n$ , if  $m^+ = n^+$  then  $m = n$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $m$  and  $n$  be natural numbers.

$\langle 1 \rangle 2$ . ASSUME:  $m^+ = n^+$

$\langle 1 \rangle 3$ .  $m \in m^+ = n^+$

$\langle 1 \rangle 4$ .  $m \in n$  or  $m = n$

$\langle 1 \rangle 5$ .  $n \in n^+ = m^+$

$\langle 1 \rangle 6$ .  $n \in m$  or  $n = m$

$\langle 1 \rangle 7$ . We cannot have  $m \in n$  and  $n \in m$

$\langle 2 \rangle 1$ . ASSUME: for a contradiction  $m \in n$  and  $n \in m$

$\langle 2 \rangle 2$ .  $m \in m$

PROOF: Since  $m$  is a transitive set (Proposition 17.4).

$\langle 2 \rangle 3$ . Q.E.D.

PROOF: This contradicts Proposition 17.2.

$\langle 1 \rangle 8$ .  $m = n$

□

**Theorem 17.6** (Recursion Theorem). *Let  $X$  be a set. Let  $a \in X$ . Let  $f : X \rightarrow X$ . There exists a function  $u : \omega \rightarrow X$  such that  $u(0) = a$  and, for all  $n \in \omega$ , we have  $u(n^+) = f(u(n))$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $\mathcal{C} = \{A \in \mathcal{P}(\omega \times X) : (0, a) \in A \wedge \forall n \in \omega. \forall x \in X. (n, x) \in A \Rightarrow (n^+, f(x)) \in A\}$

$\langle 1 \rangle 2$ .  $\mathcal{C} \neq \emptyset$

PROOF:  $\omega \times X \in \mathcal{C}$

$\langle 1 \rangle 3$ . LET:  $u = \bigcap \mathcal{C}$

$\langle 1 \rangle 4$ .  $u \in \mathcal{C}$

$\langle 1 \rangle 5$ .  $u$  is a function.

$\langle 2 \rangle 1$ . LET:  $P(n)$  be the property:  $\forall x, y \in X. (n, x) \in u \wedge (n, y) \in u \Rightarrow x = y$   
 $\langle 2 \rangle 2$ .  $P(0)$   
 $\langle 3 \rangle 1$ .  $\forall x \in X. (0, x) \in u \Rightarrow x = a$   
PROOF: If  $(0, x) \in u$  and  $x \neq a$  then  $u - \{(0, x)\} \in \mathcal{C}$  and so  $u - \{(0, x)\} \subseteq u$ , which is impossible.  
 $\langle 2 \rangle 3$ . For every natural number  $n$ , if  $P(n)$  then  $P(n^+)$ .  
 $\langle 3 \rangle 1$ . LET:  $n$  be a natural number.  
 $\langle 3 \rangle 2$ . ASSUME:  $P(n)$   
 $\langle 3 \rangle 3$ . LET:  $x, y \in X$   
 $\langle 3 \rangle 4$ . ASSUME:  $(n^+, x), (n^+, y) \in u$   
 $\langle 3 \rangle 5$ . PICK  $x', y' \in X$  such that  $(n, x') \in u, (n, y') \in u$  and  $f(x') = x$  and  $f(y') = y$   
PROOF: If no such  $x'$  exists then  $u - \{(n^+, x)\} \in \mathcal{C}$  and so  $u - \{(n^+, x)\} \subseteq u$  which is impossible. Similarly for  $y'$ .  
 $\langle 3 \rangle 6$ .  $x' = y'$   
PROOF:  $\langle 3 \rangle 2$   
 $\langle 3 \rangle 7$ .  $x = y$

□

**Proposition 17.7.** *For any natural number  $n$ , either  $n = 0$  or there exists a natural number  $m$  such that  $n = m^+$ .*

PROOF: Easy induction on  $n$ . □

**Proposition 17.8.**  *$\omega$  is a transitive set.*

PROOF:

$\langle 1 \rangle 1$ . LET:  $P(n)$  be the property  $\forall x \in n. x \in \omega$   
 $\langle 1 \rangle 2$ .  $P(0)$   
PROOF: Vacuous.  
 $\langle 1 \rangle 3$ . For any natural number  $n$ , if  $P(n)$  then  $P(n^+)$ .  
 $\langle 2 \rangle 1$ . LET:  $n$  be a natural number.  
 $\langle 2 \rangle 2$ . ASSUME:  $P(n)$   
 $\langle 2 \rangle 3$ . LET:  $x \in n^+$   
 $\langle 2 \rangle 4$ .  $x \in n$  or  $x = n$   
 $\langle 2 \rangle 5$ . CASE:  $x \in n$   
PROOF: Then  $x \in \omega$  by  $\langle 2 \rangle 2$ .  
 $\langle 2 \rangle 6$ . CASE:  $x = n$   
PROOF: Then  $x \in \omega$  by  $\langle 2 \rangle 1$ .

□

**Proposition 17.9.** *For any natural number  $n$  and any nonempty subset  $E \subseteq n$ , there exists  $k \in E$  such that  $\forall m \in E. k = m \vee k \in m$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $P(n)$  be the property: for any nonempty subset  $E \subseteq n$ , there exists  $k \in E$  such that  $\forall m \in E. k = m \vee k \in m$   
 $\langle 1 \rangle 2$ .  $P(0)$



PROOF: Vacuous as there is no nonempty subset of 0.

$\langle 1 \rangle 3$ . For any natural number  $n$ , if  $P(n)$  then  $P(n^+)$ .

$\langle 2 \rangle 1$ . LET:  $n$  be a natural number.

$\langle 2 \rangle 2$ . ASSUME:  $P(n)$

$\langle 2 \rangle 3$ . LET:  $E$  be a nonempty subset of  $n^+$

$\langle 2 \rangle 4$ . CASE:  $E - \{n\} = \emptyset$   
PROOF: Then  $E = \{n\}$  so take  $k = n$ .

$\langle 2 \rangle 5$ . CASE:  $E - \{n\} \neq \emptyset$

$\langle 3 \rangle 1$ . PICK  $k \in E - \{n\}$  such that  $\forall m \in E - \{n\}. k = m \vee k \in m$   
PROOF: By  $\langle 2 \rangle 2$ .

$\langle 3 \rangle 2$ .  $\forall m \in E. k = m \vee k \in m$   
PROOF: Since  $k \in n$ .

□

## Chapter 18

# Arithmetic

**Definition 18.1** (Addition). Define *addition*  $+$  on  $\omega$  by recursion thus:

$$\begin{aligned}m + 0 &= m \\m + n^+ &= (m + n)^+\end{aligned}$$

**Proposition 18.2.** *For all  $m, n, p \in \omega$  we have*

$$m + (n + p) = (m + n) + p .$$

PROOF:

$\langle 1 \rangle 1$ . LET:  $P(p)$  be the property  $\forall m, n \in \omega. m + (n + p) = (m + n) + p$

$\langle 1 \rangle 2$ .  $P(0)$

PROOF:  $m + (n + 0) = m + n = (m + n) + 0$ .

$\langle 1 \rangle 3$ .  $\forall p \in \omega. P(p) \Rightarrow P(p^+)$

$\langle 2 \rangle 1$ . LET:  $p \in \omega$

$\langle 2 \rangle 2$ . ASSUME:  $P(p)$

$\langle 2 \rangle 3$ . LET:  $m, n \in \omega$

$\langle 2 \rangle 4$ .  $m + (n + p^+) = (m + n) + p^+$

PROOF:

$$\begin{aligned}m + (n + p^+) &= m + (n + p)^+ \\&= (m + (n + p))^+ \\&= ((m + n) + p)^+ \\&= (m + n) + p^+\end{aligned}$$

□

**Proposition 18.3.** *For all  $m, n \in \omega$ , we have*

$$m + n = n + m .$$

PROOF:

$\langle 1 \rangle 1$ . LET:  $P(m)$  be the property  $\forall n \in \omega. m + n = n + m$

$\langle 1 \rangle 2. P(0)$   
 $\langle 2 \rangle 1. \text{ LET: } Q(n) \text{ be the property } 0 + n = n + 0$   
 $\langle 2 \rangle 2. Q(0)$   
 PROOF: Trivial.  
 $\langle 2 \rangle 3. \forall n \in \omega. Q(n) \Rightarrow Q(n^+)$   
 $\langle 3 \rangle 1. \text{ LET: } n \in \omega$   
 $\langle 3 \rangle 2. \text{ ASSUME: } Q(n)$   
 $\langle 3 \rangle 3. 0 + n^+ = n^+ + 0$   
 PROOF:  

$$\begin{aligned}
 0 + n^+ &= (0 + n)^+ \\
 &= (n + 0)^+ & (\langle 3 \rangle 2) \\
 &= n^+ \\
 &= n^+ + 0
 \end{aligned}$$

$\langle 1 \rangle 3. \forall m \in \omega. P(m) \Rightarrow P(m^+)$   
 $\langle 2 \rangle 1. \text{ LET: } m \in \omega$   
 $\langle 2 \rangle 2. \text{ ASSUME: } P(m)$   
 $\langle 2 \rangle 3. \text{ LET: } Q(n) \text{ be the property } m^+ + n = n + m^+$   
 $\langle 2 \rangle 4. Q(0)$   
 PROOF:  $\langle 1 \rangle 2$   
 $\langle 2 \rangle 5. \forall n \in \omega. Q(n) \Rightarrow Q(n^+)$   
 $\langle 3 \rangle 1. \text{ LET: } n \in \omega$   
 $\langle 3 \rangle 2. \text{ ASSUME: } Q(n)$   
 $\langle 3 \rangle 3. Q(n^+)$   
 PROOF:  

$$\begin{aligned}
 m^+ + n^+ &= (m^+ + n)^+ \\
 &= (n + m^+)^+ & (\langle 3 \rangle 2) \\
 &= (n + m)^{++} \\
 &= (m + n)^{++} & (\langle 2 \rangle 2) \\
 &= (m + n^+)^+ \\
 &= (n^+ + m)^+ & (\langle 2 \rangle 2) \\
 &= n^+ + m^+
 \end{aligned}$$

□

**Definition 18.4** (Multiplication). Define *multiplication*  $\cdot$  on  $\omega$  by

$$\begin{aligned}
 m0 &= 0 \\
 mn^+ &= mn + m
 \end{aligned}$$

**Proposition 18.5.** For all  $m, n, p \in \omega$ , we have

$$m(n + p) = mn + mp .$$

PROOF:

$\langle 1 \rangle 1. \text{ LET: } P(p) \text{ be the statement } \forall m, n \in \omega. m(n + p) = mn + mp$

⟨1⟩2.  $P(0)$

PROOF:

$$\begin{aligned} m(n+0) &= mn \\ &= mn + 0 \\ &= mn + m0 \end{aligned}$$

⟨1⟩3.  $\forall p \in \omega. P(p) \Rightarrow P(p^+)$

⟨2⟩1. LET:  $p \in \omega$

⟨2⟩2. ASSUME:  $P(p)$

⟨2⟩3. LET:  $m, n \in \omega$

⟨2⟩4.  $m(n+p^+) = mn + mp^+$

PROOF:

$$\begin{aligned} m(n+p^+) &= m(n+p)^+ \\ &= m(n+p) + m \\ &= (mn+mp) + m && (\langle 2 \rangle 2) \\ &= mn + (mp+m) && (\text{Proposition 18.2}) \\ &= mn + mp^+ \end{aligned}$$

□

**Proposition 18.6.** *For all  $m, n, p \in \omega$  we have*

$$m(np) = (mn)p .$$

PROOF:

⟨1⟩1. LET:  $P(p)$  be the statement  $\forall m, n \in \omega. m(np) = (mn)p$

⟨1⟩2.  $P(0)$

PROOF:

$$\begin{aligned} m(n0) &= m0 \\ &= 0 \\ &= (mn)0 \end{aligned}$$

⟨1⟩3.  $\forall p \in \omega. P(p) \Rightarrow P(p^+)$

⟨2⟩1. LET:  $p \in \omega$

⟨2⟩2. ASSUME:  $P(p)$

⟨2⟩3. LET:  $m, n \in \omega$

⟨2⟩4.  $m(np^+) = (mn)p^+$

PROOF:

$$\begin{aligned} m(np^+) &= m(np+n) \\ &= m(np) + mn && (\text{Proposition 18.5}) \\ &= (mn)p + mn && (\langle 2 \rangle 2) \\ &= (mn)p^+ \end{aligned}$$

□

**Proposition 18.7.** *For all  $m, n \in \omega$ , we have*

$$mn = nm .$$

PROOF:

$\langle 1 \rangle 1$ . LET:  $P(m)$  be the statement  $\forall n \in \omega. mn = nm$

$\langle 1 \rangle 2$ .  $P(0)$

$\langle 2 \rangle 1$ . LET:  $Q(n)$  be the statement  $0n = n0$

$\langle 2 \rangle 2$ .  $Q(0)$

PROOF: Trivial.

$\langle 2 \rangle 3$ .  $\forall n \in \omega. Q(n) \Rightarrow Q(n^+)$

$\langle 3 \rangle 1$ . LET:  $n \in \omega$

$\langle 3 \rangle 2$ . ASSUME:  $Q(n)$

$\langle 3 \rangle 3$ .  $Q(n^+)$

PROOF:

$$\begin{aligned}
 0n^+ &= 0n + 0 \\
 &= 0n \\
 &= n0 & (\langle 3 \rangle 2) \\
 &= 0 \\
 &= n^+0
 \end{aligned}$$

$\langle 1 \rangle 3$ .  $\forall m \in \omega. P(m) \Rightarrow P(m^+)$

$\langle 2 \rangle 1$ . LET:  $m \in \omega$

$\langle 2 \rangle 2$ . ASSUME:  $P(m)$

$\langle 2 \rangle 3$ . LET:  $Q(n)$  be the statement  $m^+n = nm^+$

$\langle 2 \rangle 4$ .  $Q(0)$

PROOF:  $\langle 1 \rangle 2$

$\langle 2 \rangle 5$ .  $\forall n \in \omega. Q(n) \Rightarrow Q(n^+)$

$\langle 3 \rangle 1$ . LET:  $n \in \omega$

$\langle 3 \rangle 2$ . ASSUME:  $Q(n)$

$\langle 3 \rangle 3$ .  $Q(n^+)$

PROOF:

$$\begin{aligned}
 m^+n^+ &= m^+n + m^+ \\
 &= (m^+n + m)^+ \\
 &= (nm^+ + m)^+ & (\langle 3 \rangle 2) \\
 &= (nm + n + m)^+ \\
 &= (mn + m + n)^+ & (\langle 2 \rangle 2, \text{Proposition 18.2, Proposition 18.3}) \\
 &= (mn^+ + n)^+ \\
 &= (n^+m + n)^+ & (\langle 2 \rangle 2) \\
 &= n^+m + n^+ \\
 &= n^+m^+
 \end{aligned}$$

□

**Definition 18.8** (Exponentiation). Define *exponentiation* on  $\omega$  by recursion:

$$\begin{aligned}
 m^0 &= 1 \\
 m^{n^+} &= m^n m
 \end{aligned}$$

**Proposition 18.9.** *For all  $m, n, p \in \omega$  we have*

$$m^{n+p} = m^n m^p .$$

PROOF:

$$\langle 1 \rangle 1. m^{n+0} = m^n m^0$$

PROOF:

$$\begin{aligned} m^{n+0} &= m^n \\ &= m^n 1 \\ &= m^n m^0 \end{aligned}$$

$$\langle 1 \rangle 2. \text{ If } m^{n+p} = m^n m^p \text{ then } m^{n+p^+} = m^n m^{p^+}$$

PROOF:

$$\begin{aligned} m^{n+p^+} &= m^{n+p} m \\ &= m^n m^p m \\ &= m^n m^{p^+} \end{aligned}$$

□

**Proposition 18.10.** *For all  $m, n, p \in \omega$  we have*

$$(m^n)^p = m^{np} .$$

PROOF:

$$\langle 1 \rangle 1. (m^n)^0 = m^{n0}$$

PROOF: Both are equal to 1.

$$\langle 1 \rangle 2. \text{ If } (m^n)^p = m^{np} \text{ then } (m^n)^{p^+} = m^{np^+}$$

PROOF:

$$\begin{aligned} (m^n)^{p^+} &= (m^n)^p m^n \\ &= m^{np} m^n \\ &= m^{np+n} && \text{(Proposition 18.9)} \\ &= m^{np^+} \end{aligned}$$

□

**Proposition 18.11.** *For any natural numbers  $m$  and  $n$ , either  $m \in n$  or  $m = n$  or  $n \in m$ .*

PROOF:

$$\langle 1 \rangle 1. \text{ LET: } P(n) \text{ be the property: for all } m \in \omega, \text{ either } m \in n \text{ or } m = n \text{ or } n \in m$$

$$\langle 1 \rangle 2. P(0)$$

$$\langle 2 \rangle 1. \text{ LET: } Q(m) \text{ be the property: either } m = 0 \text{ or } 0 \in m$$

$$\langle 2 \rangle 2. Q(0)$$

PROOF: Since  $0 = 0$ .

$$\langle 2 \rangle 3. \text{ For all } m \in \omega, \text{ if } Q(m) \text{ then } Q(m^+)$$

PROOF: If  $m = 0$  or  $0 \in m$  then  $0 \in m^+$ .

$$\langle 1 \rangle 3. \text{ For any natural number } n, \text{ if } P(n) \text{ then } P(n^+)$$

$\langle 2 \rangle 1$ . LET:  $n$  be a natural number.  
 $\langle 2 \rangle 2$ . ASSUME:  $P(n)$   
 $\langle 2 \rangle 3$ . LET:  $m \in \omega$   
 $\langle 2 \rangle 4$ .  $m \in n$  or  $m = n$  or  $n \in m$   
PROOF:  $\langle 2 \rangle 2$   
 $\langle 2 \rangle 5$ . CASE:  $m \in n$  or  $m = n$   
PROOF: Then  $m \in n^+$ .  
 $\langle 2 \rangle 6$ . CASE:  $n \in m$   
 $\langle 3 \rangle 1$ . PICK  $p$  such that  $m = p^+$   
 $\langle 3 \rangle 2$ .  $n \in p$  or  $n = p$   
 $\langle 3 \rangle 3$ . CASE:  $n \in p$   
PROOF: Then  $n^+ \in p^+ = m$  by Proposition 16.7.  
 $\langle 3 \rangle 4$ . CASE:  $n = p$   
PROOF: Then  $m = n^+$ .

□

**Corollary 18.11.1** (Trichotomy). *For any natural numbers  $m$  and  $n$ , exactly one of  $m \in n$ ,  $m = n$ ,  $n \in m$  holds.*

PROOF:

$\langle 1 \rangle 1$ . We never have  $m \in n$  and  $m = n$ .  
PROOF: By Corollary 17.2.1.  
 $\langle 1 \rangle 2$ . We never have  $m \in n$  and  $n \in m$ .  
PROOF: Since  $m$  is a transitive set this would imply  $m \in m$  contradicting Corollary 17.2.1.  
 $\langle 1 \rangle 3$ . We never have  $m = n$  and  $n \in m$ .  
PROOF: By Corollary 17.2.1.

□

**Proposition 18.12.** *For any natural numbers  $m$  and  $n$ , we have  $m \in n$  if and only if  $m \subsetneq n$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $m$  and  $n$  be natural numbers.  
 $\langle 1 \rangle 2$ . If  $m \in n$  then  $m \subsetneq n$ .  
PROOF: Since  $n$  is a transitive set, and  $m \neq n$  by Corollary 17.2.1.  
 $\langle 1 \rangle 3$ . If  $m \subsetneq n$  then  $m \in n$ .  
 $\langle 2 \rangle 1$ . ASSUME:  $m \subsetneq n$   
 $\langle 2 \rangle 2$ .  $n \notin m$   
PROOF: Proposition 17.2.  
 $\langle 2 \rangle 3$ .  $m \neq n$   
 $\langle 2 \rangle 4$ .  $m \in n$   
PROOF: Trichotomy.

□

**Definition 18.13.** Given natural numbers  $m$  and  $n$ , we write  $m < n$  iff  $m \in n$ .  
We write  $m \leq n$  iff  $m < n \vee m = n$ .

**Proposition 18.14.** *For natural numbers  $m$  and  $n$ , if  $m \leq n$  and  $n \leq m$  then  $m = n$ .*

PROOF: We cannot have  $m < n$  and  $n < m$  by trichotomy.  $\square$

**Proposition 18.15.** *For natural numbers  $m$ ,  $n$  and  $k$ , if  $m < n$  then  $m + k < n + k$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $m, n \in \omega$

$\langle 1 \rangle 2$ . ASSUME:  $m < n$

$\langle 1 \rangle 3$ .  $m + 0 < n + 0$

$\langle 1 \rangle 4$ .  $\forall k \in \omega. m + k < n + k \Rightarrow m + k^+ < n + k^+$

PROOF: By Proposition 16.7.

$\square$

**Proposition 18.16.** *For natural numbers  $m$ ,  $n$  and  $k$ , if  $m < n$  and  $k \neq 0$  then  $mk < nk$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $m, n \in \omega$

$\langle 1 \rangle 2$ . ASSUME:  $m < n$

$\langle 1 \rangle 3$ .  $m1 < n1$

$\langle 1 \rangle 4$ . For all  $k \in \omega$ , if  $k \neq 0$  and  $mk < nk$  then  $m(k + 1) < n(k + 1)$

PROOF:

$$\begin{aligned} m(k + 1) &= mk + m \\ &< mk + n && \text{(Proposition 18.15)} \\ &< nk + n && \text{(Proposition 18.15)} \\ &= n(k + 1) \end{aligned}$$

$\square$

**Proposition 18.17.** *For any nonempty set of natural numbers  $E$ , there exists  $k \in E$  such that  $\forall m \in E. k \leq m$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $E \subseteq \omega$

$\langle 1 \rangle 2$ . ASSUME: there is no  $k \in E$  such that  $\forall m \in E. k \leq m$ .

PROVE:  $E = \emptyset$

$\langle 1 \rangle 3$ .  $\forall n \in \omega. n \notin E$

$\langle 2 \rangle 1$ . LET:  $P(n)$  be the property:  $\forall m < n. m \notin E$

$\langle 2 \rangle 2$ .  $P(0)$

PROOF: Vacuous.

$\langle 2 \rangle 3$ .  $\forall n \in \omega. P(n) \Rightarrow P(n + 1)$

$\langle 3 \rangle 1$ . LET:  $n \in \omega$

$\langle 3 \rangle 2$ . ASSUME:  $\forall m < n. m \notin E$

$\langle 3 \rangle 3$ .  $n \notin E$

PROOF: From  $\langle 1 \rangle 2$ .



□  $\langle 3 \rangle 4. \forall m < n + 1. m \notin E$

**Definition 18.18** (Equivalent). Sets  $E$  and  $F$  are *equivalent*,  $E \sim F$ , iff there exists a one-to-one correspondence between them.

**Proposition 18.19.** *For any set  $X$ , equivalence is an equivalence relation on  $\mathcal{P}X$ .*

PROOF: Easy.

**Proposition 18.20.** *Let  $n$  be a natural number. Let  $X$  be a proper subset of  $n$ . Then there exists  $m < n$  such that  $X \sim m$ .*

PROOF:

$\langle 1 \rangle 1.$  LET:  $P(n)$  be the property: for every proper subset  $X \subsetneq n$ , there exists  $m < n$  such that  $X \sim m$ .

$\langle 1 \rangle 2.$   $P(0)$

PROOF: Vacuous.

$\langle 1 \rangle 3.$   $\forall n \in \omega. P(n) \Rightarrow P(n + 1)$

$\langle 2 \rangle 1.$  LET:  $n \in \omega$

$\langle 2 \rangle 2.$  ASSUME:  $P(n)$

$\langle 2 \rangle 3.$  LET:  $X$  be a proper subset of  $n + 1$

$\langle 2 \rangle 4.$  CASE:  $X - \{n\} = n$

PROOF: Then  $X = n$  so  $X \sim n < n + 1$ .

$\langle 2 \rangle 5.$  CASE:  $X - \{n\} \subsetneq n$

$\langle 3 \rangle 1.$  PICK  $m < n$  such that  $X - \{n\} \sim m$

$\langle 3 \rangle 2.$   $X \sim m$  or  $X \sim m + 1$

PROOF: If  $n \in X$  then  $X \sim m + 1$ . If  $n \notin X$  then  $X \sim m$ .

□

**Proposition 18.21.** *For every natural number  $n$ , we have  $n$  is not equivalent to a proper subset of  $n$ .*

PROOF:

$\langle 1 \rangle 1.$  LET:  $P(n)$  be the property: every one-to-one function  $n \rightarrow n$  is onto.

$\langle 1 \rangle 2.$   $P(0)$

PROOF: The only function  $0 \rightarrow 0$  is  $\emptyset$ .

$\langle 1 \rangle 3.$   $\forall n \in \omega. P(n) \Rightarrow P(n + 1)$

$\langle 2 \rangle 1.$  LET:  $n \in \omega$

$\langle 2 \rangle 2.$  ASSUME:  $P(n)$

$\langle 2 \rangle 3.$  ASSUME:  $f : n + 1 \rightarrow n + 1$  is one-to-one.

$\langle 2 \rangle 4.$  LET:  $g : n \rightarrow n$  be the function

$$g(k) = \begin{cases} f(k) & \text{if } f(k) < n \\ f(n) & \text{if } f(k) = n \end{cases}$$

PROOF: If  $k < n$  and  $f(k) = n$  then  $f(n) < n$  since  $f$  is one-to-one.

$\langle 2 \rangle 5.$   $g$  is one-to-one.

$\langle 3 \rangle 1.$  LET:  $k, l < n$

$\langle 3 \rangle 2$ . ASSUME:  $g(k) = g(l)$   
 $\langle 3 \rangle 3$ . CASE:  $f(k) < n$  and  $f(l) < n$   
 PROOF: Then  $f(k) = g(k) = g(l) = f(l)$  so  $k = l$  since  $f$  is one-to-one.  
 $\langle 3 \rangle 4$ . CASE:  $f(k) < n$  and  $f(l) = n$   
 PROOF: Then  $f(k) = g(k) = g(l) = f(n)$  contradicting the fact that  $f$  is one-to-one.  
 $\langle 3 \rangle 5$ . CASE:  $f(k) = n$  and  $f(l) < n$   
 PROOF: Similar.  
 $\langle 3 \rangle 6$ . CASE:  $f(k) = n$  and  $f(l) = n$   
 PROOF: Then  $k = l$  since  $f$  is one-to-one.  
 $\langle 2 \rangle 6$ .  $g$  maps  $n$  onto  $n$ .  
 PROOF:  $\langle 2 \rangle 2$   
 $\langle 2 \rangle 7$ .  $f$  maps  $n + 1$  onto  $n + 1$ .  
 $\langle 3 \rangle 1$ . LET:  $l < n + 1$   
 $\langle 3 \rangle 2$ . CASE:  $l < n$   
 $\langle 4 \rangle 1$ . PICK  $k < n$  such that  $g(k) = l$   
 $\langle 4 \rangle 2$ .  $f(k) = l$  or  $f(n) = l$   
 $\langle 3 \rangle 3$ . CASE:  $l = n$   
 $\langle 4 \rangle 1$ . CASE:  $f(n) = n$   
 PROOF: Then  $l \in \text{ran } f$  as required.  
 $\langle 4 \rangle 2$ . CASE:  $f(n) < n$   
 $\langle 5 \rangle 1$ . PICK  $k < n$  such that  $g(k) = f(n)$   
 $\langle 5 \rangle 2$ .  $f(k) = n$

□

**Corollary 18.21.1.** *Equivalent natural numbers are equal.*

**Definition 18.22** (Finite). A set is *finite* iff it is equivalent to a natural number; otherwise, it is *infinite*.

**Proposition 18.23.** *No finite set is equivalent to one of its proper subsets.*

PROOF: From Proposition 18.21. □

**Proposition 18.24.**  $\omega$  is infinite.

PROOF: Since the function that maps  $n$  to  $n + 1$  is a one-to-one correspondence between  $\omega$  and  $\omega - \{0\}$ . □

**Proposition 18.25.** *Every subset of a finite set is finite.*

PROOF: Proposition 18.20. □

**Definition 18.26** (Number of Elements). For any finite set  $E$ , the *number of elements* in  $E$ ,  $\sharp(E)$ , is the unique natural number such that  $E \sim \sharp(E)$ .

**Proposition 18.27.** *Let  $E$  and  $F$  be finite sets. If  $E \subseteq F$  then  $\sharp(E) \leq \sharp(F)$ .*

PROOF: Proposition 18.20. □

**Proposition 18.28.** *Let  $E$  and  $F$  be disjoint finite sets. Then  $E \cup F$  is finite and  $\sharp(E \cup F) = \sharp(E) \cup \sharp(F)$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $P(n)$  be the statement:  $n \in \omega$  and for any  $m \in \omega$ , if  $E \sim m$ ,  $F \sim n$  and  $E \cap F = \emptyset$ , then  $E \cup F \sim m + n$

$\langle 1 \rangle 2$ .  $P(0)$

$\langle 2 \rangle 1$ . LET:  $m \in \omega$

$\langle 2 \rangle 2$ . LET:  $E \sim m$  and  $F \sim 0$

$\langle 2 \rangle 3$ .  $F = \emptyset$

$\langle 2 \rangle 4$ .  $E \cup F = E \sim m = m + 0$

$\langle 1 \rangle 3$ .  $\forall n \in \omega. P(n) \Rightarrow P(n + 1)$

$\langle 2 \rangle 1$ . LET:  $n \in \omega$

$\langle 2 \rangle 2$ . ASSUME:  $P(n)$

$\langle 2 \rangle 3$ . LET:  $m \in \omega$

$\langle 2 \rangle 4$ . LET:  $E \sim m$  and  $F \sim n + 1$

$\langle 2 \rangle 5$ . ASSUME:  $E \cap F = \emptyset$

$\langle 2 \rangle 6$ . PICK  $f \in F$

$\langle 2 \rangle 7$ .  $F - \{f\} \sim n$

$\langle 2 \rangle 8$ .  $E \cap (F - \{f\}) = \emptyset$

$\langle 2 \rangle 9$ .  $E \cup (F - \{f\}) \sim m + n$

PROOF:  $\langle 2 \rangle 2$

$\langle 2 \rangle 10$ .  $E \cup F \sim m + n + 1$

□

**Corollary 18.28.1.** *The union of two finite sets is finite.*

PROOF: Since, if  $E$  and  $F$  are finite, then  $E \cup F = (E - F) \cup (E \cap F) \cup (F - E)$  and these are finite and disjoint. □

**Proposition 18.29.** *If  $E$  and  $F$  are finite sets then  $E \times F$  is finite and  $\sharp(E \times F) = \sharp(E)\sharp(F)$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $P(n)$  be the statement:  $n \in \omega$  and for all  $m \in \omega$ , if  $E \sim m$  and  $F \sim n$  then  $E \times F \sim mn$

$\langle 1 \rangle 2$ .  $P(0)$

PROOF: If  $F \sim 0$  then  $F = \emptyset$  so  $E \times F = \emptyset \sim 0$ .

$\langle 1 \rangle 3$ .  $\forall n \in \omega. P(n) \Rightarrow P(n + 1)$

$\langle 2 \rangle 1$ . LET:  $n \in \omega$

$\langle 2 \rangle 2$ . ASSUME:  $P(n)$

$\langle 2 \rangle 3$ . LET:  $m \in \omega$

$\langle 2 \rangle 4$ . ASSUME:  $E \sim m$  and  $F \sim n + 1$

$\langle 2 \rangle 5$ . PICK  $f \in F$

$\langle 2 \rangle 6$ .  $F - \{f\} \sim n$

$\langle 2 \rangle 7$ .  $E \times (F - \{f\}) \sim mn$

$\langle 2 \rangle 8$ .  $E \times F = (E \times (F - \{f\})) \cup (E \times \{f\})$

$\langle 2 \rangle 9$ .  $E \times \{f\} \sim m$

⟨2⟩10.  $E \times F \sim mn + m$

PROOF: Proposition 18.28.

□

**Proposition 18.30.** *For any finite sets  $E$  and  $F$ , we have  $E^F$  is finite and  $\sharp(E^F) = \sharp(E)^{\sharp(F)}$ .*

PROOF:

⟨1⟩1. LET:  $P(n)$  be the property:  $n \in \omega$  and for all  $m \in \omega$ , if  $E \sim m$  and  $F \sim n$  then  $E^F \sim m^n$

⟨1⟩2.  $P(0)$

PROOF: Since  $E^\emptyset = \{\emptyset\} \sim 1$

⟨1⟩3.  $\forall n \in \omega. P(n) \Rightarrow P(n+1)$

⟨2⟩1. LET:  $n \in \omega$

⟨2⟩2. ASSUME:  $P(n)$

⟨2⟩3. LET:  $m \in \omega$

⟨2⟩4. LET:  $E \sim m$  and  $F \sim n+1$

⟨2⟩5. PICK  $f \in F$

⟨2⟩6.  $F - \{f\} \sim n$

⟨2⟩7. LET:  $\phi : E^F \rightarrow E^{F-\{f\}} \times E$  be the function  $\phi(g) = (g \upharpoonright (F - \{f\}), g(f))$

⟨2⟩8.  $\phi$  is a one-to-one correspondence

⟨2⟩9.  $\sharp(E^F) = m^{n+1}$

PROOF:

$$\begin{aligned} \sharp(E^F) &= \sharp(E^{F-\{f\}} \times E) \\ &= \sharp(E^{F-\{f\}}) \sharp(E) && \text{(Proposition 18.29)} \\ &= m^n m && (\langle 2 \rangle 2, \langle 2 \rangle 4) \\ &= m^{n+1} \end{aligned}$$

□

**Corollary 18.30.1.** *If  $E$  is finite then  $\mathcal{P}E$  is finite and  $\sharp(\mathcal{P}E) = 2^{\sharp(E)}$ .*

**Proposition 18.31.** *The union of a finite set of finite sets is finite.*

PROOF:

⟨1⟩1. LET:  $P(n)$  be the property: for any set  $E$ , if  $E \sim n$  and every element of  $E$  is finite, then  $\bigcup E$  is finite.

⟨1⟩2.  $P(0)$

PROOF: Since  $\bigcup \emptyset = \emptyset$  is finite.

⟨1⟩3.  $\forall n \in \omega. P(n) \Rightarrow P(n+1)$

⟨2⟩1. LET:  $n$  be a natural number.

⟨2⟩2. ASSUME:  $P(n)$

⟨2⟩3. LET:  $E \sim n+1$

⟨2⟩4. PICK  $X \in E$

⟨2⟩5.  $E - \{X\} \sim n$

⟨2⟩6.  $\bigcup(E - \{X\})$  is finite.

PROOF: ⟨2⟩2

⟨2⟩7.  $\bigcup E = \bigcup(E - \{X\}) \cup X$

⟨2⟩8.  $\bigcup E$  is finite.

PROOF: Corollary 18.28.1.

□

**Proposition 18.32.** *Every nonempty finite set of natural numbers has a greatest element.*

PROOF:

⟨1⟩1. LET:  $P(n)$  be the property: for every  $E \subseteq \mathbb{N}$ , if  $E \sim n$  then  $E$  has a greatest element.

⟨1⟩2.  $P(1)$

PROOF: Since  $k$  is the greatest element of  $\{k\}$ .

⟨1⟩3.  $\forall n \geq 1. P(n) \Rightarrow P(n+1)$

⟨2⟩1. LET:  $n \geq 1$

⟨2⟩2. ASSUME:  $P(n)$

⟨2⟩3. ASSUME:  $E \subseteq \omega$  and  $E \sim n+1$

⟨2⟩4. PICK  $k \in E$

⟨2⟩5. LET:  $l$  be the greatest element of  $E - \{k\}$

⟨2⟩6. Either  $k$  or  $l$  is greatest in  $E$ .

□

## Chapter 19

# Order