

C1 Set Theory

Robin Adams

August 12, 2022

1 Primitive Notions

Let there be *sets*.

Let there be a binary relation called *membership*, \in . When $x \in y$ holds, we say x is a *member* or *element* of y . We write $x \notin y$ iff x is not a member of y .

2 The Axioms

Axiom 1 (Extensionality). *If two sets have exactly the same members, then they are equal.*

As a consequence of this axiom, we may identify a set A with the class $\{x : x \in A\}$. The use of the symbols \in and $=$ is consistent.

Definition 2. We say that a class \mathbf{A} is a *set* iff there exists a set A such that $A = \mathbf{A}$. That is, the class $\{x : P(x)\}$ is a set iff

$$\exists A. \forall x (x \in A \leftrightarrow P(x)) .$$

Otherwise, \mathbf{A} is a *proper class*.

Definition 3 (Subset). If A is a set and \mathbf{B} is a class, we say A is a *subset* of \mathbf{B} iff $A \subseteq \mathbf{B}$.

Axiom 4 (Empty Set). *The empty class is a set, called the empty set.*

Axiom 5 (Pairing). *For any objects a and b , the class $\{a, b\}$ is a set, called a pair set.*

Definition 6 (Union). For any class of sets \mathbf{A} , the *union* $\bigcup \mathbf{A}$ is the class $\{x : \exists A \in \mathbf{A}. x \in A\}$.

We write $\bigcup_{P[x_1, \dots, x_n]} t[x_1, \dots, x_n]$ for $\bigcup \{t[x_1, \dots, x_n] : P[x_1, \dots, x_n]\}$.

Proposition 7. *If $\mathbf{A} \subseteq \mathbf{B}$ then $\bigcup \mathbf{A} \subseteq \bigcup \mathbf{B}$.*

PROOF: Easy. \square

Axiom 8 (Union). *For any set A , the union $\bigcup A$ is a set.*

Proposition 9. *For any sets A and B , the class $A \cup B$ is a set.*

PROOF: It is $\bigcup\{A, B\}$. \square

Proposition Schema 10. *For any objects a_1, \dots, a_n , the class $\{a_1, \dots, a_n\}$ is a set.*

PROOF: By repeated application of the Pairing and Union axioms. \square

Definition 11 (Power Set). For any set A , the *power set* of A , $\mathcal{P}A$, is the class of all subsets of A .

Axiom 12 (Power Set). *For any set A , the class $\mathcal{P}A$ is a set.*

Axiom 13 (Subset, Aussonderung). *For any class \mathbf{A} and set B , if $\mathbf{A} \subseteq B$ then \mathbf{A} is a set.*

Proposition 14. *For any set A and class \mathbf{B} , the intersection $A \cap \mathbf{B}$ is a set.*

PROOF: By the Subset Axiom since it is a subclass of A . \square

Proposition 15. *For any set A and class \mathbf{B} , the relative complement $A - \mathbf{B}$ is a set.*

PROOF: By the Subset Axiom since it is a subclass of A . \square

Theorem 16. *The universal class \mathbf{V} is a proper class.*

PROOF:

$\langle 1 \rangle 1$. ASSUME: \mathbf{V} is a set.

$\langle 1 \rangle 2$. LET: $R = \{x : x \notin x\}$

$\langle 1 \rangle 3$. R is a set.

PROOF: By the Subset Axiom.

$\langle 1 \rangle 4$. $R \in R$ if and only if $R \notin R$

$\langle 1 \rangle 5$. Q.E.D.

PROOF: This is a contradiction.

\square

Definition 17 (Intersection). For any class of sets \mathbf{A} , the *intersection* $\bigcap \mathbf{A}$ is the class $\{x : \forall A \in \mathbf{A}. x \in A\}$.

We write $\bigcap_{P[x_1, \dots, x_n]} t[x_1, \dots, x_n]$ for $\bigcap \{t[x_1, \dots, x_n] : P[x_1, \dots, x_n]\}$.

Proposition 18. *For any nonempty class of sets \mathbf{A} , the class $\bigcap \mathbf{A}$ is a set.*

PROOF: Pick $A \in \mathbf{A}$. Then $\bigcap \mathbf{A} \subseteq A$. \square

Proposition 19. *If $\mathbf{A} \subseteq \mathbf{B}$ then $\bigcap \mathbf{B} \subseteq \bigcap \mathbf{A}$.*

PROOF: Easy. \square

Proposition 20. *For any set A and class of sets \mathbf{B} , we have*

$$A \cup \bigcap \mathbf{B} = \bigcap \{A \cup X \mid X \in \mathbf{B}\}$$

PROOF: Easy. \square

Proposition 21. *For any set A and class of sets \mathbf{B} , we have*

$$A \cap \bigcup \mathbf{B} = \bigcup \{A \cap X \mid X \in \mathbf{B}\}$$

PROOF: Easy. \square

Proposition 22. *For any set C and class of sets \mathbf{A} , we have*

$$C - \bigcup \mathbf{A} = \bigcap \{C - X \mid X \in \mathbf{A}\} .$$

PROOF: Easy. \square

Proposition 23. *For any set C and class of sets \mathbf{A} , we have*

$$C - \bigcap \mathbf{A} = \bigcup \{C - X \mid X \in \mathbf{A}\} .$$

PROOF: Easy. \square

3 Ordered Pairs

Definition 24 (Ordered Pair). For any objects a and b , the *ordered pair* (a, b) is $\{\{a\}, \{a, b\}\}$. We call a its *first coordinate* and b its *second coordinate*.

Theorem 25. *For any objects (a, b) , we have $(a, b) = (c, d)$ if and only if $a = c$ and $b = d$.*

PROOF:

$\langle 1 \rangle 1$. If $(a, b) = (c, d)$ then $a = c$ and $b = d$

$\langle 2 \rangle 1$. ASSUME: $(a, b) = (c, d)$

$\langle 2 \rangle 2$. $a = c$

PROOF: Since $\{a\} = \bigcap(a, b) = \bigcap(c, d) = \{c\}$.

$\langle 2 \rangle 3$. $\{a, b\} = \{c, d\}$

PROOF: $\{a, b\} = \bigcup(a, b) = \bigcup(c, d) = \{c, d\}$.

$\langle 2 \rangle 4$. $b = c$ or $b = d$

$\langle 2 \rangle 5$. CASE: $b = c$

$\langle 3 \rangle 1$. $a = b$

$\langle 3 \rangle 2$. $\{c, d\} = \{a\}$

$\langle 3 \rangle 3$. $b = d$

$\langle 2 \rangle 6$. CASE: $b = d$

PROOF: We have $a = c$ and $b = d$ as required.

$\langle 1 \rangle 2$. If $a = c$ and $b = d$ then $(a, b) = (c, d)$

PROOF: Trivial.

\square

Definition 26 (Cartesian Product). The *Cartesian product* of classes \mathbf{A} and \mathbf{B} is the class

$$\mathbf{A} \times \mathbf{B} = \{(x, y) : x \in \mathbf{A}, y \in \mathbf{B}\} .$$

Lemma 27. *For any objects x and y and set C , if $x \in C$ and $y \in C$ then $(x, y) \in \mathcal{PP}C$.*

PROOF: Easy. \square

Corollary 27.1. *For any sets A and B , the Cartesian product $A \times B$ is a set.*

PROOF: By the Subset Axiom applied to $\mathcal{PP}(A \cup B)$. \square

Lemma 28. *If $(x, y) \in \mathbf{A}$ then $x, y \in \bigcup \bigcup \mathbf{A}$.*

PROOF: Easy. \square

4 Relations

Definition 29 (Relation). A *relation* is a class of ordered pairs. It is *small* iff it is a set.

When \mathbf{R} is a relation, we write $x\mathbf{R}y$ for $(x, y) \in \mathbf{R}$.

Definition 30 (Domain). The *domain* of a class \mathbf{R} is $\text{dom } \mathbf{R} = \{x : \exists y.(x, y) \in \mathbf{R}\}$.

Definition 31 (Range). The *range* of a class \mathbf{R} is $\text{ran } \mathbf{R} = \{y : \exists x.(x, y) \in \mathbf{R}\}$.

Definition 32 (Field). The *field* of a class \mathbf{R} is $\text{fld } \mathbf{R} = \text{dom } \mathbf{R} \cup \text{ran } \mathbf{R}$.

Proposition 33. *If R is a set then $\text{dom } R$, $\text{ran } R$ and $\text{fld } R$ are sets.*

PROOF: Apply the Subset Axiom to $\bigcup \bigcup R$. \square

5 n -ary Relations

Definition 34. Given objects a, b, c , define the *ordered triple* (a, b, c) to be $((a, b), c)$.

Define $(a, b, c, d) = ((a, b, c), d)$, etc.

Define the *1-tuple* (a) to be a .

Definition 35 (n -ary Relation). Given a class \mathbf{A} , an *n -ary relation* on \mathbf{A} is a class of ordered n -tuples, all of whose components are in \mathbf{A} .