

C1 Set Theory

Robin Adams

September 2, 2022

1 Primitive Notions

Let there be *sets*.

Let there be a binary relation called *membership*, \in . When $x \in y$ holds, we say x is a *member* or *element* of y . We write $x \notin y$ iff x is not a member of y .

2 The Axioms

Axiom 1 (Extensionality). *If two sets have exactly the same members, then they are equal.*

As a consequence of this axiom, we may identify a set A with the class $\{x : x \in A\}$. The use of the symbols \in and $=$ is consistent.

Definition 2. We say that a class \mathbf{A} is a *set* iff there exists a set A such that $A = \mathbf{A}$. That is, the class $\{x : P(x)\}$ is a set iff

$$\exists A. \forall x (x \in A \leftrightarrow P(x)) .$$

Otherwise, \mathbf{A} is a *proper class*.

Definition 3 (Subset). If A is a set and \mathbf{B} is a class, we say A is a *subset* of \mathbf{B} iff $A \subseteq \mathbf{B}$.

Axiom 4 (Empty Set). *The empty class is a set, called the empty set.*

Axiom 5 (Replacement). *For any property $P(x, y)$, the following is an axiom:*

Let A be a set. Assume that, for all $x \in A$, there is at most one y such that $P(x, y)$. Then $\{y : \exists x \in A. P(x, y)\}$ is a set.

Definition 6 (Power Set). For any set A , the *power set* of A , $\mathcal{P}A$, is the class of all subsets of A .

Axiom 7 (Power Set). *For any set A , the class $\mathcal{P}A$ is a set.*

Theorem 8 (Pairing). *For any objects a and b , the class $\{a, b\}$ is a set, called a pair set.*

PROOF: Let a and b be sets. Let $P(x, y)$ be the formula $(x = \emptyset \ \& \ y = a)$ or $(x = \mathcal{P}\emptyset \ \& \ y = b)$. Then we have $(\forall x \in \mathcal{P}\mathcal{P}\emptyset) \forall y_1 \forall y_2 (P(x, y_1) \ \& \ P(x, y_2) \Rightarrow y_1 = y_2)$, hence there exists a set c such that

$$\forall y (y \in c \Leftrightarrow (\exists x \in \mathcal{P}\mathcal{P}\emptyset) P(x, y))$$

The members of c are just a and b . \square

Definition 9 (Union). For any class of sets \mathbf{A} , the *union* $\bigcup \mathbf{A}$ is the class $\{x : \exists A \in \mathbf{A}. x \in A\}$.

We write $\bigcup_{P[x_1, \dots, x_n]} t[x_1, \dots, x_n]$ for $\bigcup \{t[x_1, \dots, x_n] : P[x_1, \dots, x_n]\}$.

Proposition 10. *If $\mathbf{A} \subseteq \mathbf{B}$ then $\bigcup \mathbf{A} \subseteq \bigcup \mathbf{B}$.*

PROOF: Easy. \square

Axiom 11 (Union). *For any set A , the union $\bigcup A$ is a set.*

Proposition 12. *For any sets A and B , the class $A \cup B$ is a set.*

PROOF: It is $\bigcup \{A, B\}$. \square

Proposition Schema 13. *For any objects a_1, \dots, a_n , the class $\{a_1, \dots, a_n\}$ is a set.*

PROOF: By repeated application of the Pairing and Union axioms. \square

Theorem 14 (Subset Axioms, Aussonderung). *For any class \mathbf{A} and set B , if $\mathbf{A} \subseteq B$ then \mathbf{A} is a set.*

PROOF: Let $Q(x, y)$ be the formula $x \in \mathbf{A} \wedge y = x$. Now we reason as follows. Let c be any set. Then we have

$$(\forall x \in B) \forall y_1 \forall y_2 (Q(x, y_1) \ \& \ Q(x, y_2) \Rightarrow y_1 = y_2)$$

Then, by a Replacement Axiom, there exists a set c such that

$$\forall y (y \in c \Leftrightarrow (\exists x \in B) Q(x, y)) .$$

This is equivalent to $\forall x (x \in c \Leftrightarrow x \in \mathbf{A})$. \square

Proposition 15. *For any set A and class \mathbf{B} , the intersection $A \cap \mathbf{B}$ is a set.*

PROOF: By the Subset Axiom since it is a subclass of A . \square

Proposition 16. *For any set A and class \mathbf{B} , the relative complement $A - \mathbf{B}$ is a set.*

PROOF: By the Subset Axiom since it is a subclass of A . \square

Theorem 17. *The universal class \mathbf{V} is a proper class.*

PROOF:

$\langle 1 \rangle 1$. ASSUME: \mathbf{V} is a set.

$\langle 1 \rangle 2$. LET: $R = \{x : x \notin x\}$

$\langle 1 \rangle 3$. R is a set.

PROOF: By the Subset Axiom.

$\langle 1 \rangle 4$. $R \in R$ if and only if $R \notin R$

⟨1⟩5. Q.E.D.

PROOF: This is a contradiction.

□

Definition 18 (Intersection). For any class of sets \mathbf{A} , the *intersection* $\bigcap \mathbf{A}$ is the class $\{x : \forall A \in \mathbf{A}. x \in A\}$.

We write $\bigcap_{P[x_1, \dots, x_n]} t[x_1, \dots, x_n]$ for $\bigcap \{t[x_1, \dots, x_n] : P[x_1, \dots, x_n]\}$.

Proposition 19. For any nonempty class of sets \mathbf{A} , the class $\bigcap \mathbf{A}$ is a set.

PROOF: Pick $A \in \mathbf{A}$. Then $\bigcap \mathbf{A} \subseteq A$. □

Proposition 20. If $\mathbf{A} \subseteq \mathbf{B}$ then $\bigcap \mathbf{B} \subseteq \bigcap \mathbf{A}$.

PROOF: Easy. □

Proposition 21. For any set A and class of sets \mathbf{B} , we have

$$A \cup \bigcap \mathbf{B} = \bigcap \{A \cup X \mid X \in \mathbf{B}\}$$

PROOF: Easy. □

Proposition 22. For any set A and class of sets \mathbf{B} , we have

$$A \cap \bigcup \mathbf{B} = \bigcup \{A \cap X \mid X \in \mathbf{B}\}$$

PROOF: Easy. □

Proposition 23. For any set C and class of sets \mathbf{A} , we have

$$C - \bigcup \mathbf{A} = \bigcap \{C - X \mid X \in \mathbf{A}\} .$$

PROOF: Easy. □

Proposition 24. For any set C and class of sets \mathbf{A} , we have

$$C - \bigcap \mathbf{A} = \bigcup \{C - X \mid X \in \mathbf{A}\} .$$

PROOF: Easy. □

Axiom 25 (Regularity). For every nonempty set A , there exists $m \in A$ such that $m \cap A = \emptyset$.

Theorem 26. No set is a member of itself.

PROOF: If $A \in A$ then there is no $m \in \{A\}$ such that $m \cap \{A\} = \emptyset$. □

Theorem 27. There are no sets a and b with $a \in b$ and $b \in a$.

PROOF: If there were, then there would be no $m \in \{a, b\}$ such that $m \cap \{a, b\} = \emptyset$. □

3 Ordered Pairs

Definition 28 (Ordered Pair). For any objects a and b , the *ordered pair* (a, b) is $\{\{a\}, \{a, b\}\}$. We call a its *first coordinate* and b its *second coordinate*.

Theorem 29. For any objects (a, b) , we have $(a, b) = (c, d)$ if and only if $a = c$ and $b = d$.

PROOF:

$\langle 1 \rangle 1$. If $(a, b) = (c, d)$ then $a = c$ and $b = d$

$\langle 2 \rangle 1$. ASSUME: $(a, b) = (c, d)$

$\langle 2 \rangle 2$. $a = c$

PROOF: Since $\{a\} = \bigcap(a, b) = \bigcap(c, d) = \{c\}$.

$\langle 2 \rangle 3$. $\{a, b\} = \{c, d\}$

PROOF: $\{a, b\} = \bigcup(a, b) = \bigcup(c, d) = \{c, d\}$.

$\langle 2 \rangle 4$. $b = c$ or $b = d$

$\langle 2 \rangle 5$. CASE: $b = c$

$\langle 3 \rangle 1$. $a = b$

$\langle 3 \rangle 2$. $\{c, d\} = \{a\}$

$\langle 3 \rangle 3$. $b = d$

$\langle 2 \rangle 6$. CASE: $b = d$

PROOF: We have $a = c$ and $b = d$ as required.

$\langle 1 \rangle 2$. If $a = c$ and $b = d$ then $(a, b) = (c, d)$

PROOF: Trivial.

□

Definition 30 (Cartesian Product). The *Cartesian product* of classes \mathbf{A} and \mathbf{B} is the class

$$\mathbf{A} \times \mathbf{B} = \{(x, y) : x \in \mathbf{A}, y \in \mathbf{B}\} .$$

Lemma 31. For any objects x and y and set C , if $x \in C$ and $y \in C$ then $(x, y) \in \mathcal{PPC}$.

PROOF: Easy. □

Corollary 31.1. For any sets A and B , the Cartesian product $A \times B$ is a set.

PROOF: By the Subset Axiom applied to $\mathcal{PP}(A \cup B)$. □

Lemma 32. If $(x, y) \in \mathbf{A}$ then $x, y \in \bigcup \bigcup \mathbf{A}$.

PROOF: Easy. □

4 Relations

Definition 33 (Relation). A *relation* is a class of ordered pairs. It is *small* iff it is a set.

When \mathbf{R} is a relation, we write $x\mathbf{R}y$ for $(x, y) \in \mathbf{R}$.

Definition 34 (Domain). The *domain* of a class \mathbf{R} is $\text{dom } \mathbf{R} = \{x : \exists y.(x, y) \in \mathbf{R}\}$.

Definition 35 (Range). The *range* of a class \mathbf{R} is $\text{ran } \mathbf{R} = \{y : \exists x.(x, y) \in \mathbf{R}\}$.

Definition 36 (Field). The *field* of a class \mathbf{R} is $\text{fld } \mathbf{R} = \text{dom } \mathbf{R} \cup \text{ran } \mathbf{R}$.

Proposition 37. *If R is a set then $\text{dom } R$, $\text{ran } R$ and $\text{fld } R$ are sets.*

PROOF: Apply the Subset Axiom to $\bigcup \bigcup R$. \square

Definition 38 (Single-Rooted). A class \mathbf{R} is *single-rooted* iff, for all $y \in \text{ran } \mathbf{R}$, there is only one x such that $x\mathbf{R}y$.

Definition 39 (Inverse). The *inverse* of a class \mathbf{F} is the class $\mathbf{F}^{-1} = \{(y, x) \mid (x, y) \in \mathbf{F}\}$.

Theorem 40. *For any class \mathbf{F} , we have $\text{dom } \mathbf{F}^{-1} = \text{ran } \mathbf{F}$ and $\text{ran } \mathbf{F}^{-1} = \text{dom } \mathbf{F}$.*

PROOF: Easy. \square

Theorem 41. *For a relation \mathbf{F} , $(\mathbf{F}^{-1})^{-1} = \mathbf{F}$.*

PROOF: Easy. \square

Definition 42 (Composition). The *composition* of classes \mathbf{F} and \mathbf{G} is the class $\mathbf{G} \circ \mathbf{F} = \{(x, z) \mid \exists y.(x, y) \in \mathbf{F} \wedge (y, z) \in \mathbf{G}\}$.

Theorem 43. *For any classes \mathbf{F} and \mathbf{G} , $(\mathbf{F} \circ \mathbf{G})^{-1} = \mathbf{G}^{-1} \circ \mathbf{F}^{-1}$.*

PROOF: Easy. \square

Definition 44 (Restriction). The *restriction* of the class \mathbf{F} to the class \mathbf{A} is the class $\mathbf{F} \upharpoonright \mathbf{A} = \{(x, y) : x \in \mathbf{A} \wedge (x, y) \in \mathbf{F}\}$.

Definition 45 (Image). The *image* of the class \mathbf{A} under the class \mathbf{F} is the class $\mathbf{F}(\mathbf{A}) = \{y : \exists x \in \mathbf{A}.(x, y) \in \mathbf{F}\}$.

Theorem 46.

$$\mathbf{F}(\mathbf{A} \cup \mathbf{B}) = \mathbf{F}(\mathbf{A}) \cup \mathbf{F}(\mathbf{B})$$

PROOF: Easy. \square

Theorem 47.

$$\mathbf{F}\left(\bigcup \mathbf{A}\right) = \bigcup \{\mathbf{F}(X) : X \in \mathbf{A}\}$$

PROOF: Easy. \square

Theorem 48.

$$\mathbf{F}(\mathbf{A} \cap \mathbf{B}) \subseteq \mathbf{F}(\mathbf{A}) \cap \mathbf{F}(\mathbf{B})$$

Equality holds if \mathbf{F} is single-rooted.

PROOF: Easy. \square

Theorem 49.

$$\mathbf{F}(\bigcap \mathbf{A}) \subseteq \bigcap \{\mathbf{F}(X) : X \in \mathbf{A}\}$$

Equality holds if \mathbf{F} is single-rooted.

PROOF: Easy. \square

Theorem 50.

$$\mathbf{F}(\mathbf{A}) - \mathbf{F}(\mathbf{B}) \subseteq \mathbf{F}(\mathbf{A} - \mathbf{B})$$

Equality holds if \mathbf{F} is single-rooted.

PROOF: Easy. \square

Definition 51 (Reflexive). A binary relation \mathbf{R} on \mathbf{A} is *reflexive* on \mathbf{A} if and only if $\forall x \in \mathbf{A}. x\mathbf{R}x$.

Definition 52 (Symmetric). A binary relation \mathbf{R} is *symmetric* iff, whenever $x\mathbf{R}y$, then $y\mathbf{R}x$.

Definition 53 (Transitive). A binary relation \mathbf{R} is *transitive* iff, whenever $x\mathbf{R}y$ and $y\mathbf{R}z$, then $x\mathbf{R}z$.

5 n -ary Relations

Definition 54. Given objects a, b, c , define the *ordered triple* (a, b, c) to be $((a, b), c)$.

Define $(a, b, c, d) = ((a, b, c), d)$, etc.

Define the *1-tuple* (a) to be a .

Definition 55 (n -ary Relation). Given a class \mathbf{A} , an *n -ary relation* on \mathbf{A} is a class of ordered n -tuples, all of whose components are in \mathbf{A} .

6 Functions

Definition 56 (Function). A *function* is a relation \mathbf{F} such that, for all $x \in \text{dom } \mathbf{F}$, there is only one y such that $x\mathbf{F}y$. We call this unique y the *value* of \mathbf{F} at x and denote it by $\mathbf{F}(x)$.

We say \mathbf{F} is a function *from* \mathbf{A} *into* \mathbf{B} , or \mathbf{F} *maps* \mathbf{A} into \mathbf{B} , and write $\mathbf{F} : \mathbf{A} \rightarrow \mathbf{B}$, iff \mathbf{F} is a function, $\text{dom } \mathbf{F} = \mathbf{A}$, and $\text{ran } \mathbf{F} \subseteq \mathbf{B}$.

If, in addition, $\text{ran } \mathbf{F} = \mathbf{B}$, we say \mathbf{F} is a function from \mathbf{A} *onto* \mathbf{B} .

Theorem 57. For a class \mathbf{F} , \mathbf{F}^{-1} is a function if and only if \mathbf{F} is single-rooted.

PROOF: Easy. \square

Theorem 58. A relation \mathbf{F} is a function if and only if \mathbf{F}^{-1} is single-rooted.

PROOF: Easy. \square

Theorem 59. For any function \mathbf{G} and classes \mathbf{A} and \mathbf{B} ,

$$\begin{aligned}\mathbf{G}^{-1}(\bigcup \mathbf{A}) &= \bigcup \{\mathbf{G}^{-1}(X) : X \in \mathbf{A}\} \\ \mathbf{G}^{-1}(\bigcap \mathbf{A}) &= \bigcap \{\mathbf{G}^{-1}(X) : X \in \mathbf{A}\} \quad (\text{if } \mathbf{A} \neq \emptyset) \\ \mathbf{G}^{-1}(\mathbf{A} - \mathbf{B}) &= \mathbf{G}^{-1}(\mathbf{A}) - \mathbf{G}^{-1}(\mathbf{B})\end{aligned}$$

PROOF: Easy. \square

Theorem 60. Assume that \mathbf{F} and \mathbf{G} are functions. Then $\mathbf{F} \circ \mathbf{G}$ is a function, its domain is $\{x \in \text{dom } \mathbf{G} : \mathbf{G}(x) \in \text{dom } \mathbf{F}\}$, and for x in its domain,

$$(\mathbf{F} \circ \mathbf{G})(x) = \mathbf{F}(\mathbf{G}(x)) .$$

PROOF: Easy. \square

Definition 61 (One-to-one). A function \mathbf{F} is *one-to-one* or an *injection* iff it is single-rooted.

Theorem 62. Let \mathbf{F} be a one-to-one function. For $x \in \text{dom } \mathbf{F}$, $\mathbf{F}^{-1}(\mathbf{F}(x)) = x$.

PROOF: Easy. \square

Theorem 63. Let \mathbf{F} be a one-to-one function. For $y \in \text{ran } \mathbf{F}$, $\mathbf{F}(\mathbf{F}^{-1}(y)) = y$.

PROOF: Easy. \square

Definition 64 (Identity Function). For any class \mathbf{A} , the *identity* function on \mathbf{A} is $\text{id}_{\mathbf{A}} = \{(x, x) \mid x \in \mathbf{A}\}$.

Theorem 65. Let $F : A \rightarrow B$. Assume $A \neq \emptyset$. Then F has a left inverse (i.e. there exists $G : B \rightarrow A$ such that $G \circ F = \text{id}_A$) if and only if F is one-to-one.

PROOF:

$\langle 1 \rangle 1$. If F is one-to-one then F has a left inverse.

$\langle 2 \rangle 1$. ASSUME: F is one-to-one.

$\langle 2 \rangle 2$. $F^{-1} : \text{ran } F \rightarrow A$

$\langle 2 \rangle 3$. PICK $a \in A$

$\langle 2 \rangle 4$. Define $G : B \rightarrow A$ by:

$$G(x) = \begin{cases} F^{-1}(x) & \text{if } x \in \text{ran } F \\ a & \text{if } x \in B - \text{ran } F \end{cases}$$

$\langle 2 \rangle 5$. $\forall x \in A. G(F(x)) = x$

$\langle 1 \rangle 2$. If F has a left inverse then F is one-to-one.

$\langle 2 \rangle 1$. ASSUME: F has a left inverse G .

$\langle 2 \rangle 2$. LET: $x, y \in A$ with $F(x) = F(y)$

$\langle 2 \rangle 3$. $x = y$

PROOF: $x = G(F(x)) = G(F(y)) = y$.

\square

Definition 66 (Binary Operation). A *binary operation* on a set A is a function from $A \times A$ into A .

7 The Axiom of Choice

Axiom 67 (Choice). *For any relation R there exists a function $H \subseteq R$ with $\text{dom } H = \text{dom } R$.*

Theorem 68. *Let $F : A \rightarrow B$. Then F has a right inverse if and only if F maps A onto B .*

PROOF:

$\langle 1 \rangle 1$. If F has a right inverse then F maps A onto B .

PROOF: If $H : B \rightarrow A$ is a right inverse, then for any y in B , we have $y = F(H(y))$.

$\langle 1 \rangle 2$. If F maps A onto B then F has a right inverse.

$\langle 2 \rangle 1$. ASSUME: F maps A onto B .

$\langle 2 \rangle 2$. PICK a function H with $H \subseteq F^{-1}$ and $\text{dom } H = \text{dom } F^{-1}$

PROOF: By the Axiom of Choice.

$\langle 2 \rangle 3$. $\text{dom } H = B$

PROOF: $\text{dom } H = \text{dom } F^{-1} = \text{ran } F = B$ by $\langle 2 \rangle 1$.

$\langle 2 \rangle 4$. For all $y \in B$ we have $F(H(y)) = y$

$\langle 3 \rangle 1$. LET: $y \in B$

$\langle 3 \rangle 2$. $(y, H(y)) \in F^{-1}$

$\langle 3 \rangle 3$. $F(H(y)) = y$

□

8 Sets of Functions

Definition 69. Let A be a set and \mathbf{B} be a class. Then \mathbf{B}^A is the class of all functions $A \rightarrow \mathbf{B}$.

9 Dependent Products

Definition 70. Let I be a set and H_i a set for all $i \in I$. Define

$$\prod_{i \in I} H_i = \{f : f \text{ is a function, } \text{dom } f = I, \forall i \in I. f(i) \in H_i\} .$$

Theorem 71. *The Axiom of Choice is equivalent to the statement: For any set I and any function H with domain I , if $H(i) \neq \emptyset$ for all $i \in I$, then $\prod_{i \in I} H(i) \neq \emptyset$*

PROOF:

$\langle 1 \rangle 1$. If the Axiom of Choice is true then, for any set I and any function H with domain I , if $H(i) \neq \emptyset$ for all $i \in I$, then $\prod_{i \in I} H(i) \neq \emptyset$.

$\langle 2 \rangle 1$. ASSUME: The Axiom of Choice.

$\langle 2 \rangle 2$. LET: I be a set.

$\langle 2 \rangle 3$. LET: H be a function with domain I .

- (2)4. ASSUME: $H(i) \neq \emptyset$ for all $i \in I$.
 (2)5. LET: $R = \{(i, x) : i \in I, x \in H(i)\}$
 (2)6. PICK a function $F \subseteq R$ with $\text{dom } F = \text{dom } R$
 PROVE: $F \in \prod_{i \in I} H(i)$
 PROOF: By the Axiom of Choice.
 (2)7. $\text{dom } H = I$
 PROOF: We have $\text{dom } R = I$ since for all $i \in I$ there exists x such that $x \in H(i)$.
 (2)8. $\forall i \in I. F(i) \in H(i)$
 PROOF: Since $iRF(i)$.
 (1)2. If, for any set I and any function H with domain I , if $H(i) \neq \emptyset$ for all $i \in I$, then $\prod_{i \in I} H(i) \neq \emptyset$, then the Axiom of Choice is true.
 (2)1. ASSUME: For any set I and any function H with domain I , if $H(i) \neq \emptyset$ for all $i \in I$, then $\prod_{i \in I} H(i) \neq \emptyset$
 (2)2. LET: R be a relation
 (2)3. LET: $I = \text{dom } R$
 (2)4. Define the function H with domain I by: for $i \in I$, $H(i) = \{y : iRy\}$
 (2)5. $H(i) \neq \emptyset$ for all $i \in I$
 (2)6. PICK $F \in \prod_{i \in I} H(i)$
 PROOF: By (2)1
 (2)7. F is a function
 (2)8. $F \subseteq R$
 PROOF: For all $i \in I$ we have $F(i) \in H(i)$ and so $iRF(i)$.
 (2)9. $\text{dom } F = \text{dom } R$

□

Theorem 72. *The following are equivalent.*

1. *The Axiom of Choice.*
2. *Let \mathcal{A} be a set such that (a) every member of \mathcal{A} is a nonempty set, and (b) any two distinct members of \mathcal{A} are disjoint. Then there exists a set C such that, for all $B \in \mathcal{A}$, we have $C \cap B$ is a singleton.*
3. *For any set A , there exists a function $F : \mathcal{P}A - \{\emptyset\} \rightarrow A$ such that $F(X) \in X$ for all $X \in \mathcal{P}A - \{\emptyset\}$.*

PROOF:

- (1)1. $1 \Rightarrow 2$

PROOF: Let \mathcal{A} be a set matching the two conditions. By the Multiplicative Axiom, pick a function $f \in \prod_{B \in \mathcal{A}} B$. Let $C = \text{ran } f$. Then $C \cap B = \{f(B)\}$ for all $B \in \mathcal{A}$.

- (1)2. $2 \Rightarrow 3$

- (2)1. ASSUME: 2

- (2)2. LET: A be a set.

- (2)3. LET: $\mathcal{A} = \{\{B\} \times B : B \in \mathcal{P}A - \{\emptyset\}\}$

- (2)4. PICK a set C such that $C \cap (\{B\} \times B)$ is a singleton for all $B \in \mathcal{P}A - \{\emptyset\}$

- (2)5. LET: $F = C \cap \bigcup \mathcal{A}$

$\langle 3 \rangle 3. x \mathbf{R} z$

PROOF: Since \mathbf{R} is transitive.

$\langle 3 \rangle 4. z \in [x]_{\mathbf{R}}$

$\langle 2 \rangle 3. y \mathbf{R} x$

PROOF: Since \mathbf{R} is symmetric.

$\langle 2 \rangle 4. [x]_{\mathbf{R}} \subseteq [y]_{\mathbf{R}}$

PROOF: Similar.

□

Definition 77 (Partition). A *partition* of a set A is a set $P \subseteq \mathcal{P}A$ such that:

- Every member of P is nonempty.
- Any two distinct members of P are disjoint.
- $A = \bigcup P$

Theorem 78. Let R be an equivalence relation on the set A . Then the set of all equivalence classes is a partition of A .

PROOF:

$\langle 1 \rangle 1$. Every equivalence class is nonempty.

PROOF: For any $x \in A$ we have $x \in [x]_R$.

$\langle 1 \rangle 2$. Any two distinct equivalence classes are disjoint.

$\langle 2 \rangle 1$. LET: $x, y \in A$

$\langle 2 \rangle 2$. ASSUME: $z \in [x]_R \cap [y]_R$

PROVE: $[x]_R = [y]_R$

$\langle 2 \rangle 3. x R y$

$\langle 3 \rangle 1. x R z$

$\langle 3 \rangle 2. y R z$

$\langle 3 \rangle 3. z R y$

PROOF: By $\langle 3 \rangle 2$ and symmetry.

$\langle 3 \rangle 4. x R y$

PROOF: By $\langle 3 \rangle 1$, $\langle 3 \rangle 3$ and transitivity.

$\langle 2 \rangle 4. [x]_R = [y]_R$

PROOF: By Lemma 3N.

$\langle 1 \rangle 3$. A is the union of all the equivalence classes.

PROOF: For any $x \in A$ we have $x \in [x]_R$.

□

Definition 79 (Quotient Set). If R is an equivalence relation on the set A , then the *quotient set* A/R is the set of all equivalence classes, and the *natural map* or *canonical map* $\phi : A \rightarrow A/R$ is defined by $\phi(x) = [x]_R$.

Theorem 80. Assume that R is an equivalence relation on A and that $F : A \rightarrow B$. Assume that F is compatible with R ; that is, whenever $x R y$, then $F(x) = F(y)$. Then there exists a unique $\bar{F} : A/R \rightarrow B$ such that $F = \bar{F} \circ \phi$.

PROOF: The unique such \bar{F} is $\{([x], F(x)) : x \in A\}$. □

11 Partial Orders

Definition 81 (Strict Partial Order). A *strict partial order* is an irreflexive, transitive relation.

If $<$ is a strict partial order, we write $x \leq y$ for $x < y \vee x = y$.

Theorem 82. Assume that $<$ is a partial order. Then for any x, y and z :

1. At most one of the three alternatives,

$$x < y, x = y, y < x,$$

can hold.

2. $x \leq y \leq x \Rightarrow x = y$.

PROOF: Easy. \square

Definition 83 (Minimal). Let $<$ be a partial order on D . An element $m \in D$ is *minimal* iff there is no $x \in D$ such that $x < m$.

Definition 84 (Maximal). Let $<$ be a partial order on D . An element $m \in D$ is *maximal* iff there is no $x \in D$ such that $m < x$.

Definition 85 (Least). Let $<$ be a partial order on D . An element $m \in D$ is *least*, *smallest* or the *minimum* iff $\forall x \in D. m \leq x$.

Definition 86 (Greatest). Let $<$ be a partial order on D . An element $m \in D$ is *greatest*, *largest* or the *maximum* iff $\forall x \in D. x \leq m$.

Proposition 87. If R is a partial ordering on D then so is R^{-1} .

PROOF: Easy. \square

Definition 88 (Upper Bound). Let $<$ be a partial order on A and $C \subseteq A$. An *upper bound* for C is an element $b \in A$ such that $\forall x \in C. x \leq b$.

Definition 89 (Least Upper Bound). Let $<$ be a partial order on A and $C \subseteq A$. The *least upper bound* or *supremum* for C is the least element in the set of upper bounds for C .

Definition 90 (Lower Bound). Let $<$ be a partial order on A and $C \subseteq A$. A *lower bound* for C is an element $b \in A$ such that $\forall x \in C. b \leq x$.

Definition 91 (Greatest Lower Bound). Let $<$ be a partial order on A and $C \subseteq A$. The *greatest lower bound* or *infimum* for C is the greatest element in the set of lower bounds for C .

Definition 92 (Initial Segment). Let $<$ be a partial order on A and $t \in A$. The *initial segment* up to t is

$$\text{seg } t = \{x \in A : x < t\} .$$

Definition 93 (Isomorphism). Let A and B be posets. An *isomorphism* between A and B is a bijection f between A and B such that, for all $x, y \in A$, we have $x < y$ if and only if $f(x) < f(y)$.

Proposition 94. *Isomorphism is an equivalence relation on the class of posets.*

PROOF: Easy. \square

Proposition 95. *Let $(A, <)$ be a poset and $B \subseteq A$. Then $< \cap B^2$ is a partial order on B .*

PROOF: Easy. \square

12 Linear Orders

Definition 96 (Linear Ordering). Let \mathbf{A} be a class. A *linear ordering* or *total ordering* on \mathbf{A} is a relation \mathbf{R} on \mathbf{A} such that:

- \mathbf{R} is transitive.
- \mathbf{R} satisfies *trichotomy* on \mathbf{A} ; i.e. for any $x, y \in \mathbf{A}$, exactly one of

$$x\mathbf{R}y, x = y, y\mathbf{R}x$$

holds.

Theorem 97. *Let \mathbf{R} be a linear ordering on \mathbf{A} .*

1. *There is no x such that $x\mathbf{R}x$.*
2. *For distinct x and y in \mathbf{A} , either $x\mathbf{R}y$ or $y\mathbf{R}x$.*

PROOF: Immediate from trichotomy. \square

Definition 98 (Strictly Monotone Functions). Let A and B be linearly ordered sets. A function $f : A \rightarrow B$ is *strictly monotone* iff, for all $x, y \in A$, if $x < y$ then $f(x) < f(y)$.

Theorem 99. *Let A and B be linearly ordered sets and $f : A \rightarrow B$ be strictly monotone. For all $x, y \in A$, if $f(x) < f(y)$ then $x < y$.*

PROOF: We have $f(x) \neq f(y)$ and $f(y) \not< f(x)$ by trichotomy, hence $x \neq y$ and $y \not< x$ since f is strictly monotone, hence $x < y$ by trichotomy. \square

Theorem 100. *Every strictly monotone function is injective.*

PROOF: If $f(x) = f(y)$, then we have $f(x) \not< f(y)$ and $f(y) \not< f(x)$ by trichotomy, hence $x \not< y$ and $y \not< x$ since f is strictly monotone, hence $x = y$ by trichotomy. \square

Proposition 101. *Let $(A, <)$ be a linearly ordered set and $B \subseteq A$. Then $< \cap B^2$ is a linear order on B .*

PROOF: Easy. \square

Definition 102. Let A and B be disjoint linearly ordered sets. The *concatenation* of A and B , $A \oplus B$, is the set $A \cup B$ under the order given by: $x < y$ iff

- $x, y \in A$ and $x < y$; or
- $x, y \in B$ and $x < y$; or
- $x \in A$ and $y \in B$.

It is easy to check this is a linear ordering.

Definition 103. Let A and B be linearly ordered sets. The *lexicographic order* on $A \times B$ is defined by: $(a_1, b_1) < (a_2, b_2)$ iff $a_1 < a_2$ or $(a_1 = a_2 \text{ and } b_1 < b_2)$.

13 Well Orderings

Definition 104 (Well Ordering). A *well ordering* on a set A is a linear ordering on A such that every nonempty subset of A has a least element.

Theorem 105 (Transfinite Induction Principle). *Let $<$ be a well ordering on A . Let $B \subseteq A$. Suppose that*

$$\forall x \in A (\text{seg } x \subseteq B \Rightarrow x \in B) .$$

Then $B = A$.

PROOF:

- $\langle 1 \rangle 1$. ASSUME: for a contradiction $B \neq A$
- $\langle 1 \rangle 2$. LET: t be the least element of $A - B$
- $\langle 1 \rangle 3$. $\text{seg } t \subseteq B$
- $\langle 1 \rangle 4$. $t \notin B$
- $\langle 1 \rangle 5$. Q.E.D.

PROOF: This is a contradiction.

\square

Theorem 106. *Assume that $<$ is a linear ordering on A . Assume that the only $<$ -inductive subset of A is A itself. Then $<$ is a well ordering on A .*

PROOF:

- $\langle 1 \rangle 1$. ASSUME: for a contradiction $B \subseteq A$ has no least element.
- $\langle 1 \rangle 2$. $A - B$ is $<$ -inductive.
 - $\langle 2 \rangle 1$. LET: $t \in A$
 - $\langle 2 \rangle 2$. ASSUME: $\text{seg } t \subseteq A - B$
 - $\langle 2 \rangle 3$. $t \notin B$

PROOF: If it were, it would be the least element of B .

- $\langle 2 \rangle 4$. $t \in A - B$

$\langle 1 \rangle 3. A - B = A$

$\langle 1 \rangle 4. B = \emptyset$

□

Theorem 107 (Transfinite Recursion Theorem Schema). *For any property $P(x, y)$ the following is a theorem:*

Assume that $<$ is a well ordering on A . Assume that $\forall x \exists ! y P(x, y)$. Then there exists a unique function F with domain A such that

$$\forall t \in A. P(F \upharpoonright \text{seg } t, F(t)) .$$

PROOF:

$\langle 1 \rangle 1.$ Given $t \in A$, let us say that a function v is *P-constructed up to t* iff $\text{dom } v = \{x \in A : x \leq t\}$ and $\forall x \in \text{dom } v. P(v \upharpoonright \text{seg } x, v(x))$

$\langle 1 \rangle 2.$ Let $t_1, t_2 \in A$ with $t_1 \leq t_2$. Let v_1 be a function that is *P-constructed up to t_1* , and v_2 a function that is *P-constructed up to t_2* . Then $\forall x \leq t_1. v_1(x) = v_2(x)$

$\langle 2 \rangle 1.$ LET: $x \leq t_1$

$\langle 2 \rangle 2.$ ASSUME: $\forall y < x. v_1(y) = v_2(y)$

$\langle 2 \rangle 3.$ $v_1 \upharpoonright \text{seg } x = v_2 \upharpoonright \text{seg } x$

$\langle 2 \rangle 4.$ $P(v_1 \upharpoonright \text{seg } x, v_1(x))$

$\langle 2 \rangle 5.$ $P(v_2 \upharpoonright \text{seg } x, v_2(x))$

$\langle 2 \rangle 6.$ $v_1(x) = v_2(x)$

PROOF: Since there is only one y such that $P(v_1 \upharpoonright \text{seg } x, y)$.

$\langle 2 \rangle 7.$ Q.E.D.

PROOF: By transfinite induction.

$\langle 1 \rangle 3.$ LET: $\mathcal{H} = \{v : \exists t \in A. v \text{ is } P\text{-constructed up to } t\}$

$\langle 1 \rangle 4.$ \mathcal{H} is a set.

PROOF: By a Replacement Axiom since, if v_1 and v_2 are both *P-constructed up to t* then $v_1 = v_2$ by $\langle 1 \rangle 2$.

$\langle 1 \rangle 5.$ LET: $F = \bigcup \mathcal{H}$

$\langle 1 \rangle 6.$ F is a function

$\langle 2 \rangle 1.$ ASSUME: tFx and tFy

$\langle 2 \rangle 2.$ PICK $v_1, v_2 \in \mathcal{H}$ such that $v_1(t) = x$ and $v_2(t) = y$

$\langle 2 \rangle 3.$ PICK $t_1, t_2 \in A$ such that v_1 is *P-constructed up to t_1* and v_2 is *P-constructed up to t_2*

$\langle 2 \rangle 4.$ ASSUME: w.l.o.g. $t_1 \leq t_2$

$\langle 2 \rangle 5.$ $v_1(t) = v_2(t)$

PROOF: By $\langle 1 \rangle 2$

$\langle 2 \rangle 6.$ $x = y$

$\langle 1 \rangle 7.$ $\forall x \in \text{dom } F. P(F \upharpoonright \text{seg } x, F(x))$

$\langle 2 \rangle 1.$ LET: $x \in \text{dom } F$

$\langle 2 \rangle 2.$ PICK $v \in \mathcal{H}$ such that $x \in \text{dom } v$

$\langle 2 \rangle 3.$ $P(v \upharpoonright \text{seg } x, v(x))$

$\langle 2 \rangle 4.$ $v \upharpoonright \text{seg } x = F \upharpoonright \text{seg } x$

PROOF: $\forall y < x. (y, v(y)) \in \bigcup \mathcal{H} = F$

$\langle 2 \rangle 5.$ $v(x) = F(x)$

PROOF: $(x, v(x)) \in \bigcup \mathcal{H} = F$

$\langle 1 \rangle 8.$ $\text{dom } F = A$

$\langle 2 \rangle 1.$ LET: $x \in A$

$\langle 2 \rangle 2.$ ASSUME: $\forall y < x. y \in \text{dom } F$

$\langle 2 \rangle 3.$ LET: z be the object such that $P(F \upharpoonright \text{seg } x, z)$

$\langle 2 \rangle 4.$ $F \upharpoonright \text{seg } x \cup \{(x, z)\}$ is P -constructed up to x

$\langle 2 \rangle 5.$ $x \in \text{dom } F$

$\langle 2 \rangle 6.$ Q.E.D.

PROOF: By transfinite induction, this proves $\forall x \in A. x \in \text{dom } F$.

$\langle 1 \rangle 9.$ F is unique.

$\langle 2 \rangle 1.$ LET: G be a function with domain A such that $\forall x \in A. P(G \upharpoonright \text{seg } x, G(x))$

PROVE: $\forall x \in A. F(x) = G(x)$

$\langle 2 \rangle 2.$ LET: $x \in A$

$\langle 2 \rangle 3.$ ASSUME: $\forall y < x. F(y) = G(y)$

$\langle 2 \rangle 4.$ $F \upharpoonright \text{seg } x = G \upharpoonright \text{seg } x$

$\langle 2 \rangle 5.$ $F(x) = G(x)$

$\langle 2 \rangle 6.$ Q.E.D.

PROOF: This completes the proof by transfinite induction.

□

Proposition 108. *Let $(A, <)$ be a well ordered set and $B \subseteq A$. Then $< \cap B^2$ is a well order on B .*

PROOF: Easy. □

Theorem 109. *Let A and B be well-ordered sets. Then one of the following holds:*

- $A \cong B$
- $\exists b \in B. A \cong \text{seg } b$
- $\exists a \in A. \text{seg } a \cong B$

PROOF:

$\langle 1 \rangle 1.$ PICK e that is not a member of A or B

$\langle 1 \rangle 2.$ Define $F : A \rightarrow B \cup \{e\}$ by:

$$F(t) = \begin{cases} \text{the least element of } B - F(\text{seg } t) & \text{if } B - F(\text{seg } t) \neq \emptyset \\ e & \text{if } B - F(\text{seg } t) = \emptyset \end{cases}$$

$\langle 1 \rangle 3.$ CASE: $e \in \text{ran } F$

$\langle 2 \rangle 1.$ LET: $a \in A$ be least such that $B - F(\text{seg } a) = \emptyset$

$\langle 2 \rangle 2.$ $F \upharpoonright \text{seg } a : \text{seg } a \cong B$

$\langle 1 \rangle 4.$ CASE: $\text{ran } F = B$

PROOF: In this case $F : A \cong B$.

$\langle 1 \rangle 5.$ CASE: $\text{ran } F \subset B$

$\langle 2 \rangle 1.$ LET: $b \in B$ be least such that $b \notin \text{ran } F$

$\langle 2 \rangle 2.$ $F : A \cong \text{seg } b$

□

14 Epsilon-Images

Lemma 110. *Let $<$ be a well ordering on A . Let E be the function on A defined by transfinite recursion thus:*

$$E(t) = \{E(x) : x < t\} \quad (t \in A) .$$

Let $\alpha = \text{ran } E$. Then:

1. $\forall t \in A. E(t) \notin E(t)$
2. E is injective.
3. $\forall s, t \in A. (s < t \Leftrightarrow E(s) \in E(t))$
4. α is a transitive set.

PROOF:

$\langle 1 \rangle 1. \forall t \in A. E(t) \notin E(t)$

$\langle 2 \rangle 1. \text{ LET: } t \in A$

$\langle 2 \rangle 2. \text{ ASSUME: } \forall s < t. E(s) \notin E(s)$

$\langle 2 \rangle 3. \text{ ASSUME: for a contradiction } E(t) \in E(t)$

$\langle 2 \rangle 4. \text{ PICK } x < t \text{ such that } E(t) = E(x)$

$\langle 2 \rangle 5. E(x) \in E(x)$

$\langle 2 \rangle 6. \text{ Q.E.D.}$

PROOF: This is a contradiction. The result follows by transfinite induction.

$\langle 1 \rangle 2. E$ is injective.

$\langle 2 \rangle 1. \text{ ASSUME: for a contradiction } E(x) = E(y) \text{ where } x \neq y$

$\langle 2 \rangle 2. \text{ ASSUME: w.l.o.g. } x < y$

$\langle 2 \rangle 3. E(x) \in E(y)$

$\langle 2 \rangle 4. \text{ Q.E.D.}$

PROOF: This contradicts $\langle 1 \rangle 1.$

$\langle 1 \rangle 3. \forall s, t \in A. (s < t \Leftrightarrow E(s) \in E(t))$

$\langle 2 \rangle 1. \text{ LET: } s, t \in A$

$\langle 2 \rangle 2. \text{ If } s < t \text{ then } E(s) \in E(t)$

PROOF: Immediate from definition of E .

$\langle 2 \rangle 3. \text{ If } E(s) \in E(t) \text{ then } s < t$

$\langle 3 \rangle 1. \text{ ASSUME: } E(s) \in E(t)$

$\langle 3 \rangle 2. \text{ PICK } x < t \text{ such that } E(s) = E(x)$

$\langle 3 \rangle 3. s = x$

PROOF: $\langle 1 \rangle 2.$

$\langle 3 \rangle 4. s < t$

$\langle 1 \rangle 4. \alpha$ is a transitive set.

PROOF: From definition of E .

Corollary 110.1. *For any well-ordered set $(A, <)$, if α is its epsilon-image, then $(A, <)$ is isomorphic to (α, \in) .*

Corollary 110.2. *The epsilon-image of any well-ordered set is well ordered by \in .*

Theorem 111. *Two well-ordered sets are isomorphic iff they have the same ϵ -image.*

PROOF:

$\langle 1 \rangle 1$. LET: A and B be well-ordered sets.

$\langle 1 \rangle 2$. If A and B have the same ϵ -image then they are isomorphic.

PROOF: From Corollary 110.1.

$\langle 1 \rangle 3$. If $A \cong B$ then A and B have the same epsilon-image.

$\langle 2 \rangle 1$. LET: $f : A \cong B$

$\langle 2 \rangle 2$. LET: $E : A \cong \alpha$ and $F : B \cong \beta$ be the canonical isomorphisms between A and B and their epsilon-images.

$\langle 2 \rangle 3$. $\forall x \in A. E(x) = F(f(x))$

$\langle 3 \rangle 1$. LET: $x \in A$

$\langle 3 \rangle 2$. ASSUME: $\forall y < x. E(y) = F(f(y))$

$\langle 3 \rangle 3$. $E(x) = F(f(x))$

PROOF:

$$\begin{aligned} E(x) &= \{E(y) : y < x\} \\ &= \{F(f(y)) : y < x\} \\ &= \{F(z) : z < f(x)\} \\ &= F(f(x)) \end{aligned}$$

$\langle 2 \rangle 4$. $\alpha = \beta$

□

15 Ordinal Numbers

Definition 112 (Ordinal Number). The *ordinal number* of a well-ordered set is its epsilon-image.

Definition 113 (Well-ordered by Epsilon). A set A is *well-ordered by epsilon* iff $\{(x, y) : x, y \in A, x \in y\}$ is a well ordering on A .

Theorem 114. *A set is an ordinal number if and only if it is a transitive set that is well-ordered by epsilon.*

PROOF:

$\langle 1 \rangle 1$. Every ordinal number is a transitive set.

PROOF: Lemma 110.

$\langle 1 \rangle 2$. Every ordinal number is well-ordered by epsilon.

PROOF: Corollary 110.2.

$\langle 1 \rangle 3$. Every transitive set that is well-ordered by epsilon is an ordinal number.

$\langle 2 \rangle 1$. LET: α be a transitive set well-ordered by epsilon.

$\langle 2 \rangle 2$. LET: β be the epsilon-image of (α, \in) with $E : \alpha \cong \beta$ the canonical isomorphism.

$\langle 2 \rangle 3$. $\forall x \in \alpha. E(x) = x$

$\langle 3 \rangle 1$. LET: $x \in \alpha$

$\langle 3 \rangle 2$. ASSUME: $\forall y < x. E(y) = y$

$\langle 3 \rangle 3. E(x) = x$

PROOF:

$$\begin{aligned}
 E(x) &= \{E(y) : y \in \alpha, y \in x\} \\
 &= \{E(y) : y \in x\} && (\alpha \text{ is a transitive set}) \\
 &= \{y : y \in x\} && (\langle 3 \rangle 2) \\
 &= x
 \end{aligned}$$

$\langle 2 \rangle 4. \alpha = \beta$

□

Theorem 115. *Every member of an ordinal number is an ordinal number.*

PROOF:

$\langle 1 \rangle 1.$ LET: α be an ordinal number.

$\langle 1 \rangle 2.$ LET: $\beta \in \alpha$

$\langle 1 \rangle 3.$ β is a transitive set.

$\langle 2 \rangle 1.$ LET: $x \in y \in \beta$

$\langle 2 \rangle 2.$ $y \in \alpha$

PROOF: Since α is a transitive set.

$\langle 2 \rangle 3.$ $x \in \alpha$

PROOF: Since α is a transitive set.

$\langle 2 \rangle 4.$ $x \in \beta$

PROOF: Since α is a partially ordered by epsilon.

$\langle 1 \rangle 4.$ β is well-ordered by epsilon.

PROOF: Since $\{(x, y) : x, y \in \beta, x \in y\}$ is the restriction of $\{(x, y) : x, y \in \alpha, x \in y\}$ to β .

$\langle 1 \rangle 5.$ β is an ordinal number.

PROOF: Theorem 114.

□

Proposition 116. *The class of ordinals is well-ordered by epsilon.*

PROOF:

$\langle 1 \rangle 1.$ For any ordinals α, β, γ , if $\alpha \in \beta \in \gamma$ then $\alpha \in \gamma$.

PROOF: Since γ is a transitive set (Lemma 110).

$\langle 1 \rangle 2.$ For any ordinal α we have $\alpha \notin \alpha$.

PROOF: Since α is well-ordered by epsilon.

$\langle 1 \rangle 3.$ For any ordinals α, β , exactly one of $\alpha \in \beta, \beta \in \alpha, \alpha = \beta$ holds.

$\langle 2 \rangle 1.$ LET: α, β be ordinals.

$\langle 2 \rangle 2.$ Either $\alpha \cong \beta$ or $\exists \gamma \in \beta. \alpha \cong \gamma$ or $\exists \gamma \in \alpha. \gamma \cong \alpha$

PROOF: Theorem 109.

$\langle 2 \rangle 3.$ Either $\alpha = \beta$ or $\exists \gamma \in \beta. \alpha = \gamma$ or $\exists \gamma \in \alpha. \gamma = \alpha$

PROOF: Since any ordinal is its own epsilon-image, and isomorphic well-orderings have equal epsilon-images.

$\langle 1 \rangle 4.$ Any nonempty set of ordinals has a least element.

$\langle 2 \rangle 1.$ LET: A be a nonempty set of ordinals.

$\langle 2 \rangle 2.$ PICK $\alpha \in A$

$\langle 2 \rangle 3.$ CASE: $A \cap \alpha = \emptyset$

PROOF: In this case, α is least in A .

$\langle 2 \rangle 4$. CASE: $A \cap \alpha \neq \emptyset$

PROOF: In this case, the least element of $A \cap \alpha$ is the least element in A .

□

Corollary 116.1. *Any transitive set of ordinal numbers is an ordinal number.*

Corollary 116.2. \emptyset is an ordinal number.

We write 0 for \emptyset considered as an ordinal number.

Definition 117 (Successor). The *successor* of a set a is the set $a^+ = a \cup \{a\}$.

Corollary 117.1. *The successor of an ordinal number is an ordinal number.*

Corollary 117.2. *For any set A of ordinal numbers, the set $\bigcup A$ is an ordinal number.*

Theorem 118 (Burali-Forti). *The class of ordinal numbers is not a set.*

PROOF:

$\langle 1 \rangle 1$. ASSUME: for a contradiction the class **On** is a set.

$\langle 1 \rangle 2$. **On** is an ordinal number.

PROOF: Corollary 116.1.

$\langle 1 \rangle 3$. **On** \in **On**

$\langle 1 \rangle 4$. Q.E.D.

PROOF: This contradicts Lemma 110.

□

Theorem 119 (Hartogs). *For any set A , there exists an ordinal not dominated by A .*

PROOF:

$\langle 1 \rangle 1$. LET: A be a set.

$\langle 1 \rangle 2$. LET: $\alpha = \{\beta : \beta \text{ is an ordinal, } \beta \preceq A\}$.

$\langle 1 \rangle 3$. LET: $W = \{(B, <) : B \subseteq A, < \text{ is a well ordering on } B\}$

$\langle 1 \rangle 4$. $\forall \beta \in \alpha. \exists (B, <) \in W. \beta$ is the epsilon-image of $(B, <)$

$\langle 2 \rangle 1$. LET: $\beta \in \alpha$

$\langle 2 \rangle 2$. PICK an injection $f : \beta \rightarrow A$

$\langle 2 \rangle 3$. Define $<$ on $f(\beta)$ by: $f(\gamma) < f(\delta)$ iff $\gamma \in \delta$

$\langle 2 \rangle 4$. $<$ well orders $f(\beta)$

$\langle 2 \rangle 5$. β is the epsilon-image of $(f(\beta), <)$ with f^{-1} the canonical isomorphism.

$\langle 1 \rangle 5$. α is a set.

PROOF: By a Replacement Axiom applied to W .

$\langle 1 \rangle 6$. α is an ordinal.

$\langle 2 \rangle 1$. α is a transitive set.

$\langle 3 \rangle 1$. LET: $\beta \in \gamma \in \alpha$

$\langle 3 \rangle 2$. $\beta \subseteq \gamma \preceq A$

$\langle 3 \rangle 3$. $\beta \preceq A$

$\langle 3 \rangle 4. \beta \in \alpha$
 $\langle 2 \rangle 2. \text{ Q.E.D.}$
 PROOF: By Corollary 116.1.
 $\langle 1 \rangle 7. \alpha \not\subseteq A$
 PROOF: Because $\alpha \notin \alpha$.
 \square

Theorem 120 (Zorn's Lemma). *The following statements are equivalent:*

1. *The Axiom of Choice*

Well-Ordering Theorem For any set A , there exists a well ordering on A .

Zorn's Lemma Let \mathcal{A} be a set such that, for every chain $\mathcal{B} \subseteq \mathcal{A}$, we have $\bigcup \mathcal{B} \in \mathcal{A}$. Then \mathcal{A} has a maximal element.

PROOF:

- $\langle 1 \rangle 1.$ If the Axiom of Choice is true then the Well-Ordering Theorem is true.
 $\langle 2 \rangle 1.$ ASSUME: The Axiom of Choice
 $\langle 2 \rangle 2.$ LET: A be any set.
 $\langle 2 \rangle 3.$ PICK an ordinal α not dominated by A .
 $\langle 2 \rangle 4.$ PICK an object e such that $e \notin A$.
 $\langle 2 \rangle 5.$ PICK a choice function $G : \mathcal{P}A - \{\emptyset\} \rightarrow A$ for A .
 $\langle 2 \rangle 6.$ Define the function $F : \alpha \rightarrow A \cup \{e\}$ by transfinite recursion thus:

$$F(\gamma) = \begin{cases} G(A - \{F(\delta) : \delta < \gamma\}) & \text{if } A - \{F(\delta) : \delta < \gamma\} \neq \emptyset \\ e & \text{if } A - \{F(\delta) : \delta < \gamma\} = \emptyset \end{cases}$$
 $\langle 2 \rangle 7.$ LET: δ be least such that $F(\delta) = e$
 PROOF: There is such a δ , otherwise F would be a bijection between α and A .
 $\langle 2 \rangle 8.$ $F \upharpoonright \delta$ is a bijection between δ and A
 $\langle 2 \rangle 9.$ Define $<$ on A by: $F(\gamma) < F(\beta)$ iff $\gamma \in \beta$ for $\gamma, \beta \in \delta$
 $\langle 2 \rangle 10.$ $<$ is a well ordering on A .
 $\langle 1 \rangle 2.$ If the Well-Ordering Theorem is true then Zorn's Lemma is true.
 $\langle 2 \rangle 1.$ ASSUME: The Well-Ordering Theorem
 $\langle 2 \rangle 2.$ LET: \mathcal{A} be a set that is closed under unions of chains.
 $\langle 2 \rangle 3.$ PICK a well ordering $<$ on \mathcal{A}
 $\langle 2 \rangle 4.$ Define the function $F : \mathcal{A} \rightarrow 2$ by transfinite recursion thus:

$$F(A) = \begin{cases} 1 & \text{if } \forall B < A. F(B) = 1 \Rightarrow B \subseteq A \\ 0 & \text{otherwise} \end{cases}$$
 $\langle 2 \rangle 5.$ LET: $\mathcal{C} = \{A \in \mathcal{A} : F(A) = 1\}$
 $\langle 2 \rangle 6.$ \mathcal{C} is a chain.
 $\langle 3 \rangle 1.$ LET: $A, B \in \mathcal{C}$
 $\langle 3 \rangle 2.$ ASSUME: w.l.o.g. $A < B$
 $\langle 3 \rangle 3.$ $F(A) = 1$
 $\langle 3 \rangle 4.$ $F(B) = 1$
 $\langle 3 \rangle 5.$ $A \subseteq B$
 $\langle 2 \rangle 7.$ $\bigcup \mathcal{C} \in \mathcal{A}$

PROOF: By $\langle 2 \rangle 2$.
 $\langle 2 \rangle 8$. $\bigcup \mathcal{C}$ is maximal in \mathcal{A}
 $\langle 3 \rangle 1$. ASSUME: $\bigcup \mathcal{C} \subseteq D \in \mathcal{A}$
 $\langle 3 \rangle 2$. $\forall B < D. F(B) = 1 \Rightarrow B \subseteq D$
PROOF: If $F(B) = 1$ then $B \in \mathcal{C}$ so $B \subseteq \bigcup \mathcal{C} \subseteq D$.
 $\langle 3 \rangle 3$. $F(D) = 1$
 $\langle 3 \rangle 4$. $D \in \mathcal{C}$
 $\langle 3 \rangle 5$. $D = \bigcup \mathcal{C}$
 $\langle 1 \rangle 3$. If Zorn's Lemma is true then the Axiom of Choice is true.
 $\langle 2 \rangle 1$. ASSUME: Zorn's Lemma
 $\langle 2 \rangle 2$. LET: R be a relation.
 $\langle 2 \rangle 3$. LET: \mathcal{A} be the set of all functions that are subsets of R .
 $\langle 2 \rangle 4$. For any chain $\mathcal{B} \subseteq \mathcal{A}$ we have $\bigcup \mathcal{B} \in \mathcal{A}$
 $\langle 2 \rangle 5$. PICK $F \in \mathcal{A}$ maximal.
 $\langle 2 \rangle 6$. $\text{dom } F = \text{dom } R$

□

Theorem 121 (Well-Ordering Theorem (Choice)). *For any set A , there exists a well ordering on A .*

PROOF:

$\langle 1 \rangle 1$. LET: A be a set.
 $\langle 1 \rangle 2$. PICK an ordinal α not dominated by A
 $\langle 1 \rangle 3$. $A \preceq \alpha$
 $\langle 1 \rangle 4$. PICK an injection $f : A \rightarrow \alpha$
 $\langle 1 \rangle 5$. Define $<$ on A by: $x < y$ iff $f(x) \in f(y)$
 $\langle 1 \rangle 6$. $<$ is a well ordering on A .

□

Corollary 121.1 (Numeration Theorem (Choice)). *Any set is equinumerous to some ordinal number.*

Theorem 122 (Transfinite Recursion). *Let $\mathbf{F} : \mathbf{V} \rightarrow \mathbf{V}$. Then there exists a function $\mathbf{G} : \mathbf{On} \rightarrow \mathbf{V}$ such that*

$$\forall \alpha \in \mathbf{On}. \mathbf{G}(\alpha) = \mathbf{F}(\mathbf{G} \upharpoonright \alpha) .$$

PROOF: Define $\mathbf{G} = \{(\alpha, y) : \exists f : \alpha^+ \rightarrow \mathbf{V}. \forall \beta \in \alpha^+. f(\beta) = \mathbf{F}(f \upharpoonright \beta)\}$. □

Definition 123 (Continuous). A function $\mathbf{F} : \mathbf{On} \rightarrow \mathbf{On}$ is *continuous* iff $\mathbf{F}(\lambda) = \bigcup_{\beta \in \lambda} \mathbf{F}(\beta)$ for every limit ordinal λ .

Theorem 124. *Let $\mathbf{F} : \mathbf{On} \rightarrow \mathbf{On}$ be continuous. Suppose $\forall \alpha \in \mathbf{On}. \mathbf{F}(\alpha) < \mathbf{F}(\alpha + 1)$. Then \mathbf{F} is strictly monotone.*

PROOF:

$\langle 1 \rangle 1$. LET: $P(\beta)$ be the statement: $\forall \alpha < \beta. \mathbf{F}(\alpha) < \mathbf{F}(\beta)$
 $\langle 1 \rangle 2$. $P(0)$

PROOF: Vacuous.

$\langle 1 \rangle 3. \forall \beta \in \mathbf{On}. P(\beta) \Rightarrow P(\beta^+)$

PROOF: For $\alpha < \beta^+$ we have $\mathbf{F}(\alpha) \leq \mathbf{F}(\beta) < \mathbf{F}(\beta^+)$.

$\langle 1 \rangle 4.$ For every limit ordinal λ , if $\forall \beta < \lambda. P(\beta)$ then $P(\lambda)$

PROOF: For $\alpha < \lambda$ we have $\mathbf{F}(\alpha) < \mathbf{F}(\alpha^+) \leq \mathbf{F}(\lambda)$.

□

Definition 125 (Normal). A function $\mathbf{F} : \mathbf{On} \rightarrow \mathbf{On}$ is *normal* iff it is strictly monotone and continuous.

Theorem 126. Let $\mathbf{F} : \mathbf{On} \rightarrow \mathbf{On}$ be normal. Let $t_0 \leq \beta$. Then there exists a greatest γ such that $\mathbf{F}(\gamma) \leq \beta$.

PROOF:

$\langle 1 \rangle 1.$ LET: $\gamma = \{\alpha \in \mathbf{On} : \mathbf{F}(\alpha) \leq \beta\}$

$\langle 1 \rangle 2.$ γ is an ordinal.

$\langle 2 \rangle 1.$ γ is a set.

PROOF: We have $\alpha \leq \mathbf{F}(\alpha)$ for all α , so $\gamma \subseteq \beta$.

$\langle 2 \rangle 2.$ γ is a transitive set.

PROOF: If $\alpha < \alpha'$ and $\mathbf{F}(\alpha') \leq \beta$ then $\mathbf{F}(\alpha) < \beta$ by monotonicity.

$\langle 1 \rangle 3.$ $\gamma \neq 0$

PROOF: By hypothesis.

$\langle 1 \rangle 4.$ CASE: γ is a successor ordinal.

PROOF: Let $\gamma = \alpha^+$. Then α is greatest such that $\mathbf{F}(\alpha) \leq \beta$.

$\langle 1 \rangle 5.$ CASE: γ is a limit ordinal.

PROOF: This is impossible since then $\mathbf{F}(\gamma) = \bigcup_{\alpha \in \gamma} \mathbf{F}(\alpha) \leq \beta$ and so $\gamma \in \gamma$.

□

Theorem 127. Let $\mathbf{F} : \mathbf{On} \rightarrow \mathbf{On}$ be normal. Let S be a nonempty set of ordinals. Then $\mathbf{F}(\sup S) = \sup \mathbf{F}(S)$.

PROOF:

$\langle 1 \rangle 1.$ $\mathbf{F}(\sup S) \geq \sup \mathbf{F}(S)$

PROOF: By monotonicity.

$\langle 1 \rangle 2.$ $\mathbf{F}(\sup S) \leq \sup \mathbf{F}(S)$

$\langle 2 \rangle 1.$ CASE: $\sup S \in S$

PROOF: Immediate.

$\langle 2 \rangle 2.$ CASE: $\sup S \notin S$

$\langle 3 \rangle 1.$ $\sup S$ is a limit ordinal.

$\langle 3 \rangle 2.$ $\mathbf{F}(\sup S) = \sup\{\mathbf{F}(\beta) : \beta < \sup S\}$

$\langle 3 \rangle 3.$ $\forall \beta < \sup S. \mathbf{F}(\beta) \leq \sup \mathbf{F}(S)$

□

Theorem 128 (Veblen Fixed-Point Theorem (1907)). A normal operation on ordinals has arbitrarily large fixed points.

That is, let $\mathbf{F} : \mathbf{On} \rightarrow \mathbf{On}$ be normal. For all $\alpha \in \mathbf{On}$, there exists $\beta \geq \alpha$ such that $\mathbf{F}(\beta) = \beta$.

PROOF: Let $\beta = \sup_{n \in \omega} F^n(\alpha)$. Then $\alpha \leq \beta$ using monotonicity, and

$$\begin{aligned} F(\beta) &= \sup_{n \in \omega} F^{n+1}(\alpha) \\ &= \beta \end{aligned}$$

□

16 Natural Numbers

Definition 129 (Inductive). A class \mathbf{A} is *inductive* iff $\emptyset \in \mathbf{A}$ and $\forall a \in \mathbf{A}. a^+ \in \mathbf{A}$.

Axiom 130 (Infinity). *There exists an inductive set.*

Definition 131 (Natural Number). A *natural number* is a set that belongs to every inductive set.

We write ω for the class of all natural numbers.

Theorem 132. *The class ω is a set.*

PROOF: Pick an inductive set I (by the Axiom of Infinity), then apply a Subset Axiom to I . □

Theorem 133. *The set ω is inductive, and is a subset of every inductive set.*

PROOF: Easy. □

Corollary 133.1 (Proof by Induction). *Any inductive subclass of ω is equal to ω .*

Theorem 134. *Every natural number except 0 is the successor of some natural number.*

PROOF: Easy proof by induction. □

Definition 135 (Peano System). A *Peano system* is a triple $\langle N, S, e \rangle$ consisting of a set N , a function $S : N \rightarrow N$ and an element $e \in N$ such that:

1. $e \notin \text{ran } S$
2. S is one-to-one
3. Any subset $A \subseteq N$ that contains e and is closed under S equals N .

Definition 136 (Transitive Set). A set A is a *transitive set* iff every member of a member of A is a member of A .

Theorem 137. *For any transitive set a , $\bigcup(a^+) = a$.*

PROOF:

$$\begin{aligned} \bigcup(a^+) &= \bigcup(a \cup \{a\}) \\ &= \bigcup a \cup \bigcup \{a\} \\ &= \bigcup a \cup a \\ &= a \end{aligned}$$

since $\bigcup a \subseteq a$. □

Theorem 138. *Every natural number is a transitive set.*

PROOF:

$\langle 1 \rangle 1$. 0 is a transitive set.

PROOF: Vacuous.

$\langle 1 \rangle 2$. For any natural number n , if n is a transitive set then n^+ is a transitive set.

$\langle 2 \rangle 1$. LET: n be a natural number that is a transitive set.

$\langle 2 \rangle 2$. $\bigcup(n^+) \subseteq n^+$

PROOF: Theorem 137.

□

Theorem 139. $\langle \omega, \sigma, 0 \rangle$ is a Peano system, where $0 = \emptyset$ and $\sigma = \{ \langle n, n^+ \rangle : n \in \omega \}$.

PROOF:

$\langle 1 \rangle 1$. $0 \notin \text{ran } \sigma$

PROOF: For any $n \in \omega$ we have $0 \neq n^+$ since $n \in n^+$ and $n \notin 0$.

$\langle 1 \rangle 2$. σ is one-to-one.

PROOF: If $m^+ = n^+$ then $m = \bigcup(m^+) = \bigcup(n^+) = n$ using Theorems 137 and 138.

$\langle 1 \rangle 3$. Any subset $A \subseteq \omega$ that contains 0 and is closed under σ equals ω .

□

Theorem 140. *The set ω is a transitive set.*

PROOF:

$\langle 1 \rangle 1$. For every natural number n we have $\forall m \in n$. m is a natural number.

$\langle 2 \rangle 1$. $\forall m \in 0$. m is a natural number.

PROOF: Vacuous.

$\langle 2 \rangle 2$. If n is a natural number and $\forall m \in n$. m is a natural number, then $\forall m \in n^+$. m is a natural number.

PROOF: Since if $m \in n^+$ we have either $m \in n$ or $m = n$, and m is a natural number in either case.

□

Theorem 141 (Recursion Theorem on ω). *Let A be a set, $a \in A$ and $F : A \rightarrow A$. Then there exists a unique function $h : \omega \rightarrow A$ such that*

$$h(0) = a ,$$

and for every n in ω ,

$$h(n^+) = F(h(n)) .$$

PROOF:

$\langle 1 \rangle 1$. Let us call a function v *acceptable* iff $\text{dom } v \subseteq \omega$, $\text{ran } v \subseteq A$ and:

1. If $0 \in \text{dom } v$ then $v(0) = a$

2. For all $n \in \omega$, if $n^+ \in \text{dom } v$ then $n \in \text{dom } v$ and $v(n^+) = F(v(n))$.

PROOF: If there were then there would be no $m \in \text{ran } f$ such that $m \cap \text{ran } f = \emptyset$, contradicting the Axiom of Regularity. \square

17 Finite Sets

Definition 144 (Finite). A set is *finite* iff it is equinumerous with a natural number. Otherwise it is infinite.

Theorem 145. *No natural number is equinumerous with a proper subset of itself.*

PROOF:

$\langle 1 \rangle 1$. Any injective function $f : 0 \rightarrow 0$ has range 0 .

PROOF: Since the only such function is \emptyset .

$\langle 1 \rangle 2$. For any natural number n , if every injective function $f : n \rightarrow n$ has range n , then every injective function $f : n^+ \rightarrow n^+$ has range n^+ .

$\langle 2 \rangle 1$. LET: $n \in \omega$

$\langle 2 \rangle 2$. ASSUME: Every injective function $f : n \rightarrow n$ has range n .

$\langle 2 \rangle 3$. LET: $f : n^+ \rightarrow n^+$ be injective.

$\langle 2 \rangle 4$. Define $g : n \rightarrow n$ by

$$g(k) = \begin{cases} f(k) & \text{if } f(k) \in n \\ f(n) & \text{if } f(k) = n \end{cases}$$

PROOF: If $k \in n$ and $f(k) = n$ then $f(n) \in n$ since f is injective.

$\langle 2 \rangle 5$. g is injective.

$\langle 3 \rangle 1$. LET: $i, j \in n$

$\langle 3 \rangle 2$. ASSUME: $g(i) = g(j)$

$\langle 3 \rangle 3$. CASE: $f(i) \in n, f(j) \in n$

PROOF: Then $f(i) = f(j)$ so $i = j$

$\langle 3 \rangle 4$. CASE: $f(i) \in n, f(j) \notin n$

PROOF: Then $f(i) = f(n)$ which is impossible as f is injective.

$\langle 3 \rangle 5$. CASE: $f(i) \notin n, f(j) \in n$

PROOF: Then $f(n) = f(j)$ which is impossible as f is injective.

$\langle 3 \rangle 6$. CASE: $f(i) \notin n, f(j) \notin n$

PROOF: Then $f(i) = f(j) = n$ so $i = j$.

$\langle 2 \rangle 6$. $\text{ran } g = n$

PROOF: By $\langle 2 \rangle 2$.

$\langle 2 \rangle 7$. $\text{ran } f = n^+$

$\langle 3 \rangle 1$. $\forall k \in n. k \in \text{ran } f$

PROOF: Since $\text{ran } g \subseteq \text{ran } f$.

$\langle 3 \rangle 2$. $n \in \text{ran } f$

$\langle 4 \rangle 1$. CASE: $f(n) \in n$

$\langle 5 \rangle 1$. PICK k such that $g(k) = f(n)$

$\langle 5 \rangle 2$. $f(k) = n$

$\langle 4 \rangle 2$. CASE: $f(n) = n$

PROOF: Then $n \in \text{ran } f$.

□

Corollary 145.1. *No finite set is equinumerous with a proper subset of itself.*

Corollary 145.2. *The set ω is infinite.*

PROOF: Since the function that maps n to $n + 1$ is a bijection between ω and the proper subset $\omega - \{0\}$. □

Corollary 145.3. *Every finite set is equinumerous with a unique natural number.*

Lemma 146. *Let n be a natural number and $C \subseteq n$. Then there exists $m \in n$ such that $C \approx m$.*

PROOF:

⟨1⟩1. For all $C \subseteq 0$, there exists $m \in 0$ such that $C \approx m$.

PROOF: In this case $C = \emptyset$ and so $C \approx 0$.

⟨1⟩2. Let $n \in \omega$. Assume that, for all $C \subseteq n$, there exists $m \in n$ such that $C \approx m$.

Let $C \subseteq n^+$. Then there exists $m \in n^+$ such that $C \approx m$.

⟨2⟩1. LET: $n \in \omega$

⟨2⟩2. ASSUME: For all $C \subseteq n$, there exists $m \in n$ such that $C \approx m$.

⟨2⟩3. LET: $C \subseteq n^+$

⟨2⟩4. CASE: $n \in C$

⟨3⟩1. PICK $m \in n$ such that $C - \{n\} \approx m$

⟨3⟩2. $C \approx m^+$

⟨2⟩5. CASE: $n \notin C$

PROOF: Then $C \subseteq n$ so $C \approx m$ for some $m \in n$.

□

Corollary 146.1. *Any subset of a finite set is finite.*

18 Cardinal Numbers

Definition 147 (Cardinality (Choice)). For any set A , define the *cardinal number* of A , $|A|$, to be the least ordinal that is equinumerous with A .

Theorem 148. *For any sets A and B , $|A| = |B|$ if and only if $A \approx B$.*

PROOF: Easy. □

Theorem 149. *For any finite set A , $|A|$ is the natural number such that $A \approx |A|$.*

PROOF: Immediate from definitions. □

Definition 150. We write \aleph_0 for $|\omega|$.

19 Cardinal Arithmetic

Definition 151 (Addition). Let κ and λ be any cardinal numbers. Then $\kappa + \lambda = |K \cup L|$, where K and L are any disjoint sets of cardinality κ and λ respectively.

To show this is well-defined, we must prove that, if $K_1 \approx K_2$, $L_1 \approx L_2$, and $K_1 \cap L_1 = K_2 \cap L_2 = \emptyset$, then $K_1 \cup L_1 \approx K_2 \cup L_2$.

PROOF: Easy.

Lemma 152. For any cardinal number κ we have $\kappa + 0 = \kappa$.

PROOF: Since for any set K we have $K \cup \emptyset = K$.

Lemma 153. For any natural number n we have $n + \aleph_0 = \aleph_0$.

PROOF: Easy. \square

Lemma 154.

$$\aleph_0 + \aleph_0 = \aleph_0$$

PROOF: Define $f : (\omega \times \{0\}) \cup (\omega \times \{1\}) \rightarrow \omega$ by $f(n, 0) = 2n$ and $f(n, 1) = 2n + 1$. Then f is a bijection. \square

Theorem 155.

$$\kappa + \lambda = \lambda + \kappa$$

PROOF: Easy. \square

Theorem 156.

$$\kappa + (\lambda + \mu) = (\kappa + \lambda) + \mu$$

PROOF: Easy. \square

Definition 157 (Multiplication). Let κ and λ be any cardinal numbers. Then $\kappa\lambda = |K \times L|$, where K and L are any sets of cardinality κ and λ respectively.

It is easy to prove this well-defined.

Lemma 158. For any cardinal number κ we have $\kappa 0 = 0$.

PROOF: For any set K we have $K \times \emptyset = \emptyset$. \square

Lemma 159. For any natural number n we have $n\aleph_0 = \aleph_0$.

PROOF: Induction on n using Lemma 154. \square

Lemma 160.

$$\aleph_0 \aleph_0 = \aleph_0$$

PROOF: Define $f : \omega \times \omega \rightarrow \omega$ by $f(m, n) = 2^m(2n + 1) - 1$. Then f is a bijection. \square

Lemma 161.

$$\kappa 1 = \kappa$$

PROOF: Easy. \square

Theorem 162.

$$\kappa\lambda = \lambda\kappa$$

PROOF: Easy. \square

Theorem 163.

$$\kappa(\lambda\mu) = (\kappa\lambda)\mu$$

PROOF: Easy. \square

Theorem 164.

$$\kappa(\lambda + \mu) = \kappa\lambda + \kappa\mu$$

PROOF: Easy. \square

Definition 165 (Exponentiation). Let κ and λ be any cardinal numbers. Then $\kappa^\lambda = |K^L|$, where K and L are any sets of cardinality κ and λ respectively.

It is easy to prove this well-defined.

Theorem 166. For any cardinal κ , $\kappa^0 = 1$.

PROOF: For any set K , there is only one function $\emptyset \rightarrow K$, namely \emptyset . \square

Theorem 167. For any non-zero cardinal κ , we have $0^\kappa = 0$.

PROOF: For any nonempty set K , there is no function $K \rightarrow \emptyset$. \square

Theorem 168. For any set A , $|\mathcal{P}A| = 2^{|A|}$.

PROOF: Define the bijection $f : \mathcal{P}A \rightarrow 2^A$ by $f(S)(a) = 1$ if $a \in S$, 0 if $a \notin S$. \square

Corollary 168.1. For any cardinal κ , we have $\kappa \neq 2^\kappa$.

Theorem 169.

$$\kappa^{\lambda+\mu} = \kappa^\lambda \kappa^\mu$$

PROOF: Easy. \square

Theorem 170.

$$(\kappa\lambda)^\mu = \kappa^\mu \lambda^\mu$$

PROOF: Easy. \square

Theorem 171.

$$(\kappa^\lambda)^\mu = \kappa^{\lambda\mu}$$

PROOF: Easy. \square

Lemma 172. The union of a set of cardinal numbers is a cardinal number.

PROOF:

$\langle 1 \rangle 1$. LET: A be a set of cardinal numbers.

$\langle 1 \rangle 2$. LET: $\alpha \in \bigcup A$

$\langle 1 \rangle 3$. PICK $\kappa \in A$ such that $\alpha \in \kappa$

$\langle 1 \rangle 4$. $\alpha \prec \kappa$

$\langle 1 \rangle 5$. $\alpha \prec \bigcup A$

\square

20 Alephs

Definition 173. Define the cardinal number \aleph_α for every ordinal α by transfinite recursion thus: \aleph_α is the least infinite cardinal different from \aleph_β for every $\beta < \alpha$.

Theorem 174. *If $\alpha < \beta$ then $\aleph_\alpha < \aleph_\beta$.*

PROOF: By minimality of \aleph_α . \square

Theorem 175. *Every infinite cardinal is of the form \aleph_α for some α .*

PROOF:

$\langle 1 \rangle 1$. LET: κ be an infinite cardinal

$\langle 1 \rangle 2$. ASSUME: for every infinite cardinal $\lambda < \kappa$, there exists α such that $\lambda = \aleph_\alpha$

$\langle 1 \rangle 3$. LET: $\alpha = \{\beta : \aleph_\beta < \kappa\}$

$\langle 1 \rangle 4$. α is a set.

PROOF: The mapping $\beta \mapsto \aleph_\beta$ is an injection $\alpha \rightarrow \kappa$.

$\langle 1 \rangle 5$. α is a transitive set.

$\langle 1 \rangle 6$. α is an ordinal.

$\langle 1 \rangle 7$. \aleph_α is the least infinite cardinal different from \aleph_β for all β such that $\aleph_\beta < \kappa$.

$\langle 1 \rangle 8$. \aleph_α is the least infinite cardinal different from λ for every infinite cardinal $\lambda < \kappa$.

PROOF: By $\langle 1 \rangle 2$.

$\langle 1 \rangle 9$. $\aleph_\alpha = \kappa$

\square

21 Arithmetic

Lemma 176. *For any natural numbers m and n , we have $m + n^+ = (m + n)^+$.*

PROOF: Easy. \square

Corollary 176.1. *The union of two finite sets is finite.*

Lemma 177. *For any natural numbers m and n we have $mn^+ = mn + m$.*

PROOF: Easy. \square

Corollary 177.1. *The Cartesian product of two finite sets is finite.*

Lemma 178. *For any natural numbers m and n we have $m^{n^+} = m^n m$.*

PROOF: Easy. \square

Corollary 178.1. *If A and B are finite sets then A^B is finite.*

22 Ordering on the Natural Numbers

Lemma 179. *For any natural numbers m and n , $m \in n$ if and only if $m^+ \in n^+$.*

PROOF:

$\langle 1 \rangle 1.$ $\forall m, n \in \omega (m \in n \Rightarrow m^+ \in n^+)$

$\langle 2 \rangle 1.$ $\forall m \in \omega (m \in 0 \Rightarrow m^+ \in 0^+)$

PROOF: Vacuous.

$\langle 2 \rangle 2.$ For all $n \in \omega$, if $\forall m \in n. m^+ \in n^+$ then $\forall m \in n^+. m^+ \in n^{++}$

$\langle 3 \rangle 1.$ LET: $n \in \omega$

$\langle 3 \rangle 2.$ ASSUME: $\forall m \in n. m^+ \in n^+$

$\langle 3 \rangle 3.$ LET: $m \in n^+$

$\langle 3 \rangle 4.$ CASE: $m \in n$

$\langle 4 \rangle 1.$ $m^+ \in n^+$

PROOF: By $\langle 3 \rangle 2$

$\langle 4 \rangle 2.$ $m^+ \in n^{++}$

$\langle 3 \rangle 5.$ CASE: $m = n$

PROOF: $m^+ = n^+ \in n^{++}$

$\langle 1 \rangle 2.$ $\forall m, n \in \omega (m^+ \in n^+ \Rightarrow m \in n)$

$\langle 2 \rangle 1.$ LET: $m, n \in \omega$

$\langle 2 \rangle 2.$ ASSUME: $m^+ \in n^+$

$\langle 2 \rangle 3.$ $m \in m^+$

$\langle 2 \rangle 4.$ $m^+ \in n$ or $m^+ = n$

$\langle 2 \rangle 5.$ $m \in n$

PROOF: If $m^+ \in n$ this follows because n is transitive (Theorem 138).

□

Lemma 180. *For any natural number n we have $n \notin n$.*

PROOF:

$\langle 1 \rangle 1.$ $0 \notin 0$

$\langle 1 \rangle 2.$ For all $n \in \omega$, if $n \notin n$ then $n^+ \notin n^+$

$\langle 2 \rangle 1.$ LET: $n \in \omega$

$\langle 2 \rangle 2.$ ASSUME: $n^+ \in n^+$

PROVE: $n \in n$

$\langle 2 \rangle 3.$ $n^+ \in n$ or $n^+ = n$

$\langle 2 \rangle 4.$ $n \in n^+$

$\langle 2 \rangle 5.$ $n \in n$

PROOF: If $n^+ \in n$ this follows because n is transitive (Theorem 138).

□

Theorem 181 (Trichotomy Law for ω). *For any natural numbers m and n , exactly one of*

$$m \in n, m = n, n \in m$$

holds.

PROOF:

$\langle 1 \rangle 1$. For any $m, n \in \omega$, at most one of $m \in n$, $m = n$, $n \in m$ holds.
 PROOF: If $m \in n$ and $m = n$ then $m \in m$ contradicting Lemma 180.
 If $m \in n$ and $n \in m$ then $m \in m$ by Theorem 138, contradicting Lemma 180.
 $\langle 1 \rangle 2$. For any $m, n \in \omega$, at least one of $m \in n$, $m = n$, $n \in m$ holds.
 $\langle 2 \rangle 1$. For all $n \in \omega$, either $0 \in n$ or $0 = n$
 $\langle 3 \rangle 1$. $0 = 0$
 $\langle 3 \rangle 2$. For all $n \in \omega$, if $0 \in n$ or $0 = n$ then $0 \in n^+$
 $\langle 2 \rangle 2$. For all $m \in \omega$, if $\forall n \in \omega (m \in n \vee m = n \vee n \in m)$ then $\forall n \in \omega (m^+ \in n \vee m^+ = n \vee n \in m^+)$
 $\langle 3 \rangle 1$. LET: $m \in \omega$
 $\langle 3 \rangle 2$. ASSUME: $\forall n \in \omega (m \in n \vee m = n \vee n \in m)$
 $\langle 3 \rangle 3$. LET: $n \in \omega$
 $\langle 3 \rangle 4$. CASE: $m \in n$
 PROOF: Then $m \in n^+$
 $\langle 3 \rangle 5$. CASE: $m = n$
 PROOF: Then $m \in n^+$
 $\langle 3 \rangle 6$. CASE: $n \in m$
 PROOF: Then $n^+ \in m^+$ by Lemma 179 so $n^+ \in m$ or $n^+ = m$.

□

Corollary 181.1. *The relation \in is a linear ordering on ω .*

Corollary 181.2. *For any natural numbers m and n ,*

$$m \in n \Leftrightarrow m \subset n .$$

PROOF:

$\langle 1 \rangle 1$. LET: $m, n \in \omega$
 $\langle 1 \rangle 2$. If $m \in n$ then $m \subset n$.
 $\langle 2 \rangle 1$. ASSUME: $m \in n$
 $\langle 2 \rangle 2$. $m \subseteq n$
 PROOF: Theorem 138.
 $\langle 2 \rangle 3$. $m \neq n$
 PROOF: Lemma 180.
 $\langle 1 \rangle 3$. If $m \subset n$ then $m \in n$.
 PROOF: We have $m \neq n$ and $n \notin m$ by $\langle 1 \rangle 2$, hence $m \in n$ by trichotomy.

□

Theorem 182. *For any natural number p , the function that maps n to $n + p$ is strictly monotone. For any natural numbers m , n and p , we have $m \in n$ if and only if $m + p \in n + p$.*

PROOF: We prove that $m \in n \Rightarrow m + p \in n + p$. This is an easy induction on p using Lemma 179. □

Theorem 183. *For any non-zero natural number p , the function that maps n to np is strictly monotone.*

PROOF: Easy induction on p using Theorem 182. □

Theorem 184 (Strong Induction). *Let A be a subset of ω and suppose that, for all $n \in \omega$, we have*

$$(\forall m < n. m \in A) \Rightarrow n \in A .$$

Then $A = \omega$.

PROOF: Prove $\forall n \in \omega. \forall m < n. m \in A$ by induction on n . \square

Theorem 185 (Well-Ordering of ω). *The ordering $<$ on ω is a well-ordering.*

PROOF: If A is a subset of ω with no least element, we prove $\forall n \in \omega. n \notin A$ by strong induction on n . \square

Theorem 186 (Choice). *Let $<$ be a linear ordering on A . Then $<$ is a well-ordering on A iff there does not exist any function $f : \omega \rightarrow \omega$ such that $f(n+1) < f(n)$ for all $n \in \omega$.*

PROOF:

$\langle 1 \rangle 1$. If $<$ is a well-ordering on A then there does not exist any function $f : \omega \rightarrow \omega$ such that $f(n+1) < f(n)$ for all $n \in \omega$.

PROOF: If there is such a function f then $\text{ran } f$ is a nonempty subset of A with no least element.

$\langle 1 \rangle 2$. If there does not exist any function $f : \omega \rightarrow A$ such that $f(n+1) < f(n)$ for all $n \in \omega$ then $<$ is a well-ordering on A .

$\langle 2 \rangle 1$. LET: $X \subseteq A$ be a nonempty subset of A with no least element.

PROVE: There exists a function $f : \omega \rightarrow A$ such that $f(n+1) < f(n)$ for all $n \in \omega$

$\langle 2 \rangle 2$. PICK $a_0 \in X$

$\langle 2 \rangle 3$. $\forall x \in X. \exists y \in X. y < x$

$\langle 2 \rangle 4$. PICK a function $g : X \rightarrow X$ such that $\forall x \in X. g(x) < x$

PROOF: By the Axiom of Choice.

$\langle 2 \rangle 5$. Define $f : \omega \rightarrow A$ recursively by:

$$f(0) = a_0$$

$$f(n^+) = g(f(n))$$

$\langle 2 \rangle 6$. $\forall n \in \omega. f(n^+) < f(n)$

\square

Lemma 187. *For any natural numbers m and n , we have $m \in n$ if and only if there exists a natural number p such that $n = m + p^+$.*

PROOF:

$\langle 1 \rangle 1$. For all m, p , we have $m \in m + p^+$

PROOF: $m = m + 0 \in m + p^+$

$\langle 1 \rangle 2$. For all m, n , if $m \in n$ then there exists p such that $n = m + p^+$

$\langle 2 \rangle 1$. For all m , if $m \in 0$ then there exists p such that $0 = m + p^+$

PROOF: Vacuous.

$\langle 2 \rangle 2$. For all $n \in \omega$, if $\forall m \in n. \exists p \in \omega. n = m + p^+$ then $\forall m \in n^+. \exists p \in \omega. n^+ = m + p^+$

- $\langle 3 \rangle 1.$ LET: $n \in \omega$
 - $\langle 3 \rangle 2.$ ASSUME: $\forall m \in n. \exists p \in \omega. n = m + p^+$
 - $\langle 3 \rangle 3.$ LET: $m \in n^+$
 - $\langle 3 \rangle 4.$ CASE: $m \in n$
 - $\langle 4 \rangle 1.$ PICK p such that $n = m + p^+$
 - $\langle 4 \rangle 2.$ $n^+ = m + p^{++}$
 - $\langle 3 \rangle 5.$ CASE: $m = n$
- PROOF: $n^+ = m + 0^+$

□

Lemma 188. *For natural numbers m, n, p and q , if $m \in n$ and $p \in q$ then $mp + nq \in mq + np$.*

- $\langle 1 \rangle 1.$ PICK natural numbers a and b such that $n = m + a^+$ and $q = p + b^+$
- PROOF: Lemma 187.
- $\langle 1 \rangle 2.$ $mp + nq = mq + np + (a^+ + b)^+$
- $\langle 1 \rangle 3.$ $mp + nq \in mq + np$
- PROOF: Lemma 187.

23 The Integers

Theorem 189. *The relation \sim is an equivalence relation on $\omega \times \omega$, where $(m, n) \sim (p, q)$ iff $m + q = n + p$.*

PROOF:

- $\langle 1 \rangle 1.$ The relation \sim is reflexive on ω^2
- PROOF: For any m, n , we have $m + n = m + n$ and so $(m, n) \sim (m, n)$.
- $\langle 1 \rangle 2.$ The relation \sim is symmetric.
- PROOF: If $m + q = n + p$ then $p + n = q + m$.
- $\langle 1 \rangle 3.$ The relation \sim is transitive.
- $\langle 2 \rangle 1.$ ASSUME: $(m, n) \sim (p, q) \sim (r, s)$
- $\langle 2 \rangle 2.$ $m + q = n + p$
- $\langle 2 \rangle 3.$ $p + s = q + r$
- $\langle 2 \rangle 4.$ $m + p + q + s = n + p + q + r$
- $\langle 2 \rangle 5.$ $m + s = n + r$
- PROOF: By cancellation of addition in ω .

□

Definition 190. The set \mathbb{Z} of *integers* is the quotient set $(\omega \times \omega) / \sim$.

Lemma 191. *If $(m, n) \sim (m', n')$ and $(p, q) \sim (p', q')$ then $(m + p, n + q) \sim (m' + p', n' + q')$.*

PROOF: Assume $m + n' = m' + n$ and $p + q' = p' + q$. Then $m + p + n' + q' = m' + p' + n + q$. □

Definition 192 (Addition). Addition $+$ on \mathbb{Z} is the binary operation such that

$$[(m, n)] + [(p, q)] = [(m + p, n + q)]$$

Theorem 193. *Addition on \mathbb{Z} is commutative.*

PROOF: From the definition. \square

Theorem 194. *Addition on \mathbb{Z} is associative.*

PROOF: Easy. \square

Definition 195 (Zero). The zero in the integers is $0 = [(0, 0)]$.

Theorem 196. *For any integer a we have $a + 0 = a$.*

PROOF: Easy. \square

Theorem 197. *For any integer a , there exists an integer b such that $a + b = 0$.*

PROOF: If $a = [(m, n)]$ take $b = [(n, m)]$. \square

Lemma 198. *If $(m, n) \sim (m', n')$ and $(p, q) \sim (p', q')$ then $(mp + nq, mq + np) \sim (m'p' + n'q', m'q' + n'p')$.*

PROOF:

- $\langle 1 \rangle 1.$ ASSUME: $m + n' = m' + n$ and $p + q' = p' + q$
 - $\langle 1 \rangle 2.$ $mp + n'p = m'p + np$
 - $\langle 1 \rangle 3.$ $m'q + nq = mq + n'q$
 - $\langle 1 \rangle 4.$ $mp + mq' = m'p' + mq$
 - $\langle 1 \rangle 5.$ $n'p' + n'q = n'p + n'q'$
 - $\langle 1 \rangle 6.$ $mp + n'p + m'q + nq + mp + mq' + n'p' + n'q = m'p + np + mq + n'q + m'p' + mq + n'p + n'q'$
 - $\langle 1 \rangle 7.$ $mp + nq + m'q' + n'p' = mq + np + m'p' + n'q'$
- \square

Definition 199 (Multiplication). *Multiplication \cdot is the binary operation on \mathbb{Z} such that*

$$[(m, n)][(p, q)] = [(mp + nq, mq + np)]$$

Theorem 200. *Multiplication is commutative.*

PROOF: Easy. \square

Theorem 201. *Multiplication is associative.*

PROOF: Easy. \square

Theorem 202. *Multiplication is distributive over addition.*

PROOF: Easy. \square

Definition 203. The integer one is $1 = [(1, 0)]$.

Theorem 204. *For any integer a we have $a1 = a$.*

PROOF: Easy. \square

Theorem 205. $0 \neq 1$

PROOF: Easy. \square

Lemma 206. *If $(m, n) \sim (m', n')$ and $(p, q) \sim (p', q')$ then $m + q \in p + n$ iff $m' + q' \in p' + n'$.*

PROOF:

$$\begin{aligned} m + q \in p + n &\Leftrightarrow m + q + n' + q' \in p + n + n' + q' \\ &\Leftrightarrow m' + n + q + q' \in p' + n + n' + q \\ &\Leftrightarrow m' + q' \in p' + n' \end{aligned} \quad \square$$

Definition 207 (Ordering). The ordering $<$ on \mathbb{Z} is defined by: $[(m, n)] < [(p, q)]$ iff $m + q \in n + p$.

Theorem 208. *The relation $<$ is a linear ordering on \mathbb{Z} .*

PROOF:

- $\langle 1 \rangle 1.$ $<$ is transitive.
 - $\langle 2 \rangle 1.$ ASSUME: $[(m, n)] < [(p, q)]$ and $[(p, q)] < [(r, s)]$
 - $\langle 2 \rangle 2.$ $m + q \in n + p$ and $p + s \in q + r$
 - $\langle 2 \rangle 3.$ $m + q + s \in n + p + s$
 - $\langle 2 \rangle 4.$ $n + p + s \in n + q + r$
 - $\langle 2 \rangle 5.$ $m + q + s \in n + q + r$
 - $\langle 2 \rangle 6.$ $m + s \in n + r$
- $\langle 1 \rangle 2.$ $<$ satisfies trichotomy.

PROOF: From trichotomy on ω .

\square

Theorem 209. *For any integers a, b and c , we have $a < b$ iff $a + c < b + c$.*

PROOF: An easy consequence of the corresponding property in ω .

Corollary 209.1. *If $a + c = b + c$ then $a = b$.*

Theorem 210. *If $0 < c$, then the function that maps an integer a to ac is strictly monotone.*

PROOF:

- $\langle 1 \rangle 1.$ LET: a, b and c be integers.
- $\langle 1 \rangle 2.$ ASSUME: $0 < c$ and $a < b$
- $\langle 1 \rangle 3.$ LET: $a = [(m, n)]$
- $\langle 1 \rangle 4.$ LET: $b = [(p, q)]$
- $\langle 1 \rangle 5.$ LET: $c = [(r, s)]$
- $\langle 1 \rangle 6.$ $s \in r$
- $\langle 1 \rangle 7.$ $m + q \in p + n$
- $\langle 1 \rangle 8.$ $(m + q)r + (p + n)s \in (m + q)s + (p + n)r$

PROOF: Lemma 188.

- $\langle 1 \rangle 9.$ $ac < bc$

□

Lemma 211. For integers a and b , $a(-b) = -(ab)$

PROOF: This follows from the fact that $ab + a(-b) = a(b + (-b)) = a0 = 0$. □

Theorem 212. For integers a , b and c , if $a < b$ and $c < 0$ then $ac > bc$.

PROOF: We have $0 < -c$ so $a(-c) < b(-c)$ hence $-(ac) < -(bc)$ so $bc < ac$. □

Theorem 213. For any integers a and b , if $ab = 0$ then $a = 0$ or $b = 0$.

PROOF: We prove if $a \neq 0$ and $b \neq 0$ then $ab \neq 0$.

If $a > 0$ and $b > 0$ then $ab > 0$. Similarly for the other four cases. □

Theorem 214. If $ac = bc$ and $c \neq 0$ then $a = b$.

PROOF: We have $(a - b)c = 0$ so $a - b = 0$ hence $a = b$. □

Definition 215 (Positive). An integer a is *positive* iff $0 < a$.

Theorem 216. Define $E : \omega \rightarrow \mathbb{Z}$ by $E(n) = [(n, 0)]$. Then E maps ω one-to-one into \mathbb{Z} , and:

1. $E(m + n) = E(m) + E(n)$
2. $E(mn) = E(m)E(n)$
3. $m \in n$ if and only if $E(m) < E(n)$.

PROOF: Routine calculations. □

24 Equinumerosity

Definition 217 (Equinumerous). Two sets A and B are *equinumerous*, $A \approx B$, iff there exists a bijection between them.

Theorem 218. *Equinumerosity is an equivalence relation on the class of sets.*

PROOF: Easy. □

Theorem 219 (Cantor 1873). *No set is equinumerous with its power set.*

PROOF:

⟨1⟩1. LET: $g : A \rightarrow \mathcal{P}A$

PROVE: g is not surjective.

⟨1⟩2. LET: $B = \{x \in A : x \notin g(x)\}$

⟨1⟩3. $\forall x \in A. g(x) \neq B$

PROOF: Because $x \in B$ iff $x \notin g(x)$.

□

25 Ordering Cardinal Numbers

Definition 220 (Dominated). A set A is *dominated* by a set B , $A \preceq B$, iff there exists an injection $f : A \rightarrow B$.

Lemma 221. *Domination is a preorder on the class of sets.*

PROOF: Easy. \square

Lemma 222. *If $A \subseteq B$ then $A \preceq B$.*

PROOF: The inclusion from A to B is an injection. \square

Lemma 223. *If $A \preceq B$, $A \approx A'$ and $B \approx B'$ then $A' \preceq B'$.*

PROOF: Easy. \square

Definition 224. Given cardinal numbers κ and λ , we write $\kappa \leq \lambda$ iff $K \preceq L$, where K is any set of cardinality κ and L is any set of cardinality λ .

We write $\kappa < \lambda$ iff $\kappa \leq \lambda$ and $\kappa \neq \lambda$.

Theorem 225 (Schröder-Bernstein). *If $A \preceq B$ and $B \preceq A$ then $A \approx B$.*

PROOF:

$\langle 1 \rangle 1$. LET: $f : A \rightarrow B$ and $g : B \rightarrow A$ be one-to-one.

$\langle 1 \rangle 2$. Define the sequence of sets $C_n \subseteq A$ by:

$$C_0 = A - \text{ran } g$$

$$C_{n+1} = g(f(C_n))$$

$\langle 1 \rangle 3$. Define $h : A \rightarrow B$ by

$$h(x) = \begin{cases} f(x) & \text{if } \exists n \in \mathbb{N}. x \in C_n \\ g^{-1}(x) & \text{otherwise} \end{cases}$$

$\langle 1 \rangle 4$. h is injective.

$\langle 2 \rangle 1$. LET: $x, y \in A$

$\langle 2 \rangle 2$. ASSUME: $h(x) = h(y)$

$\langle 2 \rangle 3$. CASE: $x \in C_m, y \in C_n$

PROOF: We have $f(x) = f(y)$ so $x = y$

$\langle 2 \rangle 4$. CASE: $x \in C_m, y \notin \bigcup_n C_n$

PROOF: This case is impossible because we would have $y = g(f(x))$ and so $y \in C_{m+1}$.

$\langle 2 \rangle 5$. CASE: $x, y \notin \bigcup_n C_n$

PROOF: We have $g^{-1}(x) = g^{-1}(y)$ so $x = y$.

$\langle 1 \rangle 5$. h is surjective.

$\langle 2 \rangle 1$. LET: $y \in B$

$\langle 2 \rangle 2$. ASSUME: $y \notin f(C_n)$ for all n

$\langle 2 \rangle 3$. $g(y) \notin C_n$ for all n

$\langle 2 \rangle 4$. $y = h(g(y))$

\square

Corollary 225.1. *The relation \leq is a partial order on the class of cardinal numbers.*

Theorem 226. *Let κ , λ and μ be cardinal numbers.*

1. $\kappa \leq \lambda \Rightarrow \kappa + \mu \leq \lambda + \mu$
2. $\kappa \leq \lambda \Rightarrow \kappa\mu \leq \lambda\mu$
3. $\kappa \leq \lambda \Rightarrow \kappa^\mu \leq \lambda^\mu$
4. $\kappa \leq \lambda \Rightarrow \mu^\kappa \leq \mu^\lambda$ if κ and μ are not both zero.

PROOF: Parts 1–3 are easy. For part 4:

Let $|K| = \kappa$, $|L| = \lambda$ and $|M| = \mu$ with $K \subseteq L$.

If $M = \emptyset$ then $\kappa \neq 0$ so $\mu^\kappa = 0 \leq \mu^\lambda$.

Otherwise, pick $a \in M$. Define $\Phi : M^K \rightarrow M^L$ by:

$$\Phi(f)(x) = \begin{cases} f(x) & \text{if } x \in K \\ a & \text{if } x \notin K \end{cases}$$

Then Φ is an injection. \square

Theorem 227 (Cardinal Comparability). *The Axiom of Choice is equivalent to the statement: for any sets C and D , either $C \preceq D$ or $D \preceq C$.*

PROOF:

- $\langle 1 \rangle 1$. If Zorn's Lemma then Cardinal Comparability.
 - $\langle 2 \rangle 1$. ASSUME: Zorn's Lemma
 - $\langle 2 \rangle 2$. LET: C and D be sets.
 - $\langle 2 \rangle 3$. LET: \mathcal{A} be the set of all injective functions f with $\text{dom } f \subseteq C$ and $\text{ran } f \subseteq D$
 - $\langle 2 \rangle 4$. For every chain $\mathcal{B} \subseteq \mathcal{A}$ we have $\bigcup \mathcal{B} \in \mathcal{A}$
 - $\langle 2 \rangle 5$. LET: $f \in \mathcal{A}$ be maximal
 - $\langle 2 \rangle 6$. $\text{dom } f = C$ or $\text{ran } f = D$
 - $\langle 2 \rangle 7$. f is an injective function $C \rightarrow D$ or f^{-1} is an injective function $D \rightarrow C$
- $\langle 1 \rangle 2$. If Cardinal Comparability then the Well-Ordering Theorem.
 - $\langle 2 \rangle 1$. ASSUME: Cardinal Comparability
 - $\langle 2 \rangle 2$. LET: A be any set
 - $\langle 2 \rangle 3$. PICK an ordinal α not dominated by A

PROOF: Hartogs' Theorem.

 - $\langle 2 \rangle 4$. $A \preceq \alpha$
 - $\langle 2 \rangle 5$. PICK an injective function $f : A \rightarrow \alpha$
 - $\langle 2 \rangle 6$. Define $<$ on A by: $x < y$ iff $f(x) \in f(y)$
 - $\langle 2 \rangle 7$. $<$ is a well ordering on A .

\square

Theorem 228 (Choice). *For any infinite set A , we have $\omega \preceq A$.*

PROOF:

- $\langle 1 \rangle 1$. LET: A be an infinite set.
- $\langle 1 \rangle 2$. PICK a choice function F for A
- $\langle 1 \rangle 3$. Define $f : \omega \rightarrow A$ by recursion by: $f(n) = F(A - \{f(0), f(1), \dots, f(n-1)\})$

PROOF: $A - \{f(0), f(1), \dots, f(n-1)\}$ is nonempty because A is infinite.
 $\langle 1 \rangle 4$. f is injective.
 \square

Corollary 228.1 (Choice). *For any infinite cardinal κ we have $\aleph_0 \leq \kappa$.*

Corollary 228.2 (Choice). *A set is infinite iff it is equinumerous to a proper subset of itself.*

Proposition 229 (Choice). *If there exists a surjection $A \rightarrow B$ then $B \preceq A$.*

PROOF: Any surjection $A \rightarrow B$ has a right inverse which is an injection $B \rightarrow A$.

26 Countable Sets

Definition 230 (Countable). A set is *countable* iff it is dominated by ω .

Proposition 231. *Any subset of a countable set is countable.*

PROOF: Easy. \square

The union of two countable sets is countable.

PROOF: Because $\aleph_0 + \aleph_0 = \aleph_0$ \square

Proposition 232. *The product of two countable sets is countable.*

PROOF: Because $\aleph_0 \aleph_0 = \aleph_0$. \square

Proposition 233 (Choice). *For any infinite set A , the set $\mathcal{P}A$ is uncountable.*

PROOF: If $|A| \geq \aleph_0$ then $|\mathcal{P}A| \geq 2^{\aleph_0}$. \square

Theorem 234 (Choice). *A countable union of countable sets is countable.*

PROOF:

$\langle 1 \rangle 1$. LET: \mathcal{A} be a countable set of countable sets.

$\langle 1 \rangle 2$. ASSUME: w.l.o.g. $\mathcal{A} \neq \emptyset$ and $\emptyset \notin \mathcal{A}$

$\langle 1 \rangle 3$. PICK a surjection $G : \omega \rightarrow \mathcal{A}$

$\langle 1 \rangle 4$. PICK a function F with domain ω such that, for all m , $F(m)$ is a surjection $\omega \rightarrow G(m)$

PROOF: By the Axiom of Choice.

$\langle 1 \rangle 5$. Define $f : \omega \times \omega \rightarrow \bigcup \mathcal{A}$ by $f(m, n) = F(m)(n)$

$\langle 1 \rangle 6$. f is surjective.

$\langle 1 \rangle 7$. $A \preceq \omega \times \omega$

\square

27 Arithmetic of Infinite Cardinals

Lemma 235 (Choice). *For any infinite cardinal κ we have $\kappa \cdot \kappa = \kappa$.*

PROOF:

- ⟨1⟩1. LET: κ be an infinite cardinal.
- ⟨1⟩2. LET: B be a set of cardinality κ .
- ⟨1⟩3. LET: $\mathcal{H} = \{f : f = \emptyset \text{ or for some infinite } A \subseteq B, f \text{ is a bijection between } A \times A \text{ and } A\}$
- ⟨1⟩4. For any chain $\mathcal{C} \subseteq \mathcal{H}$, we have $\bigcup \mathcal{C} \in \mathcal{H}$
 - ⟨2⟩1. LET: $\mathcal{C} \subseteq \mathcal{H}$ be a chain.
 - ⟨2⟩2. ASSUME: w.l.o.g. \mathcal{C} has a nonempty element.
PROOF: Otherwise $\bigcup \mathcal{C} = \emptyset \in \mathcal{H}$.
 - ⟨2⟩3. $\bigcup \mathcal{C}$ is an injective function.
 - ⟨2⟩4. LET: $A = \text{ran } \bigcup \mathcal{C}$
 - ⟨2⟩5. A is infinite.
 - ⟨2⟩6. $\bigcup \mathcal{C}$ is a bijection between $A \times A$ and A .
- ⟨3⟩1. LET: $a_1, a_2 \in A$
- ⟨3⟩2. PICK $f_1, f_2 \in \mathcal{C}$ such that $a_1 \in \text{ran } f_1$ and $a_2 \in \text{ran } f_2$
- ⟨3⟩3. ASSUME: w.l.o.g. $f_1 \subseteq f_2$
- ⟨3⟩4. $\langle a_1, a_2 \rangle \in \text{dom } f_2$
- ⟨3⟩5. $\langle a_1, a_2 \rangle \in \text{dom } \bigcup \mathcal{C}$
- ⟨1⟩5. PICK a maximal $f_0 \in \mathcal{H}$
PROOF: Zorn's Lemma.
- ⟨1⟩6. $f_0 \neq \emptyset$
PROOF: B has a countable subset A , say, and $A \times A \approx A$.
- ⟨1⟩7. PICK $A_0 \subseteq B$ infinite such that f_0 is a bijection between $A_0 \times A_0$ and A_0 .
- ⟨1⟩8. LET: $\lambda = |A_0|$
- ⟨1⟩9. λ is infinite
- ⟨1⟩10. $\lambda = \lambda \cdot \lambda$
- ⟨1⟩11. $\lambda = \kappa$
 - ⟨2⟩1. $|B - A_0| < \lambda$
 - ⟨3⟩1. ASSUME: for a contradiction $\lambda \leq |B - A_0|$
 - ⟨3⟩2. PICK $D \subseteq B - A_0$ with $|D| = \lambda$
 - ⟨3⟩3. $(A_0 \cup D) \times (A_0 \cup D) = (A_0 \times A_0) \cup (A_0 \times D) \cup (D \times A_0) \cup (D \times D)$
 - ⟨3⟩4. $f_0 : A_0 \times A_0 \approx A_0$
 - ⟨3⟩5. $|(A_0 \times D) \cup (D \times A_0) \cup (D \times D)| = \lambda$
PROOF:

$$\begin{aligned} |(A_0 \times D) \cup (D \times A_0) \cup (D \times D)| &= \lambda \cdot \lambda + \lambda \cdot \lambda + \lambda \cdot \lambda \\ &= \lambda + \lambda + \lambda & (\langle 1 \rangle 10) \\ &= 3 \cdot \lambda \\ &\leq \lambda \cdot \lambda \\ &= \lambda & (\langle 1 \rangle 10) \end{aligned}$$
 - ⟨3⟩6. PICK a bijection $g : (A_0 \times D) \cup (D \times A_0) \cup (D \times D) \approx D$
 - ⟨3⟩7. $f_0 \cup g : (A_0 \cup D) \times (A_0 \cup D) \approx A_0 \cup D$
 - ⟨3⟩8. Q.E.D.
- PROOF: This contradicts the maximality of f_0 .
- ⟨2⟩2. $\lambda = \kappa$

PROOF:

$$\begin{aligned}
\kappa &= |B| \\
&= |A_0| + |B - A_0| \\
&\leq \lambda + \lambda \\
&= 2 \cdot \lambda \\
&\leq \lambda \cdot \lambda \\
&= \lambda \\
&\leq \kappa
\end{aligned}$$

□

Corollary 235.1 (Absorption Law of Cardinal Arithmetic (Choice)). *Let κ and λ be cardinal numbers, the larger of which is infinite and the smaller of which is nonzero. Then*

$$\kappa + \lambda = \kappa \cdot \lambda = \max(\kappa, \lambda) \ .$$

PROOF:

⟨1⟩1. ASSUME: w.l.o.g. $\kappa \leq \lambda$

⟨1⟩2. $\kappa + \lambda = \lambda$

PROOF:

$$\begin{aligned}
\lambda &\leq \kappa + \lambda \\
&\leq \lambda + \lambda \\
&= 2 \cdot \lambda \\
&\leq \lambda \cdot \lambda \\
&= \lambda
\end{aligned}$$

⟨1⟩3. $\kappa \cdot \lambda = \lambda$

PROOF:

$$\begin{aligned}
\lambda &= 1 \cdot \lambda \\
&\leq \kappa \cdot \lambda \\
&\leq \lambda \cdot \lambda \\
&= \lambda
\end{aligned}$$

□

28 Rank

Definition 236. Define the set V_α for every ordinal α by transfinite recursion thus:

$$V_\alpha = \bigcup \{ \mathcal{P}V_\beta : \beta \in \alpha \} \ .$$

Lemma 237. *For any ordinal α , V_α is a transitive set.*

PROOF:

⟨1⟩1. LET: α be an ordinal.

- ⟨1⟩2. LET: $x \in y \in V_\alpha$
- ⟨1⟩3. PICK $\beta \in \alpha$ such that $y \in \mathcal{P}V_\beta$
- ⟨1⟩4. $x \in V_\beta$
- ⟨1⟩5. PICK $\gamma \in \beta$ such that $x \in \mathcal{P}V_\gamma$
- ⟨1⟩6. $\gamma \in \alpha$ and $x \in \mathcal{P}V_\gamma$
- ⟨1⟩7. $x \in V_\alpha$

□

Theorem 238. For ordinals $\beta \in \alpha$ we have $V_\beta \subseteq V_\alpha$.

PROOF:

$$\begin{aligned}
 V_\beta &= \bigcup_{\gamma \in \beta} \mathcal{P}V_\gamma \\
 &\subseteq \bigcup_{\gamma \in \alpha} \mathcal{P}V_\gamma \\
 &= V_\alpha
 \end{aligned}
 \quad \square$$

Theorem 239.

$$V_0 = \emptyset$$

PROOF: Immediate from definitions. □

Theorem 240. For any ordinal α , $V_{\alpha+} = \mathcal{P}V_\alpha$.

PROOF:

$$\begin{aligned}
 V_{\alpha+} &= \bigcup_{\beta \leq \alpha} \mathcal{P}V_\beta \\
 &= \mathcal{P}V_\alpha
 \end{aligned}$$

by Theorem 238. □

Theorem 241. For λ a limit ordinal, $V_\lambda = \bigcup_{\beta < \lambda} V_\beta$.

PROOF:

$$\begin{aligned}
 V_\lambda &= \bigcup_{\beta < \lambda} \mathcal{P}V_\beta \\
 &= \bigcup_{\beta < \lambda} V_{\beta+} \\
 &= \bigcup_{\beta < \lambda} V_\beta
 \end{aligned}$$

since $\beta < \lambda$ iff $\beta^+ < \lambda$. □

Definition 242 (Grounded, Rank). A set A is *grounded* iff $\exists \alpha. A \subseteq V_\alpha$. The *rank* of a grounded set A , $\text{rank } A$, is then the least ordinal α such that $A \subseteq V_\alpha$.

Theorem 243. If A is grounded and $a \in A$ then a is grounded and $\text{rank } a < \text{rank } A$.

PROOF: We have $a \in A \subseteq V_{\text{rank } A}$. So $a \in \mathcal{P}V_\alpha$ for some $\alpha < \text{rank } A$, i.e. $a \subseteq V_\alpha$ for some $\alpha < \text{rank } A$, as required.

Theorem 244. *If every member of A is grounded then A is grounded and*

$$\text{rank } A = \sup_{a \in A} (\text{rank } a)^+ .$$

PROOF:

- $\langle 1 \rangle 1.$ LET: $\alpha = \sup_{a \in A} (\text{rank } a)^+$
- $\langle 1 \rangle 2.$ $A \subseteq V_\alpha$
 - $\langle 2 \rangle 1.$ LET: $a \in A$
 - $\langle 2 \rangle 2.$ $a \subseteq V_{\text{rank } a}$
 - $\langle 2 \rangle 3.$ $a \in V_{(\text{rank } a)^+}$
 - $\langle 2 \rangle 4.$ $a \in V_\alpha$
- $\langle 1 \rangle 3.$ If $A \subseteq V_\beta$ then $\alpha \leq \beta$
 - $\langle 2 \rangle 1.$ ASSUME: $A \subseteq V_\beta$
 - $\langle 2 \rangle 2.$ $\forall a \in A. a \in V_\beta$
 - $\langle 2 \rangle 3.$ $\forall a \in A. \exists \gamma < \beta. a \subseteq V_\gamma$
 - $\langle 2 \rangle 4.$ $\forall a \in A. \exists \gamma < \beta. \text{rank } a \leq \gamma$
 - $\langle 2 \rangle 5.$ $\forall a \in A. \text{rank } a < \beta$
 - $\langle 2 \rangle 6.$ $\forall a \in A. (\text{rank } a)^+ \leq \beta$
 - $\langle 2 \rangle 7.$ $\alpha \leq \beta$

□

Theorem 245. *Every set is grounded.*

PROOF:

- $\langle 1 \rangle 1.$ ASSUME: for a contradiction c is not grounded.
- $\langle 1 \rangle 2.$ LET: B be the transitive closure of $\{c\}$.
- $\langle 1 \rangle 3.$ LET: $A = \{x \in B : x \text{ is not grounded}\}$
- $\langle 1 \rangle 4.$ PICK $m \in A$ such that $m \cap A = \emptyset$
 - PROOF: By the Axiom of Regularity.
- $\langle 1 \rangle 5.$ Every member of m is grounded.
 - PROOF: Every member of m is in B by transitivity but not in A .
- $\langle 1 \rangle 6.$ m is grounded.
 - PROOF: Theorem 244.
- $\langle 1 \rangle 7.$ Q.E.D.
 - PROOF: This contradicts the fact that $m \in A$.

□