

Topology

Robin Adams

May 31, 2022

Contents

1	Set Theory	4
2	Order Theory	6
3	Real Analysis	10
4	Group Theory	11
5	Topological Spaces	12
6	Closed Set	13
7	Interior	14
8	Closure	15
9	Boundary	17
10	Limit Points	17
11	Basis for a Topology	18
12	Local Basis at a Point	22
13	Convergence	23
14	Locally Finite Sets	25
15	Open Maps	26
16	Continuous Functions	26
17	Homeomorphisms	31
18	The Order Topology	32

19 The nth Root Function	35
20 The Product Topology	35
20.1 Continuous in Each Variable Separately	38
21 The Subspace Topology	38
22 The Box Topology	42
23 T_1 Spaces	44
24 Hausdorff Spaces	45
25 The First Countability Axiom	48
26 Strong Continuity	50
27 Saturated Sets	50
28 Quotient Maps	51
29 Quotient Topology	54
30 Retractions	56
31 Homogeneous Spaces	56
32 Regular Spaces	56
33 Connected Spaces	56
34 Totally Disconnected Spaces	65
35 Paths and Path Connectedness	65
36 The Topologist's Sine Curve	68
37 The Long Line	69
38 Components	71
39 Path Components	72
40 Local Connectedness	73
41 Local Path Connectedness	75
42 Weak Local Connectedness	76

43 Quasicomponents	77
44 Open Coverings	78
45 Compact Spaces	78
46 Perfect Maps	85
47 Topological Groups	85
48 The Metric Topology	93
49 Real Linear Algebra	99
50 The Uniform Topology	106
51 Uniform Convergence	109
52 Isometric Imbeddings	110

1 Set Theory

Definition 1 (Cover). Let X be a set and $\mathcal{A} \subseteq \mathcal{P}X$. Then \mathcal{A} *covers* X , or is a *covering* of X , if and only if $\bigcup \mathcal{A} = X$.

Definition 2 (Finite Intersection Property). Let X be a set and $\mathcal{A} \subseteq \mathcal{P}X$. Then \mathcal{A} satisfies the *finite intersection property* if and only if every nonempty finite subset of \mathcal{A} has nonempty intersection.

Lemma 3. *Let X be a set. Let $\mathcal{A} \subseteq \mathcal{P}X$ have the finite intersection property. Then there exists a maximal set \mathcal{D} such that $\mathcal{A} \subseteq \mathcal{D} \subseteq \mathcal{P}X$ and \mathcal{D} has the finite intersection property.*

PROOF:

$\langle 1 \rangle 1$. LET: $\mathbb{F} = \{\mathcal{D} \mid \mathcal{A} \subseteq \mathcal{D} \subseteq \mathcal{P}X, \mathcal{D} \text{ has the finite intersection property}\}$

$\langle 1 \rangle 2$. Every chain in \mathbb{F} has an upper bound.

$\langle 2 \rangle 1$. LET: \mathbb{C} be a chain in \mathbb{F} .

$\langle 2 \rangle 2$. ASSUME: without loss of generality $\mathbb{C} \neq \emptyset$

PROVE: $\bigcup \mathbb{C} \in \mathbb{F}$

PROOF: If $\mathbb{C} = \emptyset$ then \mathcal{A} is an upper bound.

$\langle 2 \rangle 3$. $\mathcal{A} \subseteq \bigcup \mathbb{C} \subseteq \mathcal{P}X$

$\langle 2 \rangle 4$. LET: $C_1, \dots, C_n \in \mathbb{C}$

PROVE: $C_1 \cap \dots \cap C_n \neq \emptyset$

$\langle 2 \rangle 5$. PICK $C_1, \dots, C_n \in \mathbb{C}$ such that $C_i \in \mathbb{C}_i$ for all i .

$\langle 2 \rangle 6$. ASSUME: without loss of generality $C_1 \subseteq \dots \subseteq C_n$

$\langle 2 \rangle 7$. $C_1, \dots, C_n \in \mathbb{C}_n$

$\langle 2 \rangle 8$. \mathbb{C}_n satisfies the finite intersection property.

$\langle 2 \rangle 9$. $C_1 \cap \dots \cap C_n \neq \emptyset$

$\langle 1 \rangle 3$. Q.E.D.

PROOF: By Zorn's Lemma.

□

Lemma 4. *Let X be a set. Let $\mathcal{D} \subseteq \mathcal{P}X$ be maximal with respect to the finite intersection property. Then any finite intersection of elements of \mathcal{D} is an element of \mathcal{D} .*

PROOF:

$\langle 1 \rangle 1$. LET: $D_1, D_2 \in \mathcal{D}$

$\langle 1 \rangle 2$. $\mathcal{D} \cup \{D_1 \cap D_2\}$ has the finite intersection property.

PROOF: Any finite intersection of members of $\mathcal{D} \cup \{D_1 \cap D_2\}$ is a finite intersection of members of \mathcal{D} .

$\langle 1 \rangle 3$. $\mathcal{D} = \mathcal{D} \cup \{D_1 \cap D_2\}$

PROOF: By maximality of \mathcal{D} .

$\langle 1 \rangle 4$. $D_1 \cap D_2 \in \mathcal{D}$.

□

Lemma 5. *Let X be a set. Let $\mathcal{D} \subseteq \mathcal{P}X$ be maximal with respect to the finite intersection property. Let $A \subseteq X$. If A intersects every member of \mathcal{D} then $A \in \mathcal{D}$.*

PROOF:

$\langle 1 \rangle 1.$ $\mathcal{D} \cup \{A\}$ has the finite intersection property.

$\langle 2 \rangle 1.$ LET: $D_1, \dots, D_n \in \mathcal{D}$

PROVE: $D_1 \cap \dots \cap D_n \cap A \neq \emptyset$

$\langle 2 \rangle 2.$ $D_1 \cap \dots \cap D_n \in \mathcal{D}$

PROOF: Lemma 4.

$\langle 2 \rangle 3.$ $D_1 \cap \dots \cap D_n \cap A \neq \emptyset$

PROOF: Since A intersects every member of \mathcal{D} .

$\langle 1 \rangle 2.$ Q.E.D.

PROOF: By maximality of \mathcal{D} .

□

Proposition 6. *Let X be a set. Let $\mathcal{D} \subseteq \mathcal{P}X$ be maximal with respect to the finite intersection property. Let $A, D \in \mathcal{P}X$. If $D \in \mathcal{D}$ and $D \subseteq A$ then $A \in \mathcal{D}$.*

PROOF:

$\langle 1 \rangle 1.$ $\mathcal{D} \cup \{A\}$ satisfies the finite intersection property.

$\langle 2 \rangle 1.$ LET: $D_1, \dots, D_n \in \mathcal{D}$

$\langle 2 \rangle 2.$ $D_1 \cap \dots \cap D_n \cap D \neq \emptyset$

PROOF: Since \mathcal{D} satisfies the finite intersection property.

$\langle 2 \rangle 3.$ $D_1 \cap \dots \cap D_n \cap A \neq \emptyset$

$\langle 1 \rangle 2.$ $\mathcal{D} = \mathcal{D} \cup \{A\}$

PROOF: By the maximality of \mathcal{D} .

$\langle 1 \rangle 3.$ $A \in \mathcal{D}$

□

Definition 7 (Graph). Let $f : A \rightarrow B$. The *graph* of f is the set $\{(x, f(x)) \mid x \in A\} \subseteq A \times B$.

2 Order Theory

Definition 8 (Preorder). Let X be a set. A *preorder* on X is a binary relation \leq on X such that:

Reflexivity For all $x \in X$, we have $x \leq x$

Transitivity For all $x, y, z \in X$, if $x \leq y$ and $y \leq z$ then $x \leq z$.

Definition 9 (Preordered Set). A *preordered set* consists of a set X and a preorder \leq on X .

Proposition 10. Let X and Y be linearly ordered sets. Let $f : X \rightarrow Y$ be strictly monotone and surjective. Then f is a poset isomorphism.

PROOF:

$\langle 1 \rangle 1.$ f is injective.

$\langle 2 \rangle 1.$ LET: $x, y \in X$

$\langle 2 \rangle 2.$ ASSUME: $f(x) = f(y)$

$\langle 2 \rangle 3.$ $x \not\leq y$

PROOF: By strong monotonicity.

$\langle 2 \rangle 4.$ $y \not\leq x$

PROOF: By strong monotonicity.

$\langle 2 \rangle 5.$ $x = y$

PROOF: By trichotomy.

$\langle 1 \rangle 2.$ f^{-1} is monotone.

$\langle 2 \rangle 1.$ LET: $x, y \in X$

$\langle 2 \rangle 2.$ ASSUME: $x \leq y$

$\langle 2 \rangle 3.$ $f^{-1}(x) \not\leq f^{-1}(y)$

PROOF: By strong monotonicity.

$\langle 2 \rangle 4.$ $f^{-1}(x) < f^{-1}(y)$

PROOF: By trichotomy.

□

Definition 11 (Interval). Let X be a preordered set and $Y \subseteq X$. Then Y is an *interval* if and only if, for all $a, b \in Y$ and $c \in X$, if $a \leq c \leq b$ then $c \in Y$.

Definition 12 (Linear Continuum). A linearly ordered set L is a *linear continuum* if and only if:

1. every nonempty subset of L that is bounded above has a supremum
2. L is dense

Proposition 13. Every interval in a linear continuum is a linear continuum.

PROOF:

$\langle 1 \rangle 1.$ LET: L be a linear continuum and I an interval in L .

$\langle 1 \rangle 2.$ Every nonempty subset of I that is bounded above has a supremum in I .

$\langle 2 \rangle 1.$ LET: $X \subseteq I$ be nonempty and bounded above by $b \in I$.

$\langle 2 \rangle 2$. LET: s be the supremum of X in L .
 PROOF: Since L is a linear continuum.
 $\langle 2 \rangle 3$. $s \in I$
 $\langle 3 \rangle 1$. PICK $a \in X$
 PROOF: Since X is nonempty ($\langle 2 \rangle 1$).
 $\langle 3 \rangle 2$. $a \leq s \leq b$
 $\langle 3 \rangle 3$. $s \in I$
 PROOF: Since I is an interval ($\langle 1 \rangle 1$).
 $\langle 2 \rangle 4$. s is the supremum of X in I
 $\langle 1 \rangle 3$. I is dense.
 $\langle 2 \rangle 1$. LET: $x, y \in I$ with $x < y$
 $\langle 2 \rangle 2$. PICK $z \in L$ with $x < z < y$
 PROOF: Since L is dense.
 $\langle 2 \rangle 3$. $z \in I$
 PROOF: Since I is an interval.

□

Definition 14 (Ordered Square). The *ordered square* I_o^2 is the set $[0, 1]^2$ under the dictionary order.

Proposition 15. *The ordered square is a linear continuum.*

PROOF:

$\langle 1 \rangle 1$. Every nonempty subset of I_o^2 bounded above has a supremum.
 $\langle 2 \rangle 1$. LET: $X \subseteq I_o^2$ be nonempty and bounded above by (b, c)
 $\langle 2 \rangle 2$. LET: $s = \sup \pi_1(X)$
 PROOF: The set $\pi_1(X)$ is nonempty and bounded above by b .
 $\langle 2 \rangle 3$. CASE: $s \in \pi_1(X)$
 $\langle 3 \rangle 1$. LET: $t = \sup \{y \in [0, 1] \mid (s, y) \in X\}$
 PROOF: This set is nonempty and bounded above by c .
 $\langle 3 \rangle 2$. (s, t) is the supremum of X .
 $\langle 2 \rangle 4$. CASE: $s \notin \pi_1(X)$
 PROOF: In this case $(s, 0)$ is the supremum of X .
 $\langle 1 \rangle 2$. I_o^2 is dense.
 $\langle 2 \rangle 1$. LET: $(x_1, y_1), (x_2, y_2) \in I_o^2$ with $(x_1, y_1) < (x_2, y_2)$
 $\langle 2 \rangle 2$. CASE: $x_1 < x_2$
 $\langle 3 \rangle 1$. PICK x_3 with $x_1 < x_3 < x_2$
 $\langle 3 \rangle 2$. $(x_1, y_1) < (x_3, y_1) < (x_2, y_2)$
 $\langle 2 \rangle 3$. CASE: $x_1 = x_2$ and $y_1 < y_2$
 $\langle 3 \rangle 1$. PICK y_3 with $y_1 < y_3 < y_2$
 $\langle 3 \rangle 2$. $(x_1, y_1) < (x_1, y_3) < (x_2, y_2)$

□

Proposition 16. *If X is a well-ordered set then $X \times [0, 1)$ under the dictionary order is a linear continuum.*

PROOF:

$\langle 1 \rangle 1$. Every nonempty set $A \subseteq X \times [0, 1)$ bounded above has a supremum.

- ⟨2⟩1. LET: $A \subseteq X \times [0, 1)$ be nonempty and bounded above
- ⟨2⟩2. LET: x_0 be the supremum of $\pi_1(A)$
- ⟨2⟩3. CASE: $x_0 \in \pi_1(A)$
 - ⟨3⟩1. LET: y_0 be the supremum of $\{y \in [0, 1) \mid (x_0, y) \in A\}$
 - ⟨3⟩2. (x_0, y_0) is the supremum of A .
- ⟨2⟩4. CASE: $x_0 \notin \pi_1(A)$
 - PROOF: In this case $(x_0, 0)$ is the supremum of A .
- ⟨1⟩2. $X \times [0, 1)$ is dense.
 - ⟨2⟩1. LET: $(x_1, y_1), (x_2, y_2) \in X \times [0, 1)$ with $(x_1, y_1) < (x_2, y_2)$
 - ⟨2⟩2. CASE: $x_1 < x_2$
 - ⟨3⟩1. PICK y_3 such that $y_1 < y_3 < 1$
 - ⟨3⟩2. $(x_1, y_1) < (x_1, y_3) < (x_2, y_2)$
 - ⟨2⟩3. CASE: $x_1 = x_2$ and $y_1 < y_2$
 - ⟨3⟩1. PICK y_3 such that $y_1 < y_3 < y_2$
 - ⟨3⟩2. $(x_1, y_1) < (x_1, y_3) < (x_2, y_2)$

□

Lemma 17. For all $a, b, c, d \in \mathbb{R}$ with $a < b$ and $c < d$, we have $[a, b) \cong [c, d)$

PROOF: The map $\lambda t.c + (t - a)(d - c)/(b - a)$ is an order isomorphism.

Proposition 18. Let X be a linearly ordered set. Let $a < b < c$ in X . Then $[a, c) \cong [0, 1)$ if and only if $[a, b) \cong [b, c) \cong [0, 1)$.

PROOF:

- ⟨1⟩1. If $[a, c) \cong [0, 1)$ then $[a, b) \cong [b, c) \cong [0, 1)$
- ⟨2⟩1. ASSUME: $f : [a, c) \cong [0, 1)$ is an order isomorphism
- ⟨2⟩2. $[a, b) \cong [0, 1)$
 - PROOF:
$$\begin{aligned} [a, b) &\cong [0, f(b)) && \text{(by the restriction of } f) \\ &\cong [0, 1) && \text{(Lemma 17)} \end{aligned}$$
- ⟨2⟩3. $[b, c) \cong [0, 1)$
 - PROOF: Similar.
- ⟨1⟩2. If $[a, b) \cong [b, c) \cong [0, 1)$ then $[a, c) \cong [0, 1)$
 - PROOF:
$$\begin{aligned} [a, c) &= [a, b) * [b, c) \\ &\cong [0, 1) * [0, 1) \\ &\cong [0, 1/2) * [1/2, 1) && \text{(Lemma 17)} \\ &= 1 \end{aligned}$$

□

Proposition 19 (CC). Let X be a linearly ordered set. Let $x_0 < x_1 < \dots$ be a strictly increasing sequence in X with supremum b . Then $[x_0, b) \cong [0, 1)$ if and only if $[x_i, x_{i+1}) \cong [0, 1)$ for all i .

PROOF:

- ⟨1⟩1. If $[x_0, b) \cong [0, 1)$ then $[x_i, x_{i+1}) \cong [0, 1)$ for all i .

PROOF: By Lemma 17

$\langle 1 \rangle 2$. If $[x_i, x_{i+1}) \cong [0, 1)$ for all i then $[x_0, b) \cong [0, 1)$

$\langle 2 \rangle 1$. ASSUME: $[x_i, x_{i+1}) \cong [0, 1)$ for all i

$\langle 2 \rangle 2$. PICK an order isomorphism $f_i : [x_i, x_{i+1}) \cong [1/2^i, 2/2^{i+1})$ for each i .

PROOF: By Lemma 17

$\langle 2 \rangle 3$. The union of the f_i s is an order isomorphism $[x_0, b) \cong [0, 1)$

□

3 Real Analysis

Definition 20. Let \mathbb{R}^∞ be the subset of \mathbb{R}^ω consisting of all sequences (a_n) such that $a_n = 0$ for all but finitely many n .

4 Group Theory

Definition 21. Given a group G and sets $A, B \subseteq G$, let $AB = \{ab \mid a \in A, b \in B\}$.

Definition 22. Given a group G and a set $A \subseteq G$, let $A^{-1} = \{a^{-1} \mid a \in A\}$.

5 Topological Spaces

Definition 23 (Topology). A *topology* on a set X is a set $\mathcal{T} \subseteq \mathcal{P}X$ such that:

- $X \in \mathcal{T}$.
- For all $\mathcal{U} \subseteq \mathcal{T}$ we have $\bigcup \mathcal{U} \in \mathcal{T}$.
- For all $U, V \in \mathcal{T}$ we have $U \cap V \in \mathcal{T}$.

We call the elements of X *points* and the elements of \mathcal{T} *open sets*.

Definition 24 (Topological Space). A *topological space* X consists of a set X and a topology on X .

Definition 25 (Discrete Space). For any set X , the *discrete* topology on X is $\mathcal{P}X$.

Definition 26 (Indiscrete Space). For any set X , the *indiscrete* or *trivial* topology on X is $\{\emptyset, X\}$.

Definition 27 (Finite Complement Topology). For any set X , the *finite complement topology* on X is $\{U \in \mathcal{P}X \mid X \setminus U \text{ is finite}\} \cup \{\emptyset\}$.

Definition 28 (Countable Complement Topology). For any set X , the *countable complement topology* on X is $\{U \in \mathcal{P}X \mid X \setminus U \text{ is countable}\} \cup \{\emptyset\}$.

Definition 29 (Finer, Coarser). Suppose that \mathcal{T} and \mathcal{T}' are two topologies on a given set X . If $\mathcal{T}' \supseteq \mathcal{T}$, we say that \mathcal{T}' is *finer* than \mathcal{T} ; if \mathcal{T}' *properly* contains \mathcal{T} , we say that \mathcal{T}' is *strictly finer* than \mathcal{T} . We also say that \mathcal{T} is *coarser* than \mathcal{T}' , or *strictly coarser*, in these two respective situations. We say \mathcal{T} is *comparable* with \mathcal{T}' if either $\mathcal{T}' \supseteq \mathcal{T}$ or $\mathcal{T} \supseteq \mathcal{T}'$.

Lemma 30. *Let X be a topological space and $U \subseteq X$. Then U is open if and only if, for all $x \in U$, there exists an open set V such that $x \in V \subseteq U$.*

PROOF:

$\langle 1 \rangle 1. \Rightarrow$

PROOF: Take $V = U$

$\langle 1 \rangle 2. \Leftarrow$

PROOF: We have $U = \bigcup \{V \text{ open in } X \mid V \subseteq U\}$.

□

Lemma 31. *Let X be a set and \mathcal{T} a nonempty set of topologies on X . Then $\bigcap \mathcal{T}$ is a topology on X , and is the finest topology that is coarser than every member of \mathcal{T} .*

PROOF:

$\langle 1 \rangle 1. X \in \bigcap \mathcal{T}$

PROOF: Since X is in every member of \mathcal{T} .

$\langle 1 \rangle 2. \bigcap \mathcal{T}$ is closed under union.

- ⟨2⟩1. LET: $\mathcal{U} \subseteq \bigcap \mathcal{T}$
- ⟨2⟩2. For all $T \in \mathcal{T}$ we have $\mathcal{U} \subseteq T$
- ⟨2⟩3. For all $T \in \mathcal{T}$ we have $\bigcup \mathcal{U} \in T$
- ⟨2⟩4. $\bigcup \mathcal{U} \in \bigcap \mathcal{T}$
- ⟨1⟩3. $\bigcap \mathcal{T}$ is closed under binary intersection.
- ⟨2⟩1. LET: $U, V \in \bigcap \mathcal{T}$
- ⟨2⟩2. For all $T \in \mathcal{T}$ we have $U, V \in T$
- ⟨2⟩3. For all $T \in \mathcal{T}$ we have $U \cap V \in T$
- ⟨2⟩4. $U \cap V \in \bigcap \mathcal{T}$

□

Lemma 32. *Let X be a set and \mathcal{T} a set of topologies on X . Then there exists a unique coarsest topology that is finer than every member of \mathcal{T} .*

PROOF: The required topology is given by

$$\bigcap \{T \in \mathcal{P}X \mid T \text{ is a topology on } X \text{ that is finer than every member of } \mathcal{T}\},$$

The set is nonempty since it contains the discrete topology. □

Definition 33 (Neighbourhood). A *neighbourhood* of a point x is an open set that contains x .

6 Closed Set

Definition 34 (Closed Set). Let X be a topological space and $A \subseteq X$. Then A is *closed* if and only if $X \setminus A$ is open.

Lemma 35. *The empty set is closed.*

PROOF: Since the whole space X is always open. □

Lemma 36. *The topological space X is closed.*

PROOF: Since \emptyset is open. □

Lemma 37. *The intersection of a nonempty set of closed sets is closed.*

PROOF: Let \mathcal{C} be a nonempty set of closed sets. Then $X \setminus \bigcap \mathcal{C} = \bigcup \{X \setminus C \mid C \in \mathcal{C}\}$ is open. □

Lemma 38. *The union of two closed sets is closed.*

PROOF: Let C and D be closed. Then $X \setminus (C \cup D) = (X \setminus C) \cap (X \setminus D)$ is open. □

Proposition 39. *Let X be a set and $\mathcal{C} \subseteq \mathcal{P}X$ a set such that:*

1. $\emptyset \in \mathcal{C}$
2. $X \in \mathcal{C}$
3. For all $\mathcal{A} \subseteq \mathcal{C}$ nonempty we have $\bigcap \mathcal{A} \in \mathcal{C}$

4. For all $C, D \in \mathcal{C}$ we have $C \cup D \in \mathcal{C}$.

Then there exists a unique topology \mathcal{T} such that \mathcal{C} is the set of closed sets, namely

$$\mathcal{T} = \{X \setminus C \mid C \in \mathcal{C}\}$$

PROOF:

$\langle 1 \rangle 1$. LET: $\mathcal{T} = \{X \setminus C \mid C \in \mathcal{C}\}$

$\langle 1 \rangle 2$. \mathcal{T} is a topology

$\langle 2 \rangle 1$. $X \in \mathcal{T}$

PROOF: Since $\emptyset \in \mathcal{C}$

$\langle 2 \rangle 2$. For all $\mathcal{U} \subseteq \mathcal{T}$ we have $\bigcup \mathcal{U} \in \mathcal{T}$

$\langle 3 \rangle 1$. LET: $\mathcal{U} \subseteq \mathcal{T}$

$\langle 3 \rangle 2$. CASE: $\mathcal{U} = \emptyset$

PROOF: In this case $\bigcup \mathcal{U} = \emptyset \in \mathcal{T}$ since $X \in \mathcal{C}$

$\langle 3 \rangle 3$. CASE: $\mathcal{U} \neq \emptyset$

PROOF: In this case $X \setminus \bigcup \mathcal{U} = \bigcap \{X \setminus U \mid U \in \mathcal{U}\} \in \mathcal{C}$.

$\langle 2 \rangle 3$. For all $U, V \in \mathcal{T}$ we have $U \cap V \in \mathcal{T}$

PROOF: Since $X \setminus (U \cap V) = (X \setminus U) \cup (X \setminus V) \in \mathcal{C}$.

$\langle 1 \rangle 3$. \mathcal{C} is the set of all closed sets in \mathcal{T}

PROOF:

C is closed in \mathcal{T}

$$\Leftrightarrow X \setminus C \in \mathcal{T}$$

$$\Leftrightarrow C \in \mathcal{C}$$

$\langle 1 \rangle 4$. If \mathcal{T}' is a topology and \mathcal{C} is the set of closed sets in \mathcal{T}' then $\mathcal{T}' = \mathcal{T}$

PROOF: We have

$$U \in \mathcal{T}$$

$$\Leftrightarrow X \setminus U \in \mathcal{C}$$

$$\Leftrightarrow X \setminus U \text{ is closed in } \mathcal{T}'$$

$$\Leftrightarrow U \in \mathcal{T}'$$

□

Proposition 40. If U is open and A is closed then $U \setminus A$ is open.

PROOF: $U \setminus A = U \cap (X \setminus A)$ is the intersection of two open sets. □

Proposition 41. If U is open and A is closed then $A \setminus U$ is closed.

PROOF: $A \setminus U = A \cap (X \setminus U)$ is the intersection of two closed sets. □

7 Interior

Definition 42 (Interior). Let X be a topological space and $A \subseteq X$. The *interior* of A , $\text{Int } A$, is the union of all the open subsets of A .

Lemma 43. The interior of a set is open.

PROOF: It is a union of open sets. \square

Lemma 44.

$$\text{Int } A \subseteq A$$

PROOF: Immediate from definition. \square

Lemma 45. *If U is open and $U \subseteq A$ then $U \subseteq \text{Int } A$*

PROOF: Immediate from definition. \square

Lemma 46. *A set A is open if and only if $A = \text{Int } A$.*

PROOF: If $A = \text{Int } A$ then A is open by Lemma 43. Conversely if A is open then $A \subseteq \text{Int } A$ by the definition of interior and so $A = \text{Int } A$.

8 Closure

Definition 47 (Closure). Let X be a topological space and $A \subseteq X$. The *closure* of A , \overline{A} , is the intersection of all the closed sets that include A .

This intersection exists since X is a closed set that includes A (Lemma 36).

Lemma 48. *The closure of a set is closed.*

PROOF: Dual to Lemma 43. \square

Lemma 49.

$$A \subseteq \overline{A}$$

PROOF: Immediate from definition. \square

Lemma 50. *If C is closed and $A \subseteq C$ then $\overline{A} \subseteq C$.*

PROOF: Immediate from definition. \square

Lemma 51. *A set A is closed if and only if $A = \overline{A}$.*

PROOF: Dual to Lemma 46. \square

Theorem 52. *Let X be a topological space, $A \subseteq X$ and $x \in X$. Then $x \in \overline{A}$ if and only if every neighbourhood of x intersects A .*

PROOF: We have

$$x \in \overline{A}$$

$$\Leftrightarrow \forall C. C \text{ closed} \wedge A \subseteq C \Rightarrow x \in C$$

$$\Leftrightarrow \forall U. U \text{ open} \wedge A \cap U = \emptyset \Rightarrow x \notin U$$

$$\Leftrightarrow \forall U. U \text{ open} \wedge x \in U \Rightarrow U \text{ intersects } A$$

\square

Proposition 53. *If $A \subseteq B$ then $\overline{A} \subseteq \overline{B}$.*

PROOF: This holds because \overline{B} is a closed set that includes A . \square

Proposition 54.

$$\overline{A \cup B} = \overline{A} \cup \overline{B}$$

PROOF:

$\langle 1 \rangle 1. \overline{A} \subseteq \overline{A \cup B}$

PROOF: By Proposition 53.

$\langle 1 \rangle 2. \overline{B} \subseteq \overline{A \cup B}$

PROOF: By Proposition 53.

$\langle 1 \rangle 3. \overline{A \cup B} \subseteq \overline{A \cup B}$

$\langle 2 \rangle 1. \text{ LET: } x \in \overline{A \cup B}$

$\langle 2 \rangle 2. \text{ ASSUME: } x \notin \overline{A}$

PROVE: $x \in \overline{B}$

$\langle 2 \rangle 3. \text{ PICK a neighbourhood } U \text{ of } x \text{ that does not intersect } A$

$\langle 2 \rangle 4. \text{ LET: } V \text{ be any neighbourhood of } x$

$\langle 2 \rangle 5. U \cap V \text{ is a neighbourhood of } x$

$\langle 2 \rangle 6. U \cap V \text{ intersects } A \cup B$

PROOF: From $\langle 2 \rangle 1$ and Theorem 52.

$\langle 2 \rangle 7. U \cap V \text{ intersects } B$

PROOF: From $\langle 2 \rangle 3$

$\langle 2 \rangle 8. V \text{ intersects } B$

$\langle 2 \rangle 9. \text{ Q.E.D.}$

PROOF: We have $x \in \overline{B}$ from Theorem 52.

\square

Proposition 55. *Let X be a topological space. Let \mathcal{D} be a set of subsets of X that is maximal with respect to the finite intersection property. Let $x \in X$. Then the following are equivalent:*

1. *For all $D \in \mathcal{D}$ we have $x \in \overline{D}$*
2. *Every neighbourhood of x is in \mathcal{D} .*

PROOF:

$\langle 1 \rangle 1. 1 \Rightarrow 2$

$\langle 2 \rangle 1. \text{ For all } D \in \mathcal{D} \text{ we have } x \in \overline{D}$

$\langle 2 \rangle 2. \text{ LET: } U \text{ be a neighbourhood of } x$

$\langle 2 \rangle 3. \mathcal{D} \cup \{U\} \text{ satisfies the finite intersection property.}$

$\langle 3 \rangle 1. \text{ LET: } D_1, \dots, D_n \in \mathcal{D}$

$\langle 3 \rangle 2. D_1 \cap \dots \cap D_n \in \mathcal{D}$

PROOF: Lemma 4.

$\langle 3 \rangle 3. x \in \overline{D_1 \cap \dots \cap D_n}$

PROOF: $\langle 2 \rangle 1, \langle 3 \rangle 2$

$\langle 3 \rangle 4. D_1 \cap \dots \cap D_n \cap U \neq \emptyset$

PROOF: Theorem 52, $\langle 2 \rangle 2, \langle 3 \rangle 3$.

$\langle 2 \rangle 4. \mathcal{D} = \mathcal{D} \cup \{U\}$

PROOF: By the maximality of \mathcal{D} .

$\langle 2 \rangle 5. U \in \mathcal{D}$
 $\langle 1 \rangle 2. 2 \Rightarrow 1$
 $\langle 2 \rangle 1. \text{ ASSUME: Every neighbourhood of } x \text{ is in } \mathcal{D}.$
 $\langle 2 \rangle 2. \text{ LET: } D \in \mathcal{D}$
 $\langle 2 \rangle 3. \text{ Every neighbourhood of } x \text{ intersects } D.$
 PROOF: From $\langle 2 \rangle 1$, $\langle 2 \rangle 2$ and the fact that \mathcal{D} satisfies the finite intersection property.
 $\langle 2 \rangle 4. x \in \overline{D}$
 PROOF: Theorem 52, $\langle 2 \rangle 3$.
 \square

9 Boundary

Definition 56 (Boundary). The *boundary* of a set A is the set $\partial A = \overline{A} \cap \overline{X \setminus A}$.

Proposition 57.

$$\text{Int } A \cap \partial A = \emptyset$$

PROOF: Since $\overline{X \setminus A} = X \setminus \text{Int } A$. \square

Proposition 58.

$$\overline{A} = \text{Int } A \cup \partial A$$

PROOF:

$$\begin{aligned}
 \text{Int } A \cup \partial A &= \text{Int } A \cup (\overline{A} \cap \overline{X \setminus A}) \\
 &= (\text{Int } A \cup \overline{A}) \cap (\text{Int } A \cup \overline{X \setminus A}) \\
 &= \overline{A} \cap X \\
 &= \overline{A}
 \end{aligned}$$

Proposition 59. $\partial A = \emptyset$ if and only if A is open and closed.

PROOF: If $\partial A = \emptyset$ then $\overline{A} = \text{Int } A$ by Proposition 58.

Proposition 60. A set U is open if and only if $\partial U = \overline{U} \setminus U$.

PROOF:

$$\begin{aligned}
 \partial U &= \overline{U} \setminus U \\
 \Leftrightarrow \overline{U} \setminus \text{Int } U &= \overline{U} \setminus U && (\text{Propositions 57, 58}) \\
 \Leftrightarrow \text{Int } U &= U && \square
 \end{aligned}$$

10 Limit Points

Definition 61 (Limit Point). Let X be a topological space, $a \in X$ and $A \subseteq X$. Then a is a *limit point*, *cluster point* or *point of accumulation* for A if and only if every neighbourhood of a intersects A at a point other than a .

Lemma 62. *The point a is an accumulation point for A if and only if $a \in \overline{A} \setminus \{a\}$.*

PROOF: From Theorem 52. \square

Theorem 63. *Let X be a topological space and $A \subseteq X$. Let A' be the set of all limit points of A . Then $\overline{A} = A \cup A'$.*

PROOF:

$\langle 1 \rangle 1.$ For all $x \in \overline{A}$, if $x \notin A$ then $x \in A'$

PROOF: From Theorem 52.

$\langle 1 \rangle 2.$ $A \subseteq \overline{A}$

PROOF: Lemma 49.

$\langle 1 \rangle 3.$ $A' \subseteq \overline{A}$

PROOF: From Theorem 52.

\square

Corollary 63.1. *A set is closed if and only if it contains all its limit points.*

Proposition 64. *In an indiscrete topology, every point is a limit point of any set that has more than one point.*

PROOF: Let X be an indiscrete space. Let A be a set with more than one point and x be a point. The only neighbourhood of x is X , which must intersect A at a point other than x . \square

Lemma 65. *Let X be a topological space and $A \subseteq B \subseteq X$. Then every limit point of A is a limit point of B .*

PROOF: Immediate from definitions. \square

11 Basis for a Topology

Definition 66 (Basis). If X is a set, a *basis* for a topology on X is a set $\mathcal{B} \subseteq \mathcal{P}X$ called *basis elements* such that

1. For all $x \in X$, there exists $B \in \mathcal{B}$ such that $x \in B$.
2. For all $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$.

If \mathcal{B} satisfies these two conditions, then we define the topology *generated* by \mathcal{B} to be $\mathcal{T} = \{U \in \mathcal{P}X \mid \forall x \in U. \exists B \in \mathcal{B}. x \in B \subseteq U\}$.

We prove this is a topology.

PROOF:

$\langle 1 \rangle 1.$ $X \in \mathcal{T}$

PROOF: For all $x \in X$ there exists $B \in \mathcal{B}$ such that $x \in B \subseteq X$ by condition 1.

- ⟨1⟩2. For all $\mathcal{U} \subseteq \mathcal{T}$ we have $\bigcup \mathcal{U} \in \mathcal{T}$
 - ⟨2⟩1. LET: $\mathcal{U} \subseteq \mathcal{T}$
 - ⟨2⟩2. LET: $x \in \bigcup \mathcal{U}$
 - ⟨2⟩3. PICK $U \in \mathcal{U}$ such that $x \in U$
 - ⟨2⟩4. PICK $B \in \mathcal{B}$ such that $x \in B \subseteq U$
 - PROOF: Since $U \in \mathcal{T}$ by ⟨2⟩1 and ⟨2⟩3.
 - ⟨2⟩5. $x \in B \subseteq \bigcup \mathcal{U}$
- ⟨1⟩3. For all $U, V \in \mathcal{T}$ we have $U \cap V \in \mathcal{T}$
 - ⟨2⟩1. LET: $U, V \in \mathcal{T}$
 - ⟨2⟩2. LET: $x \in U \cap V$
 - ⟨2⟩3. PICK $B_1 \in \mathcal{B}$ such that $x \in B_1 \subseteq U$
 - ⟨2⟩4. PICK $B_2 \in \mathcal{B}$ such that $x \in B_2 \subseteq V$
 - ⟨2⟩5. PICK $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$
 - PROOF: By condition 2.
 - ⟨2⟩6. $x \in B_3 \subseteq U \cap V$

□

Lemma 67. *Let X be a set. Let \mathcal{B} be a basis for a topology \mathcal{T} on X . Then \mathcal{T} is the set of all unions of subsets of \mathcal{B} .*

PROOF:

- ⟨1⟩1. For all $U \in \mathcal{T}$, there exists $\mathcal{A} \subseteq \mathcal{B}$ such that $U = \bigcup \mathcal{A}$
 - ⟨2⟩1. LET: $U \in \mathcal{T}$
 - ⟨2⟩2. LET: $\mathcal{A} = \{B \in \mathcal{B} \mid B \subseteq U\}$
 - ⟨2⟩3. $U \subseteq \bigcup \mathcal{A}$
 - ⟨3⟩1. LET: $x \in U$
 - ⟨3⟩2. PICK $B \in \mathcal{B}$ such that $x \in B \subseteq U$
 - PROOF: Since \mathcal{B} is a basis for \mathcal{T} .
 - ⟨3⟩3. $x \in B \in \mathcal{A}$
 - ⟨2⟩4. $\bigcup \mathcal{A} \subseteq U$
 - PROOF: From the definition of \mathcal{A} (⟨2⟩2).
- ⟨1⟩2. For all $\mathcal{A} \subseteq \mathcal{B}$ we have $\bigcup \mathcal{A} \in \mathcal{T}$
 - ⟨2⟩1. $\mathcal{B} \subseteq \mathcal{T}$
 - PROOF: If $B \in \mathcal{B}$ and $x \in B$, then there exists $B' \in \mathcal{B}$ such that $x \in B' \subseteq B$, namely $B' = B$.
 - ⟨2⟩2. Q.E.D.
 - PROOF: Since \mathcal{T} is closed under union.

□

Corollary 67.1. *Let X be a set. Let \mathcal{B} be a basis for a topology \mathcal{T} on X . Then \mathcal{T} is the coarsest topology that includes \mathcal{B} .*

PROOF: Since every topology that includes \mathcal{B} includes all unions of subsets of \mathcal{B} . □

Lemma 68. *Let X be a topological space. Suppose that \mathcal{C} is a set of open sets such that, for every open set U and every point $x \in U$, there exists $C \in \mathcal{C}$ such that $x \in C \subseteq U$. Then \mathcal{C} is a basis for the topology on X .*

PROOF:

$\langle 1 \rangle 1$. For all $x \in X$, there exists $C \in \mathcal{C}$ such that $x \in C$

PROOF: Immediate from hypothesis.

$\langle 1 \rangle 2$. For all $C_1, C_2 \in \mathcal{C}$ and $x \in C_1 \cap C_2$, there exists $C_3 \in \mathcal{C}$ such that $x \in C_3 \subseteq C_1 \cap C_2$

PROOF: Since $C_1 \cap C_2$ is open.

$\langle 1 \rangle 3$. Every open set is open in the topology generated by \mathcal{C}

PROOF: Immediate from hypothesis.

$\langle 1 \rangle 4$. Every union of a subset of \mathcal{C} is open.

PROOF: Since every member of \mathcal{C} is open.

□

Lemma 69. *Let \mathcal{B} and \mathcal{B}' be bases for the topologies \mathcal{T} and \mathcal{T}' respectively on the set X . Then the following are equivalent.*

1. $\mathcal{T} \subseteq \mathcal{T}'$

2. For all $B \in \mathcal{B}$ and $x \in B$, there exists $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$.

PROOF:

$\langle 1 \rangle 1$. $1 \Rightarrow 2$

$\langle 2 \rangle 1$. ASSUME: $\mathcal{T} \subseteq \mathcal{T}'$

$\langle 2 \rangle 2$. LET: $B \in \mathcal{B}$ and $x \in B$

$\langle 2 \rangle 3$. $B \in \mathcal{T}$

PROOF: Corollary 67.1.

$\langle 2 \rangle 4$. $B \in \mathcal{T}'$

PROOF: By $\langle 2 \rangle 1$

$\langle 2 \rangle 5$. There exists $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$

PROOF: Since \mathcal{B}' is a basis for \mathcal{T}' .

$\langle 1 \rangle 2$. $2 \Rightarrow 1$

$\langle 2 \rangle 1$. ASSUME: 2

$\langle 2 \rangle 2$. LET: $U \in \mathcal{T}$

PROVE: $U \in \mathcal{T}'$

$\langle 2 \rangle 3$. LET: $x \in U$

PROVE: There exists $B' \in \mathcal{B}'$ such that $x \in B' \subseteq U$

$\langle 2 \rangle 4$. PICK $B \in \mathcal{B}$ such that $x \in B \subseteq U$

PROOF: Since \mathcal{B} is a basis for \mathcal{T} .

$\langle 2 \rangle 5$. PICK $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$

PROOF: By $\langle 2 \rangle 1$.

$\langle 2 \rangle 6$. $x \in B' \subseteq U$

□

Theorem 70. *Let X be a topological space, $A \subseteq X$ and $x \in X$. Let \mathcal{B} be a basis for X . Then $x \in \overline{A}$ if and only if, for all $B \in \mathcal{B}$, if $x \in B$ then B intersects A .*

PROOF:

$\langle 1 \rangle 1$. If $x \in \overline{A}$ then, for all $B \in \mathcal{B}$, if $x \in B$ then B intersects A .

PROOF: This follows from Theorem 52 since every element of \mathcal{B} is open (Corollary 67.1).

- ⟨1⟩2. Suppose that, for all $B \in \mathcal{B}$, if $x \in B$ then B intersects A . Then $x \in \overline{A}$.
 ⟨2⟩1. ASSUME: For all $B \in \mathcal{B}$, if $x \in B$ then B intersects A .
 ⟨2⟩2. LET: U be an open set that contains x
 PROVE: U intersects A .
 ⟨2⟩3. PICK $B \in \mathcal{B}$ such that $x \in B \subseteq U$.
 ⟨2⟩4. B intersects A .
 PROOF: From ⟨2⟩1.
 ⟨2⟩5. U intersects A .
 ⟨2⟩6. Q.E.D.
 PROOF: By Theorem 52.

□

Definition 71 (Lower Limit Topology on the Real Line). The *lower limit topology on the real line* is the topology on \mathbb{R} generated by the basis consisting of all half-open intervals of the form $[a, b)$.

We write \mathbb{R}_l for the topological space \mathbb{R} under the lower limit topology.

We prove this is a basis for a topology.

PROOF:

- ⟨1⟩1. For all $x \in \mathbb{R}$ there exists an interval $[a, b)$ such that $x \in [a, b)$.
 PROOF: Take $[a, b) = [x, x + 1)$.
 ⟨1⟩2. For any open intervals $[a, b)$, $[c, d)$ if $x \in [a, b) \cap [c, d)$, then there exists an interval $[e, f)$ such that $x \in [e, f) \subseteq [a, b) \cap [c, d)$
 PROOF: Take $[e, f) = [\max(a, c), \min(b, d))$.

□

Definition 72 (K -topology on the Real Line). Let $K = \{1/n \mid n \in \mathbb{Z}^+\}$.

The *K -topology on the real line* is the topology on \mathbb{R} generated by the basis consisting of all open intervals (a, b) and all sets of the form $(a, b) \setminus K$.

We write \mathbb{R}_K for the topological space \mathbb{R} under the K -topology.

We prove this is a basis for a topology.

PROOF:

- ⟨1⟩1. For all $x \in \mathbb{R}$ there exists an open interval (a, b) such that $x \in (a, b)$.
 PROOF: Take $(a, b) = (x - 1, x + 1)$.
 ⟨1⟩2. For any basic open sets B_1, B_2 if $x \in B_1 \cap B_2$, then there exists a basic open set B_3 such that $x \in B_3 \subseteq B_1 \cap B_2$
 ⟨2⟩1. CASE: $B_1 = (a, b)$, $B_2 = (c, d)$
 PROOF: Take $B_3 = (\max(a, c), \min(b, d))$.
 ⟨2⟩2. CASE: $B_1 = (a, b)$ or $(a, b) \setminus K$, $B_2 = (c, d)$ or $(c, d) \setminus K$, and they are not both open intervals.
 PROOF: Take $B_3 = (\max(a, c), \min(b, d)) \setminus K$.

□

Lemma 73. The lower limit topology and the K -topology are incomparable.

PROOF:

⟨1⟩1. The interval $[10, 11)$ is not open in the K -topology.

PROOF: There is no open interval (a, b) such that $10 \in (a, b) \subseteq [10, 11)$ or $10 \in (a, b) \setminus K \subseteq [10, 11)$.

⟨1⟩2. The set $(-1, 1) \setminus K$ is not open in the lower limit topology.

PROOF: There is no half-open interval $[a, b)$ such that $0 \in [a, b) \subseteq (-1, 1) \setminus K$, since there must be a positive integer n with $1/n \in [a, b)$.

□

Definition 74 (Subbasis). A *subbasis* \mathcal{S} for a topology on X is a set $\mathcal{S} \subseteq \mathcal{P}X$ such that $\bigcup \mathcal{S} = X$.

The topology *generated* by the subbasis \mathcal{S} is the set of all unions of finite intersections of elements of \mathcal{S} .

We prove this is a topology.

PROOF:

⟨1⟩1. The set \mathcal{B} of all finite intersections of elements of \mathcal{S} forms a basis for a topology on X .

⟨2⟩1. $\bigcup \mathcal{B} = X$

PROOF: Since $\mathcal{S} \subseteq \mathcal{B}$.

⟨2⟩2. \mathcal{B} is closed under binary intersection.

PROOF: By definition.

⟨1⟩2. Q.E.D.

PROOF: By Lemma 67.

□

We have simultaneously proved:

Proposition 75. *Let \mathcal{S} be a subbasis for the topology on X . Then the set of all finite intersections of elements of \mathcal{S} is a basis for the topology on X .*

Proposition 76. *Let X be a set. Let \mathcal{S} be a subbasis for a topology \mathcal{T} on X . Then \mathcal{T} is the coarsest topology that includes \mathcal{S} .*

PROOF: Since every topology that includes \mathcal{S} includes every union of finite intersections of elements of \mathcal{S} . □

12 Local Basis at a Point

Definition 77 (Local Basis). Let X be a topological space and $a \in X$. A (*local*) *basis at a* is a set \mathcal{B} of neighbourhoods of a such that every neighbourhood of a includes some member of \mathcal{B} .

Lemma 78. *If there exists a countable local basis at a point a , then there exists a countable local basis $\{B_n \mid n \geq 1\}$ such that $B_1 \supseteq B_2 \supseteq \cdots$.*

PROOF: Given a countable local basis $\{C_n \mid n \geq 1\}$, take $B_n = C_1 \cap \cdots \cap C_n$. □

13 Convergence

Definition 79 (Convergence). Let X be a topological space. Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of points in X and $l \in X$. Then the sequence $(a_n)_{n \in \mathbb{N}}$ *converges* to the *limit* l , $a_n \rightarrow l$ as $n \rightarrow \infty$, if and only if, for every neighbourhood U of l , there exists N such that, for all $n \geq N$, we have $a_n \in U$.

Lemma 80. *Let X be a topological space. Let $A \subseteq X$ and $l \in X$. If there is a sequence of points in A that converges to l then $l \in \bar{A}$.*

PROOF:

- $\langle 1 \rangle 1$. LET: (a_n) be a sequence of points in A that converges to l .
- $\langle 1 \rangle 2$. LET: U be a neighbourhood of l .
- $\langle 1 \rangle 3$. PICK N such that, for all $n \geq N$, we have $a_n \in U$.
- $\langle 1 \rangle 4$. $a_N \in U \cap A$
- $\langle 1 \rangle 5$. Q.E.D.

PROOF: Theorem 52.

□

Proposition 81. *Let X be a topological space. Let \mathcal{B} be a basis for the topology on X . Let (a_n) be a sequence in X and $l \in X$. Then $a_n \rightarrow l$ as $n \rightarrow \infty$ if and only if, for every $B \in \mathcal{B}$ with $l \in B$, there exists N such that, for all $n \geq N$, we have $a_n \in B$.*

PROOF:

- $\langle 1 \rangle 1$. If $a_n \rightarrow l$ as $n \rightarrow \infty$ then, for every $B \in \mathcal{B}$ with $l \in B$, there exists N such that, for all $n \geq N$, we have $a_n \in B$.

PROOF: Since every element of \mathcal{B} is open (Corollary 67.1).

- $\langle 1 \rangle 2$. If, for every $B \in \mathcal{B}$ with $l \in B$, there exists N such that, for all $n \geq N$, we have $a_n \in B$, then $a_n \rightarrow l$ as $n \rightarrow \infty$.
 - $\langle 2 \rangle 1$. ASSUME: for every $B \in \mathcal{B}$ with $l \in B$, there exists N such that, for all $n \geq N$, we have $a_n \in B$
 - $\langle 2 \rangle 2$. LET: U be a neighbourhood of l .
 - $\langle 2 \rangle 3$. PICK $B \in \mathcal{B}$ such that $l \in B \subseteq U$
 - $\langle 2 \rangle 4$. PICK N such that, for all $n \geq N$, we have $a_n \in B$
- PROOF: From $\langle 2 \rangle 1$.
- $\langle 2 \rangle 5$. For all $n \geq N$ we have $a_n \in U$

□

Lemma 82. *If a sequence (a_n) is constant with $a_n = l$ for all n , then $a_n \rightarrow l$ as $n \rightarrow \infty$.*

PROOF: Immediate from definitions. □

Theorem 83. *Let X be a linearly ordered set. Let (s_n) be a monotone increasing sequence in X with a supremum s . Then $s_n \rightarrow s$ as $n \rightarrow \infty$.*

PROOF:

- $\langle 1 \rangle 1$. ASSUME: s is not least in X .

PROOF: Otherwise (s_n) is the constant sequence s and the result follows from Lemma 82.

- ⟨1⟩2. LET: U be a neighbourhood of s .
- ⟨1⟩3. PICK $a < s$ such that $(a, s] \subseteq U$
- ⟨1⟩4. PICK N such that $a < a_N$.
- ⟨1⟩5. For all $n \geq N$ we have $a_n \in (a, s]$
- ⟨1⟩6. For all $n \geq N$ we have $a_n \in U$.

□

Theorem 84. If $\sum_{i=0}^{\infty} a_i = s$ and $\sum_{i=0}^{\infty} b_i = t$ then $\sum_{i=0}^{\infty} (ca_i + b_i) = cs + t$.

PROOF: $\sum_{i=0}^N (ca_i + b_i) = c \sum_{i=0}^N a_i + \sum_{i=0}^N b_i \rightarrow cs + t$ as $n \rightarrow \infty$. □

Theorem 85 (Comparison Test). If $|a_i| \leq b_i$ for all i and $\sum_{i=0}^{\infty} b_i$ converges then $\sum_{i=0}^{\infty} a_i$ converges.

PROOF:

- ⟨1⟩1. $\sum_{i=0}^{\infty} |a_i|$ converges

PROOF: The partial sums $\sum_{i=0}^N |a_i|$ form an increasing sequence bounded above by $\sum_{i=0}^{\infty} b_i$.

- ⟨1⟩2. LET: $c_i = |a_i| + a_i$ for all i

- ⟨1⟩3. $\sum_{i=0}^{\infty} c_i$ converges

PROOF: Each c_i is either $2|a_i|$ or 0. So the partial sums $\sum_{i=0}^N c_i$ form an increasing sequence bounded above by $2 \sum_{i=0}^{\infty} b_i$.

- ⟨1⟩4. Q.E.D.

PROOF: Since $a_i = c_i - |a_i|$.

□

Corollary 85.1. If $\sum_{i=0}^{\infty} |a_i|$ converges then $\sum_{i=0}^{\infty} a_i$ converges.

Theorem 86 (Weierstrass M-test). Let X be a set and $(f_n : X \rightarrow \mathbb{R})$ be a sequence of functions. Let

$$s_n(x) = \sum_{i=0}^n f_i(x)$$

for all n, x . Suppose $|f_i(x)| \leq M_i$ for all $i \geq 0$ and $x \in X$. If the series $\sum_{i=0}^{\infty} M_i$ converges, then the sequence (s_n) converges uniformly to

$$s(x) = \sum_{i=0}^{\infty} f_i(x) .$$

PROOF:

- ⟨1⟩1. LET: $r_n = \sum_{i=n+1}^{\infty} M_i$ for all n

- ⟨1⟩2. Given $0 \leq n < k$, we have $|s_k(x) - s_n(x)| \leq r_n$

PROOF:

$$\begin{aligned}
|s_k(x) - s_n(x)| &= \left| \sum_{i=n+1}^k f_i(x) \right| \\
&\leq \sum_{i=n+1}^k |f_i(x)| \\
&\leq \sum_{i=n+1}^k M_i \\
&\leq r_n
\end{aligned}$$

⟨1⟩3. Given $n \geq 0$ we have $|s(x) - s_n(x)| \leq r_n$

PROOF: By taking the limit $k \rightarrow \infty$ in ⟨1⟩2.

⟨1⟩4. Q.E.D.

PROOF: Since $r_n \rightarrow 0$ as $n \rightarrow \infty$.

□

14 Locally Finite Sets

Definition 87 (Locally Finite). Let X be a topological space and $\{A_\alpha\}$ a family of subsets of X . Then \mathcal{A} is *locally finite* if and only if every point in X has a neighbourhood that intersects A_α for only finitely many α .

Theorem 88 (Pasting Lemma). Let X and Y be topological spaces and $f : X \rightarrow Y$. Let $\{A_\alpha\}$ be a locally finite family of closed subsets of X that cover X . Suppose $f \upharpoonright A_\alpha$ is continuous for all α . Then f is continuous.

PROOF:

⟨1⟩1. Let X and Y be topological spaces and $f : X \rightarrow Y$. Let A and B be closed subsets of X such that $X = A \cup B$. Suppose $f \upharpoonright A$ and $f \upharpoonright B$ are continuous. Then f is continuous.

⟨2⟩1. LET: $C \subseteq Y$ be closed.

⟨2⟩2. $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$

⟨2⟩3. $f^{-1}(C)$ and $g^{-1}(C)$ are closed in X .

PROOF: Theorems 98 and 148.

⟨2⟩4. $h^{-1}(C)$ is closed in X .

PROOF: Lemma 38.

⟨2⟩5. Q.E.D.

PROOF: Theorem 98.

⟨1⟩2. Let X and Y be topological spaces and $f : X \rightarrow Y$. Let $\{A_\alpha\}$ be a finite family of closed subsets of X that cover X . Suppose $f \upharpoonright A_\alpha$ is continuous for all α . Then f is continuous.

PROOF: From ⟨1⟩1 by induction.

⟨1⟩3. Let X and Y be topological spaces and $f : X \rightarrow Y$. Let $\{A_\alpha\}$ be a locally finite family of closed subsets of X that cover X . Suppose $f \upharpoonright A_\alpha$ is continuous for all α . Then f is continuous.

- (2)1. LET: $x \in X$
 PROVE: f is continuous at x
 (2)2. PICK a neighbourhood U of x that intersects A_α for only finitely many α .
 (2)3. $f \upharpoonright U$ is continuous
 PROOF: By (1)2.
 (2)4. Q.E.D.
 PROOF: Lemma 108.

□

The following example shows that we cannot remove the assumption of local finiteness.

Example 89. Define $f : [-1, 1] \rightarrow \mathbb{R}$ by: $f(x) = 1$ if $x < -1$, $f(x) = 0$ if $x > 1$. Let $C_n = [-1, -1/n]$ for $n \geq 1$, and $D = [0, 1]$. Then $[-1, 1] = \bigcup_{n=1}^{\infty} C_n \cup D$ and f is continuous on each C_n and each D , but f is not continuous on $[-1, 1]$.

15 Open Maps

Definition 90 (Open Map). Let X and Y be topological spaces and $f : X \rightarrow Y$. Then f is an *open map* if and only if, for every open set U in X , the set $f(U)$ is open in Y .

Lemma 91. Let X and Y be topological spaces and $f : X \rightarrow Y$. Let \mathcal{B} be a basis for the topology on X . If $f(B)$ is open in Y for all $B \in \mathcal{B}$, then f is an open map.

PROOF: From Lemma 67. □

Proposition 92. Let X and Y be topological spaces. Let \mathcal{B} be a basis for the topology on X . Let $f : X \rightarrow Y$. Suppose that, for all $B \in \mathcal{B}$, we have $f(B)$ is open in Y . Then f is an open map.

PROOF: For any $\mathcal{A} \subseteq \mathcal{B}$, we have $f(\bigcup \mathcal{A}) = \bigcup_{B \in \mathcal{A}} f(B)$ is open in Y . The result follows from Lemma 67. □

16 Continuous Functions

Definition 93 (Continuous). Let X and Y be topological spaces. A function $f : X \rightarrow Y$ is *continuous* if and only if, for every open set V in Y , the set $f^{-1}(V)$ is open in X .

Proposition 94. Let X and Y be topological spaces and $f : X \rightarrow Y$. Let \mathcal{B} be a basis for Y . Then f is continuous if and only if, for all $B \in \mathcal{B}$, we have $f^{-1}(B)$ is open in X .

PROOF:

- (1)1. If f is continuous then, for all $B \in \mathcal{B}$, we have $f^{-1}(B)$ is open in X .

PROOF: Since every element of \mathcal{B} is open (Lemma 67).

⟨1⟩2. Suppose that, for all $B \in \mathcal{B}$, we have $f^{-1}(B)$ is open in X . Then f is continuous.

⟨2⟩1. ASSUME: For all $B \in \mathcal{B}$, we have $f^{-1}(B)$ is open in X .

⟨2⟩2. LET: V be open in Y .

⟨2⟩3. PICK $\mathcal{A} \subseteq \mathcal{B}$ such that $V = \bigcup \mathcal{A}$

PROOF: By Lemma 67.

⟨2⟩4. $f^{-1}(V)$ is open in X .

PROOF:

$$\begin{aligned} f^{-1}(V) &= f^{-1}\left(\bigcup \mathcal{A}\right) \\ &= \bigcup_{B \in \mathcal{A}} f^{-1}(B) \end{aligned}$$

□

Proposition 95. *Let X and Y be topological spaces and $f : X \rightarrow Y$. Let \mathcal{S} be a subbasis for Y . Then f is continuous if and only if, for all $S \in \mathcal{S}$, we have $f^{-1}(S)$ is open in X .*

PROOF:

⟨1⟩1. If f is continuous then, for all $S \in \mathcal{S}$, we have $f^{-1}(S)$ is open in X .

PROOF: Since every element of \mathcal{S} is open.

⟨1⟩2. Suppose that, for all $S \in \mathcal{S}$, we have $f^{-1}(S)$ is open in X . Then f is continuous.

⟨2⟩1. ASSUME: For all $S \in \mathcal{S}$, we have $f^{-1}(S)$ is open in X .

⟨2⟩2. LET: $S_1, \dots, S_n \in \mathcal{S}$

⟨2⟩3. $f^{-1}(S_1 \cap \dots \cap S_n)$ is open in X

PROOF: Since $f^{-1}(S_1 \cap \dots \cap S_n) = f^{-1}(S_1) \cap \dots \cap f^{-1}(S_n)$.

⟨2⟩4. Q.E.D.

PROOF: By Propositions 94 and 75.

□

Proposition 96. *Let X and Y be topological spaces and $f : X \rightarrow Y$. Let \mathcal{S} be a basis for Y . Then f is continuous if and only if, for all $V \in \mathcal{S}$, we have $f^{-1}(V)$ is open in X .*

PROOF:

⟨1⟩1. If f is continuous then, for all $V \in \mathcal{S}$, we have $f^{-1}(V)$ is open in X .

PROOF: Since every element of \mathcal{S} is open.

⟨1⟩2. Suppose that, for all $V \in \mathcal{S}$, we have $f^{-1}(V)$ is open in X . Then f is continuous.

⟨2⟩1. ASSUME: For all $V \in \mathcal{S}$, we have $f^{-1}(V)$ is open in X .

⟨2⟩2. For every set B that is the finite intersection of elements of \mathcal{S} , we have $f^{-1}(B)$ is open in X .

PROOF: Because $f^{-1}(V_1 \cap \dots \cap V_n) = f^{-1}(V_1) \cap \dots \cap f^{-1}(V_n)$.

⟨2⟩3. Q.E.D.

PROOF: From Propositions 75 and 94.

□

Definition 97 (Continuous at a Point). Let X and Y be topological spaces. Let $f : X \rightarrow Y$ and $x \in X$. Then f is *continuous at x* if and only if, for every neighbourhood V of $f(x)$, there exists a neighbourhood U of x such that $f(U) \subseteq V$.

Theorem 98. Let X and Y be topological spaces and $f : X \rightarrow Y$. Then the following are equivalent:

1. f is continuous.
2. For all $A \subseteq X$, we have $f(\overline{A}) \subseteq \overline{f(A)}$
3. For all $B \subseteq Y$ closed, we have $f^{-1}(B)$ is closed in X .
4. f is continuous at every point of X .

PROOF:

$\langle 1 \rangle 1. 1 \Rightarrow 2$

$\langle 2 \rangle 1.$ ASSUME: f is continuous.

$\langle 2 \rangle 2.$ LET: $A \subseteq X$

$\langle 2 \rangle 3.$ LET: $x \in \overline{A}$

PROVE: $f(x) \in \overline{f(A)}$

$\langle 2 \rangle 4.$ LET: V be a neighbourhood of $f(x)$

$\langle 2 \rangle 5.$ $f^{-1}(V)$ is a neighbourhood of x

$\langle 2 \rangle 6.$ PICK $y \in A \cap f^{-1}(V)$

PROOF: By Theorem 52.

$\langle 2 \rangle 7.$ $f(y) \in V \cap f(A)$

$\langle 2 \rangle 8.$ Q.E.D.

PROOF: By Theorem 52.

$\langle 1 \rangle 2. 2 \Rightarrow 3$

$\langle 2 \rangle 1.$ ASSUME: 2

$\langle 2 \rangle 2.$ LET: B be closed in Y

$\langle 2 \rangle 3.$ LET: $x \in \overline{f^{-1}(B)}$

PROVE: $x \in f^{-1}(B)$

$\langle 2 \rangle 4.$ $f(x) \in B$

PROOF:

$$f(x) \in \overline{f(f^{-1}(B))}$$

$$\subseteq \overline{f(f^{-1}(B))}$$

$((2)1)$

$$\subseteq \overline{B}$$

(Proposition 53)

$$= B$$

$\langle 1 \rangle 3. 3 \Rightarrow 1$

$\langle 2 \rangle 1.$ ASSUME: 3

$\langle 2 \rangle 2.$ LET: V be open in Y

$\langle 2 \rangle 3.$ $Y \setminus V$ is closed in Y

$\langle 2 \rangle 4.$ $f^{-1}(Y \setminus V)$ is closed in X

$\langle 2 \rangle 5.$ $X \setminus f^{-1}(V)$ is closed in X

$\langle 2 \rangle 6.$ $f^{-1}(V)$ is open in X

⟨1⟩4. $1 \Rightarrow 4$

PROOF: For any neighbourhood V of $f(x)$, the set $U = f^{-1}(V)$ is a neighbourhood of x such that $f(U) \subseteq V$.

⟨1⟩5. $4 \Rightarrow 1$

⟨2⟩1. ASSUME: 4

⟨2⟩2. LET: V be open in Y

⟨2⟩3. LET: $x \in f^{-1}(V)$

⟨2⟩4. V is a neighbourhood of $f(x)$

⟨2⟩5. PICK a neighbourhood U of x such that $f(U) \subseteq V$

⟨2⟩6. $x \in U \subseteq f^{-1}(V)$

⟨2⟩7. Q.E.D.

PROOF: By Lemma 30.

□

Theorem 99. *A constant function is continuous.*

PROOF: Let X and Y be topological spaces. Let $b \in Y$, and let $f : X \rightarrow Y$ be the constant function with value b . For any open $V \subseteq Y$, the set $f^{-1}(V)$ is either X (if $b \in V$) or \emptyset (if $b \notin V$). □

Theorem 100. *If A is a subspace of X then the inclusion $j : A \rightarrow X$ is continuous.*

PROOF: For any V open in X , we have $j^{-1}(V) = V \cap A$ is open in A . □

Theorem 101. *The composite of two continuous functions is continuous.*

PROOF: Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be continuous. For any V open in Z , we have $(g \circ f)^{-1}(V) = f^{-1}(g^{-1}(V))$ is open in X . □

Theorem 102. *Let $f : X \rightarrow Y$ be a continuous function and A be a subspace of X . Then the restriction $f \upharpoonright A : A \rightarrow Y$ is continuous.*

PROOF: Let V be open in Y . Then $(f \upharpoonright A)^{-1}(V) = f^{-1}(V) \cap A$ is open in A . □

Theorem 103. *Let $f : X \rightarrow Y$ be continuous. Let Z be a subspace of Y such that $f(X) \subseteq Z$. Then the corestriction $f : X \rightarrow Z$ is continuous.*

PROOF:

⟨1⟩1. LET: V be open in Z .

⟨1⟩2. PICK U open in Y such that $V = U \cap Z$.

⟨1⟩3. $f^{-1}(V) = f^{-1}(U)$

⟨1⟩4. $f^{-1}(V)$ is open in X .

□

Theorem 104. *Let $f : X \rightarrow Y$ be continuous. Let Z be a space such that Y is a subspace of Z . Then the expansion $f : X \rightarrow Z$ is continuous.*

PROOF: Let V be open in Z . Then $f^{-1}(V) = f^{-1}(V \cap Y)$ is open in X . □

Theorem 105. *Let X and Y be topological spaces. Let $f : X \rightarrow Y$. Suppose \mathcal{U} is a set of open sets in X such that $X = \bigcup \mathcal{U}$ and, for all $U \in \mathcal{U}$, we have $f \upharpoonright U : U \rightarrow Y$ is continuous. Then f is continuous.*

PROOF:

- $\langle 1 \rangle 1$. LET: V be open in Y
- $\langle 1 \rangle 2$. $f^{-1}(V) = \bigcup_{U \in \mathcal{U}} (f \upharpoonright U)^{-1}(V)$
- $\langle 1 \rangle 3$. For all $U \in \mathcal{U}$, we have $(f \upharpoonright U)^{-1}(V)$ is open in U .
- $\langle 1 \rangle 4$. For all $U \in \mathcal{U}$, we have $(f \upharpoonright U)^{-1}(V)$ is open in X .

PROOF: Lemma 147.

□

Proposition 106. *Let X and X' be the same set X under two topologies \mathcal{T} and \mathcal{T}' . Let $i : X \rightarrow X'$ be the identity function. Then i is continuous if and only if $\mathcal{T}' \subseteq \mathcal{T}$.*

PROOF: Immediate from definitions. □

Proposition 107. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and $a \in \mathbb{R}$. Then f is continuous on the right at a if and only if f is continuous at a as a function $\mathbb{R}_l \rightarrow \mathbb{R}$.*

PROOF:

- $\langle 1 \rangle 1$. If f is continuous on the right at a then f is continuous at a as a function $\mathbb{R}_l \rightarrow \mathbb{R}$.
 - $\langle 2 \rangle 1$. ASSUME: f is continuous on the right at a .
 - $\langle 2 \rangle 2$. LET: V be a neighbourhood of $f(a)$
 - $\langle 2 \rangle 3$. PICK b, c such that $f(a) \in (b, c) \subseteq V$.
 - $\langle 2 \rangle 4$. LET: $\epsilon = \min(c - f(a), f(a) - b)$
 - $\langle 2 \rangle 5$. PICK $\delta > 0$ such that, for all x , if $a < x < a + \delta$ then $|f(x) - f(a)| < \epsilon$
 - $\langle 2 \rangle 6$. LET: $U = [a, a + \delta)$
 - $\langle 2 \rangle 7$. $f(U) \subseteq V$
- $\langle 1 \rangle 2$. If f is continuous at a as a function $\mathbb{R}_l \rightarrow \mathbb{R}$ then f is continuous on the right at a .
 - $\langle 2 \rangle 1$. ASSUME: f is continuous at a as a function $\mathbb{R}_l \rightarrow \mathbb{R}$
 - $\langle 2 \rangle 2$. LET: $\epsilon > 0$
 - $\langle 2 \rangle 3$. PICK a neighbourhood U of a such that $f(U) \subseteq (f(a) - \epsilon, f(a) + \epsilon)$
 - $\langle 2 \rangle 4$. PICK b, c such that $a \in [b, c) \subset U$
 - $\langle 2 \rangle 5$. LET: $\delta = c - a$
 - $\langle 2 \rangle 6$. For all x , if $a < x < a + \delta$ then $|f(x) - f(a)| < \epsilon$

□

Lemma 108. *Let $f : X \rightarrow Y$. Let Z be an open subspace of X and $a \in Z$. If $f \upharpoonright Z$ is continuous at a then f is continuous at a .*

PROOF:

- $\langle 1 \rangle 1$. LET: V be a neighbourhood of $f(a)$
- $\langle 1 \rangle 2$. PICK a neighbourhood W of a in Z such that $f(W) \subseteq V$
- $\langle 1 \rangle 3$. W is a neighbourhood of a in X such that $f(W) \subseteq V$

PROOF: Lemma 147.

□

Proposition 109. *Let $f : A \rightarrow B$ and $g : C \rightarrow D$ be continuous. Define $f \times g : A \times C \rightarrow B \times D$ by*

$$(f \times g)(a, c) = (f(a), g(c)) .$$

Then $f \times g$ is continuous.

PROOF: $\pi_1 \circ (f \times g) = f \circ \pi_1$ and $\pi_2 \circ (f \times g) = g \circ \pi_2$ are continuous by Theorem 101. The result follows by Theorem 137.

Proposition 110. *Let X and Y be topological spaces and $f : X \rightarrow Y$ be continuous. If $a_n \rightarrow l$ as $n \rightarrow \infty$ in X then $f(a_n) \rightarrow f(l)$ as $n \rightarrow \infty$.*

PROOF:

⟨1⟩1. LET: V be a neighbourhood of $f(l)$

⟨1⟩2. PICK a neighbourhood U of l such that $f(U) \subseteq V$

⟨1⟩3. PICK N such that, for all $n \geq N$, we have $a_n \in U$

⟨1⟩4. For all $n \geq N$ we have $f(a_n) \in V$

□

17 Homeomorphisms

Definition 111 (Homeomorphism). Let X and Y be topological spaces. A *Homeomorphism* f between X and Y , $f : X \cong Y$, is a bijection $f : X \rightarrow Y$ such that both f and f^{-1} are continuous.

Lemma 112. *Let X and Y be topological spaces and $f : X \rightarrow Y$ a bijection. Then the following are equivalent:*

1. f is a homeomorphism.
2. f is continuous and an open map.
3. f is continuous and a closed map.
4. For any $U \subseteq X$, we have U is open if and only if $f(U)$ is open.

PROOF: Immediate from definitions. □

Proposition 113. *Let X and X' be the same set X under two topologies \mathcal{T} and \mathcal{T}' . Let $i : X \rightarrow X'$ be the identity function. Then i is a homeomorphism if and only if $\mathcal{T} = \mathcal{T}'$.*

PROOF: Immediate from definitions. □

Definition 114 (Topological Property). Let P be a property of topological spaces. Then P is a *topological* property if and only if, for any spaces X and Y , if P holds of X and $X \cong Y$ then P holds of Y .

Definition 115 (Topological Imbedding). Let X and Y be topological spaces and $f : X \rightarrow Y$. Then f is a *topological imbedding* if and only if the corestriction $f : X \rightarrow f(X)$ is a homeomorphism.

Proposition 116. Let X and Y be topological spaces and $a \in X$. The function $i : Y \rightarrow X \times Y$ that maps y to (a, y) is an imbedding.

PROOF:

$\langle 1 \rangle 1$. i is injective

$\langle 1 \rangle 2$. i is continuous.

PROOF: For U open in X and V open in Y , we have $i^{-1}(U \times V)$ is V if $a \in U$, and \emptyset if $a \notin U$.

$\langle 1 \rangle 3$. $i : Y \rightarrow i(Y)$ is an open map.

PROOF: For V open in Y we have $i(V) = (X \times V) \cap i(Y)$.

□

18 The Order Topology

Definition 117 (Order Topology). Let X be a linearly ordered set with at least two points. The *order topology* on X is the topology generated by the basis \mathcal{B} consisting of:

- all open intervals (a, b) ;
- all intervals of the form $[\perp, b)$ where \perp is least in X ;
- all intervals of the form $(a, \top]$ where \top is greatest in X .

We prove this is a basis for a topology.

PROOF:

$\langle 1 \rangle 1$. For all $x \in X$ there exists $B \in \mathcal{B}$ such that $x \in B$.

$\langle 2 \rangle 1$. LET: $x \in X$

$\langle 2 \rangle 2$. CASE: x is greatest in X .

$\langle 3 \rangle 1$. PICK $y \in X$ with $y \neq x$

$\langle 3 \rangle 2$. $x \in (y, x] \in \mathcal{B}$

$\langle 2 \rangle 3$. CASE: x is least in X .

$\langle 3 \rangle 1$. PICK $y \in X$ with $y \neq x$

$\langle 3 \rangle 2$. $x \in [x, y) \in \mathcal{B}$

$\langle 2 \rangle 4$. CASE: x is neither greatest nor least in X .

$\langle 3 \rangle 1$. PICK $a, b \in X$ with $a < x$ and $x < b$

$\langle 3 \rangle 2$. $x \in (a, b) \in \mathcal{B}$

$\langle 1 \rangle 2$. For all $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$

$\langle 2 \rangle 1$. LET: $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$

$\langle 2 \rangle 2$. CASE: $B_1 = (a, b)$, $B_2 = (c, d)$

PROOF: Take $B_3 = (\max(a, c), \min(b, d))$.

$\langle 2 \rangle 3$. CASE: $B_1 = (a, b)$, $B_2 = [\perp, d)$

PROOF: Take $B_3 = (a, \min(b, d))$.
 $\langle 2 \rangle 4$. CASE: $B_1 = (a, b)$, $B_2 = (c, \top]$
PROOF: Take $B_3 = (\max(a, c), b)$.
 $\langle 2 \rangle 5$. CASE: $B_1 = [\perp, b)$, $B_2 = [\perp, d)$
PROOF: Take $B_3 = [\perp, \min(b, d))$.
 $\langle 2 \rangle 6$. CASE: $B_1 = [\perp, b)$, $B_2 = (c, \top]$
PROOF: Take $B_3 = (c, b)$.

□

Lemma 118. *Let X be a linearly ordered set. Then the open rays form a subbasis for the order topology on X .*

PROOF:

$\langle 1 \rangle 1$. Every open ray is open.
 $\langle 2 \rangle 1$. For all $a \in X$, the ray $(-\infty, a)$ is open.
 $\langle 3 \rangle 1$. LET: $x \in (-\infty, a)$
 $\langle 3 \rangle 2$. CASE: x is least in X
PROOF: $x \text{ in } [x, a) = (-\infty, a)$.
 $\langle 3 \rangle 3$. CASE: x is not least in X
 $\langle 4 \rangle 1$. PICK $y < x$
 $\langle 4 \rangle 2$. $x \in (y, a) \subseteq (-\infty, a)$
 $\langle 2 \rangle 2$. For all $a \in X$, the ray $(a, +\infty)$ is open.
PROOF: Similar.
 $\langle 1 \rangle 2$. Every basic open set is a finite intersection of open rays.
PROOF: We have $(a, b) = (a, +\infty) \cap (-\infty, b)$, $[\perp, b) = (-\infty, b)$ and $(a, \top] = (a, +\infty)$.

□

Definition 119 (Standard Topology on the Real Line). The *standard topology on the real line* is the order topology on \mathbb{R} generated by the standard order.

Lemma 120. *The lower limit topology is strictly finer than the standard topology on \mathbb{R} .*

PROOF:

$\langle 1 \rangle 1$. Every open interval is open in the lower limit topology.
PROOF: If $x \in (a, b)$ then $x \in [x, b) \subseteq (a, b)$.
 $\langle 1 \rangle 2$. The half-open interval $[0, 1)$ is not open in the standard topology.
PROOF: There is no open interval (a, b) such that $0 \in (a, b) \subseteq [0, 1)$.

□

Lemma 121. *The K -topology is strictly finer than the standard topology on \mathbb{R} .*

PROOF:

$\langle 1 \rangle 1$. Every open interval is open in the K -topology.
PROOF: Corollary 67.1.
 $\langle 1 \rangle 2$. The set $(-1, 1) \setminus K$ is not open in the standard topology.
PROOF: There is no open interval (a, b) such that $0 \in (a, b) \subseteq (-1, 1) \setminus K$, since there must be a positive integer n with $1/n \in (a, b)$.

□

Lemma 122. *Let X be a topological space. Let Y be a linearly ordered set in the order topology. Let $f, g : X \rightarrow Y$ be continuous. Then $C = \{x \in X \mid f(x) \leq g(x)\}$ is closed.*

PROOF:

⟨1⟩1. LET: $x \in X \setminus C$

⟨1⟩2. $f(x) > g(x)$

PROVE: There exists a neighbourhood U of x such that $U \subseteq X \setminus C$

⟨1⟩3. CASE: There exists y such that $g(x) < y < f(x)$

PROOF: Take $U = g^{-1}((-\infty, y)) \cup f^{-1}(y, +\infty)$.

⟨1⟩4. CASE: There is no y such that $g(x) < y < f(x)$

PROOF: Take $U = g^{-1}((-\infty, f(x))) \cup f^{-1}(g(x), +\infty)$.

□

Proposition 123. *Let X be a topological space. Let Y be a linearly ordered set in the order topology. Let $f, g : X \rightarrow Y$ be continuous. Define $h : X \rightarrow Y$ by $h(x) = \min(f(x), g(x))$. Then h is continuous.*

PROOF: By the Pasting Lemma applied to $\{x \in X \mid f(x) \leq g(x)\}$ and $\{x \in X \mid g(x) \leq f(x)\}$, which are closed by Lemma 122.

Proposition 124. *Let X and Y be linearly ordered sets in the order topology. Let $f : X \rightarrow Y$ be strictly monotone and surjective. Then f is a homeomorphism.*

PROOF:

⟨1⟩1. f is bijective.

PROOF: Proposition 10.

⟨1⟩2. f is continuous.

⟨2⟩1. For all $y \in Y$ we have $f^{-1}((y, +\infty))$ is open.

⟨3⟩1. LET: $y \in Y$

⟨3⟩2. PICK $x \in X$ such that $f(x) = y$

PROOF: Since f is surjective.

⟨3⟩3. $f^{-1}((y, +\infty)) = (x, +\infty)$

PROOF: By strict monotonicity.

⟨2⟩2. For all $y \in Y$ we have $f^{-1}((-\infty, y))$ is open.

PROOF: Similar.

⟨1⟩3. f^{-1} is continuous.

⟨2⟩1. For all $x \in X$ we have $f((x, +\infty))$ is open.

PROOF: $f((x, +\infty)) = (f(x), +\infty)$.

⟨2⟩2. For all $x \in X$ we have $f((-\infty, x))$ is open.

PROOF: $f((-\infty, x)) = (-\infty, f(x))$.

□

19 The n th Root Function

Proposition 125. For all $n \geq 1$, the function $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ defined by $f(x) = x^n$ is a homeomorphism.

PROOF:

$\langle 1 \rangle 1$. f is strictly monotone.

$\langle 2 \rangle 1$. LET: $x, y \in \mathbb{R}$ with $0 \leq x < y$

$\langle 2 \rangle 2$. $x^n < y^n$

$$y^n - x^n = (y - x)(y^{n-1} + y^{n-2}x + y^{n-3}x^2 + \cdots + x^{n-1}) > 0$$

$\langle 1 \rangle 2$. f is surjective.

$\langle 2 \rangle 1$. LET: $y \in \mathbb{R}_{\geq 0}$

$\langle 2 \rangle 2$. PICK $x \in \mathbb{R}$ such that $y \leq x^n$

PROOF: If $y \leq 1$ take $x = 1$, otherwise take $x = y$.

$\langle 2 \rangle 3$. There exists $x' \in [0, x]$ such that $(x')^n = y$

PROOF: By the Intermediate Value Theorem.

$\langle 1 \rangle 3$. Q.E.D.

PROOF: Proposition 124.

□

Definition 126. For $n \geq 1$, the n th root function is the function $\mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ that is the inverse of $\lambda x.x^n$.

20 The Product Topology

Definition 127 (Product Topology). Let $\{A_i\}_{i \in I}$ be a family of topological spaces. The *product topology* on $\prod_{i \in I} A_i$ is the topology generated by the subbasis consisting of the sets of the form $\pi_i^{-1}(U)$ where $i \in I$ and U is open in A_i .

Proposition 128. The product topology on $\prod_{i \in I} A_i$ is generated by the basis consisting of all sets of the form $\prod_{i \in I} U_i$ where $\{U_i\}_{i \in I}$ is a family such that each U_i is an open set in A_i and $U_i = A_i$ for all but finitely many i .

PROOF: From Proposition 75. □

Proposition 129. If A_i is closed in X_i for all $i \in I$ then $\prod_{i \in I} A_i$ is closed in $\prod_{i \in I} X_i$.

PROOF:

$$\left(\prod_{i \in I} X_i\right) \setminus \left(\prod_{i \in I} A_i\right) = \bigcup_{j \in I} \left(\prod_{i \in I} X_i \setminus \pi_j^{-1}(A_j)\right) \square$$

Proposition 130. Let $\{A_i\}_{i \in I}$ be a family of topological spaces. For $i \in I$, let \mathcal{B}_i be a basis for the topology on A_i . Then $\mathcal{B} = \{\prod_{i \in I} B_i \mid \forall i \in I. B_i \in \mathcal{B}_i, B_i = A_i \text{ for all but finitely many } i\}$ is a basis for the box topology on $\prod_{i \in I} A_i$.

PROOF:

- ⟨1⟩1. Every set in \mathcal{B} is open.
- ⟨1⟩2. For every point $a \in \prod_{i \in I} A_i$ and every open set U with $a \in U$, there exists $B \in \mathcal{B}$ such that $a \in B \subseteq U$.
 - ⟨2⟩1. LET: U be open and $a \in U$
 - ⟨2⟩2. PICK a family $\{U_i\}_{i \in I}$ such that each U_i is open in A_i , such that $U_i = A_i$ except for $i = i_1, \dots, i_n$, and such that $a \in \prod_{i \in I} U_i \subseteq U$.
 - ⟨2⟩3. For $j = 1, \dots, n$, PICK $B_{i_j} \in \mathcal{B}_{i_j}$ such that $a_{i_j} \in B_{i_j} \subseteq U_{i_j}$
 - ⟨2⟩4. LET: $B = \prod_{i \in I} B_i$ where $B_i = A_i$ for $i \neq i_1, \dots, i_n$
 - ⟨2⟩5. $B \in \mathcal{B}$
 - ⟨2⟩6. $a \in B \subseteq U$
- ⟨1⟩3. Q.E.D.

PROOF: Lemma 68.

□

Proposition 131. *Let $\{A_i\}_{i \in I}$ be a family of topological spaces. Then the projections $\pi_i : \prod_{i \in I} A_i \rightarrow A_i$ are open maps.*

PROOF: From Lemma 91. □

Example 132. The projections are not always closed maps. For example, $\pi_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$ maps the closed set $\{(x, 1/x) \mid x > 0\}$ to $(0, +\infty)$.

Proposition 133. *Let $\{X_i\}_{i \in I}$ be a family of sets. For $i \in I$, let \mathcal{T}_i and \mathcal{U}_i be topologies on X_i . Let \mathcal{P} be the product topology on $\prod_{i \in I} X_i$ generated by the topologies \mathcal{T}_i , and \mathcal{Q} the product topology on the same set generated by the topologies \mathcal{U}_i . Then $\mathcal{P} \subseteq \mathcal{Q}$ if and only if $\mathcal{T}_i \subseteq \mathcal{U}_i$ for all i .*

PROOF:

- ⟨1⟩1. If $\mathcal{T}_i \subseteq \mathcal{U}_i$ for all i then $\mathcal{P} \subseteq \mathcal{Q}$

PROOF: By Corollary 67.1.
- ⟨1⟩2. If $\mathcal{P} \subseteq \mathcal{Q}$ then $\mathcal{T}_i \subseteq \mathcal{U}_i$ for all i
 - ⟨2⟩1. ASSUME: $\mathcal{P} \subseteq \mathcal{Q}$
 - ⟨2⟩2. LET: $i \in I$
 - ⟨2⟩3. LET: $U \in \mathcal{T}_i$
 - ⟨2⟩4. LET: $U_i = U$ and $U_j = X_j$ for $j \neq i$
 - ⟨2⟩5. $\prod_{i \in I} U_i \in \mathcal{P}$
 - ⟨2⟩6. $\prod_{i \in I} U_i \in \mathcal{Q}$
 - ⟨2⟩7. $U \in \mathcal{U}_i$

PROOF: From Proposition 131.

□

Proposition 134 (AC). *Let $\{X_i\}_{i \in I}$ be a family of topological spaces. Let $A_i \subseteq X_i$ for all $i \in I$. Then*

$$\prod_{i \in I} \overline{A_i} = \overline{\prod_{i \in I} A_i} .$$

PROOF:

- (1)1. $\overline{\prod_{i \in I} A_i} \subseteq \prod_{i \in I} \overline{A_i}$
 (2)1. For all $i \in I$ we have $A_i \subseteq \overline{A_i}$
 PROOF: Lemma 49.
 (2)2. $\prod_{i \in I} A_i \subseteq \prod_{i \in I} \overline{A_i}$
 (2)3. Q.E.D.
 PROOF: Since $\prod_{i \in I} A_i$ is closed by Proposition 129.
 (1)2. $\prod_{i \in I} \overline{A_i} \subseteq \overline{\prod_{i \in I} A_i}$
 (2)1. LET: $x \in \prod_{i \in I} \overline{A_i}$
 (2)2. LET: U be a neighbourhood of x
 (2)3. PICK V_i open in X_i such that $x \in \prod_{i \in I} V_i \subseteq U$ with $V_i = X_i$ except for
 $i = i_1, \dots, i_n$
 (2)4. For $i \in I$, pick $a_i \in V_i \cap A_i$
 PROOF: By Theorem 52 and (2)1 using the Axiom of Choice.
 (2)5. U intersects $\prod_{i \in I} A_i$
 (2)6. Q.E.D.
 PROOF: $a \in U \cap \prod_{i \in I} A_i$

□

Example 135. The closure of \mathbb{R}^∞ in \mathbb{R}^ω is \mathbb{R}^ω

PROOF:

- (1)1. LET: $a \in \mathbb{R}^\omega$
 (1)2. LET: U be any neighbourhoods of a .
 (1)3. PICK U_n open in \mathbb{R} for all n such that $a \in \prod_{n \geq 0} U_n \subseteq U$ and $U_n = \mathbb{R}$ for
 all n except n_1, \dots, n_k
 (1)4. LET: $b_n = a_n$ for $n = n_1, \dots, n_k$ and $b_n = 0$ for all other n
 (1)5. $b \in \mathbb{R}^\infty \cap U$
 (1)6. Q.E.D.

PROOF: From Theorem 52.

□

Proposition 136. Let $\{X_i\}_{i \in I}$ be a family of topological spaces. Let (a_n) be a sequence in $\prod_{i \in I} X_i$ and $l \in \prod_{i \in I} X_i$. Then $a_n \rightarrow l$ as $n \rightarrow \infty$ if and only if, for all $i \in I$, we have $\pi_i(a_n) \rightarrow \pi_i(l)$ as $n \rightarrow \infty$.

PROOF:

- (1)1. If $a_n \rightarrow l$ as $n \rightarrow \infty$ then, for all $i \in I$, we have $\pi_i(a_n) \rightarrow \pi_i(l)$ as $n \rightarrow \infty$
 PROOF: Proposition 110.
 (1)2. If, for all $i \in I$, we have $\pi_i(a_n) \rightarrow \pi_i(l)$ as $n \rightarrow \infty$, then $a_n \rightarrow l$ as $n \rightarrow \infty$
 (2)1. ASSUME: For all $i \in I$, we have $\pi_i(a_n) \rightarrow \pi_i(l)$ as $n \rightarrow \infty$
 (2)2. LET: V be a neighbourhood of l
 (2)3. PICK open sets U_i in X_i such that $l \in \prod_{i \in I} U_i \subseteq V$ and $U_i = X_i$ for all
 i except $i = i_1, \dots, i_k$
 (2)4. For $j = 1, \dots, k$, PICK N_j such that, for all $n \geq N_j$, we have $\pi_{i_j}(a_n) \in$
 U_{i_j}
 (2)5. LET: $N = \max(N_1, \dots, N_k)$
 (2)6. For all $n \geq N$ we have $a_n \in V$

□

Theorem 137. *Let A be a topological space and $\{X_i\}_{i \in I}$ be a family of topological spaces. Let $f : A \rightarrow \prod_{i \in I} X_i$ be a function. If $\pi_i \circ f$ is continuous for all $i \in I$ then f is continuous.*

PROOF:

⟨1⟩1. LET: $i \in I$ and U be open in X_i

⟨1⟩2. $f^{-1}(\pi_i^{-1}(U))$ is open in A

⟨1⟩3. Q.E.D.

PROOF: Proposition 95.

□

20.1 Continuous in Each Variable Separately

Definition 138 (Continuous in Each Variable Separately). Let $F : X \times Y \rightarrow Z$. Then F is *continuous in each variable separately* if and only if:

- for every $a \in X$ the function $\lambda y \in Y. F(a, y)$ is continuous;
- for every $b \in Y$ the function $\lambda x \in X. F(x, b)$ is continuous.

Proposition 139. *Let $F : X \times Y \rightarrow Z$. If F is continuous then F is continuous in each variable separately.*

PROOF: For $a \in X$, the function $\lambda y \in Y. F(a, y)$ is $F \circ i$ where $i : Y \rightarrow X \times Y$ maps y to (a, y) . We have i is continuous by Proposition 116, hence $F \circ i$ is continuous by Theorem 101.

Similarly for $\lambda x \in X. F(x, b)$ for $b \in Y$. □

Example 140. Define $F : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$F(x, y) = \begin{cases} xy/(x^2 + y^2) & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

Then F is continuous in each variable separately but not continuous.

Proposition 141. *Let $f : A \rightarrow C$ and $g : B \rightarrow D$ be open maps. Then $f \times g : A \times B \rightarrow C \times D$ is an open map.*

PROOF: Given U open in A and V open in B . Then $(f \times g)(U \times V) = f(U) \times g(V)$ is open in $C \times D$. The result follows from Proposition 92. □

21 The Subspace Topology

Definition 142 (Subspace Topology). Let X be a topological space and $Y \subseteq X$. The *subspace topology* on Y is $\mathcal{T} = \{U \cap Y \mid U \text{ is open in } X\}$.

We prove this is a topology.

PROOF:

⟨1⟩1. $Y \in \mathcal{T}$

PROOF: Since $Y = X \cap Y$

⟨1⟩2. For all $\mathcal{U} \subseteq \mathcal{T}$, we have $\bigcup \mathcal{U} \in \mathcal{T}$

⟨2⟩1. LET: $\mathcal{U} \subseteq \mathcal{T}$

⟨2⟩2. LET: $\mathcal{V} = \{V \text{ open in } X \mid V \cap Y \in \mathcal{U}\}$

⟨2⟩3. $\bigcup \mathcal{U} = (\bigcup \mathcal{V}) \cap Y$

⟨1⟩3. For all $U, V \in \mathcal{T}$, we have $U \cap V \in \mathcal{T}$

⟨2⟩1. LET: $U, V \in \mathcal{T}$

⟨2⟩2. PICK U', V' open in X such that $U = U' \cap Y$ and $V = V' \cap Y$

⟨2⟩3. $(U \cap V) = (U' \cap V') \cap Y$

□

Theorem 143. Let X be a topological space and Y a subspace of X . Let $A \subseteq Y$. Then A is closed in Y if and only if there exists a closed set C in X such that $A = C \cap Y$.

PROOF: We have

$$\begin{aligned} & A \text{ is closed in } Y \\ \Leftrightarrow & Y \setminus A \text{ is open in } Y \\ \Leftrightarrow & \exists U \text{ open in } X. Y \setminus A = Y \cap U \\ \Leftrightarrow & \exists C \text{ closed in } X. Y \setminus A = Y \cap (X \setminus U) \\ \Leftrightarrow & \exists C \text{ closed in } X. A = Y \cap U \end{aligned} \quad \square$$

Theorem 144. Let Y be a subspace of X . Let $A \subseteq Y$. Let \bar{A} be the closure of A in X . Then the closure of A in Y is $\bar{A} \cap Y$.

PROOF: The closure of A in Y is

$$\begin{aligned} & \bigcap \{C \text{ closed in } Y \mid A \subseteq C\} \\ &= \bigcap \{D \cap Y \mid D \text{ closed in } X, A \subseteq D \cap Y\} \quad (\text{Theorem 143}) \\ &= \bigcap \{D \mid D \text{ closed in } X, A \subseteq D\} \cap Y \\ &= \bar{A} \cap Y \end{aligned} \quad \square$$

Lemma 145. Let X be a topological space and $Y \subseteq X$. Let \mathcal{B} be a basis for the topology on X . Then $\mathcal{B}' = \{B \cap Y \mid B \in \mathcal{B}\}$ is a basis for the subspace topology on Y .

PROOF:

⟨1⟩1. Every element in \mathcal{B}' is open in Y

⟨1⟩2. For every open set U in Y and point $y \in U$, there exists $B' \in \mathcal{B}'$ such that $y \in B' \subseteq U$

⟨2⟩1. LET: U be open in Y and $y \in U$

⟨2⟩2. PICK V open in X such that $U = V \cap Y$

⟨2⟩3. PICK $B \in \mathcal{B}$ such that $y \in B \subseteq V$

⟨2⟩4. LET: $B' = B \cap Y$

⟨2⟩5. $B' \in \mathcal{B}'$

⟨2⟩6. $y \in B' \subseteq U$

⟨1⟩3. Q.E.D.

PROOF: By Lemma 68.

□

Lemma 146. *Let X be a topological space and $Y \subseteq X$. Let \mathcal{S} be a basis for the topology on X . Then $\mathcal{S}' = \{S \cap Y \mid S \in \mathcal{S}\}$ is a subbasis for the subspace topology on Y .*

PROOF: The set $\{B \cap Y \mid B \text{ is a finite intersection of elements of } \mathcal{S}\}$ is a basis for the subspace topology by Lemma 145, and this is the set of all finite intersections of elements of \mathcal{S}' . □

Lemma 147. *Let Y be a subspace of X . If U is open in Y and Y is open in X then U is open in X .*

PROOF:

⟨1⟩1. PICK V open in X such that $U = V \cap Y$

⟨1⟩2. U is open in X

PROOF: Since it is the intersection of two open sets V and Y .

□

Theorem 148. *Let Y be a subspace of X and $A \subseteq Y$. If A is closed in Y and Y is closed in X then A is closed in X .*

PROOF: Pick a closed set C in X such that $A = C \cap Y$ (Theorem 143). Then A is the intersection of two sets closed in X , hence A is closed in X (Lemma 37).

□

Theorem 149. *Let $\{X_i\}_{i \in I}$ be a family of topological spaces. Let A_i be a subspace of X_i for all $i \in I$. Then the product topology on $\prod_{i \in I} A_i$ is the same as the topology it inherits as a subspace of $\prod_{i \in I} X_i$.*

PROOF: The product topology is generated by the subbasis

$$\begin{aligned} & \{\pi_i^{-1}(U) \mid i \in I, U \text{ open in } A_i\} \\ &= \{\pi_i^{-1}(V) \cap A_i \mid i \in I, V \text{ open in } X_i\} \\ &= \{\pi_i^{-1}(V) \mid i \in I, V \text{ open in } X_i\} \cap \prod_{i \in I} A_i \end{aligned}$$

and this is a subbasis for the subspace topology by Lemma 146. □

Theorem 150. *Let X be an ordered set in the order topology. Let $Y \subseteq X$ be an interval. Then the order topology on Y is the same as the subspace topology on Y .*

PROOF:

⟨1⟩1. The order topology is finer than the subspace topology.

⟨2⟩1. For every open ray R in X , the set $R \cap Y$ is open in the order topology.

⟨3⟩1. For all $a \in X$, we have $(-\infty, a) \cap Y$ is open in the order topology.

⟨4⟩1. CASE: For all $y \in Y$ we have $y < a$
 PROOF: In this case $(-\infty, a) \cap Y = Y$.
 ⟨4⟩2. CASE: For all $y \in Y$ we have $a < y$
 PROOF: In this case $(-\infty, a) \cap Y = \emptyset$.
 ⟨4⟩3. CASE: There exists $y \in Y$ such that $y \leq a$ and $y \in Y$ such that
 $a \leq y$
 ⟨5⟩1. $a \in Y$
 PROOF: Because Y is an interval.
 ⟨5⟩2. $(-\infty, a) \cap Y = \{y \in Y \mid y < a\}$
 ⟨3⟩2. For all $a \in X$, we have $(a, +\infty) \cap Y$ is open in the order topology.
 PROOF: Similar.
 ⟨2⟩2. Q.E.D.
 PROOF: By Lemmas 118 and 146 and Proposition 76.
 ⟨1⟩2. The subspace topology is finer than the order topology.
 ⟨2⟩1. Every open ray in Y is open in the subspace topology.
 PROOF: For any $a \in Y$ we have $(-\infty, a)_Y = (-\infty, a)_X \cap Y$ and $(a, +\infty)_Y = (a, +\infty)_X \cap Y$.
 ⟨2⟩2. Q.E.D.
 PROOF: By Lemma 118 and Proposition 76
 □

This example shows that we cannot remove the hypothesis that Y is an interval:

Example 151. The order topology on I_o^2 is different from the subspace topology as a subspace of \mathbb{R}^2 under the dictionary order topology.

PROOF: The set $\{1/2\} \times (1/2, 1)$ is open in the subspace topology but not in the order topology. □

Proposition 152. Let X be a topological space, Y a subspace of X , and Z a subspace of Y . Then the subspace topology on Z inherited from X is the same as the subspace topology on Z inherited from Y .

PROOF: The subspace topology inherited from Y is

$$\begin{aligned}
 & \{V \cap Z \mid V \text{ open in } Y\} \\
 &= \{U \cap Y \cap Z \mid U \text{ open in } X\} \\
 &= \{U \cap Z \mid U \text{ open in } X\}
 \end{aligned}$$

which is the subspace topology inherited from X . □

Definition 153 (Unit Circle). The *unit circle* S^1 is $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ as a subspace of \mathbb{R}^2 .

Definition 154 (Unit 2-sphere). The *unit 2-sphere* is $S^2 = \{(x, y, z) \mid x^2 + y^2 + z^2 \leq 1\}$ as a subspace of \mathbb{R}^3 .

Proposition 155. Let $f : X \rightarrow Y$ be an open map and $A \subseteq X$ be open. Then the restriction $f \upharpoonright A : A \rightarrow f(A)$ is an open map.

PROOF:

$\langle 1 \rangle 1$. LET: U be open in A

$\langle 1 \rangle 2$. U is open in X

PROOF: Lemma 147.

$\langle 1 \rangle 3$. $f(U)$ is open in Y

$\langle 1 \rangle 4$. $f(U)$ is open in $f(A)$

PROOF: Since $f(U) = f(U) \cap f(A)$.

□

Example 156. This example shows that we cannot remove the hypothesis that A is open.

Let $A = \{(x, y) \in \mathbb{R}^2 \mid (x > 0 \text{ and } y = 1/x) \text{ or } x = y = 0\}$. Then $\pi_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$ is an open map, but $\pi_1 \upharpoonright A : A \rightarrow [0, +\infty)$ is not, because it maps the set $\{0, 0\}$ which is open in A to $\{0\}$ which is not open in $[0, +\infty)$.

Proposition 157. *Let Y be a subspace of X . Let $A \subseteq Y$ and $l \in Y$. Then l is a limit point of A in Y if and only if l is a limit point of A in X .*

PROOF: Both are equivalent to the condition that any neighbourhood of l in X intersects A in a point other than l . □

22 The Box Topology

Definition 158 (Box Topology). Let $\{A_i\}_{i \in I}$ be a family of topological spaces. The *box topology* on $\prod_{i \in I} A_i$ is the topology generated by the set of all sets of the form $\prod_{i \in I} U_i$ where $\{U_i\}_{i \in I}$ is a family such that each U_i is open in A_i .

This is a basis since it covers $\prod_{i \in I} A_i$ and is closed under intersection.

Proposition 159. *The box topology is finer than the product topology.*

PROOF: From Proposition 128. □

Corollary 159.1. *If A_i is closed in X_i for all $i \in I$ then $\prod_{i \in I} A_i$ is closed in $\prod_{i \in I} X_i$ under the box topology.*

PROOF: From Proposition 129.

Proposition 160 (AC). *Let $\{A_i\}_{i \in I}$ be a family of topological spaces. For $i \in I$, let \mathcal{B}_i be a basis for the topology on A_i . Then $\mathcal{B} = \{\prod_{i \in I} B_i \mid \forall i \in I, B_i \in \mathcal{B}_i\}$ is a basis for the box topology on $\prod_{i \in I} A_i$.*

PROOF:

$\langle 1 \rangle 1$. Every set of the form $\prod_{i \in I} B_i$ is open.

$\langle 1 \rangle 2$. For every point $a \in \prod_{i \in I} A_i$ and every open set U with $a \in U$, there exists $B \in \mathcal{B}$ such that $a \in B \subseteq U$.

$\langle 2 \rangle 1$. LET: U be open and $a \in U$

$\langle 2 \rangle 2$. PICK a family $\{U_i\}_{i \in I}$ such that each U_i is open in A_i and $a \in \prod_{i \in I} U_i \subseteq U$.

⟨2⟩3. For $i \in I$, PICK $B_i \in \mathcal{B}_i$ such that $a_i \in B_i \subseteq U_i$

PROOF: Using the Axiom of Choice.

⟨2⟩4. $a \in \prod_{i \in I} B_i \subseteq U$

⟨1⟩3. Q.E.D.

PROOF: Lemma 68.

□

Theorem 161. *Let $\{X_i\}_{i \in I}$ be a family of topological spaces. Let A_i be a subspace of X_i for all $i \in I$. Give $\prod_{i \in I} X_i$ the box topology. Then the box topology on $\prod_{i \in I} A_i$ is the same as the topology it inherits as a subspace of $\prod_{i \in I} X_i$.*

PROOF: The box topology is generated by the basis

$$\begin{aligned} & \left\{ \prod_{i \in I} U_i \mid \forall i \in I, U_i \text{ open in } A_i \right\} \\ &= \left\{ \prod_{i \in I} (V_i \cap A_i) \mid \forall i \in I, V_i \text{ open in } X_i \right\} \\ &= \left\{ \prod_{i \in I} V_i \mid \forall i \in I, V_i \text{ open in } X_i \right\} \cap \prod_{i \in I} A_i \end{aligned}$$

and this is a basis for the subspace topology by Lemma 145. □

Proposition 162 (AC). *Let $\{X_i\}_{i \in I}$ be a family of topological spaces. Give $\prod_{i \in I} X_i$ the box topology. Let $A_i \subseteq X_i$ for all $i \in I$. Then*

$$\prod_{i \in I} \overline{A_i} = \overline{\prod_{i \in I} A_i}.$$

PROOF:

⟨1⟩1. $\overline{\prod_{i \in I} A_i} \subseteq \prod_{i \in I} \overline{A_i}$

⟨2⟩1. For all $i \in I$ we have $A_i \subseteq \overline{A_i}$

PROOF: Lemma 49.

⟨2⟩2. $\prod_{i \in I} A_i \subseteq \prod_{i \in I} \overline{A_i}$

⟨2⟩3. Q.E.D.

PROOF: Since $\prod_{i \in I} A_i$ is closed by Corollary 159.1.

⟨1⟩2. $\prod_{i \in I} \overline{A_i} \subseteq \overline{\prod_{i \in I} A_i}$

⟨2⟩1. LET: $x \in \prod_{i \in I} \overline{A_i}$

⟨2⟩2. LET: U be a neighbourhood of x

⟨2⟩3. PICK V_i open in X_i such that $x \in \prod_{i \in I} V_i \subseteq U$

⟨2⟩4. For $i \in I$, pick $a_i \in V_i \cap A_i$

PROOF: By Theorem 52 and ⟨2⟩1 using the Axiom of Choice.

⟨2⟩5. U intersects $\prod_{i \in I} A_i$

⟨2⟩6. Q.E.D.

PROOF: $a \in U \cap \prod_{i \in I} A_i$.

□

The following example shows that Theorem 137 fails in the box topology.

Example 163. Define $f : \mathbb{R} \rightarrow \mathbb{R}^\omega$ by $f(t) = (t, t, \dots)$. Then $\pi_n \circ f = \text{id}_{\mathbb{R}}$ is continuous for all n . But f is not continuous when \mathbb{R}^ω is given the box topology because the inverse image of

$$(-1, 1) \times (-1/2, 1/2) \times (-1/3, 1/3) \times \dots$$

is $\{0\}$ which is not open.

The following example shows that Proposition 136 fails in the box topology.

Example 164. Give \mathbb{R}^ω the box topology. Let $a_n = (1/n, 1/n, \dots)$ for $n \geq 1$ and $l = (0, 0, \dots)$. Then $\pi_i(a_n) \rightarrow \pi_i(l)$ as $n \rightarrow \infty$ for all i , but $a_n \not\rightarrow l$ as $n \rightarrow \infty$ since the open set

$$(-1, 1) \times (-1/2, 1/2) \times (-1/3, 1/3) \times \dots$$

contains l but does not contain any a_n .

Example 165. The set \mathbb{R}^∞ is closed in \mathbb{R}^ω under the box topology. For let (a_n) be any sequence not in \mathbb{R}^∞ . Let U_n be an open interval around a_n that does not contain 0 if $a_n \neq 0$, and $U_n = \mathbb{R}$ if $a_n = 0$. Then $\prod_{n \geq 0} U_n$ is a neighbourhood of (a_n) that does not intersect \mathbb{R}^∞ .

23 T_1 Spaces

Definition 166 (T_1 Space). A topological space is T_1 if and only if every singleton is closed.

Lemma 167. A space is T_1 if and only if every finite set is closed.

PROOF: From Lemma 38. \square

Theorem 168. In a T_1 space, a point a is a limit point of a set A if and only if every neighbourhood of a contains infinitely many points of A .

PROOF:

$\langle 1 \rangle 1$. If a is a limit point of A then every neighbourhood of a contains infinitely many points of A .

$\langle 2 \rangle 1$. ASSUME: a is a limit point of A .

$\langle 2 \rangle 2$. LET: U be a neighbourhood of a .

$\langle 2 \rangle 3$. ASSUME: for a contradiction U contains only finitely many points of A .

$\langle 2 \rangle 4$. $(U \cap A) \setminus \{a\}$ is closed.

PROOF: By the T_1 axiom.

$\langle 2 \rangle 5$. $(U \setminus A) \cup \{a\}$ is open.

PROOF: It is $U \setminus ((U \cap A) \setminus \{a\})$.

$\langle 2 \rangle 6$. $(U \setminus A) \cup \{a\}$ intersects A in a point other than a .

PROOF: From $\langle 2 \rangle 1$.

$\langle 2 \rangle 7$. Q.E.D.

\square

⟨1⟩2. If every neighbourhood of a contains infinitely many points of A then a is a limit point of A .

PROOF: Immediate from definitions.

□

(To see this does not hold in every space, see Proposition 64.)

Proposition 169. *A space is T_1 if and only if, for any two distinct points x and y , there exist neighbourhoods U of x and V of y such that $x \notin V$ and $y \notin U$.*

PROOF:

⟨1⟩1. LET: X be a topological space.

⟨1⟩2. If X is T_1 then, for any two distinct points x and y , there exist neighbourhoods U of x and V of y such that $x \notin V$ and $y \notin U$.

PROOF: This holds because $\{x\}$ and $\{y\}$ are closed.

⟨1⟩3. Suppose, for any two distinct points x and y , there exist neighbourhoods U of x and V of y such that $x \notin V$ and $y \notin U$. Then X is T_1 .

⟨2⟩1. ASSUME: For any two distinct points x and y , there exist neighbourhoods U of x and V of y such that $x \notin V$ and $y \notin U$.

⟨2⟩2. LET: $a \in X$

⟨2⟩3. $\{a\}$ is closed.

PROOF: For all $b \neq a$ there exists a neighbourhood U of b such that $U \subseteq X \setminus \{a\}$.

□

Proposition 170. *A subspace of a T_1 space is T_1 .*

PROOF: From Proposition 148.

24 Hausdorff Spaces

Definition 171 (Hausdorff Space). A topological space is *Hausdorff* if and only if, for any points x, y with $x \neq y$, there exist disjoint open sets U and V such that $x \in U$ and $y \in V$.

Theorem 172. *Every Hausdorff space is T_1 .*

PROOF:

⟨1⟩1. LET: X be a Hausdorff space.

⟨1⟩2. LET: $b \in X$

PROVE: $\overline{\{b\}} = \{b\}$

⟨1⟩3. ASSUME: $a \in \overline{\{b\}}$ and $a \neq b$

⟨1⟩4. PICK disjoint neighbourhoods U of a and V of b .

⟨1⟩5. U intersects $\{b\}$

PROOF: Theorem 52.

⟨1⟩6. $b \in U$

⟨1⟩7. Q.E.D.

PROOF: This contradicts the fact that U and V are disjoint (⟨1⟩4).

□

Proposition 173. *An infinite set under the finite complement topology is T_1 but not Hausdorff.*

PROOF:

⟨1⟩1. LET: X be an infinite set under the finite complement topology.

⟨1⟩2. Every singleton is closed.

PROOF: By definition.

⟨1⟩3. PICK $a, b \in X$ with $a \neq b$

⟨1⟩4. There are no disjoint neighbourhoods U of a and V of b .

⟨2⟩1. LET: U be a neighbourhood of a and V a neighbourhood of b .

⟨2⟩2. $X \setminus U$ and $X \setminus V$ are finite.

⟨2⟩3. PICK $c \in X$ that is not in $X \setminus U$ or $X \setminus V$.

⟨2⟩4. $c \in U \cap V$

□

Proposition 174. *The product of a family of Hausdorff spaces is Hausdorff.*

PROOF:

⟨1⟩1. LET: $\{X_i\}_{i \in I}$ be a family of Hausdorff spaces.

⟨1⟩2. LET: $a, b \in \prod_{i \in I} X_i$ with $a \neq b$

⟨1⟩3. PICK $i \in I$ such that $a_i \neq b_i$

⟨1⟩4. PICK U, V disjoint open sets in X_i with $a_i \in U$ and $b_i \in V$

⟨1⟩5. $\pi_i^{-1}(U)$ and $\pi_i^{-1}(V)$ are disjoint open sets in $\prod_{i \in I} X_i$ with $a \in \pi_i^{-1}(U)$ and $b \in \pi_i^{-1}(V)$

□

Theorem 175. *Every linearly ordered set under the order topology is Hausdorff.*

PROOF:

⟨1⟩1. LET: X be a linearly ordered set under the order topology.

⟨1⟩2. LET: $a, b \in X$ with $a \neq b$

⟨1⟩3. ASSUME: w.l.o.g. $a < b$

⟨1⟩4. CASE: There exists c such that $a < c < b$

PROOF: The sets $(-\infty, c)$ and $(c, +\infty)$ are disjoint neighbourhoods of a and b respectively.

⟨1⟩5. CASE: There is no c such that $a < c < b$

PROOF: The sets $(-\infty, b)$ and $(a, +\infty)$ are disjoint neighbourhoods of a and b respectively.

□

Theorem 176. *A subspace of a Hausdorff space is Hausdorff.*

PROOF:

⟨1⟩1. LET: X be a Hausdorff space and Y a subspace of X .

⟨1⟩2. LET: $x, y \in Y$ with $x \neq y$

⟨1⟩3. PICK disjoint neighbourhoods U of x and V of y in X .

⟨1⟩4. $U \cap Y$ and $V \cap Y$ are disjoint neighbourhoods of x and y respectively in Y .

□

Proposition 177. *A space X is Hausdorff if and only if the diagonal $\Delta = \{(x, x) \mid x \in X\}$ is closed in X^2 .*

PROOF:

X is Hausdorff

$$\Leftrightarrow \forall x, y \in X. x \neq y \Rightarrow \exists V, W \text{ open. } x \in V \wedge y \in W \wedge V \cap W = \emptyset$$

$$\Leftrightarrow \forall (x, y) \in X^2 \setminus \Delta. \exists V, W \text{ open. } (x, y) \subseteq V \times W \subseteq X^2 \setminus \Delta$$

$$\Leftrightarrow \Delta \text{ is closed}$$

□

Theorem 178. *In a Hausdorff space, a sequence has at most one limit.*

PROOF:

⟨1⟩1. LET: X be a Hausdorff space.

⟨1⟩2. ASSUME: for a contradiction $a_n \rightarrow l$ as $n \rightarrow \infty$, $a_n \rightarrow m$ as $n \rightarrow \infty$, and $l \neq m$

⟨1⟩3. PICK disjoint neighbourhoods U of l and V of m

PROOF: By the Hausdorff axiom.

⟨1⟩4. PICK M and N such that $a_n \in U$ for $n \geq M$ and $a_n \in V$ for $n \geq N$

⟨1⟩5. $a_{\max(M, N)} \in U \cap V$

⟨1⟩6. Q.E.D.

PROOF: This contradicts the fact that U and V are disjoint (⟨1⟩3).

□

To see this is not always true in spaces that are T_1 but not Hausdorff:

Proposition 179. *Let X be an infinite set under the finite complement topology. Let $(a_n)_{n \in \mathbb{N}}$ be a sequence with all points distinct. Then for every $l \in X$ we have $a_n \rightarrow l$ as $n \rightarrow \infty$.*

PROOF: Let U be any neighbourhood of l . Since $X \setminus U$ is finite, there must exist N such that, for all $n \geq N$, we have $a_n \in U$. □

Proposition 180. *Let X be a topological space. Let Y a Hausdorff space. Let $A \subseteq X$. Let $f, g : \overline{A} \rightarrow Y$ be continuous. If f and g agree on A then $f = g$.*

PROOF:

⟨1⟩1. LET: $x \in \overline{A}$

⟨1⟩2. ASSUME: $f(x) \neq g(x)$

⟨1⟩3. PICK disjoint neighbourhoods V of $f(x)$ and W of $g(x)$.

⟨1⟩4. PICK $y \in f^{-1}(V) \cap g^{-1}(W) \cap A$

PROOF: Since $f^{-1}(V) \cap g^{-1}(W)$ is a neighbourhood of x and hence intersects A .

⟨1⟩5. $f(y) = g(y) \in V \cap W$

⟨1⟩6. Q.E.D.

PROOF: This contradicts the fact that V and W are disjoint ($\langle 1 \rangle 3$).

□

Proposition 181. *Let $\{X_i\}_{i \in I}$ be a family of Hausdorff spaces. Then $\prod_{i \in I} X_i$ under the box topology is Hausdorff.*

PROOF:

$\langle 1 \rangle 1$. LET: $\{X_i\}_{i \in I}$ be a family of Hausdorff spaces.

$\langle 1 \rangle 2$. LET: $a, b \in \prod_{i \in I} X_i$ with $a \neq b$

$\langle 1 \rangle 3$. PICK $i \in I$ such that $a_i \neq b_i$

$\langle 1 \rangle 4$. PICK U, V disjoint open sets in X_i with $a_i \in U$ and $b_i \in V$

$\langle 1 \rangle 5$. $\pi_i^{-1}(U)$ and $\pi_i^{-1}(V)$ are disjoint open sets in $\prod_{i \in I} X_i$ with $a \in \pi_i^{-1}(U)$ and $b \in \pi_i^{-1}(V)$

□

Proposition 182. *Let \mathcal{T} and \mathcal{T}' be two topologies on the same set X with $\mathcal{T} \subseteq \mathcal{T}'$. If \mathcal{T} is Hausdorff then \mathcal{T}' is Hausdorff.*

PROOF: Immediate from definitions.

Proposition 183. *Let X be a Hausdorff space. Let $\mathcal{D} \subseteq \mathcal{P}X$ be maximal with respect to the finite intersection property. Then $\bigcap_{D \in \mathcal{D}} \overline{D}$ contains at most one point.*

PROOF:

$\langle 1 \rangle 1$. LET: $x, y \in \bigcap_{D \in \mathcal{D}} \overline{D}$

$\langle 1 \rangle 2$. ASSUME: for a contradiction $x \neq y$

$\langle 1 \rangle 3$. PICK disjoint open subsets U and V of x and y respectively.

$\langle 1 \rangle 4$. $U, V \in \mathcal{D}$

PROOF: Proposition 55.

$\langle 1 \rangle 5$. Q.E.D.

PROOF: This contradicts the fact that \mathcal{D} satisfies the finite intersection property.

□

25 The First Countability Axiom

Definition 184 (First Countability Axiom). A topological space X satisfies the *first countability axiom*, or is *first countable*, if and only if every point has a countable local basis.

Lemma 185 (Sequence Lemma (CC)). *Let X be a first countable space. Let $A \subseteq X$ and $l \in \overline{A}$. Then there exists a sequence in A that converges to l .*

PROOF:

$\langle 1 \rangle 1$. PICK a countable local basis $\{B_n \mid n \in \mathbb{Z}^+\}$ at l such that $B_1 \supseteq B_2 \supseteq \dots$.

PROOF: Lemma 78.

$\langle 1 \rangle 2$. For all $n \geq 1$, PICK $a_n \in A \cap B_n$.

PROVE: $a_n \rightarrow l$ as $n \rightarrow \infty$
 <1>3. LET: U be a neighbourhood of A
 <1>4. PICK N such that $B_N \subseteq U$
 <1>5. For $n \geq N$ we have $a_n \in U$
 PROOF: $a_n \in B_n \subseteq B_N \subseteq U$
 \square

Theorem 186 (CC). *Let X be a first countable space and Y a topological space. Let $f : X \rightarrow Y$. Suppose that, for every sequence (x_n) in X and $l \in X$, if $x_n \rightarrow l$ as $n \rightarrow \infty$, then $f(x_n) \rightarrow f(l)$ as $n \rightarrow \infty$. Then f is continuous.*

PROOF:
 <1>1. LET: $A \subseteq X$
 <1>2. LET: $a \in A$
 PROVE: $f(a) \in \overline{f(A)}$
 <1>3. PICK a sequence (x_n) in A that converges to a .
 PROOF: By the Sequence Lemma.
 <1>4. $f(x_n) \rightarrow f(a)$
 <1>5. $f(a) \in \overline{f(A)}$
 PROOF: By Lemma 80.
 <1>6. Q.E.D.
 PROOF: By Theorem 98.
 \square

Example 187 (CC). The space \mathbb{R}^ω under the box product is not first countable.

PROOF: Let $\{B_n \mid n \geq 0\}$ be a countable set of neighbourhoods of $\vec{0}$. We will construct a neighbourhood of $\vec{0}$ that does not include any of these.
 For $n \geq 0$, pick a neighbourhood U_n of 0 such that $U_n \subset \pi_n(B_p)$. Then $\prod_{n=0}^\infty U_n$ is a neighbourhood of $\vec{0}$ that does not include any B_n . \square

Example 188. If J is an uncountable set then \mathbb{R}^J is not first countable.

PROOF:
 <1>1. LET: $\{B_n \mid n \geq 0\}$ be any countable set of neighbourhoods of $\vec{0}$.
 <1>2. For $n \geq 0$, PICK a basis element $\prod_{\alpha \in J} U_{n\alpha}$ that contains $\vec{0}$ and is included in B_n .
 PROOF: Using the Axiom of Countable Choice.
 <1>3. For $n \geq 0$,
 LET: $J_n = \{\alpha \in J \mid U_{n\alpha} \neq \mathbb{R}\}$
 <1>4. PICK $\beta \in J$ such that $\beta \notin J_n$ for any n .
 PROOF: Since each J_n is finite so $\bigcup_n J_n$ is countable.
 <1>5. $\pi_\beta((-1, 1))$ is a neighbourhood of $\vec{0}$ that does not include any B_n .
 \square

Example 189. The space \mathbb{R}_l is first countable.

PROOF: For any $a \in \mathbb{R}$, the set $\{[a, a + 1/n) \mid n \geq 1\}$ is a countable local basis.

Example 190. The ordered square is first countable.

PROOF: For any $(a, b) \in I_o^2$ with $b \neq 0, 1$, the set $\{(\{a\} \times (b - 1/n, b + 1/n)) \cap I_o^2 \mid n \geq 1\}$ is a countable local basis.

26 Strong Continuity

Definition 191 (Strongly Continuous). Let X and Y be topological spaces and $f : X \rightarrow Y$ be a function. Then f is *strongly continuous* if and only if, for every subset $U \subseteq Y$, we have U is open in Y if and only if $f^{-1}(U)$ is open in X .

Proposition 192. Let X and Y be topological spaces and $f : X \rightarrow Y$ be a function. Then f is strongly continuous if and only if, for every subset $C \subseteq Y$, we have C is closed in Y if and only if $f^{-1}(C)$ is closed in X .

PROOF: Since $X \setminus f^{-1}(C) = f^{-1}(Y \setminus C)$. \square

Proposition 193. Let X, Y and Z be topological spaces. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. If f and g are strongly continuous then so is $g \circ f$.

PROOF: Since $(g \circ f)^{-1}(U) = f^{-1}(g^{-1}(U))$. \square

Proposition 194. Let X, Y and Z be topological spaces. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. If $g \circ f$ is continuous and f is strongly continuous then g is continuous.

PROOF:

$\langle 1 \rangle 1$. LET: $V \subseteq Z$ be open.

$\langle 1 \rangle 2$. $f^{-1}(g^{-1}(V))$ is open in X .

PROOF: Since $g \circ f$ is continuous.

$\langle 1 \rangle 3$. $f^{-1}(V)$ is open in Y .

PROOF: Since g is strongly continuous.

\square

Proposition 195. Let X, Y and Z be topological spaces. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. If $g \circ f$ is strongly continuous and f is strongly continuous then g is strongly continuous.

PROOF: For $V \subseteq Z$, we have V is open iff $f^{-1}(g^{-1}(V))$ is open iff $g^{-1}(V)$ is open.

27 Saturated Sets

Definition 196. Let X and Y be sets and $p : X \rightarrow Y$ a surjective function. Let $C \subseteq X$. Then C is *saturated* with respect to p if and only if, for all $x, y \in X$, if $x \in C$ and $p(x) = p(y)$ then $y \in C$.

Proposition 197. Let X and Y be sets and $p : X \rightarrow Y$ a surjective function. Let $C \subseteq X$. Then the following are equivalent:

1. C is saturated with respect to p .
2. There exists $D \subseteq Y$ such that $C = p^{-1}(D)$
3. $C = p^{-1}(p(C))$.

PROOF:

$\langle 1 \rangle 1. 1 \Rightarrow 3$

$\langle 2 \rangle 1.$ ASSUME: C is saturated with respect to p .

$\langle 2 \rangle 2. C \subseteq p^{-1}(p(C))$

PROOF: Trivial.

$\langle 2 \rangle 3. p^{-1}(p(C)) \subseteq C$

$\langle 3 \rangle 1.$ LET: $x \in p^{-1}(p(C))$

$\langle 3 \rangle 2. p(x) \in p(C)$

$\langle 3 \rangle 3.$ There exists $y \in C$ such that $p(x) = p(y)$

$\langle 3 \rangle 4. x \in C$

PROOF: From $\langle 2 \rangle 1.$

$\langle 1 \rangle 2. 3 \Rightarrow 2$

PROOF: Trivial.

$\langle 1 \rangle 3. 2 \Rightarrow 1$

PROOF: This follows because if $p(x) \in D$ and $p(x) = p(y)$ then $p(y) \in D$.

□

28 Quotient Maps

Definition 198 (Quotient Map). Let X and Y be topological spaces and $p : X \rightarrow Y$. Then p is a *quotient map* if and only if p is surjective and strongly continuous.

Proposition 199. Let X and Y be topological spaces and $p : X \rightarrow Y$ be a surjective function. Then the following are equivalent.

1. p is a quotient map.
2. p is continuous and maps saturated open sets to open sets.
3. p is continuous and maps saturated closed sets to closed sets.

PROOF:

$\langle 1 \rangle 1. 1 \Rightarrow 2$

$\langle 2 \rangle 1.$ ASSUME: p is a quotient map.

$\langle 2 \rangle 2.$ LET: U be a saturated open set in X .

$\langle 2 \rangle 3. p^{-1}(p(U))$ is open in X .

PROOF: Since $U = p^{-1}(p(U))$ be Proposition 197.

$\langle 2 \rangle 4. p(U)$ is open in Y .

PROOF: From $\langle 2 \rangle 1.$

$\langle 1 \rangle 2. 1 \Rightarrow 3$

PROOF: Similar.

- ⟨1⟩3. $2 \Rightarrow 1$
 - ⟨2⟩1. ASSUME: p is continuous and maps saturated open sets to open sets.
 - ⟨2⟩2. LET: $U \subseteq Y$
 - ⟨2⟩3. ASSUME: $p^{-1}(U)$ is open in X
 - ⟨2⟩4. $p^{-1}(U)$ is saturated.
 - PROOF: Proposition 197.
 - ⟨2⟩5. U is open in Y .
 - ⟨1⟩4. $3 \Rightarrow 1$
 - PROOF: Similar.
-

Corollary 199.1. *Every surjective continuous open map is a quotient map.*

Corollary 199.2. *Every surjective continuous closed map is a quotient map.*

Example 200. The converses of these corollaries do not hold.

Let $A = \{(x, y) \mid x \geq 0\} \cup \{(x, y) \mid y = 0\}$. Then $\pi_1 : A \rightarrow \mathbb{R}$ is a quotient map, but not an open map or a closed map.

We prove that π_1 maps saturated open sets to open sets:

- ⟨1⟩1. LET: $\pi_1^{-1}(U)$ be a saturated open set in A
- PROVE: U is open in \mathbb{R}
- ⟨1⟩2. LET: $x \in U$
- ⟨1⟩3. $(x, 0) \in \pi_1(U)^{-1}$
- ⟨1⟩4. PICK W, V open in \mathbb{R} such that $(x, 0) \subseteq W \times V \subseteq \pi_1(U)^{-1}$
- ⟨1⟩5. $x \in W \subseteq U$

It is not an open map because it maps $((-1, 1) \times (1, 2)) \cap A$ to $[0, 1)$.

It is not a closed map because it maps $\{(x, 1/x) \mid x > 0\}$ to $(0, +\infty)$.

Proposition 201. *Let $p : X \twoheadrightarrow Y$ be a quotient map. Let $A \subseteq X$ be saturated with respect to p . Let $q : A \twoheadrightarrow p(A)$ be the restriction of p .*

- 1. *If A is either open or closed in X then q is a quotient map.*
- 2. *If p is either an open map or a closed map then q is a quotient map.*

PROOF:

- ⟨1⟩1. LET: $p : X \twoheadrightarrow Y$ be a quotient map.
- ⟨1⟩2. LET: $A \subseteq X$ be saturated with respect to p .
- ⟨1⟩3. LET: $q : A \twoheadrightarrow p(A)$ be the restriction of p .
- ⟨1⟩4. q is continuous.
- PROOF: Theorem 102.
- ⟨1⟩5. If A is open in X then q is a quotient map.
 - ⟨2⟩1. ASSUME: A is open in X .
 - ⟨2⟩2. q maps saturated open sets to open sets.
 - ⟨3⟩1. LET: $U \subseteq A$ be saturated with respect to q and open in A
 - ⟨3⟩2. U is saturated with respect to p
 - ⟨4⟩1. LET: $x, y \in X$

1

Define $f : [0, 1] \rightarrow [2, 3] \rightarrow [0, 2]$ by $f(x) = x$ if $x \leq 1$, $f(x) = x - 1$ if $x \geq 2$. Then f is a quotient map but its restriction f' to $[0, 1) \cup [2, 3]$ is not, because

$f'^{-1}([1, 2])$ is open but $[1, 2]$ is not.

For a counterexample where A is saturated, see Example 208.

Proposition 203. *Let $p : A \twoheadrightarrow C$ and $q : B \twoheadrightarrow D$ be open quotient maps. Then $p \times q : A \times B \rightarrow C \times D$ is an open quotient map.*

PROOF: From Corollary 199.1, Proposition 141 and Theorem 137. \square

Theorem 204. *Let $p : X \twoheadrightarrow Y$ be a quotient map. Let Z be a topological space and $f : Y \rightarrow Z$ be a function. Then*

1. $f \circ p$ is continuous if and only if f is continuous.
2. $f \circ p$ is a quotient map if and only if f is a quotient map.

PROOF:

$\langle 1 \rangle 1$. If $f \circ p$ is continuous then f is continuous.

PROOF: Proposition 194.

$\langle 1 \rangle 2$. If f is continuous then $f \circ p$ is continuous.

PROOF: Theorem 101.

$\langle 1 \rangle 3$. If $f \circ p$ is a quotient map then f is a quotient map.

PROOF: Proposition 195.

$\langle 1 \rangle 4$. If f is a quotient map then $f \circ p$ is a quotient map.

PROOF: From Proposition 193.

\square

Proposition 205. *Let X and Y be topological spaces. Let $p : X \rightarrow Y$ and $f : Y \rightarrow X$ be continuous maps such that $p \circ f = \text{id}_Y$. Then p is a quotient map.*

PROOF:

$\langle 1 \rangle 1$. LET: $V \subseteq Y$

$\langle 1 \rangle 2$. ASSUME: $p^{-1}(V)$ is open in X .

$\langle 1 \rangle 3$. $f^{-1}(p^{-1}(V))$ is open in Y .

PROOF: Because f is continuous.

$\langle 1 \rangle 4$. V is open in Y .

PROOF: Because $f^{-1}(p^{-1}(V)) = V$.

\square

29 Quotient Topology

Definition 206 (Quotient Topology). Let X be a topological space, Y a set and $p : X \twoheadrightarrow Y$ be a surjective function. Then the *quotient topology* on Y is the unique topology on Y with respect to which p is a quotient map, namely $\mathcal{T} = \{U \in \mathcal{P}Y \mid p^{-1}(U) \text{ is open in } X\}$.

We prove this is a topology.

PROOF:

⟨1⟩1. $Y \in \mathcal{T}$

PROOF: Since $p^{-1}(Y) = X$ by surjectivity.

⟨1⟩2. For all $\mathcal{A} \subseteq \mathcal{T}$ we have $\bigcup \mathcal{A} \in \mathcal{T}$

PROOF: Since $p^{-1}(\bigcup \mathcal{A}) = \bigcup_{U \in \mathcal{A}} p^{-1}(U)$

⟨1⟩3. For all $U, V \in \mathcal{T}$ we have $U \cap V \in \mathcal{T}$

PROOF: Since $p^{-1}(U \cap V) = p^{-1}(U) \cap p^{-1}(V)$.

□

Definition 207 (Quotient Space). Let X be a topological space and \sim an equivalence relation on X . Let $p : X \rightarrow X/\sim$ be the canonical surjection. Then X/\sim under the quotient topology is called a *quotient space*, *identification space* or *decomposition space* of X .

Here is a counterexample showing we cannot remove all the hypotheses of Proposition 201 except that A is saturated.

Example 208. Let $X = (0, 1/2] \cup \{1\} \cup \{1 + 1/n : n \geq 2\}$ as a subspace of \mathbb{R} . Define R to be the equivalence relation on X where xRy iff $(x = y \text{ or } |x - y| = 1)$, so we identify $1/n$ with $1 + 1/n$ for all $n \geq 2$. Let Y be the resulting quotient space X/R in the quotient topology and $p : X \rightarrow Y$ the canonical surjection.

Let $A = \{1\} \cup (0, 1/2] \setminus \{1/n : n \geq 2\} \subseteq X$. Then A is saturated under p but the restriction q of p to A is not a quotient map because it maps the saturated open set $\{1\}$ to $\{1\}$ which is not open in $p(A)$.

Proposition 209. Let X, Y and Z be topological spaces. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. If f and g are quotient maps then so is $g \circ f$.

PROOF: From Proposition 193. □

Example 210. The product of two quotient maps is not necessarily a quotient map.

Let $X = \mathbb{R}$ and X^* the quotient space formed by identifying all positive integers. Let $p : X \rightarrow X^*$ be the canonical surjection.

We prove $p \times \text{id}_{\mathbb{Q}} : X \times \mathbb{Q} \rightarrow X^* \times \mathbb{Q}$ is not a quotient map.

PROOF:

⟨1⟩1. For $n \geq 1$,

LET: $c_n = \sqrt{2}/n$

⟨1⟩2. For $n \geq 1$,

LET: $U_n = \{(x, y) \in X \times \mathbb{Q} \mid n - 1/4 < x < n + 1/4, (y + n > x + c_n \text{ and } y + n > -x + c_n) \text{ or } (y + n < x + c_n \text{ and } y + n < -x + c_n)\}$

⟨1⟩3. For $n \geq 1$, we have U_n is open in $X \times \mathbb{Q}$

⟨1⟩4. For $n \geq 1$, we have $\{n\} \times \mathbb{Q} \subseteq U_n$

⟨1⟩5. LET: $U = \bigcup_{n=1}^{\infty} U_n$

⟨1⟩6. U is open in $X \times \mathbb{Q}$

⟨1⟩7. U is saturated with respect to $p \times \text{id}_{\mathbb{Q}}$

⟨1⟩8. LET: $U' = (p \times \text{id}_{\mathbb{Q}})(U)$

⟨1⟩9. ASSUME: for a contradiction U' is open in $X^* \times \mathbb{Q}$

⟨1⟩10. $(1, 0) \in U'$

- ⟨1⟩11. PICK a neighbourhood W of 1 in X^* and $\delta > 0$ such that $W \times (-\delta, \delta) \subseteq U'$
- ⟨1⟩12. $p^{-1}(W) \times (-\delta, \delta) \subseteq U$
- ⟨1⟩13. PICK n such that $c_n < \delta$
- ⟨1⟩14. $n \in p^{-1}(W)$
- ⟨1⟩15. PICK $\epsilon > 0$ such that $\epsilon < \delta - c_n$ and $\epsilon < 1/4$ and $(n - \epsilon, n + \epsilon) \subseteq p^{-1}(W)$
- ⟨1⟩16. $(n - \epsilon, n + \epsilon) \times (-\delta, \delta) \subseteq U$
- ⟨1⟩17. PICK a rational y such that $c_n - \epsilon/2 < y < c_n + \epsilon/2$
- ⟨1⟩18. $(n + \epsilon/2, y) \notin U$
- ⟨1⟩19. Q.E.D.

PROOF: This contradicts ⟨1⟩16.

□

Proposition 211. *Let X be a topological space and \sim an equivalence relation on X . Then X/\sim is T_1 if and only if every equivalence class is closed in X .*

PROOF: Immediate from definitions. □

30 Retractions

Definition 212 (Retraction). Let X be a topological space and $A \subseteq X$. A *retraction* of X onto A is a continuous map $r : X \rightarrow A$ such that, for all $a \in A$, we have $r(a) = a$.

Proposition 213. *Every retraction is a quotient map.*

PROOF: Proposition 205 with f the inclusion $A \hookrightarrow X$. □

31 Homogeneous Spaces

Definition 214 (Homogeneous). A topological space X is *homogeneous* if and only if, for any points $a, b \in X$, there exists a homeomorphism $\phi : X \cong X$ such that $\phi(a) = b$.

32 Regular Spaces

Definition 215 (Regular Space). A topological space X is *regular* if and only if, for any closed set A and point $a \notin A$, there exist disjoint open sets U, V such that $A \subseteq U$ and $a \in V$.

33 Connected Spaces

Definition 216 (Separation). A *separation* of a topological space X is a pair of disjoint open sets U, V such that $U \cup V = \emptyset$.

Definition 217 (Connected). A topological space is *connected* if and only if it has no separation; otherwise it is *disconnected*.

Proposition 218. *A topological space X is connected if and only if the only sets that are both open and closed are X and \emptyset .*

Immediate from definitions.

Lemma 219. *If Y is a subspace of X , a separation of Y is a pair of disjoint nonempty sets A and B whose union is Y , neither of which contains a limit point of the other.*

PROOF:

- $\langle 1 \rangle 1.$ LET: $A, B \subseteq Y$
- $\langle 1 \rangle 2.$ If A and B form a separation of Y then A and B are disjoint and nonempty, $A \cup B = Y$, and neither of A and B contains a limit point of the other.
 - $\langle 2 \rangle 1.$ ASSUME: A and B form a separation of Y
 - $\langle 2 \rangle 2.$ A and B are disjoint and nonempty and $A \cup B = Y$
 - PROOF: From $\langle 2 \rangle 1$ and the definition of separation.
 - $\langle 2 \rangle 3.$ A does not contain a limit point of B
 - $\langle 3 \rangle 1.$ ASSUME: for a contradiction $l \in A$ and l is a limit point of B in X .
 - $\langle 3 \rangle 2.$ l is a limit point of B in Y
 - PROOF: Proposition 157.
 - $\langle 3 \rangle 3.$ $l \in B$
 - $\langle 4 \rangle 1.$ B is closed in Y
 - PROOF: Since A is open in Y and $B = Y \setminus A$ from $\langle 2 \rangle 1$.
 - $\langle 4 \rangle 2.$ Q.E.D.
 - PROOF: Corollary 63.1.
 - $\langle 3 \rangle 4.$ Q.E.D.
 - PROOF: This contradicts the fact that $A \cap B = \emptyset$ ($\langle 2 \rangle 1$).
 - $\langle 2 \rangle 4.$ B does not contain a limit point of A
 - PROOF: Similar.
- $\langle 1 \rangle 3.$ If A and B are disjoint and nonempty, $A \cup B = Y$, and neither of A and B contains a limit point of the other, then A and B form a separation of Y .
 - $\langle 2 \rangle 1.$ ASSUME: A and B are disjoint and nonempty, $A \cup B = Y$, and neither of A and B contains a limit point of the other.
 - $\langle 2 \rangle 2.$ A is open in Y
 - $\langle 3 \rangle 1.$ B is closed in Y
 - $\langle 4 \rangle 1.$ LET: l be a limit point of B in Y
 - $\langle 4 \rangle 2.$ l is a limit point of B in X
 - PROOF: Proposition 157.
 - $\langle 4 \rangle 3.$ $l \notin A$
 - PROOF: By $\langle 2 \rangle 1$
 - $\langle 4 \rangle 4.$ $l \in B$
 - PROOF: By $\langle 2 \rangle 1$ since $A \cup B = Y$
 - $\langle 4 \rangle 5.$ Q.E.D.
 - PROOF: Corollary 63.1.
 - $\langle 3 \rangle 2.$ Q.E.D.
 - PROOF: Since $A = Y \setminus B$.

⟨2⟩3. B is open in Y

PROOF: Similar.

□

Example 220. Every set under the indiscrete topology is connected.

Example 221. The discrete topology on a set X is connected if and only if $|X| \leq 1$.

Example 222. The finite complement topology on a set X is connected if and only if either $|X| \leq 1$ or X is infinite.

Example 223. The countable complement topology on a set X is connected if and only if either $|X| \leq 1$ or X is uncountable.

Example 224. The rationals \mathbb{Q} are disconnected. For any irrational a , the sets $(-\infty, a) \cap \mathbb{Q}$ and $(a, +\infty) \cap \mathbb{Q}$ form a separation of \mathbb{Q} .

Lemma 225. Let X be a topological space. If C and D form a separation of X , and Y is a connected subspace of X , then either $Y \subseteq C$ or $Y \subseteq D$.

PROOF: Otherwise $Y \cap C$ and $Y \cap D$ would form a separation of Y . □

Theorem 226. The union of a set of connected subspaces of a space X that have a point in common is connected.

PROOF:

⟨1⟩1. LET: \mathcal{A} be a set of connected subspaces of the space X that have the point a in common.

⟨1⟩2. ASSUME: for a contradiction C and D form a separation of $\bigcup \mathcal{A}$

⟨1⟩3. ASSUME: without loss of generality $a \in C$

⟨1⟩4. For all $A \in \mathcal{A}$ we have $A \subseteq C$

PROOF: Lemma 225.

⟨1⟩5. $D = \emptyset$

⟨1⟩6. Q.E.D.

PROOF: This contradicts ⟨1⟩2.

□

Theorem 227. Let X be a topological space and A a connected subspace of X . If $A \subseteq B \subseteq \overline{A}$ then B is connected.

PROOF:

⟨1⟩1. ASSUME: for a contradiction C and D form a separation of B .

⟨1⟩2. ASSUME: without loss of generality $A \subseteq C$

PROOF: Lemma 225.

⟨1⟩3. $B \subseteq C$

⟨2⟩1. LET: $x \in B$

⟨2⟩2. $x \in \overline{A}$

⟨2⟩3. Either $x \in A$ or x is a limit point of A .

PROOF: Theorem 63.

⟨2⟩4. Either $x \in A$ or x is a limit point of C .

PROOF: Lemma 65, ⟨1⟩2.

⟨2⟩5. $x \in C$

PROOF: Lemma 219.

⟨1⟩4. $D = \emptyset$

⟨1⟩5. Q.E.D.

PROOF: This contradicts ⟨1⟩1.

□

Theorem 228. *The image of a connected space under a continuous map is connected.*

PROOF:

⟨1⟩1. LET: $f : X \rightarrow Y$ be a surjective continuous map where X is connected.

⟨1⟩2. ASSUME: for a contradiction C and D form a separation of Y .

⟨1⟩3. $f^{-1}(C)$ and $f^{-1}(D)$ form a separation of X .

□

Theorem 229. *The product of a family of connected spaces is connected.*

PROOF:

⟨1⟩1. The product of two connected spaces is connected.

⟨2⟩1. LET: X and Y be connected spaces.

⟨2⟩2. PICK $a \in X$ and $b \in Y$

PROOF: We may assume X and Y are nonempty since otherwise $X \times Y = \emptyset$ which is connected.

⟨2⟩3. $X \times \{b\}$ is connected.

PROOF: It is homeomorphic to X .

⟨2⟩4. For all $x \in X$ we have $\{x\} \times Y$ is connected.

PROOF: It is homeomorphic to Y .

⟨2⟩5. For any $x \in X$

LET: $T_x = (X \times \{b\}) \cup (\{x\} \times Y)$

⟨2⟩6. For all $x \in X$, T_x is connected.

PROOF: Theorem 226 since $(x, b) \in (X \times \{b\}) \cap (\{x\} \times Y)$.

⟨2⟩7. $X \times Y$ is connected.

PROOF: Theorem 226 since $X \times Y = \bigcup_{x \in X} T_x$ and (a, b) is a point in every T_x .

⟨1⟩2. The product of a finite family of connected spaces is connected.

PROOF: From ⟨1⟩1 by induction.

⟨1⟩3. The product of any family of connected spaces is connected.

⟨2⟩1. LET: $\{X_\alpha\}_{\alpha \in J}$ be a family of connected spaces.

⟨2⟩2. LET: $X = \prod_{\alpha \in J} X_\alpha$

⟨2⟩3. PICK $a \in X$

PROOF: We may assume $X \neq \emptyset$ as the empty space is connected.

⟨2⟩4. For every finite subset K of J ,

LET: $X_K = \{x \in X \mid \forall \alpha \in J \setminus K. x_\alpha = a_\alpha\}$

⟨2⟩5. For every finite $K \subseteq J$, we have X_K is connected.

PROOF: From $\langle 1 \rangle 2$ since $X_K \cong \prod_{\alpha \in K} X_K$.

$\langle 2 \rangle 6$. LET: $Y = \bigcup_K X_K$

$\langle 2 \rangle 7$. Y is connected

PROOF: Theorem 226 since a is a common point.

$\langle 2 \rangle 8$. $X = \bar{Y}$

$\langle 3 \rangle 1$. LET: $x \in X$

$\langle 3 \rangle 2$. LET: $U = \prod_{\alpha \in J} U_\alpha$ be a basic neighbourhood of x where $U_\alpha = X_\alpha$ for all α except $\alpha \in K$ for some finite $K \subseteq J$

$\langle 3 \rangle 3$. LET: $y \in X$ be the point with $y_\alpha = x_\alpha$ for $\alpha \in K$ and $y_\alpha = a_\alpha$ for all other α

$\langle 3 \rangle 4$. $y \in U \cap X_K$

$\langle 3 \rangle 5$. $y \in U \cap Y$

$\langle 2 \rangle 9$. X is connected.

PROOF: Theorem 227.

□

Example 230. The set \mathbb{R}^ω is disconnected under the box topology. The set of bounded sequences and the set of unbounded sequences form a separation.

Proposition 231. Let \mathcal{T} and \mathcal{T}' be two topologies on the same set X . If $\mathcal{T} \subseteq \mathcal{T}'$ and (X, \mathcal{T}') is connected then (X, \mathcal{T}) is connected.

PROOF: If U and V form a separation of (X, \mathcal{T}) then they form a separation of (X, \mathcal{T}') . □

Proposition 232. Let X be a topological space and (A_n) a sequence of connected subspaces of X . If $A_n \cap A_{n+1} \neq \emptyset$ for all n then $\bigcup_n A_n$ is connected.

PROOF:

$\langle 1 \rangle 1$. ASSUME: for a contradiction C and D form a separation of $\bigcup_n A_n$

$\langle 1 \rangle 2$. ASSUME: without loss of generality $A_0 \subseteq C$

PROOF: Lemma 225.

$\langle 1 \rangle 3$. For all n we have $A_n \subseteq C$

PROOF:

$\langle 2 \rangle 1$. ASSUME: $A_n \subseteq C$

$\langle 2 \rangle 2$. PICK $x \in A_n \cap A_{n+1}$

$\langle 2 \rangle 3$. $x \in C$

$\langle 2 \rangle 4$. $A_{n+1} \subseteq C$

PROOF: Lemma 225.

$\langle 2 \rangle 5$. Q.E.D.

PROOF: The result follows by induction.

$\langle 1 \rangle 4$. $D = \emptyset$

$\langle 1 \rangle 5$. Q.E.D.

PROOF: This contradicts $\langle 1 \rangle 1$.

□

Proposition 233. Let X be a topological space. Let $A, C \subseteq X$. If C is connected and intersects both A and $X \setminus A$ then C intersects ∂A .

PROOF: Otherwise $C \cap A^\circ$ and $C \setminus \overline{A}$ would form a separation of C . \square

Example 234. The space \mathbb{R}_l is disconnected. For any real x , the sets $(-\infty, x)$ and $[x, +\infty)$ form a separation.

Proposition 235. *Let X and Y be connected spaces. Let A be a proper subset of X and B a proper subset of Y . Then $(X \times Y) \setminus (A \times B)$ is connected.*

PROOF:

$\langle 1 \rangle 1$. PICK $a \in X \setminus A$ and $b \in Y \setminus B$

$\langle 1 \rangle 2$. For $x \in X \setminus A$ we have $(X \times \{b\}) \cup (\{x\} \times Y)$ is connected.

PROOF: Theorem 226 since (x, b) is a common point.

$\langle 1 \rangle 3$. For $y \in Y \setminus B$ we have $(X \times \{y\}) \cup (\{a\} \times Y)$ is connected.

PROOF: Theorem 226 since (a, y) is a common point.

$\langle 1 \rangle 4$. $(X \times Y) \setminus (A \times B)$ is connected.

PROOF: Theorem 226 since it is the union of the sets in $\langle 1 \rangle 2$ and $\langle 1 \rangle 3$ with (a, b) as a common point.

\square

Proposition 236. *Let $p : X \rightarrow Y$ be a quotient map. If Y is connected and $p^{-1}(y)$ is connected for all $y \in Y$, then X is connected.*

PROOF:

$\langle 1 \rangle 1$. ASSUME: for a contradiction C and D form a separation of X .

$\langle 1 \rangle 2$. C is saturated.

$\langle 2 \rangle 1$. LET: $x \in C$, $y \in X$ with $p(x) = p(y) = a$, say

$\langle 2 \rangle 2$. $y \notin D$

PROOF: Otherwise $p^{-1}(a) \cap C$ and $p^{-1}(a) \cap D$ form a separation of $p^{-1}(a)$.

$\langle 2 \rangle 3$. $y \in C$

$\langle 1 \rangle 3$. D is saturated.

PROOF: Similar.

$\langle 1 \rangle 4$. $p(C)$ and $p(D)$ form a separation of Y .

\square

Proposition 237. *Let X be a connected space and Y a connected subspace of X . Suppose A and B form a separation of $X \setminus Y$. Then $Y \cup A$ and $Y \cup B$ are both connected.*

PROOF:

$\langle 1 \rangle 1$. $Y \cup A$ is connected.

$\langle 2 \rangle 1$. ASSUME: for a contradiction C and D form a separation of $Y \cup A$

$\langle 2 \rangle 2$. ASSUME: without loss of generality $Y \subseteq C$

$\langle 2 \rangle 3$. PICK open sets A_1, B_1, C_1, D_1 in X with

$$A = A_1 \setminus Y$$

$$B = B_1 \setminus Y$$

$$C = C_1 \cap (Y \cup A)$$

$$D = D_1 \cap (Y \cup A)$$

$\langle 2 \rangle 4$. $B_1 \cup C_1$ and $A_1 \cap D_1$ form a separation of X

⟨1⟩2. $Y \cup B$ is connected.

PROOF: Similar.

□

Theorem 238. *Let L be a linearly ordered set under the order topology. Then L is connected if and only if L is a linear continuum.*

PROOF:

⟨1⟩1. If L is a linear continuum then L is connected.

⟨2⟩1. LET: L be a linear continuum under the order topology.

⟨2⟩2. ASSUME: for a contradiction C and D form a separation of L .

⟨2⟩3. PICK $a \in C$ and $b \in D$.

⟨2⟩4. ASSUME: without loss of generality $a < b$.

⟨2⟩5. LET: $S = \{x \in L \mid a < x \text{ and } [a, x) \subseteq C\}$

⟨2⟩6. S is nonempty.

PROOF: Since $a \in C$ and C is open.

⟨2⟩7. S is bounded above by b .

PROOF: Since $b \notin C$.

⟨2⟩8. LET: $s = \sup S$

⟨2⟩9. $s \in S$

⟨3⟩1. LET: $y \in [a, s)$

PROVE: $y \in C$

⟨3⟩2. PICK z with $y < z \in S$

PROOF: By minimality of s .

⟨3⟩3. $y \in [a, z) \subseteq C$

⟨2⟩10. CASE: $s \in C$

⟨3⟩1. PICK x such that $s < x$ and $[s, x) \subseteq C$

PROOF: Since C is open and s is not greatest in L because $s < b$.

⟨3⟩2. $x \in S$

PROOF: Since $[a, x) = [a, s) \cup [s, x) \subseteq C$.

⟨3⟩3. Q.E.D.

PROOF: This contradicts the fact that s is an upper bound for S .

⟨2⟩11. CASE: $s \in D$

⟨3⟩1. PICK $x < s$ such that $(x, s] \subseteq D$

⟨3⟩2. PICK y with $x < y < s$

PROOF: Since L is dense.

⟨3⟩3. $y \in C$

PROOF: From ⟨2⟩9.

⟨3⟩4. $y \in D$

PROOF: From ⟨3⟩1.

⟨3⟩5. Q.E.D.

⟨3⟩6. LET: L be a linear continuum under the order topology.

⟨3⟩7. ASSUME: for a contradiction C and D form a separation of L .

⟨3⟩8. PICK $a \in C$ and $b \in D$.

⟨3⟩9. ASSUME: without loss of generality $a < b$.

⟨3⟩10. LET: $S = \{x \in L \mid a < x \text{ and } [a, x) \subseteq C\}$

⟨3⟩11. S is nonempty.

PROOF: Since $a \in C$ and C is open.

⟨3⟩12. S is bounded above by b .
PROOF: Since $b \notin C$.

⟨3⟩13. LET: $s = \sup S$

⟨3⟩14. $s \in S$
⟨4⟩1. LET: $y \in [a, s)$
PROVE: $y \in C$

⟨4⟩2. PICK z with $y < z \in S$
PROOF: By minimality of s .

⟨4⟩3. $y \in [a, z) \subseteq C$

⟨3⟩15. CASE: $s \in C$
⟨4⟩1. PICK x such that $s < x$ and $[s, x) \subseteq C$
PROOF: Since C is open and s is not greatest in L because $s < b$.

⟨4⟩2. $x \in S$
PROOF: Since $[a, x) = [a, s) \cup [s, x) \subseteq C$.

⟨4⟩3. Q.E.D.
PROOF: This contradicts the fact that s is an upper bound for S .

⟨3⟩16. CASE: $s \in D$
⟨4⟩1. PICK $x < s$ such that $(x, s] \subseteq D$
⟨4⟩2. PICK y with $x < y < s$
PROOF: Since L is dense.

⟨4⟩3. $y \in C$
PROOF: From ⟨2⟩9.

⟨4⟩4. $y \in D$
PROOF: From ⟨3⟩1.

⟨4⟩5. Q.E.D.
PROOF: This contradicts ⟨2⟩2.

⟨1⟩2. If L is connected then L is a linear continuum.

⟨2⟩1. ASSUME: L is connected.

⟨2⟩2. Every nonempty subset of L that is bounded above has a supremum.

⟨3⟩1. LET: X be a nonempty subset of L bounded above by b .

⟨3⟩2. ASSUME: for a contradiction X has no supremum.

⟨3⟩3. LET: U be the set of upper bounds of X ,

⟨3⟩4. U is nonempty.
PROOF: Since $b \in U$.

⟨3⟩5. U is open.
⟨4⟩1. LET: $x \in U$
⟨4⟩2. PICK an upper bound y for X such that $y < x$
⟨4⟩3. Either x is greatest in L and $(y, x] \subseteq U$, or there exists $z > x$ such that $(y, z) \subseteq U$

⟨3⟩6. LET: V be the set of lower bounds of U .

⟨3⟩7. V is nonempty.
PROOF: Since $X \subseteq V$

⟨3⟩8. V is open.
⟨4⟩1. LET: $x \in V$
⟨4⟩2. PICK $y \in X$ with $x < y$

PROOF: x cannot be an upper bound for X , because it would be the supremum of X .

$\langle 4 \rangle 3$. Either x least in L and $[x, y) \subseteq V$, or there exists $z < x$ such that $(z, y) \subseteq V$

$\langle 3 \rangle 9$. $L = U \cup V$

$\langle 4 \rangle 1$. LET: $x \in L \setminus U$

$\langle 4 \rangle 2$. PICK $y \in X$ such that $x < y$

$\langle 4 \rangle 3$. For all $u \in U$ we have $x < y \leq u$

$\langle 4 \rangle 4$. $x \in V$

$\langle 3 \rangle 10$. $U \cap V = \emptyset$

PROOF: Any element of $U \cap V$ would be a supremum of X .

$\langle 3 \rangle 11$. U and V form a separation of L .

$\langle 3 \rangle 12$. Q.E.D.

PROOF: This contradicts $\langle 2 \rangle 1$.

$\langle 2 \rangle 3$. L is dense.

$\langle 3 \rangle 1$. LET: $x, y \in L$ with $x < y$

$\langle 3 \rangle 2$. There exists $z \in L$ such that $x < z < y$

PROOF: Otherwise $(-\infty, y)$ and $(x, +\infty)$ would form a separation of L .

□

Corollary 238.1. *The real line \mathbb{R} is connected.*

Corollary 238.2. *Every interval in \mathbb{R} is connected.*

Corollary 238.3. *The ordered square is connected.*

Theorem 239 (Intermediate Value Theorem). *Let X be a connected space. Let Y be a linearly ordered set under the order topology. Let $f : X \rightarrow Y$ be continuous. Let $a, b \in X$ and $r \in Y$. Suppose $f(a) < r < f(b)$. Then there exists $c \in X$ such that $f(c) = r$.*

PROOF: Otherwise $f^{-1}((-\infty, r))$ and $f^{-1}((r, +\infty))$ would form a separation of X . □

Proposition 240. *Every function $f : [0, 1] \rightarrow [0, 1]$ has a fixed point.*

PROOF:

$\langle 1 \rangle 1$. LET: $g : [0, 1] \rightarrow [-1, 1]$ be the function $g(x) = f(x) - x$

PROVE: there exists $x \in [0, 1]$ such that $g(x) = 0$

$\langle 1 \rangle 2$. ASSUME: without loss of generality $g(0) \neq 0$ and $g(1) \neq 0$

$\langle 1 \rangle 3$. $g(0) > 0$

$\langle 1 \rangle 4$. $g(1) < 0$

$\langle 1 \rangle 5$. There exists $x \in (0, 1)$ such that $g(x) = 0$

PROOF: By the Intermediate Value Theorem.

Proposition 241. *Give \mathbb{R}^ω the box topology. Let $x, y \in \mathbb{R}^\omega$. Then x and y lie in the same component if and only if $x - y$ is eventually zero, i.e. there exists N such that, for all $n \geq N$, we have $x_n = y_n$.*

PROOF:

- ⟨1⟩1. The component containing 0 is the set of sequences that are eventually zero.
- ⟨2⟩1. LET: B be the set of sequences that are eventually zero.
- ⟨2⟩2. B is path-connected.
- ⟨3⟩1. LET: $x, y \in B$
- ⟨3⟩2. PICK N such that, for all $n \geq N$, we have $x_n = y_n = 0$
- ⟨3⟩3. LET: $p : [0, 1] \rightarrow \mathbb{R}^\omega$, $p(t) = (1 - t)x + ty$
PROVE: p is continuous.
- ⟨3⟩4. LET: $t \in [0, 1]$ and $\prod_j U_j$ be a basic open neighbourhood of $p(t)$, where each U_j is open in \mathbb{R}
- ⟨3⟩5. PICK δ such that, for all $n < N$ and all $s \in [0, 1]$, if $|s - t| < \delta$ then $p(s)_n \in U_n$
- ⟨3⟩6. For all $s \in [0, 1]$, if $|s - t| < \delta$ then $p(s) \in \prod_j U_j$
- ⟨2⟩3. B is connected.
PROOF: Proposition 247.
- ⟨2⟩4. If C is connected and $B \subseteq C$ then $B = C$.
- ⟨3⟩1. ASSUME: C is connected and $B \subseteq C$
- ⟨3⟩2. ASSUME: for a contradiction $x \in C \setminus B$
- ⟨3⟩3. For $n \geq 1$,
LET: $c_n = 1$ if $x_n = 0$, $c_n = n/x_n$ otherwise
- ⟨3⟩4. LET: $h : \mathbb{R}^\omega \rightarrow \mathbb{R}^\omega$ be the function $h(x) = (c_n x_n)_{n \geq 1}$
- ⟨3⟩5. h is a homeomorphism of \mathbb{R}^ω with itself.
- ⟨3⟩6. $h(x)$ is unbounded.
PROOF: For any $b > 0$, pick $N > b$ such that $x_N \neq 0$. Then $h(x)_N > b$.
- ⟨3⟩7. $h^{-1}(\{\text{bounded sequences}\}) \cap C$ and $h^{-1}(\{\text{unbounded sequences}\}) \cap C$ form a separation of C
- ⟨3⟩8. Q.E.D.
PROOF: This contradicts ⟨3⟩1.
- ⟨1⟩2. Q.E.D.
PROOF: Since $\lambda x.x - y$ is a homeomorphism of \mathbb{R}^ω with itself.
-

34 Totally Disconnected Spaces

Definition 242 (Totally Disconnected). A topological space X is *totally disconnected* if and only if the only connected subspaces are the singletons.

Example 243. Every discrete space is totally disconnected.

Example 244. The rationals \mathbb{Q} are totally disconnected.

35 Paths and Path Connectedness

Definition 245 (Path). Let X be a topological space and $a, b \in X$. A *path* from a to b is a continuous function $p : [0, 1] \rightarrow X$ such that $p(0) = a$ and $p(1) = b$.

Definition 246 (Path Connected). A topological space is *path connected* if and only if there exists a path between any two points.

Proposition 247. *Every path connected space is connected.*

PROOF:

- ⟨1⟩1. LET: X be a path connected space.
- ⟨1⟩2. ASSUME: for a contradiction C and D form a separation of X .
- ⟨1⟩3. PICK $a \in C$ and $b \in D$.
- ⟨1⟩4. PICK a path $p : [0, 1] \rightarrow X$ from a to b .
- ⟨1⟩5. $p^{-1}(C)$ and $p^{-1}(D)$ form a separation of $[0, 1]$.
- ⟨1⟩6. Q.E.D.

PROOF: This contradicts Corollary 238.2.

□

An example that shows the converse does not hold:

Example 248. The ordered square is not path connected.

PROOF:

- ⟨1⟩1. ASSUME: for a contradiction $p : [0, 1] \rightarrow I_o^2$ is a path from $(0, 0)$ to $(1, 1)$.
- ⟨1⟩2. p is surjective.

PROOF: By the Intermediate Value Theorem.

- ⟨1⟩3. For $x \in [0, 1]$, PICK a rational $q_x \in p^{-1}((x, 0), (x, 1))$

PROOF: Since $p^{-1}((x, 0), (x, 1))$ is open and nonempty by ⟨1⟩2.

- ⟨1⟩4. For $x, y \in [0, 1]$, if $x \neq y$ then $q_x \neq q_y$

PROOF: We have $p(q_x) \neq p(q_y)$ because $((x, 0), (x, 1))$ and $((y, 0), (y, 1))$ are disjoint.

- ⟨1⟩5. $\{q_x \mid x \in [0, 1]\}$ is an uncountable set of rationals.

- ⟨1⟩6. Q.E.D.

PROOF: This contradicts the fact that the rationals are countable.

□

Proposition 249. *The continuous image of a path connected space is path connected.*

PROOF:

- ⟨1⟩1. LET: X be a path connected space, Y a topological space, and $f : X \rightarrow Y$ be continuous and surjective.
- ⟨1⟩2. LET: $a, b \in Y$
- ⟨1⟩3. PICK $c, d \in X$ with $f(c) = a$ and $f(d) = b$
- ⟨1⟩4. PICK a path $p : [0, 1] \rightarrow X$ from c to d .
- ⟨1⟩5. $f \circ p$ is a path from a to b in Y .

□

Proposition 250 (AC). *The product of a family of path-connected spaces is path-connected.*

PROOF:

- ⟨1⟩1. LET: $\{X_\alpha\}_{\alpha \in J}$ be a family of path-connected spaces.

- <1>2. LET: $a, b \in \prod_{\alpha \in J} X_\alpha$
 <1>3. For $\alpha \in J$, PICK a path $p_\alpha : [0, 1] \rightarrow X_\alpha$ from a_α to b_α
 PROOF: Using the Axiom of Choice.
 <1>4. Define $p : [0, 1] \rightarrow \prod_{\alpha \in J} X_\alpha$ by $p(t)_\alpha = p_\alpha(t)$
 <1>5. p is a path from a to b .
 PROOF: Theorem 137.

□

Proposition 251. *The continuous image of a path-connected space is path-connected.*

PROOF:

- <1>1. LET: $f : X \rightarrow Y$ be continuous and surjective where X is path-connected.
 <1>2. LET: $a, b \in Y$
 <1>3. PICK $a', b' \in X$ with $f(a') = a$ and $f(b') = b$.
 <1>4. PICK a path $p : [0, 1] \rightarrow X$ from a' to b' .
 <1>5. $f \circ p$ is a path from a to b .

□

Proposition 252. *Let X be a topological space. The union of a set of path-connected subspaces of X that have a point in common is path-connected.*

PROOF:

- <1>1. LET: \mathcal{A} be a set of path-connected subspaces of X with the point a in common.
 <1>2. LET: $b, c \in \bigcup \mathcal{A}$
 <1>3. PICK $B, C \in \mathcal{A}$ with $b \in B$ and $c \in C$.
 <1>4. PICK a path p in B from b to a .
 <1>5. PICK a path q in C from a to c .
 <1>6. The concatenation of p and q is a path from b to c in $\bigcup \mathcal{A}$.

□

Proposition 253. *Let $A \subseteq \mathbb{R}^2$ be countable. Then $\mathbb{R}^2 \setminus A$ is path-connected.*

PROOF:

- <1>1. LET: $a, b \in \mathbb{R}^2 \setminus A$
 <1>2. PICK a line l in \mathbb{R}^2 with a on one side and b on the other.
 <1>3. For every point x on l ,
 LET: p_x be the path in \mathbb{R}^2 consisting of a line from a to x then a line from x to b
 <1>4. For $x \neq y$ we have p_x and p_y have no points in common except a and b
 <1>5. There are only countably many x such that a point of A lies on p_x .
 <1>6. There exists x such that p_x is a path from a to b in $\mathbb{R}^2 \setminus A$.

□

Proposition 254. *Every open connected subspace of \mathbb{R}^2 is path-connected.*

PROOF:

- <1>1. LET: U be an open connected subspace of \mathbb{R}^2 .

$\langle 1 \rangle 2$. For all $x_0 \in U$,
 LET: $PC(x_0) = \{y \in U \mid \text{there exists a path from } x \text{ to } y\}$
 $\langle 1 \rangle 3$. For all $x_0 \in U$, the set $PC(x_0)$ is open and closed in U .
 $\langle 2 \rangle 1$. LET: $x_0 \in U$
 $\langle 2 \rangle 2$. $PC(x_0)$ is open in U
 $\langle 3 \rangle 1$. LET: $y \in PC(x_0)$
 $\langle 3 \rangle 2$. PICK $\epsilon > 0$ such that $B(y, \epsilon) \subseteq U$
 PROOF: Since U is open.
 $\langle 3 \rangle 3$. $B(y, \epsilon) \subseteq PC(x_0)$
 PROOF: For all $z \in B(y, \epsilon)$, pick a path from x_0 to y then concatenate the straight line from y to z .
 $\langle 2 \rangle 3$. $PC(x_0)$ is closed in U
 $\langle 3 \rangle 1$. LET: $y \in U$ be a limit point of $PC(x_0)$
 $\langle 3 \rangle 2$. PICK $\epsilon > 0$ such that $B(y, \epsilon) \subseteq U$
 $\langle 3 \rangle 3$. PICK $z \in PC(x_0) \cap B(y, \epsilon)$
 $\langle 3 \rangle 4$. $y \in PC(x_0)$
 PROOF: Pick a path from x_0 to z then concatenate the straight line from z to y .
 $\langle 1 \rangle 4$. $PC(x_0) = U$
 PROOF: Proposition 218.
 \square

Example 255. If A is a connected subspace of X , then A° is not necessarily connected.

Take two closed circles in \mathbb{R}^2 that touch at one point. The interior of this space is two open circles, and these two circles form a separation.

Example 256. If A is a connected subspace of X then ∂A is not necessarily connected.

We have $[0, 1]$ is connected but $\partial[0, 1] = \{0, 1\}$ is not.

Example 257. If A is a subspace of X and A° and ∂A are connected, then A is not necessarily connected.

We have $\mathbb{Q}^\circ = \emptyset$ and $\partial\mathbb{Q} = \mathbb{R}$ are connected but \mathbb{Q} is not connected.

36 The Topologist's Sine Curve

Definition 258 (The Topologist's Sine Curve). Let $S = \{(x, \sin 1/x) \mid 0 < x \leq 1\}$, The *topologist's sine curve* is the closure \overline{S} of S in \mathbb{R}^2 .

Proposition 259. The topologist's sine curve is connected.

PROOF:

$\langle 1 \rangle 1$. LET: $S = \{(x, \sin 1/x) \mid 0 < x \leq 1\}$

$\langle 1 \rangle 2$. S is connected.

PROOF: Theorem 228.

$\langle 1 \rangle 3$. \overline{S} is connected.

PROOF: Theorem 227.

□

Proposition 260. *The topologist's sine curve is $\{(x, \sin 1/x) \mid 0 < x \leq 1\} \cup (\{0\} \times [-1, 1])$.*

PROOF: Sketch proof: Given a point $(0, y)$ with $-1 \leq y \leq 1$, pick a such that $\sin a = y$. Then $((1/a, y), (1/(a + 2\pi), y), (1/(a + 4\pi), y), \dots)$ is a sequence in S that converges to $(0, y)$.

Conversely, let (x, y) be any point not in $S \cup (\{0\} \times [-1, 1])$. If $x < 0$ or $y > 1$ or $y < -1$ then we can easily find a neighbourhood that does not intersect $S \cup (\{0\} \times [-1, 1])$. If $x > 0$ and $-1 \leq y \leq 1$, then we have $y \neq \sin 1/x$. Hence pick a neighbourhood that does not intersect S .

Proposition 261. *Every closed subset of \mathbb{R} that is bounded above has a greatest element.*

PROOF: It has a supremum, which is a limit point of the set and hence an element. □

Proposition 262 (CC). *The topologist's sine curve is not path connected.*

PROOF:

⟨1⟩1. ASSUME: For a contradiction $p : [0, 1] \rightarrow \bar{S}$ is a path from $(0, 0)$ to $(1, \sin 1)$.

⟨1⟩2. $\{t \in [0, 1] \mid p(t) \in \{0\} \times [-1, 1]\}$ is closed.

PROOF: Since p is continuous and $\{0\} \times [-1, 1]$ is closed.

⟨1⟩3. LET: b be the largest number in $[0, 1]$ such that $p(b) \in \{0\} \times [-1, 1]$.

PROOF: Proposition 261.

⟨1⟩4. LET: $x : [b, 1] \rightarrow \bar{S}$ be the function $\pi_1 \circ p$

⟨1⟩5. LET: $y : [b, 1] \rightarrow \bar{S}$ be the function $\pi_2 \circ p$

⟨1⟩6. PICK a sequence t_n in $(b, 1]$ such that $t_n \rightarrow b$ and $y(t_n) = (-1)^n$ for all n

⟨2⟩1. LET: $n \geq 1$

⟨2⟩2. PICK u with $0 < u < x(1/n)$ and $\sin(1/u) = (-1)^n$

⟨2⟩3. PICK t_n with $b < t_n < 1/n$ and $x(t_n) = u$

PROOF: By the Intermediate Value Theorem

⟨1⟩7. Q.E.D.

PROOF: This contradicts Proposition 110 since y is continuous and $y(t_n)$ does not converge.

□

Corollary 262.1. *The closure of a path-connected subspace of a space is not necessarily path-connected.*

37 The Long Line

Definition 263 (The Long Line). The *long line* is the space $\omega_1 \times [0, 1]$ in the dictionary order under the order topology, where ω_1 is the first uncountable ordinal.

Lemma 264. *For any ordinal α with $0 < \alpha < \omega_1$ we have $[(0, 0), (\alpha, 0)) \cong [0, 1)$*

$\langle 1 \rangle 1.$ $[(0, 0), (1, 0)) \cong [0, 1)$

PROOF: The map π_2 is a homeomorphism.

$\langle 1 \rangle 2.$ If $[(0, 0), (\alpha, 0)) \cong [0, 1)$ then $[(0, 0), (\alpha + 1, 0)) \cong [0, 1)$

PROOF: Proposition 18.

$\langle 1 \rangle 3.$ If λ is a limit ordinal with $\lambda < \omega_1$ and $[(0, 0), (\alpha, 0)) \cong [0, 1)$ for all α with $0 < \alpha < \lambda$ then $[(0, 0), (\lambda, 0)) \cong [0, 1)$

$\langle 2 \rangle 1.$ LET: λ be a limit ordinal $< \omega_1$

$\langle 2 \rangle 2.$ ASSUME: $[(0, 0), (\alpha, 0)) \cong [0, 1)$ for all α with $0 < \alpha < \lambda$

$\langle 2 \rangle 3.$ PICK a sequence of ordinals $\alpha_0 < \alpha_1 < \dots$ with limit λ

PROOF: Since λ is countable.

$\langle 2 \rangle 4.$ $[(\alpha_i, 0), (\alpha_{i+1}, 0)) \cong [0, 1)$ for all i

PROOF: Lemma 17.

$\langle 2 \rangle 5.$ Q.E.D.

PROOF: By Proposition 19.

$\langle 1 \rangle 4.$ Q.E.D.

PROOF: By transfinite induction.

Proposition 265 (CC). *The long line is path-connected.*

PROOF:

$\langle 1 \rangle 1.$ LET: $(\alpha, i), (\beta, j) \in \omega_1 \times [0, 1)$

$\langle 1 \rangle 2.$ ASSUME: without loss of generality $(\alpha, i) < (\beta, j)$

$\langle 1 \rangle 3.$ $[(0, 0), (\beta + 1, 0)) \cong [0, 1)$

PROOF: By Lemma 264

$\langle 1 \rangle 4.$ $[(\alpha, i), (\beta, j)) \cong [0, 1)$

PROOF: Lemma 17.

$\langle 1 \rangle 5.$ PICK a homeomorphism $q : [0, 1) \rightarrow [(\alpha, i), (\beta, j))$

$\langle 1 \rangle 6.$ $q \cup \{(1, (\beta, j))\}$ is a path from (α, i) to (β, j)

□

Proposition 266. *Every point in the long line has a neighbourhood homeomorphic to an interval in \mathbb{R} .*

PROOF: For any (α, i) in the long line, the neighbourhood $[(0, 0), (\alpha + 1, 0))$ satisfies the condition by Lemma 264.

Proposition 267. *The long line L is not second countable.*

PROOF:

$\langle 1 \rangle 1.$ LET: \mathcal{B} be a basis for L .

$\langle 1 \rangle 2.$ For $\alpha < \omega_1$, PICK $B_\alpha \in \mathcal{B}$ such that $(\alpha, 1/2) \in B_\alpha \subseteq ((\alpha, 0), (\alpha + 1, 0))$

$\langle 1 \rangle 3.$ \mathcal{B} is uncountable.

PROOF: The mapping $\alpha \mapsto B_\alpha$ is an injection $\omega_1 \rightarrow \mathcal{B}$.

Corollary 267.1. *The long line cannot be imbedded into \mathbb{R}^n for any n .*

38 Components

Proposition 268. *Let X be a topological space. Define the relation \sim on X by $a \sim b$ if and only if there exists a connected subspace $A \subseteq X$ with $a, b \in A$. Then \sim is an equivalence relation on X .*

PROOF:

$\langle 1 \rangle 1.$ \sim is reflexive.

PROOF: For any $a \in X$ we have $\{a\}$ is a connected subspace that contains a .

$\langle 1 \rangle 2.$ \sim is symmetric.

PROOF: Trivial.

$\langle 1 \rangle 3.$ \sim is transitive.

$\langle 2 \rangle 1.$ LET: $a, b, c \in X$

$\langle 2 \rangle 2.$ ASSUME: $a \sim b$ and $b \sim c$

$\langle 2 \rangle 3.$ PICK connected subspaces A and B with $a, b \in A$ and $b, c \in B$

$\langle 2 \rangle 4.$ $A \cup B$ is a connected subspace that contains a and c

PROOF: Theorem 226.

□

Definition 269 ((Connected) Component). Let X be a topological space. The *(connected) components* of X are the equivalence classes under the above \sim .

Lemma 270. *Let X be a topological space. If $A \subseteq X$ is connected and nonempty then there exists a unique component C of X such that $A \subseteq C$.*

PROOF:

$\langle 1 \rangle 1.$ PICK $a \in A$

$\langle 1 \rangle 2.$ LET: C be the \sim -equivalence class of a .

$\langle 1 \rangle 3.$ $A \subseteq C$

PROOF: For all $x \in A$ we have $x \sim a$.

$\langle 1 \rangle 4.$ If C' is a component and $A \subseteq C'$ then $C = C'$

PROOF: Since we have $a \in C'$.

□

Theorem 271. *Let X be a topological space. The components of X are connected disjoint subspaces of X whose union is X such that each nonempty connected subspace of X intersects only one of them.*

PROOF:

$\langle 1 \rangle 1.$ Every component of X is connected.

PROOF: For $a \in X$, the \sim -equivalence class of a is $\bigcup \{A \subseteq X \mid A \text{ is connected, } a \in A\}$ which is connected by Theorem 226.

$\langle 1 \rangle 2.$ The components form a partition of X .

PROOF: Immediate from the definition.

$\langle 1 \rangle 3.$ Every nonempty connected subspace of X intersects a unique component of X .

$\langle 2 \rangle 1.$ LET: $A \subseteq X$ be connected and nonempty.

$\langle 2 \rangle 2.$ LET: C be the component such that $A \subseteq C$

PROOF: Lemma 270.

$\langle 2 \rangle 3$. A intersects C

$\langle 2 \rangle 4$. If A intersects the component C' then $C' = C$

$\langle 3 \rangle 1$. LET: C' be a component that intersects A

$\langle 3 \rangle 2$. PICK $b \in A \cap C'$

$\langle 3 \rangle 3$. $A \subseteq C'$

PROOF: For all $x \in A$ we have $x \sim b$.

$\langle 3 \rangle 4$. $C = C'$

PROOF: By uniqueness in $\langle 2 \rangle 2$.

□

Proposition 272. *Every component of a space is closed.*

PROOF:

$\langle 1 \rangle 1$. LET: X be a topological space and C a component of X .

$\langle 1 \rangle 2$. \overline{C} is connected.

PROOF: Theorem 227.

$\langle 1 \rangle 3$. $C = \overline{C}$

PROOF: Lemma 225.

$\langle 1 \rangle 4$. C is closed.

PROOF: Lemma 51.

□

Proposition 273. *If a topological space has finitely many components then every component is open.*

PROOF: Each component is the complement of a finite union of closed sets. □

39 Path Components

Proposition 274. *Let X be a topological space. Define the relation \sim on X by: $a \sim b$ if and only if there exists a path in X from a to b . Then \sim is an equivalence relation on X .*

PROOF:

$\langle 1 \rangle 1$. \sim is reflexive.

PROOF: For $a \in X$, the constant function $[0, 1] \rightarrow X$ with value a is a path from a to a .

$\langle 1 \rangle 2$. \sim is symmetric.

PROOF: If $p : [0, 1] \rightarrow X$ is a path from a to b , then $\lambda t.p(1-t)$ is a path from b to a .

$\langle 1 \rangle 3$. \sim is transitive.

PROOF: Concatenate paths.

□

Definition 275 (Path Component). Let X be a topological space. The *path components* of X are the equivalence relations under \sim .

Theorem 276. *The path components of X are path-connected disjoint subspaces of X whose union is X such that every nonempty path-connected subspace of X intersects exactly one path component.*

PROOF:

⟨1⟩1. Every path component is path-connected.

PROOF: If a and b are in the same path component then $a \sim b$, i.e. there exists a path from a to b .

⟨1⟩2. The path components are disjoint and their union is X .

PROOF: Immediate from the definition.

⟨1⟩3. Every non-empty path-connected subspace of X intersects exactly one path component.

⟨2⟩1. LET: A be a nonempty path-connected subspace of X .

⟨2⟩2. PICK $a \in A$

⟨2⟩3. A intersects the \sim -equivalence class of a .

⟨2⟩4. LET: C be any path component that intersects A .

⟨2⟩5. PICK $b \in A \cap C$

⟨2⟩6. $a \sim b$

PROOF: Since A is path-connected.

⟨2⟩7. C is the \sim -equivalence class of a .

□

Proposition 277. *Every path component is included in a component.*

PROOF:

⟨1⟩1. LET: X be a topological space and C a path component of X .

⟨1⟩2. C is path-connected.

PROOF: Theorem 276.

⟨1⟩3. C is connected.

PROOF: Proposition 247.

⟨1⟩4. C is included in a component.

PROOF: Lemma 270.

□

40 Local Connectedness

Definition 278 (Locally Connected). Let X be a topological space and $a \in X$. Then X is *locally connected* at a if and only if every neighbourhood of a includes a connected neighbourhood of a .

The space X is *locally connected* if and only if it is locally connected at every point.

Example 279. The real line is both connected and locally connected.

Example 280. The space $\mathbb{R} \setminus \{0\}$ is disconnected but locally connected.

Example 281. The topologist's sine curve is connected but not locally connected.

Example 282. The rationals \mathbb{Q} are neither connected nor locally connected.

Theorem 283. *A topological space X is locally connected if and only if, for every open set U in X , every component of U is open in X .*

PROOF:

⟨1⟩1. If X is locally connected then, for every open set U in X , every component of U is open in X .

⟨2⟩1. ASSUME: X is locally connected.

⟨2⟩2. LET: U be open in X .

⟨2⟩3. LET: C be a component of U .

⟨2⟩4. LET: $a \in C$

⟨2⟩5. LET: V be a connected neighbourhood of a such that $V \subseteq U$

⟨2⟩6. $V \subseteq C$

PROOF: Lemma 270.

⟨2⟩7. Q.E.D.

PROOF: Lemma 30.

⟨1⟩2. If, for every open set U in X , every component of U is open in X , then X is locally connected.

⟨2⟩1. ASSUME: for every open set U in X , every component of U is open in X .

⟨2⟩2. LET: $a \in X$

⟨2⟩3. LET: U be a neighbourhood of a

⟨2⟩4. The component of U that contains a is a connected neighbourhood of a included in U .

□

Example 284. The ordered square is locally connected.

PROOF: Every neighbourhood of a point includes an interval around that point, which is connected by Theorem 238.

Example 285. Let X be the set of all rational points on the line segment $[0, 1] \times \{0\}$, and Y the set of all rational points on the line segment $[0, 1] \times \{1\}$. Let A be the space consisting of all line segments joining the point $(0, 1)$ to a point of X , and all line segments joining the point $(1, 0)$ to a point of Y . Then A is path-connected but is not locally connected at any point,

Proposition 286. *Let X and Y be topological spaces and $p : X \twoheadrightarrow Y$ be a quotient map. If X is locally connected then so is Y .*

PROOF:

⟨1⟩1. LET: U be an open set in Y .

⟨1⟩2. LET: C be a component of U .

⟨1⟩3. $p^{-1}(C)$ is a union of components of $p^{-1}(U)$

⟨2⟩1. LET: $x \in p^{-1}(C)$

$\langle 2 \rangle 2$. LET: D be the component of $p^{-1}(U)$ that contains x .
 $\langle 2 \rangle 3$. $p(D)$ is connected.
 PROOF: Theorem 228.
 $\langle 2 \rangle 4$. $p(D) \subseteq C$.
 PROOF: From $\langle 1 \rangle 2$ since $p(x) \in p(D) \cap C$ ($\langle 2 \rangle 1$, $\langle 2 \rangle 2$).
 $\langle 2 \rangle 5$. $D \subseteq p^{-1}(C)$
 $\langle 1 \rangle 4$. $p^{-1}(C)$ is open in $p^{-1}(U)$
 PROOF: Theorem 283.
 $\langle 1 \rangle 5$. C is open in U
 PROOF: Since the restriction of p to $p : p^{-1}(U) \rightarrow U$ is a quotient map by Proposition 201.
 $\langle 1 \rangle 6$. Q.E.D.
 PROOF: Theorem 283.
 \square

41 Local Path Connectedness

Definition 287 (Locally Path-Connected). Let X be a topological space and $a \in X$. Then X is *locally path-connected* at a if and only if every neighbourhood of a includes a path-connected neighbourhood of a .

The space X is *locally path-connected* if and only if it is locally path-connected at every point.

Theorem 288. *A topological space X is locally path-connected if and only if, for every open set U in X , every path component of U is open in X .*

PROOF:

$\langle 1 \rangle 1$. If X is locally path-connected then, for every open set U in X , every path component of U is open in X .
 $\langle 2 \rangle 1$. ASSUME: X is locally path-connected.
 $\langle 2 \rangle 2$. LET: U be open in X .
 $\langle 2 \rangle 3$. LET: C be a path component of U .
 $\langle 2 \rangle 4$. LET: $a \in C$
 $\langle 2 \rangle 5$. LET: V be a path-connected neighbourhood of a such that $V \subseteq U$
 $\langle 2 \rangle 6$. $V \subseteq C$
 PROOF: Lemma 270.
 $\langle 2 \rangle 7$. Q.E.D.
 PROOF: Lemma 30.
 $\langle 1 \rangle 2$. If, for every open set U in X , every component of U is open in X , then X is locally connected.
 $\langle 2 \rangle 1$. ASSUME: for every open set U in X , every component of U is open in X .
 $\langle 2 \rangle 2$. LET: $a \in X$
 $\langle 2 \rangle 3$. LET: U be a neighbourhood of a
 $\langle 2 \rangle 4$. The component of U that contains a is a connected neighbourhood of a included in U .

□

Theorem 289. *If a space is locally path connected then its components and its path components are the same.*

PROOF:

⟨1⟩1. LET: X be a locally path connected space.

⟨1⟩2. LET: C be a component of X .

⟨1⟩3. LET: $x \in C$

⟨1⟩4. LET: P be the path component of x

PROVE: $P = C$

⟨1⟩5. $P \subseteq C$

PROOF: Proposition 277.

⟨1⟩6. LET: Q be the union of the other path components included in C

⟨1⟩7. $C = P \cup Q$

PROOF: Proposition 277.

⟨1⟩8. P and Q are open in C

⟨2⟩1. C is open.

PROOF: Theorem 283.

⟨2⟩2. Q.E.D.

PROOF: Theorem 288.

⟨1⟩9. $Q = \emptyset$

PROOF: Otherwise P and Q would form a separation of C .

□

Example 290. The ordered square is not locally path connected, since it is connected but not path connected.

Proposition 291. *Let X be a locally path-connected space. Then every connected open subspace of X is path-connected.*

PROOF:

⟨1⟩1. LET: U be a connected open subspace of X .

⟨1⟩2. LET: P be a path component of U .

⟨1⟩3. LET: Q be the union of the other path components of U .

⟨1⟩4. P and Q are open in U .

PROOF: Theorem 288.

⟨1⟩5. $Q = \emptyset$

PROOF: Otherwise P and Q form a separation of U .

□

42 Weak Local Connectedness

Definition 292 (Weakly Locally Connected). Let X be a topological space and $a \in X$. Then X is *weakly locally connected* at a if and only if every neighbourhood of a includes a connected subspace that includes a neighbourhood of a .

Proposition 293. *Let X be a topological space. If X is weakly locally connected at every point then X is locally connected.*

PROOF:

⟨1⟩1. ASSUME: X is weakly locally connected at every point.

⟨1⟩2. LET: U be open in X .

⟨1⟩3. LET: C be a component of U .

⟨1⟩4. C is open in X .

⟨2⟩1. LET: $x \in C$

⟨2⟩2. PICK a connected subspace D of U that includes a neighbourhood V of x .

⟨2⟩3. $D \subseteq C$

PROOF: Lemma 270.

⟨2⟩4. $x \in V \subseteq C$

⟨2⟩5. Q.E.D.

PROOF: Lemma 30.

⟨1⟩5. Q.E.D.

PROOF: Theorem 283.

□

Example 294. The 'infinite broom' (Munkres p.163) is an example of a space that is weakly locally connected at a point p but not locally connected at p .

43 Quasicomponents

Proposition 295. *Let X be a topological space. Define \sim on X by $x \sim y$ if and only if there exists no separation U and V of X such that $x \in U$ and $y \in V$. Then \sim is an equivalence relation on X .*

PROOF:

⟨1⟩1. \sim is reflexive.

PROOF: Since the two sets that make up a separation are disjoint.

⟨1⟩2. \sim is symmetric.

PROOF: Immediate from the definition.

⟨1⟩3. \sim is transitive.

⟨2⟩1. ASSUME: $x \sim y$ and $y \sim z$

⟨2⟩2. ASSUME: for a contradiction there is a separation U and V of X with $x \in U$ and $z \in V$

⟨2⟩3. $y \in U$ or $y \in V$

⟨2⟩4. Q.E.D.

PROOF: Either case contradicts ⟨2⟩1.

□

Definition 296 (Quasicomponents). For X a topological space, the *quasicomponents* of X are the equivalence classes under \sim .

Proposition 297. *Let X be a topological space. Then every component of X is included in a quasicomponent of X .*

PROOF:

$\langle 1 \rangle 1$. LET: C be a component of X .

$\langle 1 \rangle 2$. LET: $x, y \in C$

PROVE: $x \sim y$

$\langle 1 \rangle 3$. ASSUME: for a contradiction there exists a separation U and V of X with
 $x \in U$ and $y \in V$

$\langle 1 \rangle 4$. $C \cap U$ and $C \cap V$ form a separation of C .

$\langle 1 \rangle 5$. Q.E.D.

PROOF: This contradicts $\langle 1 \rangle 1$.

Proposition 298. *In a locally connected space, the components and the quasi-components are the same.*

PROOF:

$\langle 1 \rangle 1$. LET: X be a locally connected space and Q a quasicomponent of X .

$\langle 1 \rangle 2$. PICK a component C of X such that $C \subseteq Q$

$\langle 1 \rangle 3$. LET: D be the union of the components of X

$\langle 1 \rangle 4$. C and D are open in X .

PROOF: Theorem 283.

$\langle 1 \rangle 5$. D cannot contain any points of Q .

PROOF: If it did, then C and D would form a separation of X and there would be points $x, y \in Q$ with $x \in C$ and $y \in D$.

$\langle 1 \rangle 6$. $C = Q$

□

44 Open Coverings

Definition 299 (Open Covering). Let X be a topological space. An *open covering* of X is a covering of X whose elements are all open sets.

45 Compact Spaces

Definition 300 (Compact). A topological space is *compact* if and only if every open covering has a finite subcovering.

Lemma 301. *Let X be a topological space and Y a subspace of X . Then Y is compact if and only if every covering of Y by sets open in X has a finite subcovering.*

PROOF:

$\langle 1 \rangle 1$. If Y is compact then every covering of Y by sets open in X has a finite subcovering.

$\langle 2 \rangle 1$. ASSUME: Y is compact.

$\langle 2 \rangle 2$. LET: \mathcal{U} be a covering of Y by sets open in X .

$\langle 2 \rangle 3$. $\{U \cap Y \mid U \in \mathcal{U}\}$ is an open covering of Y .

$\langle 2 \rangle 4$. PICK a finite subcovering $\{U_1 \cap Y, \dots, U_n \cap Y\}$

- ⟨2⟩5. $\{U_1, \dots, U_n\}$ is a finite subcovering of \mathcal{U} .
- ⟨1⟩2. If every covering of Y by sets open in X has a finite subcovering then Y is compact.
- ⟨2⟩1. LET: \mathcal{U} be an open covering of Y .
- ⟨2⟩2. LET: $\mathcal{V} = \{V \text{ open in } X \mid V \cap Y \in \mathcal{U}\}$.
- ⟨2⟩3. \mathcal{V} is a covering of Y by sets open in X .
- ⟨2⟩4. PICK a finite subcovering $\{V_1, \dots, V_n\}$
- ⟨2⟩5. $\{V_1 \cap Y, \dots, V_n \cap Y\}$ is a finite subcovering of \mathcal{U} .

□

Proposition 302. *Every closed subspace of a compact space is compact.*

PROOF:

- ⟨1⟩1. LET: X be a compact space and $Y \subseteq X$ be closed.
- ⟨1⟩2. LET: \mathcal{U} be a covering of Y by sets open in X .
- ⟨1⟩3. $\mathcal{U} \cup \{X \setminus Y\}$ is an open covering of X .
- ⟨1⟩4. PICK a finite subcovering \mathcal{U}_0
- ⟨1⟩5. $\mathcal{U}_0 \cap \mathcal{U}$ is a finite subset of \mathcal{U} that covers Y .

□

Theorem 303. *The continuous image of a compact space is compact.*

PROOF:

- ⟨1⟩1. LET: $f : X \rightarrow Y$ be continuous and surjective.
- ⟨1⟩2. LET: \mathcal{V} be an open covering of Y
- ⟨1⟩3. $\{p^{-1}(V) \mid V \in \mathcal{V}\}$ is an open covering of X .
- ⟨1⟩4. PICK a finite subcovering $\{p^{-1}(V_1), \dots, p^{-1}(V_n)\}$
- ⟨1⟩5. $\{V_1, \dots, V_n\}$ covers Y .

□

Theorem 304. *Let A and B be compact subspaces of X and Y respectively. Let N be an open set in $X \times Y$ that includes $A \times B$. Then there exist open sets U and V in X and Y respectively such that $A \subseteq U$, $B \subseteq V$ and $U \times V \subseteq N$.*

PROOF:

- ⟨1⟩1. For all $x \in A$, there exist neighbourhoods U of x and V of B such that $U \times V \subseteq N$.
- ⟨2⟩1. LET: $x \in A$
- ⟨2⟩2. For all $y \in B$, there exist neighbourhoods U of x and V of y such that $U \times V \subseteq N$
- ⟨2⟩3. $\{V \text{ open in } Y \mid \exists \text{ neighbourhood } U \text{ of } x, U \times V \subseteq N\}$ covers B .
- ⟨2⟩4. PICK a finite subcover $\{V_1, \dots, V_n\}$
- ⟨2⟩5. For $i = 1, \dots, n$, PICK a neighbourhood U_i of x such that $U_i \times V_i \subseteq N$
- ⟨2⟩6. LET: $U = U_1 \cap \dots \cap U_n$
- ⟨2⟩7. LET: $V = V_1 \cup \dots \cup V_n$
- ⟨2⟩8. U is a neighbourhood of x .
- ⟨2⟩9. V is a neighbourhood of B .
- ⟨2⟩10. $U \times V \subseteq N$

- ⟨1⟩2. $\{U \text{ open in } X \mid \exists \text{ neighbourhood } V \text{ of } B. U \times V \subseteq N\}$ covers A .
 - ⟨1⟩3. PICK a finite subcover $\{U_1, \dots, U_n\}$
 - ⟨1⟩4. For $i = 1, \dots, n$, PICK a neighbourhood V_i of B such that $U_i \times V_i \subseteq N$
 - ⟨1⟩5. LET: $U = U_1 \cup \dots \cup U_n$
 - ⟨1⟩6. LET: $V = V_1 \cap \dots \cap V_n$
 - ⟨1⟩7. U and V are open.
 - ⟨1⟩8. $A \subseteq U$
 - ⟨1⟩9. $B \subseteq V$
 - ⟨1⟩10. $U \times V \subseteq N$
-

Corollary 304.1 (Tube Lemma). *Let X and Y be topological spaces with Y compact. Let $a \in X$ and N be an open set in $X \times Y$ that includes $\{a\} \times Y$. Then there exists a neighbourhood W of a such that N includes the tube $W \times Y$.*

Theorem 305. *Let X be a topological space. Then X is compact if and only if every set of closed sets that has the finite intersection property has nonempty intersection.*

PROOF: The following are equivalent.

1. X is compact.
2. Every open covering of X has a finite subcovering.
3. For any set \mathcal{C} of closed sets, if $\{X \setminus C \mid C \in \mathcal{C}\}$ covers X then there is a finite subset \mathcal{C}_0 such that $\{X \setminus C \mid C \in \mathcal{C}_0\}$ covers X
4. For any set \mathcal{C} of closed sets, if $\bigcap \mathcal{C} = \emptyset$ then there is a finite subset \mathcal{C}_0 with empty intersection.
5. Any set of closed sets with the finite intersection property has nonempty intersection.

□

Corollary 305.1. *Let X be a topological space and $C_1 \supseteq C_2 \supseteq \dots$ a nested sequence of nonempty closed sets. Then $\bigcap_n C_n$ is nonempty.*

Proposition 306. *Let \mathcal{T} and \mathcal{T}' be two topologies on the same set X with $\mathcal{T} \subseteq \mathcal{T}'$. If \mathcal{T}' is compact then \mathcal{T} is compact.*

PROOF:

- ⟨1⟩1. LET: $\mathcal{U} \subseteq \mathcal{T}$ cover X
- ⟨1⟩2. $\mathcal{U} \subseteq \mathcal{T}'$
- ⟨1⟩3. A finite subset of \mathcal{U} covers X .

□

Corollary 306.1. *If \mathcal{T} and \mathcal{T}' are two compact Hausdorff topologies on the same set X , then either $\mathcal{T} = \mathcal{T}'$ or \mathcal{T} and \mathcal{T}' are incomparable.*

PROOF: From the Proposition and Proposition 182. \square

Example 307. Any set under the finite complement topology is compact.

Proposition 308. *Let X be a topological space. A finite union of compact subspaces of X is compact.*

PROOF:

$\langle 1 \rangle 1$. LET: A and B be compact subspaces of X .

$\langle 1 \rangle 2$. LET: \mathcal{U} be a set of open sets in X that covers $A \cup B$

$\langle 1 \rangle 3$. PICK a finite subset \mathcal{U}_1 that covers A .

PROOF: Lemma 301.

$\langle 1 \rangle 4$. PICK a finite subset \mathcal{U}_2 that covers B .

PROOF: Lemma 301.

$\langle 1 \rangle 5$. $\mathcal{U}_1 \cup \mathcal{U}_2$ is a finite subset that covers $A \cup B$.

$\langle 1 \rangle 6$. Q.E.D.

PROOF: Lemma 301.

\square

Proposition 309. *Let A and B be disjoint compact subspaces of the Hausdorff space X . Then there exist disjoint open sets U and V that include A and B respectively.*

PROOF: From Theorem 304 with $N = X^2 \setminus \{(x, x) \mid x \in X\}$. \square

Corollary 309.1. *Every compact subspace of a Hausdorff space is closed.*

Theorem 310. *Let $f : X \rightarrow Y$ be a bijective continuous function. If X is compact and Y is Hausdorff then f is a homeomorphism.*

PROOF:

$\langle 1 \rangle 1$. LET: $C \subseteq X$ be closed.

$\langle 1 \rangle 2$. C is compact.

PROOF: Proposition 302.

$\langle 1 \rangle 3$. $f(C)$ is compact.

PROOF: Theorem 303.

$\langle 1 \rangle 4$. $f(C)$ is closed.

PROOF: Corollary 309.1.

$\langle 1 \rangle 5$. Q.E.D.

PROOF: Lemma 112.

\square

Proposition 311. *Let X be a compact space, Y a Hausdorff space, and $f : X \rightarrow Y$ a continuous map. Then f is a closed map.*

PROOF:

$\langle 1 \rangle 1$. LET: $C \subseteq X$ be closed.

$\langle 1 \rangle 2$. C is compact.

PROOF: Proposition 302.

⟨1⟩3. $f(C)$ is compact.

PROOF: Theorem 303.

⟨1⟩4. $f(C)$ is closed.

PROOF: Corollary 309.1.

□

Proposition 312. *If Y is compact then the projection $\pi_1 : X \times Y \rightarrow X$ is a closed map.*

PROOF:

⟨1⟩1. LET: $A \subseteq X \times Y$ be closed.

⟨1⟩2. LET: $x \in X \setminus \pi_1(A)$

⟨1⟩3. PICK a neighbourhood U of x such that $U \times Y \subseteq (X \times Y) \setminus A$

PROOF: By the Tube Lemma.

⟨1⟩4. $x \in U \subseteq X \setminus \pi_1(A)$

⟨1⟩5. Q.E.D.

PROOF: So $X \setminus \pi_1(A)$ is open by Lemma 30.

□

Theorem 313. *Let X be a topological space and Y a compact Hausdorff space. Let $f : X \rightarrow Y$ be a function. Then f is continuous if and only if the graph of f is closed in $X \times Y$.*

PROOF:

⟨1⟩1. LET: G_f be the graph of f .

⟨1⟩2. If f is continuous then G_f is closed.

⟨2⟩1. ASSUME: f is continuous.

⟨2⟩2. LET: $(x, y) \in (X \times Y) \setminus G_f$

⟨2⟩3. PICK disjoint neighbourhoods U and V of y and $f(x)$ respectively.

⟨2⟩4. $f^{-1}(V) \times U$ is a neighbourhood of (x, y) disjoint from G_f .

⟨1⟩3. If G_f is closed then f is continuous.

⟨2⟩1. ASSUME: G_f is closed.

⟨2⟩2. LET: $x \in X$ and V be a neighbourhood of $f(x)$.

⟨2⟩3. $G_f \cap (X \times (Y \setminus V))$ is closed.

⟨2⟩4. $\pi_1(G_f \cap (X \times (Y \setminus V)))$ is closed.

PROOF: Proposition 312.

⟨2⟩5. LET: $U = X \setminus \pi_1(G_f \cap (X \times (Y \setminus V)))$

⟨2⟩6. U is a neighbourhood of x

⟨2⟩7. $f(U) \subseteq V$

□

Theorem 314. *Let X be a compact topological space. Let $(f_n : X \rightarrow \mathbb{R})$ be a monotone increasing sequence of continuous functions and $f : X \rightarrow \mathbb{R}$ a continuous function. If (f_n) converges pointwise to f , then (f_n) converges uniformly to f .*

PROOF:

⟨1⟩1. LET: $\epsilon > 0$

- (1)2. For all $x \in X$, there exists N such that, for all $n \geq N$, we have $|f_n(x) - f(x)| < \epsilon$
 (1)3. For $n \geq 1$,
 LET: $U_n = \{x \in X \mid |f_n(x) - f(x)| < \epsilon\}$
 (1)4. For $n \geq 1$, we have U_n is open in X .
 (2)1. LET: $x \in X$
 (2)2. LET: $\delta = \epsilon - |f_n(x) - f(x)|$
 (2)3. PICK a neighbourhood U of x such that $f(U) \subseteq B(f(x), \delta/2)$
 (2)4. PICK a neighbourhood V of x such that $f_n(V) \subseteq B(f_n(x), \delta/2)$
 (2)5. $f(U \cap V) \subseteq U_n$
 PROOF: For $y \in U \cap V$ we have
 $|f_n(y) - f(y)| \leq |f_n(y) - f_n(x)| + |f_n(x) - f(x)| + |f(x) - y(y)|$
 $< \delta/2 + |f_n(x) - f(x)| + \delta/2$
 $= \epsilon$
 (1)5. $\{U_n \mid n \geq 1\}$ covers X
 PROOF: From (1)2
 (1)6. PICK N such that $X = U_N$
 (2)1. PICK n_1, \dots, n_k such that U_{n_1}, \dots, U_{n_k} cover X .
 (2)2. LET: $N = \max(n_1, \dots, n_k)$
 (2)3. For all i we have $U_{n_i} \subseteq U_N$
 PROOF: Since (f_n) is monotone increasing.
 (2)4. $X = U_N$
 (1)7. For all $x \in X$ and $n \geq N$ we have $|f_n(x) - f(x)| < \epsilon$
 □

An example to show that we cannot remove the hypothesis that X is compact:

Example 315. Let $X = (0, 1)$, $f_n(x) = -x^n$ and $f(x) = 0$ for $x \in X$ and $n \geq 1$. Then $f_n \rightarrow f$ pointwise and (f_n) is monotone increasing but the convergence is not uniform since, for all $N \geq 1$, there exists $x \in (0, 1)$ such that $-x^N < -1/2$.

An example to show that we cannot remove the hypothesis that (f_n) is monotone increasing:

Example 316. Let $X = [0, 1]$, $f_n(x) = 1/(n^3(x - 1/n)^2 + 1)$ and $f(x) = 0$ for $x \in X$ and $n \geq 1$. Then X is compact and $f_n \rightarrow f$ pointwise but the convergence is not uniform since, for all $N \geq 1$, there exists $x \in [0, 1]$ such that $f_N(x) = 1$, namely $x = 1/N$.

Theorem 317. Let X be a compact Hausdorff space. Let \mathcal{A} be a chain of closed connected subsets of X . Then $\bigcap \mathcal{A}$ is connected.

PROOF:

- (1)1. ASSUME: for a contradiction C and D form a separation of $\bigcap \mathcal{A}$.
 (1)2. PICK disjoint open sets U and V that include C and D respectively.
 PROOF: Proposition 309.

- (1)3. $\{A \setminus (U \cup V) \mid A \in \mathcal{A}\}$ is a set of closed sets with the finite intersection property.
 (2)1. For all $A \in \mathcal{A}$ we have $A \setminus (U \cup V)$ is closed.
 (2)2. For all $A_1, \dots, A_n \in \mathcal{A}$ we have $(A_1 \cap \dots \cap A_n) \setminus (U \cup V)$ is nonempty.
 PROOF:
 (3)1. LET: $A_1, \dots, A_n \in \mathcal{A}$
 (3)2. ASSUME: without loss of generality $A_1 \subseteq A_2, \dots, A_n$
 PROOF: Since \mathcal{A} is a chain.
 (3)3. $A_1 \setminus (U \cup V)$ is nonempty
 PROOF: Otherwise $(A_1 \cap \dots \cap A_n \cap U)$ and $(A_1 \cap \dots \cap A_n \cap V)$ would form a separation of A_n .
 (1)4. $\bigcap \mathcal{A} \setminus (U \cup V)$ is nonempty.
 PROOF: Theorem 305.
 (1)5. Q.E.D.
 PROOF: This contradicts (1)1 since $\bigcap \mathcal{A} \setminus (U \cup V) = \bigcap \mathcal{A} \setminus (C \cup D)$.

□

Theorem 318 (Tychonoff Theorem (AC)). *The product of a family of compact spaces is compact.*

PROOF:

- (1)1. LET: $\{X_\alpha\}_{\alpha \in J}$ be a family of compact spaces.
 (1)2. LET: $X = \prod_{\alpha \in J} X_\alpha$
 (1)3. For any $\mathcal{A} \subseteq \mathcal{P}X$, we have $\bigcap_{A \in \mathcal{A}} \overline{A} \neq \emptyset$
 (2)1. LET: $\mathcal{A} \subseteq \mathcal{P}X$
 (2)2. PICK $\mathcal{D} \supseteq \mathcal{A}$ that is maximal with respect to the finite intersection property.
 PROVE: $\bigcap_{D \in \mathcal{D}} \overline{D} \neq \emptyset$
 PROOF: Lemma 3.
 (2)3. For $\alpha \in J$, PICK $x_\alpha \in X_\alpha$ such that $x_\alpha \in \bigcap_{D \in \mathcal{D}} \overline{\pi_\alpha(D)}$
 PROOF: Theorem 305 since $\{\overline{\pi_\alpha(D)} \mid D \in \mathcal{D}\}$ is a set of closed sets in X_α with the finite intersection property.
 (2)4. LET: $x = (x_\alpha)_{\alpha \in J}$
 PROVE: $x \in \bigcap_{D \in \mathcal{D}} \overline{D}$
 (2)5. For any $\beta \in J$ and neighbourhood U of x_β in X_β , we have $\pi_\beta^{-1}(U)$ intersects every element of \mathcal{D}
 (3)1. LET: $\beta \in J$
 (3)2. LET: U be a neighbourhood of x_β in X_β .
 (3)3. LET: $D \in \mathcal{D}$
 (3)4. $x_\beta \in \overline{\pi_\beta(D)}$
 PROOF: From (2)3
 (3)5. U intersects $\pi_\beta(D)$.
 (3)6. $\pi_\beta^{-1}(U)$ intersects D .
 (2)6. For any $\beta \in J$ and neighbourhood U of x_β in X_β , we have $\pi_\beta^{-1}(U) \in \mathcal{D}$
 PROOF: Lemma 5.
 (2)7. Every basic neighbourhood of x is an element of \mathcal{D}

PROOF: Lemma 4.

⟨2⟩8. Every basic neighbourhood of x intersects every element of \mathcal{D}

PROOF: Since \mathcal{D} satisfies the finite intersection property.

⟨2⟩9. For all $D \in \mathcal{D}$ we have $x \in \overline{D}$

⟨1⟩4. Q.E.D.

PROOF: Theorem 305.

□

46 Perfect Maps

Definition 319 (Perfect Map). Let X and Y be topological spaces and $f : X \rightarrow Y$. Then f is a *perfect map* if and only if f is a closed map, continuous, surjective and $p^{-1}(y)$ is compact for all $y \in Y$.

Proposition 320. Let X be a topological space, Y a compact space, and $p : X \rightarrow Y$ a closed map such that, for all $y \in Y$, we have $p^{-1}(y)$ is compact. Then X is compact.

PROOF:

⟨1⟩1. LET: \mathcal{A} be a set of closed sets in X with the finite intersection property.

⟨1⟩2. $\mathcal{B} = \{p(A_1 \cap \cdots \cap A_n) \mid A_1, \dots, A_n \in \mathcal{A}\}$ is a set of closed sets in Y with the finite intersection property.

PROOF: Since p is a closed map.

⟨1⟩3. PICK $y \in \bigcap \mathcal{B}$

PROOF: Theorem 305 since Y is compact.

⟨1⟩4. $\{A \cap p^{-1}(y) \mid A \in \mathcal{A}\}$ is a set of closed sets in $p^{-1}(y)$ with the finite intersection property.

⟨1⟩5. PICK $x \in \bigcap_{A \in \mathcal{A}} (A \cap p^{-1}(y))$

PROOF: Theorem 305 since $p^{-1}(y)$ is compact.

⟨1⟩6. $x \in \bigcap \mathcal{A}$

⟨1⟩7. Q.E.D.

PROOF: Theorem 305.

□

47 Topological Groups

Definition 321 (Topological Group). A *topological group* G consists of a T_1 space G and continuous maps $\cdot : G^2 \rightarrow G$ and $(\)^{-1} : G \rightarrow G$ such that $(G, \cdot, (\)^{-1})$ is a group.

Example 322. 1. The integers \mathbb{Z} under addition are a topological group.

2. The real numbers \mathbb{R} under addition are a topological group.

3. The positive reals under multiplication are a topological group.

4. The set $\{z \in \mathbb{C} \mid |z| = 1\}$ under multiplication and given the topology of S^1 is a topological group.
5. For any $n \geq 0$, the general linear group $GL_n(\mathbb{R})$ is a topological group under matrix multiplication, considered as a subspace of \mathbb{R}^{n^2} .

Lemma 323. *Let G be a T_1 space and $\cdot : G^2 \rightarrow G$, $(\)^{-1} : G \rightarrow G$ be functions such that $(G, \cdot, (\)^{-1})$ is a group. Then G is a topological group if and only if the function $f : G^2 \rightarrow G$ that maps (x, y) to xy^{-1} is continuous.*

PROOF:

$\langle 1 \rangle 1$. If G is a topological group then f is continuous.

PROOF: From Theorem 101.

$\langle 1 \rangle 2$. If f is continuous then G is a topological group.

$\langle 2 \rangle 1$. ASSUME: f is continuous.

$\langle 2 \rangle 2$. $(\)^{-1}$ is continuous.

PROOF: Since $x^{-1} = f(e, x)$.

$\langle 2 \rangle 3$. \cdot is continuous.

PROOF: Since $xy = f(x, y^{-1})$.

□

Lemma 324. *Let G be a topological group and H a subgroup of G . Then H is a topological group under the subspace topology.*

PROOF:

$\langle 1 \rangle 1$. H is T_1 .

PROOF: From Proposition 170.

$\langle 1 \rangle 2$. multiplication and inverse on H are continuous.

PROOF: From Theorem 102.

□

Lemma 325. *Let G be a topological group and H a subgroup of G . Then \overline{H} is a subgroup of G .*

PROOF:

$\langle 1 \rangle 1$. LET: $x, y \in \overline{H}$

PROVE: $xy^{-1} \in \overline{H}$

$\langle 1 \rangle 2$. LET: U be any neighbourhood of xy^{-1}

$\langle 1 \rangle 3$. LET: $f : G^2 \rightarrow G$, $f(a, b) = ab^{-1}$

$\langle 1 \rangle 4$. $f^{-1}(U)$ is a neighbourhood of (x, y)

$\langle 1 \rangle 5$. PICK neighbourhoods V, W of x and y respectively such that $f(V \times W) \subseteq U$.

$\langle 1 \rangle 6$. PICK $a \in V \cap H$ and $b \in W \cap H$

PROOF: Theorem 52.

$\langle 1 \rangle 7$. $ab^{-1} \in U \cap H$

$\langle 1 \rangle 8$. Q.E.D.

PROOF: By Theorem 52.

□

Proposition 326. *Let G be a topological group and $\alpha \in G$. Then the maps $l_\alpha, r_\alpha : G \rightarrow G$ defined by $l_\alpha(x) = \alpha x$, $r_\alpha(x) = x\alpha$ are homeomorphisms of G with itself.*

PROOF: They are continuous with continuous inverses $l_{\alpha^{-1}}$ and $r_{\alpha^{-1}}$. \square

Corollary 326.1. *Every topological group is homogeneous.*

PROOF: Given a topological group G and $a, b \in G$, we have $l_{ba^{-1}}$ is a homeomorphism that maps a to b . \square

Proposition 327. *Let G be a topological group and H a subgroup of G . Give G/H the quotient topology. For all $\alpha \in G$, the map $\overline{f_\alpha}$ that sends xH to αxH is a homeomorphism $G/H \cong G/H$.*

PROOF:

$\langle 1 \rangle 1.$ $\overline{f_\alpha}$ is well-defined.

PROOF: If $xy^{-1} \in H$ then $(\alpha x)(\alpha y)^{-1} = xy^{-1} \in H$.

$\langle 1 \rangle 2.$ $\overline{f_\alpha}$ is continuous.

PROOF: Theorem 204 since $\overline{f_\alpha} \circ p = p \circ f_\alpha$ is continuous, where $p : G \twoheadrightarrow G/H$ is the canonical surjection.

$\langle 1 \rangle 3.$ $\overline{f_\alpha}^{-1}$ is continuous.

PROOF: Similar since $\overline{f_\alpha}^{-1} = \overline{f_{\alpha^{-1}}}$.

\square

Corollary 327.1. *Let G be a topological group and H a subgroup of G . Give G/H the quotient topology. Then G/H is homogeneous.*

Proposition 328. *Let G be a topological group and H a subgroup of G . Give G/H the quotient topology. If H is closed in G then G/H is T_1 .*

PROOF:

$\langle 1 \rangle 1.$ LET: $p : G \twoheadrightarrow G/H$ be the canonical surjection

$\langle 1 \rangle 2.$ LET: $x \in G$

$\langle 1 \rangle 3.$ $p^{-1}(xH) = f_x(H)$

$\langle 1 \rangle 4.$ $p^{-1}(xH)$ is closed in G

PROOF: Since H is closed and f_x is a homomorphism of G with itself.

$\langle 1 \rangle 5.$ $\{xH\}$ is closed in G/H

\square

Proposition 329. *Let G be a topological group and H a subgroup of G . Give G/H the quotient topology. Then the canonical surjection $p : G \twoheadrightarrow G/H$ is an open map.*

PROOF:

$\langle 1 \rangle 1.$ LET: $U \subseteq G$ be open.

$\langle 1 \rangle 2.$ $p^{-1}(p(U)) = \bigcup_{h \in H} r_h(U)$

$\langle 1 \rangle 3.$ $p^{-1}(p(U))$ is open.

$\langle 1 \rangle 4.$ $p(U)$ is open.

□

Proposition 330. *Let G be a topological group and H a closed normal subgroup of G . Then G/H is a topological group under the quotient topology.*

PROOF:

⟨1⟩1. G/H is T_1

PROOF: Proposition 328.

⟨1⟩2. The map $\bar{m} : (xH, yH) \mapsto xy^{-1}H$ is continuous.

⟨2⟩1. $p^2 : G^2 \rightarrow (G/H)^2$ is a quotient map.

PROOF: Propositions 203, 329.

⟨2⟩2. $\bar{m} \circ p^2$ is continuous.

PROOF: As it is $p^2 \circ m$ where $m : G^2 \rightarrow G$ with $m(x, y) = xy^{-1}$

□

Lemma 331. *Let G be a topological group and $A, B \subseteq G$. If either A or B is open then AB is open.*

PROOF: If A is open we have $AB = \bigcup_{b \in B} r_b(A)$. Similarly if B is open. □

Definition 332 (Symmetric Neighbourhood). Let G be a topological group. A neighbourhood V of e is *symmetric* if and only if $V = V^{-1}$.

Lemma 333. *Let G be a topological group. Let V be a neighbourhood of e . Then V is symmetric if and only if, for all $x \in V$, we have $x^{-1} \in V$.*

PROOF:

⟨1⟩1. If V is symmetric then, for all $x \in V$, we have $x^{-1} \in V$

PROOF: Immediate from definitions.

⟨1⟩2. If, for all $x \in V$, we have $x^{-1} \in V$, then V is symmetric.

⟨2⟩1. ASSUME: for all $x \in V$ we have $x^{-1} \in V$

⟨2⟩2. $V \subseteq V^{-1}$

PROOF: If $x \in V$ then there exists $y \in V$ such that $x = y^{-1}$, namely $y = x^{-1}$

⟨2⟩3. $V^{-1} \subseteq V$

PROOF: Immediate from ⟨2⟩1.

□

Lemma 334. *Let G be a topological group. For every neighbourhood U of e , there exists a symmetric neighbourhood V of e such that $V^2 \subseteq U$.*

PROOF:

⟨1⟩1. LET: U be a neighbourhood of e .

⟨1⟩2. PICK a neighbourhood V' of e such that $V'V' \subseteq U$

PROOF: Such a neighbourhood exists because multiplication in G is continuous.

⟨1⟩3. PICK a neighbourhood W of e such that $WW^{-1} \subseteq V'$

PROOF: Such a neighbourhood exists because the function that maps (x, y) to xy^{-1} is continuous.

- ⟨1⟩4. LET: $V = WW^{-1}$
- ⟨1⟩5. V is a neighbourhood of e
 - ⟨2⟩1. $e \in V$
 - PROOF: Since $e \in W$ so $e = ee^{-1} \in V$.
 - ⟨2⟩2. V is open
 - PROOF: Lemma 331.
- ⟨1⟩6. V is symmetric
 - ⟨2⟩1. For all $x \in V$ we have $x^{-1} \in V$
 - ⟨3⟩1. LET: $x \in V$
 - ⟨3⟩2. PICK $y, z \in W$ such that $x = yz^{-1}$
 - ⟨3⟩3. $x^{-1} = zy^{-1}$
 - ⟨3⟩4. $x^{-1} \in V$
 - ⟨3⟩5. $x \in V^{-1}$
 - ⟨2⟩2. Q.E.D.
 - PROOF: Lemma 333
- ⟨1⟩7. $V^2 \subseteq U$
 - PROOF: We have $V^2 \subseteq (V')^2 \subseteq U$

□

Proposition 335. *Every topological group is Hausdorff.*

PROOF:

- ⟨1⟩1. LET: G be a topological group.
- ⟨1⟩2. LET: $x, y \in G$ with $x \neq y$
- ⟨1⟩3. LET: $U = G \setminus \{x[^{-1}y]\}$
- ⟨1⟩4. PICK a symmetric neighbourhood V of e such that $VV \subseteq U$
 - ⟨2⟩1. U is open
 - PROOF: Since G is T_1 .
 - ⟨2⟩2. $e \in U$
 - PROOF: Since $x \neq y$
 - ⟨2⟩3. Q.E.D.
 - PROOF: Lemma 334.
- ⟨1⟩5. Vx and Vy are disjoint neighbourhoods of x and y respectively.
 - ⟨2⟩1. Vx is open
 - PROOF: Since $Vx = r_x(V)$
 - ⟨2⟩2. Vy is open
 - PROOF: Similar.
 - ⟨2⟩3. $Vx \cap Vy = \emptyset$
 - ⟨3⟩1. ASSUME: for a contradiction $z \in Vx \cap Vy$
 - ⟨3⟩2. PICK $a, b \in V$ such that $z = ax = by$
 - ⟨3⟩3. $xy^{-1} \in VV$
 - PROOF: Since $xy^{-1} = a^{-1}b$
 - ⟨3⟩4. $xy^{-1} \in U$
 - ⟨3⟩5. Q.E.D.
 - PROOF: From ⟨1⟩3.

□

Proposition 336. *Every topological group is regular.*

PROOF:

- ⟨1⟩1. LET: G be a topological group.
- ⟨1⟩2. LET: $A \subseteq G$ be a closed set and $a \notin A$.
- ⟨1⟩3. LET: $U = G \setminus Aa^{-1}$
- ⟨1⟩4. PICK a symmetric neighbourhood V of e such that $VV \subseteq U$
 - ⟨2⟩1. U is open
PROOF: Since $Aa^{-1} = r_{a^{-1}}(A)$ is closed.
 - ⟨2⟩2. $e \in U$
PROOF: Since $a \notin A$.
 - ⟨2⟩3. Q.E.D.
PROOF: Lemma 334.
- ⟨1⟩5. VA and Va are disjoint open sets with $A \subseteq VA$ and $a \in Va$
 - ⟨2⟩1. VA is open
PROOF: Lemma 331
 - ⟨2⟩2. Va is open
PROOF: Lemma 331
 - ⟨2⟩3. $VA \cap Va = \emptyset$
 - ⟨3⟩1. ASSUME: for a contradiction $z \in VA \cap Va$
 - ⟨3⟩2. PICK $b, c \in V$ and $d \in A$ with $z = bd = ca$
 - ⟨3⟩3. $da^{-1} \in U$
PROOF: Since $da^{-1} = b^{-1}c \in VV \subseteq U$
 - ⟨3⟩4. Q.E.D.
PROOF: This contradicts ⟨1⟩3

□

Proposition 337. *Let G be a topological group and H a subgroup of G . Give G/H the quotient topology. If H is closed in G then G/H is regular.*

PROOF:

- ⟨1⟩1. LET: $p : G \rightarrow G/H$ be the canonical surjection.
- ⟨1⟩2. LET: A be a closed set in G/H and $aH \in (G/H) \setminus A$.
- ⟨1⟩3. LET: $B = p^{-1}(A)$
- ⟨1⟩4. B is a closed saturated set in G .
- ⟨1⟩5. $B \cap aH = \emptyset$
- ⟨1⟩6. $B = BH$
- ⟨1⟩7. PICK a symmetric neighbourhood V of e such that VB does not intersect Va
 - ⟨2⟩1. LET: $U = G \setminus Ba^{-1}$
 - ⟨2⟩2. PICK a symmetric neighbourhood V of e such that $VV \subseteq U$
 - ⟨3⟩1. U is open
PROOF: Since $Ba^{-1} = r_{a^{-1}}(B)$ is closed.
 - ⟨3⟩2. $e \in U$
PROOF: If $e \in Ba^{-1}$ then $a \in B$
 - ⟨3⟩3. Q.E.D.
PROOF: Lemma 334

- $\langle 2 \rangle 3.$ $VB \cap Va = \emptyset$
 PROOF: If $vb = v'a$ for $v, v' \in V$ and $b \in B$ then we have $ba^{-1} = v^{-1}v' \in Ba \cap VV \subseteq Ba \cap U$.
 $\langle 1 \rangle 8.$ $p(VB)$ and $p(Va)$ are disjoint open sets
 $\langle 2 \rangle 1.$ $p(VB)$ and $p(Va)$ are open.
 PROOF: Proposition 329.
 $\langle 2 \rangle 2.$ $p(VB) \cap p(Va) = \emptyset$
 PROOF: If $vbH = v'aH$ for $v, v' \in V, b \in B$ then $v'a = vbh$ for some $h \in H$.
 Hence $v'a \in Va \cap VBH = Va \cap VB$.
 $\langle 1 \rangle 9.$ $A \subseteq p(VB)$
 $\langle 1 \rangle 10.$ $aH \in p(Va)$
 \square

Proposition 338. *Let G be a topological group. The component of G that contains e is a normal subgroup of G .*

PROOF:

- $\langle 1 \rangle 1.$ LET: C be the component of G that contains e .
 $\langle 1 \rangle 2.$ For all $x \in G$, xC is the component of G that contains x .
 $\langle 2 \rangle 1.$ LET: $x \in G$
 $\langle 2 \rangle 2.$ LET: D be the component of G that contains x .
 $\langle 2 \rangle 3.$ $xC \subseteq D$
 PROOF: Since xC is connected by Theorem 228.
 $\langle 2 \rangle 4.$ $D \subseteq xC$
 PROOF: Since $x^{-1}D \subseteq C$ similarly.
 $\langle 1 \rangle 3.$ For all $x \in G$, Cx is the component of G that contains x .
 PROOF: Similar.
 $\langle 1 \rangle 4.$ For all $x \in C$ we have $xC = Cx = C$
 $\langle 1 \rangle 5.$ For all $x \in C$ we have $x^{-1}C = C$
 $\langle 1 \rangle 6.$ For all $x \in C$ we have $x^{-1} \in C$
 $\langle 1 \rangle 7.$ For all $x, y \in C$ we have $xy \in C$
 PROOF: Since $xyC = xC = x$.
 $\langle 1 \rangle 8.$ For all $x \in G$ we have $xC = Cx$.
 PROOF: From $\langle 1 \rangle 2$ and $\langle 1 \rangle 3$.
 \square

Lemma 339. *Let G be a topological group. Let A be a closed set in G and B a compact subspace of G such that $A \cap B = \emptyset$. Then there exists a symmetric neighbourhood U of e such that $AU \cap BU = \emptyset$.*

PROOF:

- $\langle 1 \rangle 1.$ For all $b \in B$ there exists a symmetric neighbourhood V of e such that
 $bV^2 \cap A = \emptyset$
 $\langle 2 \rangle 1.$ LET: $b \in B$
 $\langle 2 \rangle 2.$ LET: $W = b^{-1}(G \setminus A)$
 $\langle 2 \rangle 3.$ W is a neighbourhood of e and $bW \cap A = \emptyset$
 $\langle 2 \rangle 4.$ PICK a symmetric neighbourhood V of e such that $V^2 \subseteq W$

- ⟨1⟩2. $\{bV^2 \mid b \in B, V \text{ is a symmetric neighbourhood of } e, bV^2 \cap A = \emptyset\}$ is an open cover of B
- ⟨1⟩3. PICK a finite subcover $b_1V_1^2, \dots, b_nV_n^2$, say.
- ⟨1⟩4. LET: $U = V_1 \cap \dots \cap V_n$
- ⟨1⟩5. $BU^2 \cap A = \emptyset$
- ⟨1⟩6. $AU \cap BU = \emptyset$

PROOF: If $av \in BU$ where $a \in A$ and $v \in V$ then $a = avv^{-1} \in BU^2 \cap A$.

□

Proposition 340 (AC). *Let G be a topological group. Let A be a closed set in G , and B a compact subspace of G . Then AB is closed.*

PROOF:

- ⟨1⟩1. LET: $x \in G \setminus AB$
- ⟨1⟩2. $A^{-1}x \cap B = \emptyset$
- ⟨1⟩3. $A^{-1}x$ is closed.
- ⟨1⟩4. PICK a symmetric neighbourhood U of e such that $A^{-1}xU \cap BU = \emptyset$
- ⟨1⟩5. xU^2 is open

PROOF: Lemma 331.

- ⟨1⟩6. $x \in xU^2 \subseteq G \setminus AB$

□

Corollary 340.1. *Let G be a topological group and $H \leq G$. Let $p : G \twoheadrightarrow G/H$ be the quotient map. If H is compact then p is a closed map.*

PROOF: For A closed in G , we have $p^{-1}(p(A)) = AH$ is closed, and so $p(A)$ is closed. □

Corollary 340.2. *Let G be a topological group and $H \leq G$. If H and G/H are compact then G is compact.*

PROOF: From Proposition 320 since, for all $aH \in G/H$, we have $p^{-1}(aH) = aH$ is compact because it is homomorphic to H . □

48 The Metric Topology

Definition 341 (Metric). Let X be a set. A *metric* on X is a function $d : X^2 \rightarrow \mathbb{R}$ such that:

1. For all $x, y \in X$, $d(x, y) \geq 0$
2. For all $x, y \in X$, $d(x, y) = 0$ if and only if $x = y$
3. For all $x, y \in X$, $d(x, y) = d(y, x)$
4. (*Triangle Inequality*) For all $x, y, z \in X$, $d(x, z) \leq d(x, y) + d(y, z)$

We call $d(x, y)$ the *distance* between x and y .

Definition 342 (Open Ball). Let X be a metric space. Let $a \in X$ and $\epsilon > 0$. The *open ball* with *centre* a and *radius* ϵ is

$$B(a, \epsilon) = \{x \in X \mid d(a, x) < \epsilon\} .$$

Definition 343 (Metric Topology). Let X be a metric space. The *metric topology* on X is the topology generated by the basis consisting of all the open balls.

We prove this is a basis for a topology.

PROOF:

$\langle 1 \rangle 1$. For every point a , there exists a ball B such that $a \in B$

PROOF: We have $a \in B(a, 1)$.

$\langle 1 \rangle 2$. For any balls B_1, B_2 and point $a \in B_1 \cap B_2$, there exists a ball B_3 such that $a \in B_3 \subseteq B_1 \cap B_2$

$\langle 2 \rangle 1$. LET: $B_1 = B(c_1, \epsilon_1)$ and $B_2 = B(c_2, \epsilon_2)$

$\langle 2 \rangle 2$. LET: $\delta = \min(\epsilon_1 - d(c_1, a), \epsilon_2 - d(c_2, a))$

PROVE: $B(a, \delta) \subseteq B_1 \cap B_2$

$\langle 2 \rangle 3$. LET: $x \in B(a, \delta)$

$\langle 2 \rangle 4$. $x \in B_1$

PROOF:

$$\begin{aligned} d(x, c_1) &= d(x, a) + d(a, c_1) \\ &< \delta + d(a, c_1) \\ &\leq \epsilon_1 \end{aligned}$$

$\langle 2 \rangle 5$. $x \in B_2$

PROOF: Similar.

□

Proposition 344. Let X be a metric space and $U \subseteq X$. Then U is open if and only if, for every $x \in U$, there exists $\epsilon > 0$ such that $B(x, \epsilon) \subseteq U$.

PROOF:

$\langle 1 \rangle 1$. If U is open then, for all $x \in U$, there exists $\epsilon > 0$ such that $B(x, \epsilon) \subseteq U$.

$\langle 2 \rangle 1$. ASSUME: U is open.

- ⟨2⟩2. LET: $x \in U$
 ⟨2⟩3. PICK $a \in X$ and $\delta > 0$ such that $x \in B(a, \delta) \subseteq U$
 ⟨2⟩4. LET: $\epsilon = \delta - d(a, x)$
 PROVE: $B(x, \epsilon) \subseteq U$
 ⟨2⟩5. LET: $y \in B(x, \epsilon)$
 ⟨2⟩6. $d(y, a) < \delta$
 PROOF:

$$\begin{aligned}
 d(y, a) &\leq d(a, x) + d(x, y) \\
 &< \delta + d(x, y) \\
 &= \epsilon
 \end{aligned}$$

- ⟨2⟩7. $y \in U$
 ⟨1⟩2. If, for all $x \in U$, there exists $\epsilon > 0$ such that $B(x, \epsilon) \subseteq U$, then U is open.
 PROOF: Immediate from definitions.

□

Definition 345 (Discrete Metric). Let X be a set. The *discrete metric* on X is defined by

$$d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

Proposition 346. *The discrete metric induces the discrete topology.*

PROOF: For any (open) set U and point $a \in U$, we have $a \in B(a, 1) \subseteq U$. □

Definition 347 (Standard Metric on \mathbb{R}). The *standard metric* on \mathbb{R} is defined by $d(x, y) = |x - y|$.

Proposition 348. *The standard metric on \mathbb{R} induces the standard topology on \mathbb{R} .*

PROOF:

- ⟨1⟩1. Every open ball is open in the standard topology on \mathbb{R} .
 PROOF: $B(a, \epsilon) = (a - \epsilon, a + \epsilon)$
 ⟨1⟩2. For every open set U and point $a \in U$, there exists $\epsilon > 0$ such that $B(a, \epsilon) \subseteq U$
 ⟨2⟩1. LET: U be an open set and $a \in U$
 ⟨2⟩2. PICK an open interval b, c such that $a \in (b, c) \subseteq U$
 ⟨2⟩3. LET: $\epsilon = \min(a - b, c - a)$
 ⟨2⟩4. $B(a, \epsilon) \subseteq U$

□

Definition 349 (Metriizable). A topological space X is *metrizable* if and only if there exists a metric on X that induces the topology.

Definition 350 (Bounded). Let X be a metric space and $A \subseteq X$. Then A is *bounded* if and only if there exists M such that, for all $x, y \in A$, we have $d(x, y) \leq M$.

Definition 351 (Diameter). Let X be a metric space and $A \subseteq X$. The *diameter* of A is

$$\text{diam } A = \sup_{x,y \in A} d(x,y) .$$

Definition 352 (Standard Bounded Metric). Let d be a metric on X . The *standard bounded metric* corresponding to d is the metric \bar{d} defined by

$$\bar{d}(x,y) = \min(d(x,y), 1) .$$

We prove this is a metric.

PROOF:

$\langle 1 \rangle 1. \bar{d}(x,y) \geq 0$

PROOF: Since $d(x,y) \geq 0$

$\langle 1 \rangle 2. \bar{d}(x,y) = 0$ if and only if $x = y$

PROOF: $\bar{d}(x,y) = 0$ if and only if $d(x,y) = 0$ if and only if $x = y$

$\langle 1 \rangle 3. \bar{d}(x,y) = \bar{d}(y,x)$

PROOF: Since $d(x,y) = d(y,x)$

$\langle 1 \rangle 4. \bar{d}(x,z) \leq \bar{d}(x,y) + \bar{d}(y,z)$

PROOF:

$$\begin{aligned} \bar{d}(x,y) + \bar{d}(y,z) &= \min(d(x,y), 1) + \min(d(y,z), 1) \\ &= \min(d(x,y) + d(y,z), d(x,y) + 1, d(y,z) + 1, 2) \\ &\geq \min(d(x,z), 1) \\ &= \bar{d}(x,z) \end{aligned}$$

□

Lemma 353. In any metric space X , the set $\mathcal{B} = \{B(a, \epsilon) \mid a \in X, \epsilon < 1\}$ is a basis for the metric topology.

PROOF:

$\langle 1 \rangle 1.$ Every element of \mathcal{B} is open.

PROOF: From Lemma 67.

$\langle 1 \rangle 2.$ For every open set U and point $a \in U$, there exists $B \in \mathcal{B}$ such that $a \in B \subseteq U$

$\langle 2 \rangle 1.$ LET: U be an open set and $a \in U$

$\langle 2 \rangle 2.$ PICK $\epsilon > 0$ such that $B(a, \epsilon) \subseteq U$

$\langle 2 \rangle 3.$ $B(a, \min(\epsilon, 1/2)) \subseteq U$

$\langle 1 \rangle 3.$ Q.E.D.

PROOF: Lemma 68.

□

Proposition 354. Let d be a metric on the set X . Then the standard bounded metric \bar{d} induces the same metric as d .

PROOF: This follows from Lemma 353 since the open balls with radius < 1 are the same under both metrics. □

Lemma 355. *Let d and d' be two metrics on the same set X . Let \mathcal{T} and \mathcal{T}' be the topologies they induce. Then $\mathcal{T} \subseteq \mathcal{T}'$ if and only if, for all $x \in X$ and $\epsilon > 0$, there exists $\delta > 0$ such that*

$$B_{d'}(x, \delta) \subseteq B_d(x, \epsilon) .$$

PROOF:

$\langle 1 \rangle 1$. If $\mathcal{T} \subseteq \mathcal{T}'$ then, for all $x \in X$ and $\epsilon > 0$, there exists $\delta > 0$ such that $B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$

PROOF: From Proposition 344 since $x \in B_d(x, \epsilon) \in \mathcal{T}'$.

$\langle 1 \rangle 2$. If, for all $x \in X$ and $\epsilon > 0$, there exists $\delta > 0$ such that $B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$, then $\mathcal{T} \subseteq \mathcal{T}'$

$\langle 2 \rangle 1$. ASSUME: For all $x \in X$ and $\epsilon > 0$, there exists $\delta > 0$ such that $B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$

$\langle 2 \rangle 2$. LET: $U \in \mathcal{T}$

$\langle 2 \rangle 3$. For all $x \in U$ there exists $\delta > 0$ such that $B_{d'}(x, \delta) \subseteq U$.

$\langle 3 \rangle 1$. LET: $x \in U$

$\langle 3 \rangle 2$. PICK $\epsilon > 0$ such that $B_d(x, \epsilon) \subseteq U$

PROOF: Proposition 344

$\langle 3 \rangle 3$. PICK $\delta > 0$ such that $B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$

PROOF: By $\langle 2 \rangle 1$

$\langle 3 \rangle 4$. $B_{d'}(x, \delta) \subseteq U$

$\langle 2 \rangle 4$. $U \in \mathcal{T}'$

PROOF: Proposition 344.

□

Proposition 356. \mathbb{R}^2 under the dictionary order topology is metrizable.

PROOF: Define $d : \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$d((x, y), (x, z)) = \max(|y - z|, 1)$$

$$d((x, y), (x', y')) = 1$$

$$\text{if } x \neq x' \square$$

$\langle 1 \rangle 1$. $x \in \bigcap_{i=1}^N \pi_i^{-1}() \subseteq B_D(a, \epsilon)$

Proposition 357. *Let $d : X^2 \rightarrow \mathbb{R}$ be a metric on X . Then the metric topology on X is the coarsest topology such that d is continuous.*

PROOF:

$\langle 1 \rangle 1$. d is continuous.

$\langle 2 \rangle 1$. LET: $a, b \in X$

$\langle 2 \rangle 2$. LET: $\epsilon > 0$

$\langle 2 \rangle 3$. LET: $\delta = \epsilon/2$

$\langle 2 \rangle 4$. LET: $x, y \in X$

$\langle 2 \rangle 5$. ASSUME: $\rho((a, b), (x, y)) < \delta$

$\langle 2 \rangle 6$. $|d(a, b) - d(x, y)| < \epsilon$

$\langle 3 \rangle 1$. $d(a, b) - d(x, y) < \epsilon$

PROOF:

$$\begin{aligned}
d(a, b) &\leq d(a, x) + d(x, y) + d(y, b) \\
&\leq d(x, y) + 2\rho((a, b), (x, y)) \\
&< d(x, y) + 2\delta \\
&= d(x, y) + \epsilon
\end{aligned}$$

$$\langle 3 \rangle 2. \ d(a, b) - d(x, y) > -\epsilon$$

PROOF: Similar.

$\langle 2 \rangle 7.$ Q.E.D.

$\langle 1 \rangle 2.$ If \mathcal{T} is any topology under which d is continuous then \mathcal{T} is finer than the metric topology.

PROOF: Since $B(a, \epsilon) = d_a^{-1}((-\infty, \epsilon))$

□

Proposition 358. *Let X be a metric space with metric d and $A \subseteq X$. The restriction of d to A is a metric on A that induces the subspace topology.*

PROOF:

$\langle 1 \rangle 1.$ The restriction of d to A is a metric on A .

$\langle 1 \rangle 2.$ Every open ball under $d \upharpoonright A$ is open under the subspace topology.

PROOF: $B_{d \upharpoonright A}(a, \epsilon) = B_d(a, \epsilon) \cap A$.

$\langle 1 \rangle 3.$ If U is open in the subspace topology and $x \in U$, then there exists a $d \upharpoonright A$ -ball B such that $x \in B \subseteq U$.

$\langle 2 \rangle 1.$ PICK V open in X such that $U = V \cap A$

$\langle 2 \rangle 2.$ PICK $\epsilon > 0$ such that $B_d(x, \epsilon) \subseteq V$

$\langle 2 \rangle 3.$ Take $B = B_{d \upharpoonright A}(x, \epsilon)$

□

Corollary 358.1. *A subspace of a metrizable space is metrizable.*

Proposition 359. *Every metrizable space is Hausdorff.*

PROOF:

$\langle 1 \rangle 1.$ LET: X be a metric space

$\langle 1 \rangle 2.$ LET: $a, b \in X$ with $a \neq b$

$\langle 1 \rangle 3.$ LET: $\epsilon = d(a, b)/2$

$\langle 1 \rangle 4.$ LET: $U = B(a, \epsilon)$ and $V = B(b, \epsilon)$

$\langle 1 \rangle 5.$ U and V are disjoint neighbourhoods of a and b respectively.

□

Proposition 360 (CC). *The product of a countable family of metrizable spaces is metrizable.*

PROOF:

$\langle 1 \rangle 1.$ LET: (X_n, d_n) be a sequence of metric spaces.

$\langle 1 \rangle 2.$ ASSUME: w.l.o.g. each d_n is bounded above by 1.

PROOF: By Proposition 354.

$\langle 1 \rangle 3.$ LET: D be the metric on \mathbb{R}^ω defined by $D(x, y) = \sup_i (d_i(x_i, y_i)/i)$.

- ⟨2⟩1. $D(x, y) \geq 0$
- ⟨2⟩2. $D(x, y) = 0$ if and only if $x = y$
- ⟨2⟩3. $D(x, y) = D(y, x)$
- ⟨2⟩4. $D(x, z) \leq D(x, y) + D(y, z)$

PROOF:

$$\begin{aligned}
 D(x, z) &= \sup_i \frac{d_i(x_i, z_i)}{i} \\
 &\leq \sup_i \frac{d_i(x_i, y_i) + d_i(y_i, z_i)}{i} \\
 &\leq \sup_i \frac{d_i(x_i, y_i)}{i} + \sup_i \frac{d_i(y_i, z_i)}{i} \\
 &= D(x, y) + D(y, z)
 \end{aligned}$$

- ⟨1⟩4. Every open ball $B_D(a, \epsilon)$ is open in the product topology.
- ⟨2⟩1. PICK N such that $1/\epsilon < N$
- ⟨2⟩2. $B_D(a, \epsilon) = \prod_{i=1}^{\infty} U_i$ where $U_i = B(a_i, i\epsilon)$ if $i \leq N$, and $U_i = X_i$ if $i > N$
- ⟨1⟩5. For any open set U and $a \in U$, there exists $\epsilon > 0$ such that $B_D(a, \epsilon) \subseteq U$.
- ⟨2⟩1. LET: $n \geq 1$, V be an open set in \mathbb{R} and $a \in \pi_n^{-1}(V)$
- ⟨2⟩2. PICK $\epsilon > 0$ such that $B_{d_n}(a, \epsilon) \subseteq V$
- ⟨2⟩3. $B_D(a, \epsilon/n) \subseteq \pi_n^{-1}(V)$

□

Theorem 361. *Let X and Y be metric spaces and $f : X \rightarrow Y$. Then f is continuous if and only if, for all $x \in X$ and $\epsilon > 0$, there exists $\delta > 0$ such that, for all $y \in X$, if $d(x, y) < \delta$ then $d(f(x), f(y)) < \epsilon$.*

PROOF:

- ⟨1⟩1. If f is continuous then, for all $x \in X$ and $\epsilon > 0$, there exists $\delta > 0$ such that, for all $y \in X$, if $d(x, y) < \delta$ then $d(f(x), f(y)) < \epsilon$
- ⟨2⟩1. ASSUME: f is continuous.
- ⟨2⟩2. LET: $x \in X$ and $\epsilon > 0$
- ⟨2⟩3. PICK a neighbourhood U of x such that $f(U) \subseteq B(f(x), \epsilon)$
- PROOF: Theorem 98.
- ⟨2⟩4. PICK $\delta > 0$ such that $B(x, \delta) \subseteq U$
- PROOF: Proposition 344.
- ⟨2⟩5. For all $y \in X$, if $d(x, y) < \delta$ then $d(f(x), f(y)) < \epsilon$
- ⟨1⟩2. If for all $x \in X$ and $\epsilon > 0$, there exists $\delta > 0$ such that, for all $y \in X$, if $d(x, y) < \delta$ then $d(f(x), f(y)) < \epsilon$, then f is continuous.
- ⟨2⟩1. ASSUME: for all $x \in X$ and $\epsilon > 0$, there exists $\delta > 0$ such that, for all $y \in X$, if $d(x, y) < \delta$ then $d(f(x), f(y)) < \epsilon$
- ⟨2⟩2. LET: $x \in X$ and V be a neighbourhood of $f(x)$
- ⟨2⟩3. PICK $\epsilon > 0$ such that $B(f(x), \epsilon) \subseteq V$
- PROOF: Proposition 344.
- ⟨2⟩4. PICK $\delta > 0$ such that, for all $y \in X$, if $d(x, y) < \delta$ then $d(f(x), f(y)) < \epsilon$
- PROOF: By ⟨2⟩1
- ⟨2⟩5. LET: $U = B(x, \delta)$
- ⟨2⟩6. U is a neighbourhood of x with $f(U) \subseteq V$

⟨2⟩7. Q.E.D.

PROOF: Theorem 98.

□

Proposition 362. *Let X be a metric space. Let (a_n) be a sequence in X and $l \in X$. Then $a_n \rightarrow l$ as $n \rightarrow \infty$ if and only if, for all $\epsilon > 0$, there exists N such that, for all $n \geq N$, we have $d(a_n, l) < \epsilon$.*

PROOF: From Proposition 81. □

Proposition 363. *Every metrizable space is first countable.*

PROOF: In any metric space X , the open balls $B(a, 1/n)$ for $n \geq 1$ form a local basis at a .

Example 364. \mathbb{R}^ω under the box topology is not metrizable.

Example 365. If J is uncountable then \mathbb{R}^J under the product topology is not metrizable.

Proposition 366. *A compact subspace of a metric space is bounded.*

PROOF:

⟨1⟩1. LET: X be a metric space and $A \subseteq X$ be compact.

⟨1⟩2. PICK $a \in A$

⟨1⟩3. $\{B(a, n) \mid n \in \mathbb{Z}^+\}$ covers A

⟨1⟩4. PICK a finite subcover $\{B(a, n_1), \dots, B(a, n_k)\}$

⟨1⟩5. LET: $N = \max(n_1, \dots, n_k)$

⟨1⟩6. For all $x, y \in A$ we have $d(x, y) < 2N$

PROOF:

$$\begin{aligned} d(x, y) &\leq d(x, a) + d(a, y) \\ &< N + N \end{aligned}$$

□

This example shows the converse does not hold:

Example 367. The space \mathbb{R} under the standard bounded metric is bounded but not compact.

49 Real Linear Algebra

Definition 368 (Square Metric). The *square metric* ρ on \mathbb{R}^n is defined by

$$\rho(\vec{x}, \vec{y}) = \max(|x_1 - y_1|, \dots, |x_n - y_n|)$$

We prove this is a metric.

PROOF:

⟨1⟩1. $\rho(\vec{x}, \vec{y}) \geq 0$

PROOF: Immediate from definition.

- $\langle 1 \rangle 2.$ $\rho(\vec{x}, \vec{y}) = 0$ if and only if $\vec{x} = \vec{y}$
 PROOF: Immediate from definition.
 $\langle 1 \rangle 3.$ $\rho(\vec{x}, \vec{y}) = \rho(\vec{y}, \vec{x})$
 PROOF: Immediate from definition.
 $\langle 1 \rangle 4.$ $\rho(\vec{x}, \vec{z}) \leq \rho(\vec{x}, \vec{y}) + \rho(\vec{y}, \vec{z})$
 PROOF: Since $|x_i - z_i| \leq |x_i - y_i| + |y_i - z_i|$.
 \square

Proposition 369. *The square metric induces the standard topology on \mathbb{R}^n .*

PROOF:

- $\langle 1 \rangle 1.$ For every $a \in X$ and $\epsilon > 0$, we have $B_\rho(a, \epsilon)$ is open in the standard product topology.

PROOF:

$$B_\rho(a, \epsilon) = (a_1 - \epsilon, a_1 + \epsilon) \times \cdots \times (a_n - \epsilon, a_n + \epsilon)$$

- $\langle 1 \rangle 2.$ For any open sets U_1, \dots, U_n in \mathbb{R} , we have $U_1 \times \cdots \times U_n$ is open in the square metric topology.
 $\langle 2 \rangle 1.$ LET: $\vec{a} \in U_1 \times \cdots \times U_n$
 $\langle 2 \rangle 2.$ For $i = 1, \dots, n$, PICK $\epsilon_i > 0$ such that $(a_i - \epsilon_i, a_i + \epsilon_i) \subseteq U_i$
 $\langle 2 \rangle 3.$ LET: $\epsilon = \min(\epsilon_1, \dots, \epsilon_n)$
 $\langle 2 \rangle 4.$ $B_\rho(\vec{a}, \epsilon) \subseteq U$
 \square

Definition 370. Given $\vec{x}, \vec{y} \in \mathbb{R}^n$, define the *sum* $\vec{x} + \vec{y}$ by

$$(x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n) .$$

Definition 371. Given $\lambda \in \mathbb{R}$ and $\vec{x} \in \mathbb{R}^n$, define the *scalar product* $\lambda \vec{x} \in \mathbb{R}^n$ by

$$\lambda(x_1, \dots, x_n) = (\lambda x_1, \dots, \lambda x_n)$$

Definition 372 (Inner Product). Given $\vec{x}, \vec{y} \in \mathbb{R}^n$, define the *inner product* $\vec{x} \cdot \vec{y} \in \mathbb{R}$ by

$$(x_1, \dots, x_n) \cdot (y_1, \dots, y_n) = x_1 y_1 + \cdots + x_n y_n .$$

We write \vec{x}^2 for $\vec{x} \cdot \vec{x}$.

Definition 373 (Norm). Let $n \geq 1$. The *norm* on \mathbb{R}^n is the function $\| \cdot \| : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$\|(x_1, \dots, x_n)\| = \sqrt{x_1^2 + \cdots + x_n^2}$$

Lemma 374.

$$\|\vec{x}\|^2 = \vec{x}^2$$

PROOF: Immediate from definitions. \square

Lemma 375.

$$\vec{x} \cdot (\vec{y} + \vec{z}) = \vec{x} \cdot \vec{y} + \vec{x} \cdot \vec{z}$$

PROOF: Each is equal to $(x_1y_1 + x_1z_1, \dots, x_ny_n + x_nz_n)$. \square

Lemma 376.

$$|\vec{x} \cdot \vec{y}| \leq \|\vec{x}\| \|\vec{y}\|$$

PROOF:

$\langle 1 \rangle 1$. ASSUME: $\vec{x} \neq \vec{0} \neq \vec{y}$

PROOF: Otherwise both sides are 0.

$\langle 1 \rangle 2$. LET: $a = 1/\|\vec{x}\|$

$\langle 1 \rangle 3$. LET: $b = 1/\|\vec{y}\|$

$\langle 1 \rangle 4$. $(a\vec{x} + b\vec{y})^2 \geq 0$ and $(a\vec{x} - b\vec{y})^2 \geq 0$

$\langle 1 \rangle 5$. $a^2\|\vec{x}\|^2 + 2ab\vec{x} \cdot \vec{y} + b^2\|\vec{y}\|^2 \geq 0$ and $a^2\|\vec{x}\|^2 - 2ab\vec{x} \cdot \vec{y} + b^2\|\vec{y}\|^2 \geq 0$

$\langle 1 \rangle 6$. $2ab\vec{x} \cdot \vec{y} + 2 \geq 0$ and $-2ab\vec{x} \cdot \vec{y} + 2 \geq 0$

$\langle 1 \rangle 7$. $\vec{x} \cdot \vec{y} \geq -1/ab$ and $\vec{x} \cdot \vec{y} \leq 1/ab$

$\langle 1 \rangle 8$. $\vec{x} \cdot \vec{y} \geq -\|\vec{x}\| \|\vec{y}\|$ and $\vec{x} \cdot \vec{y} \leq \|\vec{x}\| \|\vec{y}\|$

\square

Lemma 377 (Triangle Inequality).

$$\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$$

PROOF:

$$\begin{aligned} \|\vec{x} + \vec{y}\|^2 &= \|\vec{x}\|^2 + 2\vec{x} \cdot \vec{y} + \|\vec{y}\|^2 \\ &\leq \|\vec{x}\|^2 + 2\|\vec{x}\| \|\vec{y}\| + \|\vec{y}\|^2 && \text{(Lemma 376)} \\ &= (\|\vec{x}\| + \|\vec{y}\|)^2 && \square \end{aligned}$$

Definition 378 (Euclidean Metric). Let $n \geq 1$. The *Euclidean metric* on \mathbb{R}^n is defined by

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

We prove this is a metric.

$\langle 1 \rangle 1$. $d(\vec{x}, \vec{y}) \geq 0$

PROOF: Immediate from definition.

$\langle 1 \rangle 2$. $d(\vec{x}, \vec{y}) = 0$ if and only if $\vec{x} = \vec{y}$

PROOF: $d(\vec{x}, \vec{y}) = 0$ if and only if $\vec{x} - \vec{y} = \vec{0}$.

$\langle 1 \rangle 3$. $d(\vec{x}, \vec{y}) = d(\vec{y}, \vec{x})$

PROOF: Immediate from definition.

$\langle 1 \rangle 4$. $d(\vec{x}, \vec{z}) \leq d(\vec{x}, \vec{y}) + d(\vec{y}, \vec{z})$

PROOF:

$$\begin{aligned} \|\vec{x} - \vec{z}\| &= \|(\vec{x} - \vec{y}) + (\vec{y} - \vec{z})\| \\ &\leq \|\vec{x} - \vec{y}\| + \|\vec{y} - \vec{z}\| && \text{(Lemma 377)} \end{aligned}$$

\square

Proposition 379. The Euclidean metric induces the standard topology on \mathbb{R}^n .

PROOF:

$\langle 1 \rangle 1$. LET: ρ be the square metric.

- (1)2. For all $\vec{a} \in \mathbb{R}^n$ and $\epsilon > 0$, we have $B_d(\vec{a}, \epsilon) \subseteq B_\rho(\vec{a}, \epsilon)$
 (2)1. LET: $\vec{x} \in B_d(\vec{a}, \epsilon)$
 (2)2. $\sqrt{(x_1 - a_1)^2 + \cdots + (x_n - a_n)^2} < \epsilon$
 (2)3. $(x_1 - a_1)^2 + \cdots + (x_n - a_n)^2 < \epsilon^2$
 (2)4. For all i we have $(x_i - a_i)^2 < \epsilon^2$
 (2)5. For all i we have $|x_i - a_i| < \epsilon$
 (2)6. $\rho(\vec{x}, \vec{a}) < \epsilon$
 (1)3. For all $\vec{a} \in \mathbb{R}^n$ and $\epsilon > 0$, we have $B_\rho(\vec{a}, \epsilon/\sqrt{n}) \subseteq B_d(\vec{a}, \epsilon)$
 (2)1. LET: $\vec{x} \in B_\rho(\vec{a}, \epsilon/\sqrt{n})$
 (2)2. $\rho(\vec{x}, \vec{a}) < \epsilon/\sqrt{n}$
 (2)3. For all i we have $|x_i - a_i| < \epsilon/\sqrt{n}$
 (2)4. For all i we have $(x_i - a_i)^2 < \epsilon^2/n$
 (2)5. $d(\vec{x}, \vec{a}) < \epsilon$
 (1)4. Q.E.D.
 PROOF: By Lemma 355.

□

Proposition 380. *Let $n \geq 0$. For all $c \in \mathbb{R}^n$ and $\epsilon > 0$, the open ball $B(c, \epsilon)$ is path connected.*

PROOF:

- (1)1. LET: $a, b \in B(c, \epsilon)$
 (1)2. LET: $p : [0, 1] \rightarrow B(c, \epsilon)$ be the function $p(t) = (1 - t)a + tb$
 PROOF: We have $p(t) \in B(c, \epsilon)$ for all t because

$$\begin{aligned}
 d(p(t), c) &= \|(1 - t)a + tb - c\| \\
 &= \|(1 - t)(a - c) + t(b - c)\| \\
 &\leq (1 - t)\|a - c\| + t\|b - c\| \\
 &< (1 - t)\epsilon + t\epsilon \\
 &= \epsilon
 \end{aligned}$$

- (1)3. p is a path from a to b .

□

Proposition 381. *Let $n \geq 0$. For all $c \in \mathbb{R}^n$ and $\epsilon > 0$, the closed ball $\overline{B(c, \epsilon)}$ is path connected.*

PROOF:

- (1)1. LET: $a, b \in \overline{B(c, \epsilon)}$
 (1)2. LET: $p : [0, 1] \rightarrow \overline{B(c, \epsilon)}$ be the function $p(t) = (1 - t)a + tb$
 PROOF: We have $p(t) \in \overline{B(c, \epsilon)}$ for all t because

$$\begin{aligned}
 d(p(t), c) &= \|(1 - t)a + tb - c\| \\
 &= \|(1 - t)(a - c) + t(b - c)\| \\
 &\leq (1 - t)\|a - c\| + t\|b - c\| \\
 &\leq (1 - t)\epsilon + t\epsilon \\
 &= \epsilon
 \end{aligned}$$

- (1)3. p is a path from a to b .

□

Lemma 382. *If $\sum_{i=0}^{\infty} x_i^2$ and $\sum_{i=0}^{\infty} y_i^2$ converge then $\sum_{i=0}^{\infty} |x_i y_i|$ converges.*

PROOF:

⟨1⟩1. For all $N \geq 0$ we have $\sum_{i=0}^N |x_i y_i| \leq \sqrt{\sum_{i=0}^N |x_i|^2} \sqrt{\sum_{i=0}^N |y_i|^2}$

PROOF: By the Cauchy-Schwarz inequality

⟨1⟩2. Q.E.D.

PROOF: Since $\sum_{i=0}^N |x_i y_i|$ is an increasing sequence bounded above by $(\sum_{i=0}^{\infty} x_i^2)(\sum_{i=0}^{\infty} y_i^2)$.

□

Corollary 382.1. *If $\sum_{i=0}^{\infty} x_i^2$ and $\sum_{i=0}^{\infty} y_i^2$ converge then $\sum_{i=0}^{\infty} (x_i + y_i)^2$ converges.*

PROOF: Since $\sum_{i=0}^{\infty} x_i^2$, $\sum_{i=0}^{\infty} y_i^2$ and $2 \sum_{i=0}^{\infty} x_i y_i$ all converge.

Definition 383 (l^2 -metric). The l^2 -metric on

$$\left\{ (x_n) \in \mathbb{R}^{\omega} \mid \sum_{n=0}^{\infty} x_n^2 \text{ converges} \right\}$$

is defined by

$$d(x, y) = \left(\sum_{n=0}^{\infty} (x_n - y_n)^2 \right)^{1/2}$$

We prove this is a metric.

PROOF:

⟨1⟩1. d is well-defined.

PROOF: By Corollary 382.1.

⟨1⟩2. $d(x, y) \geq 0$

⟨1⟩3. $d(x, y) = 0$ if and only if $x = y$

⟨1⟩4. $d(x, y) = d(y, x)$

⟨1⟩5. $d(x, z) \leq d(x, y) + d(y, z)$

PROOF: By Lemma 377.

□

Theorem 384. *Addition is a continuous function $\mathbb{R}^2 \rightarrow \mathbb{R}$.*

PROOF:

⟨1⟩1. LET: $a, b \in \mathbb{R}$

⟨1⟩2. LET: $\epsilon > 0$

⟨1⟩3. LET: $\delta = \epsilon/2$

⟨1⟩4. LET: $x, y \in \mathbb{R}$

⟨1⟩5. ASSUME: $\rho((a, b), (x, y)) < \delta$

⟨1⟩6. $|(a + b) - (x + y)| < \epsilon$

PROOF:

$$\begin{aligned}
|(a+b) - (x+y)| &= |a-x| + |b-y| \\
&\leq 2\rho((a,b), (x,y)) \\
&< 2\delta \\
&= \epsilon
\end{aligned}$$

⟨1⟩7. Q.E.D.

PROOF: Theorem 361

□

Theorem 385. *Multiplication is a continuous function $\mathbb{R}^2 \rightarrow \mathbb{R}$.*

PROOF:

⟨1⟩1. LET: $a, b \in \mathbb{R}$

⟨1⟩2. LET: $\epsilon > 0$

⟨1⟩3. LET: $\delta = \min(\epsilon/(|a| + |b| + 1), 1)$

⟨1⟩4. LET: $x, y \in \mathbb{R}$

⟨1⟩5. ASSUME: $\rho((a,b), (x,y)) < \delta$

⟨1⟩6. $|ab - xy| < \epsilon$

PROOF:

$$\begin{aligned}
|ab - xy| &= |a(b-y) + (a-x)b - (a-x)(b-y)| \\
&\leq |a||b-y| + |b||a-x| + |a-x||b-y| \\
&< |a|\delta + |b|\delta + \delta^2 && (\langle 1 \rangle 5) \\
&\leq |a|\delta + |b|\delta + \delta && (\langle 1 \rangle 3) \\
&\leq \epsilon && (\langle 1 \rangle 3)
\end{aligned}$$

⟨1⟩7. Q.E.D.

PROOF: Theorem 361

□

Theorem 386. *The function $f : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ defined by $f(x) = x^{-1}$ is continuous.*

PROOF:

⟨1⟩1. For all $a \in \mathbb{R}$ we have $f^{-1}((a, +\infty))$ is open.

PROOF: The set is

$$(a^{-1}, +\infty) \text{ if } a > 0$$

$$(0, +\infty) \text{ if } a = 0$$

$$(-\infty, a^{-1}) \cup (0, +\infty) \text{ if } a < 0$$

⟨1⟩2. For all $a \in \mathbb{R}$ we have $f^{-1}((-\infty, a))$ is open.

PROOF: Similar.

⟨1⟩3. Q.E.D.

PROOF: By Proposition 95 and Lemma 118.

□

Definition 387. For $n \geq 0$, the *unit ball* B^n is the space $\{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$.

Proposition 388. *For all $n \geq 0$, the unit ball B^n is path connected.*

PROOF:

⟨1⟩1. LET: $a, b \in B^n$

⟨1⟩2. LET: $p : [0, 1] \rightarrow B^n$ be the function $p(t) = (1 - t)a + tb$

PROOF: We have $p(t) \in B^n$ for all t because

$$\begin{aligned}\|(1 - t)a + tb\| &\leq (1 - t)\|a\| + t\|b\| \\ &\leq (1 - t) + t \\ &= 1\end{aligned}$$

⟨1⟩3. p is a path from a to b .

□

Definition 389 (Punctured Euclidean Space). For $n \geq 0$, defined *punctured Euclidean space* to be $\mathbb{R}^n \setminus \{0\}$.

Proposition 390. For $n > 1$, *punctured Euclidean space* $\mathbb{R}^n \setminus \{0\}$ is path connected.

PROOF:

⟨1⟩1. LET: $a, b \in \mathbb{R}^n \setminus \{0\}$

⟨1⟩2. CASE: 0 is on the line from a to b

⟨2⟩1. PICK a point c not on the line from a to b

⟨2⟩2. The path consisting of a straight line from a to c followed by a straight line from c to b is a path from a to b .

⟨1⟩3. CASE: 0 is not on the line from a to b

PROOF: The straight line from a to b is a path from a to b .

Corollary 390.1. For $n > 1$, the spaces \mathbb{R} and \mathbb{R}^n are not homeomorphic.

PROOF: For any point a , the space $\mathbb{R} \setminus \{a\}$ is disconnected.

Definition 391 (Unit Sphere). For $n \geq 1$, the *unit sphere* S^{n-1} is the space

$$S^{n-1} = \{x \in \mathbb{R}^n \mid \|x\| = 1\} .$$

Proposition 392. For $n > 1$, the unit sphere S^{n-1} is path connected.

PROOF: The map $g : \mathbb{R}^n \setminus \{0\} \rightarrow S^{n-1}$ defined by $g(x) = x/\|x\|$ is continuous and surjective. The result follows by Proposition 249. □

Proposition 393. Let $f : S^1 \rightarrow \mathbb{R}$ be continuous. Then there exists $x \in S^1$ such that $f(x) = f(-x)$.

PROOF:

⟨1⟩1. LET: $g : S^1 \rightarrow \mathbb{R}$ be the function $g(x) = f(x) - f(-x)$

PROVE: There exists $x \in S^1$ such that $g(x) = 0$

⟨1⟩2. ASSUME: without loss of generality $g((1, 0)) > 0$

⟨1⟩3. $g((-1, 0)) < 0$

⟨1⟩4. There exists x such that $g(x) = 0$

PROOF: By the Intermediate Value Theorem.

□

Definition 394 (Topologist's Sine Curve). Let $S = \{(x, \sin 1/x) \mid 0 < x \leq 1\}$. The *topologist's sine curve* is the closure \bar{S} of S .

Proposition 395.

$$\bar{S} = S \cup (\{0\} \times [-1, 1])$$

Proposition 396. *The topologist's sine curve is connected.*

PROOF:

$\langle 1 \rangle 1$. LET: $S = \{(x, \sin 1/x) \mid 0 < x \leq 1\}$

$\langle 1 \rangle 2$. S is connected.

PROOF: Theorem 228.

$\langle 1 \rangle 3$. \bar{S} is connected.

PROOF: Theorem 227.

□

Proposition 397 (CC). *The topologist's sine curve is not path connected.*

PROOF:

$\langle 1 \rangle 1$. ASSUME: for a contradiction $p : [0, 1] \rightarrow \bar{S}$ is a path from $(0, 0)$ to $(1, \sin 1)$.

$\langle 1 \rangle 2$. $p^{-1}(\{0\} \times [0, 1])$ is closed.

$\langle 1 \rangle 3$. LET: b be the greatest element of $p^{-1}(\{0\} \times [0, 1])$.

$\langle 1 \rangle 4$. $b < 1$

PROOF: Since $p(1) = (1, \sin 1)$.

$\langle 1 \rangle 5$. PICK a sequence $(t_n)_{n \geq 1}$ in $(b, 1]$ such that $t_n \rightarrow b$ and $\pi_2(p(t_n)) = (-1)^n$

$\langle 2 \rangle 1$. LET: $n \geq 1$

$\langle 2 \rangle 2$. PICK u with $0 < u < \pi_1(p(1/n))$ such that $\sin(1/u) = (-1)^n$

$\langle 2 \rangle 3$. PICK t_n such that $0 < t_n < 1/n$ and $\pi_1(p(t_n)) = u$

PROOF: One exists by the Intermediate Value Theorem.

$\langle 1 \rangle 6$. Q.E.D.

PROOF: This contradicts 110.

□

50 The Uniform Topology

Definition 398 (Uniform Metric). Let J be a set. The *uniform metric* $\bar{\rho}$ on \mathbb{R}^J is defined by

$$\bar{\rho}(a, b) = \sup_{j \in J} \bar{d}(a_j, b_j)$$

where \bar{d} is the standard bounded metric on \mathbb{R} .

The *uniform topology* on \mathbb{R}^J is the topology induced by the uniform metric.

We prove this is a metric.

PROOF:

$\langle 1 \rangle 1$. $\bar{\rho}(a, b) \geq 0$

PROOF: Immediate from definitions.

$\langle 1 \rangle 2$. $\bar{\rho}(a, b) = 0$ if and only if $a = b$

PROOF: Immediate from definitions.

⟨1⟩3. $\bar{\rho}(a, b) = \bar{\rho}(b, a)$

PROOF: Immediate from definitions.

⟨1⟩4. $\bar{\rho}(a, c) \leq \bar{\rho}(a, b) + \bar{\rho}(b, c)$

PROOF:

$$\begin{aligned}\bar{\rho}(a, c) &= \sup_{j \in J} \bar{d}(a_j, c_j) \\ &\leq \sup_{j \in J} (\bar{d}(a_j, b_j) + \bar{d}(b_j, c_j)) \\ &\leq \sup_{j \in J} \bar{d}(a_j, b_j) + \sup_{j \in J} \bar{d}(b_j, c_j) \\ &= \bar{\rho}(a, b) + \bar{\rho}(b, c)\end{aligned}$$

□

Proposition 399. *The uniform topology on \mathbb{R}^J is finer than the product topology.*

PROOF:

⟨1⟩1. LET: $j \in J$ and U be open in \mathbb{R}

PROVE: $\pi_j^{-1}(U)$ is open in the uniform topology.

⟨1⟩2. LET: $a \in \pi_j^{-1}(U)$

⟨1⟩3. PICK $\epsilon > 0$ such that $(a_j - \epsilon, a_j + \epsilon) \subseteq U$

⟨1⟩4. $B_{\bar{\rho}}(a, \epsilon) \subseteq \pi_j^{-1}(U)$

□

Proposition 400. *The uniform topology on \mathbb{R}^J is coarser than the box topology.*

PROOF:

⟨1⟩1. LET: $a \in \mathbb{R}^J$ and $\epsilon > 0$

PROVE: $B(a, \epsilon)$ is open in the box topology.

⟨1⟩2. LET: $b \in B(a, \epsilon)$

⟨1⟩3. For $j \in J$ we have $|a_j - b_j| < \epsilon$

⟨1⟩4. For $j \in J$,

LET: $\delta_j = (\epsilon - |a_j - b_j|)/2$

⟨1⟩5. $\prod_{j \in J} (b_j - \delta_j, b_j + \delta_j) \subseteq B(a, \epsilon)$

□

Proposition 401. *The uniform topology on \mathbb{R}^J is strictly finer than the product topology if and only if J is infinite.*

PROOF:

⟨1⟩1. If J is finite then the uniform and product topologies coincide.

PROOF: The uniform, box and product topologies are all the same.

⟨1⟩2. If J is infinite then the uniform and product topologies are different.

PROOF: The set $B(\vec{0}, 1)$ is open in the uniform topology but not the product topology.

□

Proposition 402 (DC). *The uniform topology on \mathbb{R}^J is strictly coarser than the box topology if and only if J is infinite.*

PROOF:

⟨1⟩1. If J is finite then the uniform and box topologies coincide.

PROOF: The uniform, box and product topologies are all the same.

⟨1⟩2. If J is infinite then the uniform and box topologies are different.

PROOF: Pick an ω -sequence (j_1, j_2, \dots) in J . Let $U = \prod_{j \in J} U_j$ where $U_{j_i} = (-1/i, 1/i)$ and $U_j = (-1, 1)$ for all other j . Then $\vec{0} \in U$ but there is no $\epsilon > 0$ such that $B(\vec{0}, \epsilon) \subseteq U$.

□

Proposition 403. *The closure of \mathbb{R}^∞ in \mathbb{R}^ω under the uniform topology is \mathbb{R}^ω .*

PROOF: Given any open ball $B(a, \epsilon)$, pick an integer N such that $1/\epsilon < N$. Then $B(a, \epsilon)$ includes sequences whose n th entry is 0 for all $n \geq N$. □

Example 404. The space \mathbb{R}^ω is disconnected under the uniform topology. The set of bounded sequences and the set of unbounded sequences form a separation.

Proposition 405. *Give \mathbb{R}^ω the uniform topology. Let $x, y \in \mathbb{R}^\omega$. Then x and y are in the same component if and only if $x - y$ is bounded.*

PROOF:

⟨1⟩1. The component containing 0 is the set of bounded sequences.

⟨2⟩1. LET: B be the set of bounded sequences.

⟨2⟩2. B is path-connected.

⟨3⟩1. LET: $x, y \in B$

⟨3⟩2. PICK $b > 0$ such that $|x_j|, |y_j| \leq b$ for all j

⟨3⟩3. LET: $p : [0, 1] \rightarrow B$ be the function $p(t) = (1 - t)x + ty$

PROVE: p is continuous.

⟨3⟩4. LET: $t \in [0, 1]$ and $\epsilon > 0$

⟨3⟩5. LET: $\delta = \epsilon/2b$

⟨3⟩6. LET: $s \in [0, 1]$ with $|s - t| < \delta$

⟨3⟩7. $\bar{p}(p(s), p(t)) < \epsilon$

PROOF:

$$\begin{aligned} \bar{p}(p(s), p(t)) &= \sup_j \bar{d}((1 - s)x_j + sy_j, (1 - t)x_j + ty_j) \\ &\leq |(s - t)x_j + (t - s)y_j| \\ &\leq |s - t||x_j - y_j| \\ &< 2b\delta \\ &= \epsilon \end{aligned}$$

⟨2⟩3. B is connected.

PROOF: Proposition 247.

⟨2⟩4. If C is connected and $B \subseteq C$ then $B = C$.

PROOF: Otherwise $B \cap C$ and $C \setminus B$ form a separation of C .

⟨1⟩2. Q.E.D.

PROOF: Since $\lambda x. x - y$ is a Homeomorphism of \mathbb{R}^ω with itself.

□

51 Uniform Convergence

Definition 406 (Uniform Convergence). Let X be a set and Y a metric space. Let $(f_n : X \rightarrow Y)$ be a sequence of functions and $f : X \rightarrow Y$ be a function. Then f_n converges uniformly to f as $n \rightarrow \infty$ if and only if, for all $\epsilon > 0$, there exists N such that, for all $n \geq N$ and $x \in X$, we have $d(f_n(x), f(x)) < \epsilon$.

Example 407. Define $f_n : [0, 1] \rightarrow \mathbb{R}$ by $f_n(x) = x^n$ for $n \geq 1$, and $f : [0, 1] \rightarrow \mathbb{R}$ by $f(x) = 0$ if $x < 1$, $f(1) = 1$. Then f_n converges to f pointwise but not uniformly.

Theorem 408 (Uniform Limit Theorem). Let X be a topological space and Y a metric space. Let $(f_n : X \rightarrow Y)$ be a sequence of continuous functions and $f : X \rightarrow Y$ be a function. If f_n converges uniformly to f as $n \rightarrow \infty$, then f is continuous.

PROOF:

$\langle 1 \rangle 1$. LET: $x \in X$ and $\epsilon > 0$

$\langle 1 \rangle 2$. PICK N such that, for all $n \geq N$ and $y \in X$, we have $d(f_n(y), f(y)) < \epsilon/3$

$\langle 1 \rangle 3$. PICK a neighbourhood U of x such that $f_N(U) \subseteq B(f_N(x), \epsilon/3)$

PROVE: $f(U) \subseteq B(f(x), \epsilon)$

$\langle 1 \rangle 4$. LET: $y \in U$

$\langle 1 \rangle 5$. $d(f(y), f(x)) < \epsilon$

PROOF:

$$\begin{aligned} d(f(y), f(x)) &\leq d(f(y), f_N(y)) + d(f_N(y), f_N(x)) + d(f_N(x), f(x)) \quad (\text{Triangle Inequality}) \\ &< \epsilon/3 + \epsilon/3 + \epsilon/3 \quad (\langle 1 \rangle 2, \langle 1 \rangle 3) \\ &= \epsilon \end{aligned}$$

□

Proposition 409. Let X be a topological space and Y a metric space. Let $(f_n : X \rightarrow Y)$ be a sequence of continuous functions and $f : X \rightarrow Y$ be a function. Let (a_n) be a sequence of points in X and $a \in X$. If f_n converges uniformly to f and a_n converges to a in X then $f_n(a_n)$ converges to $f(a)$ uniformly in Y .

PROOF:

$\langle 1 \rangle 1$. LET: $\epsilon > 0$

$\langle 1 \rangle 2$. PICK N_1 such that, for all $n \geq N_1$ and $x \in X$, we have $d(f_n(x), f(x)) < \epsilon/2$

$\langle 1 \rangle 3$. PICK N_2 such that, for all $n \geq N_2$, we have $a_n \in f^{-1}(B(a, \epsilon/2))$

PROOF: Using the fact that f is continuous from the Uniform Limit Theorem.

$\langle 1 \rangle 4$. LET: $N = \max(N_1, N_2)$

$\langle 1 \rangle 5$. LET: $n \geq N$

$\langle 1 \rangle 6$. $d(f_n(a_n), f(a)) < \epsilon$

PROOF:

$$\begin{aligned} d(f_n(a_n), f(a)) &\leq d(f_n(a_n), f(a_n)) + d(f(a_n), f(a)) \quad (\text{Triangle Inequality}) \\ &< \epsilon/2 + \epsilon/2 \quad (\langle 1 \rangle 2, \langle 1 \rangle 3) \\ &= \epsilon \end{aligned}$$

□

Proposition 410. *Let X be a set. Let $(f_n : X \rightarrow \mathbb{R})$ be a sequence of functions and $f : X \rightarrow \mathbb{R}$ be a function. Then f_n converges uniformly to f as $n \rightarrow \infty$ if and only if $f_n \rightarrow f$ as $n \rightarrow \infty$ in \mathbb{R}^X under the uniform topology.*

PROOF:

- ⟨1⟩1. If f_n converges uniformly to f then f_n converges to f under the uniform topology.
- ⟨2⟩1. ASSUME: f_n converges uniformly to f
- ⟨2⟩2. LET: $\epsilon > 0$
- ⟨2⟩3. PICK N such that, for all $n \geq N$ and $x \in X$, we have $d(f_n(x), f(x)) < \epsilon/2$
- ⟨2⟩4. For all $n \geq N$ we have $\bar{\rho}(f_n, f) \leq \epsilon/2$
- ⟨2⟩5. For all $n \geq N$ we have $\bar{\rho}(f_n, f) < \epsilon$
- ⟨1⟩2. If f_n converges to f under the uniform topology then f_n converges uniformly to f .
- ⟨2⟩1. ASSUME: f_n converges to f under the uniform topology.
- ⟨2⟩2. LET: $\epsilon > 0$
- ⟨2⟩3. PICK N such that, for all $n \geq N$, we have $\bar{\rho}(f_n, f) < \min(\epsilon, 1/2)$
- ⟨2⟩4. LET: $n \geq N$
- ⟨2⟩5. LET: $x \in X$
- ⟨2⟩6. $\bar{\rho}(f_n, f) < \min(\epsilon, 1/2)$
- PROOF: From ⟨2⟩3.
- ⟨2⟩7. $d(f_n(x), f(x)) < \min(\epsilon, 1/2)$
- ⟨2⟩8. $d(f_n(x), f(x)) < \epsilon$

□

52 Isometric Imbeddings

Definition 411. Let X and Y be metric spaces. An *isometric imbedding* $f : X \rightarrow Y$ is a function such that, for all $x, y \in X$, we have $d(f(x), f(y)) = d(x, y)$.

Proposition 412. *Every isometric imbedding is an imbedding.*

PROOF:

- ⟨1⟩1. LET: $f : X \rightarrow Y$ be an isometric imbedding.
- ⟨1⟩2. f is injective.
- PROOF: If $f(x) = f(y)$ then $d(f(x), f(y)) = 0$ hence $d(x, y) = 0$ hence $x = y$.
- ⟨1⟩3. f is continuous.
- PROOF: For all $\epsilon > 0$, if $d(x, y) < \epsilon$ then $d(f(x), f(y)) < \epsilon$.
- ⟨1⟩4. $f : X \rightarrow f(X)$ is an open map.
- PROOF: $f(B(a, \epsilon)) = B(f(a), \epsilon) \cap f(X)$.

□