

# Topology

Robin Adams

May 26, 2022

## Contents

<b>1</b>	<b>Set Theory</b>	<b>4</b>
<b>2</b>	<b>Order Theory</b>	<b>5</b>
<b>3</b>	<b>Real Analysis</b>	<b>9</b>
<b>4</b>	<b>Group Theory</b>	<b>10</b>
<b>5</b>	<b>Topological Spaces</b>	<b>11</b>
<b>6</b>	<b>Closed Set</b>	<b>12</b>
<b>7</b>	<b>Interior</b>	<b>13</b>
<b>8</b>	<b>Closure</b>	<b>14</b>
<b>9</b>	<b>Boundary</b>	<b>15</b>
<b>10</b>	<b>Limit Points</b>	<b>16</b>
<b>11</b>	<b>Basis for a Topology</b>	<b>16</b>
<b>12</b>	<b>Local Basis at a Point</b>	<b>21</b>
<b>13</b>	<b>Convergence</b>	<b>21</b>
<b>14</b>	<b>Locally Finite Sets</b>	<b>23</b>
<b>15</b>	<b>Open Maps</b>	<b>24</b>
<b>16</b>	<b>Continuous Functions</b>	<b>24</b>
<b>17</b>	<b>Homeomorphisms</b>	<b>29</b>
<b>18</b>	<b>The Order Topology</b>	<b>30</b>

<b>19 The <math>n</math>th Root Function</b>	<b>33</b>
<b>20 The Product Topology</b>	<b>33</b>
20.1 Continuous in Each Variable Separately . . . . .	36
<b>21 The Subspace Topology</b>	<b>36</b>
<b>22 The Box Topology</b>	<b>40</b>
<b>23 <math>T_1</math> Spaces</b>	<b>42</b>
<b>24 Hausdorff Spaces</b>	<b>43</b>
<b>25 The First Countability Axiom</b>	<b>46</b>
<b>26 Strong Continuity</b>	<b>47</b>
<b>27 Saturated Sets</b>	<b>48</b>
<b>28 Quotient Maps</b>	<b>49</b>
<b>29 Quotient Topology</b>	<b>52</b>
<b>30 Retractions</b>	<b>54</b>
<b>31 Homogeneous Spaces</b>	<b>54</b>
<b>32 Regular Spaces</b>	<b>54</b>
<b>33 Connected Spaces</b>	<b>54</b>
<b>34 Totally Disconnected Spaces</b>	<b>63</b>
<b>35 Paths and Path Connectedness</b>	<b>63</b>
<b>36 The Topologist's Sine Curve</b>	<b>66</b>
<b>37 The Long Line</b>	<b>67</b>
<b>38 Components</b>	<b>68</b>
<b>39 Path Components</b>	<b>70</b>
<b>40 Local Connectedness</b>	<b>71</b>
<b>41 Local Path Connectedness</b>	<b>73</b>
<b>42 Weak Local Connectedness</b>	<b>74</b>

43 Quasicomponents	75
44 Open Coverings	76
45 Compact Spaces	76
46 Perfect Maps	82
47 Topological Groups	82
48 The Metric Topology	90
49 Real Linear Algebra	96
50 The Uniform Topology	103
51 Uniform Convergence	106
52 Isometric Imbeddings	107

## 1 Set Theory

**Definition 1** (Cover). Let  $X$  be a set and  $\mathcal{A} \subseteq \mathcal{P}X$ . Then  $\mathcal{A}$  *covers*  $X$ , or is a *covering* of  $X$ , if and only if  $\bigcup \mathcal{A} = X$ .

**Definition 2** (Finite Intersection Property). Let  $X$  be a set and  $\mathcal{A} \subseteq \mathcal{P}X$ . Then  $\mathcal{A}$  satisfies the *finite intersection property* if and only if every nonempty finite subset of  $\mathcal{A}$  has nonempty intersection.

**Definition 3** (Graph). Let  $f : A \rightarrow B$ . The *graph* of  $f$  is the set  $\{(x, f(x)) \mid x \in A\} \subseteq A \times B$ .

## 2 Order Theory

**Definition 4** (Preorder). Let  $X$  be a set. A *preorder* on  $X$  is a binary relation  $\leq$  on  $X$  such that:

**Reflexivity** For all  $x \in X$ , we have  $x \leq x$

**Transitivity** For all  $x, y, z \in X$ , if  $x \leq y$  and  $y \leq z$  then  $x \leq z$ .

**Definition 5** (Preordered Set). A *preordered set* consists of a set  $X$  and a preorder  $\leq$  on  $X$ .

**Proposition 6.** Let  $X$  and  $Y$  be linearly ordered sets. Let  $f : X \rightarrow Y$  be strictly monotone and surjective. Then  $f$  is a poset isomorphism.

PROOF:

$\langle 1 \rangle 1.$   $f$  is injective.

$\langle 2 \rangle 1.$  LET:  $x, y \in X$

$\langle 2 \rangle 2.$  ASSUME:  $f(x) = f(y)$

$\langle 2 \rangle 3.$   $x \not\leq y$

PROOF: By strong monotonicity.

$\langle 2 \rangle 4.$   $y \not\leq x$

PROOF: By strong monotonicity.

$\langle 2 \rangle 5.$   $x = y$

PROOF: By trichotomy.

$\langle 1 \rangle 2.$   $f^{-1}$  is monotone.

$\langle 2 \rangle 1.$  LET:  $x, y \in X$

$\langle 2 \rangle 2.$  ASSUME:  $x \leq y$

$\langle 2 \rangle 3.$   $f^{-1}(x) \not\leq f^{-1}(y)$

PROOF: By strong monotonicity.

$\langle 2 \rangle 4.$   $f^{-1}(x) < f^{-1}(y)$

PROOF: By trichotomy.

□

**Definition 7** (Interval). Let  $X$  be a preordered set and  $Y \subseteq X$ . Then  $Y$  is an *interval* if and only if, for all  $a, b \in Y$  and  $c \in X$ , if  $a \leq c \leq b$  then  $c \in Y$ .

**Definition 8** (Linear Continuum). A linearly ordered set  $L$  is a *linear continuum* if and only if:

1. every nonempty subset of  $L$  that is bounded above has a supremum
2.  $L$  is dense

**Proposition 9.** Every interval in a linear continuum is a linear continuum.

PROOF:

$\langle 1 \rangle 1.$  LET:  $L$  be a linear continuum and  $I$  an interval in  $L$ .

$\langle 1 \rangle 2.$  Every nonempty subset of  $I$  that is bounded above has a supremum in  $I$ .

$\langle 2 \rangle 1.$  LET:  $X \subseteq I$  be nonempty and bounded above by  $b \in I$ .

$\langle 2 \rangle 2$ . LET:  $s$  be the supremum of  $X$  in  $L$ .  
 PROOF: Since  $L$  is a linear continuum.  
 $\langle 2 \rangle 3$ .  $s \in I$   
 $\langle 3 \rangle 1$ . PICK  $a \in X$   
 PROOF: Since  $X$  is nonempty ( $\langle 2 \rangle 1$ ).  
 $\langle 3 \rangle 2$ .  $a \leq s \leq b$   
 $\langle 3 \rangle 3$ .  $s \in I$   
 PROOF: Since  $I$  is an interval ( $\langle 1 \rangle 1$ ).  
 $\langle 2 \rangle 4$ .  $s$  is the supremum of  $X$  in  $I$   
 $\langle 1 \rangle 3$ .  $I$  is dense.  
 $\langle 2 \rangle 1$ . LET:  $x, y \in I$  with  $x < y$   
 $\langle 2 \rangle 2$ . PICK  $z \in L$  with  $x < z < y$   
 PROOF: Since  $L$  is dense.  
 $\langle 2 \rangle 3$ .  $z \in I$   
 PROOF: Since  $I$  is an interval.

□

**Definition 10** (Ordered Square). The *ordered square*  $I_o^2$  is the set  $[0, 1]^2$  under the dictionary order.

**Proposition 11.** *The ordered square is a linear continuum.*

PROOF:

$\langle 1 \rangle 1$ . Every nonempty subset of  $I_o^2$  bounded above has a supremum.  
 $\langle 2 \rangle 1$ . LET:  $X \subseteq I_o^2$  be nonempty and bounded above by  $(b, c)$   
 $\langle 2 \rangle 2$ . LET:  $s = \sup \pi_1(X)$   
 PROOF: The set  $\pi_1(X)$  is nonempty and bounded above by  $b$ .  
 $\langle 2 \rangle 3$ . CASE:  $s \in \pi_1(X)$   
 $\langle 3 \rangle 1$ . LET:  $t = \sup \{y \in [0, 1] \mid (s, y) \in X\}$   
 PROOF: This set is nonempty and bounded above by  $c$ .  
 $\langle 3 \rangle 2$ .  $(s, t)$  is the supremum of  $X$ .  
 $\langle 2 \rangle 4$ . CASE:  $s \notin \pi_1(X)$   
 PROOF: In this case  $(s, 0)$  is the supremum of  $X$ .  
 $\langle 1 \rangle 2$ .  $I_o^2$  is dense.  
 $\langle 2 \rangle 1$ . LET:  $(x_1, y_1), (x_2, y_2) \in I_o^2$  with  $(x_1, y_1) < (x_2, y_2)$   
 $\langle 2 \rangle 2$ . CASE:  $x_1 < x_2$   
 $\langle 3 \rangle 1$ . PICK  $x_3$  with  $x_1 < x_3 < x_2$   
 $\langle 3 \rangle 2$ .  $(x_1, y_1) < (x_3, y_1) < (x_2, y_2)$   
 $\langle 2 \rangle 3$ . CASE:  $x_1 = x_2$  and  $y_1 < y_2$   
 $\langle 3 \rangle 1$ . PICK  $y_3$  with  $y_1 < y_3 < y_2$   
 $\langle 3 \rangle 2$ .  $(x_1, y_1) < (x_1, y_3) < (x_2, y_2)$

□

**Proposition 12.** *If  $X$  is a well-ordered set then  $X \times [0, 1)$  under the dictionary order is a linear continuum.*

PROOF:

$\langle 1 \rangle 1$ . Every nonempty set  $A \subseteq X \times [0, 1)$  bounded above has a supremum.

- ⟨2⟩1. LET:  $A \subseteq X \times [0, 1)$  be nonempty and bounded above
- ⟨2⟩2. LET:  $x_0$  be the supremum of  $\pi_1(A)$
- ⟨2⟩3. CASE:  $x_0 \in \pi_1(A)$ 
  - ⟨3⟩1. LET:  $y_0$  be the supremum of  $\{y \in [0, 1) \mid (x_0, y) \in A\}$
  - ⟨3⟩2.  $(x_0, y_0)$  is the supremum of  $A$ .
- ⟨2⟩4. CASE:  $x_0 \notin \pi_1(A)$ 
  - PROOF: In this case  $(x_0, 0)$  is the supremum of  $A$ .
- ⟨1⟩2.  $X \times [0, 1)$  is dense.
  - ⟨2⟩1. LET:  $(x_1, y_1), (x_2, y_2) \in X \times [0, 1)$  with  $(x_1, y_1) < (x_2, y_2)$
  - ⟨2⟩2. CASE:  $x_1 < x_2$ 
    - ⟨3⟩1. PICK  $y_3$  such that  $y_1 < y_3 < 1$
    - ⟨3⟩2.  $(x_1, y_1) < (x_1, y_3) < (x_2, y_2)$
  - ⟨2⟩3. CASE:  $x_1 = x_2$  and  $y_1 < y_2$ 
    - ⟨3⟩1. PICK  $y_3$  such that  $y_1 < y_3 < y_2$
    - ⟨3⟩2.  $(x_1, y_1) < (x_1, y_3) < (x_2, y_2)$

□

**Lemma 13.** For all  $a, b, c, d \in \mathbb{R}$  with  $a < b$  and  $c < d$ , we have  $[a, b) \cong [c, d)$

PROOF: The map  $\lambda t.c + (t - a)(d - c)/(b - a)$  is an order isomorphism.

**Proposition 14.** Let  $X$  be a linearly ordered set. Let  $a < b < c$  in  $X$ . Then  $[a, c) \cong [0, 1)$  if and only if  $[a, b) \cong [b, c) \cong [0, 1)$ .

PROOF:

- ⟨1⟩1. If  $[a, c) \cong [0, 1)$  then  $[a, b) \cong [b, c) \cong [0, 1)$
- ⟨2⟩1. ASSUME:  $f : [a, c) \cong [0, 1)$  is an order isomorphism
- ⟨2⟩2.  $[a, b) \cong [0, 1)$ 
  - PROOF:
$$\begin{aligned} [a, b) &\cong [0, f(b)) && \text{(by the restriction of } f) \\ &\cong [0, 1) && \text{(Lemma 13)} \end{aligned}$$
- ⟨2⟩3.  $[b, c) \cong [0, 1)$ 
  - PROOF: Similar.
- ⟨1⟩2. If  $[a, b) \cong [b, c) \cong [0, 1)$  then  $[a, c) \cong [0, 1)$ 
  - PROOF:
$$\begin{aligned} [a, c) &= [a, b) * [b, c) \\ &\cong [0, 1) * [0, 1) \\ &\cong [0, 1/2) * [1/2, 1) && \text{(Lemma 13)} \\ &= 1 \end{aligned}$$

□

**Proposition 15 (CC).** Let  $X$  be a linearly ordered set. Let  $x_0 < x_1 < \dots$  be a strictly increasing sequence in  $X$  with supremum  $b$ . Then  $[x_0, b) \cong [0, 1)$  if and only if  $[x_i, x_{i+1}) \cong [0, 1)$  for all  $i$ .

PROOF:

- ⟨1⟩1. If  $[x_0, b) \cong [0, 1)$  then  $[x_i, x_{i+1}) \cong [0, 1)$  for all  $i$ .

PROOF: By Lemma 13

$\langle 1 \rangle 2$ . If  $[x_i, x_{i+1}) \cong [0, 1)$  for all  $i$  then  $[x_0, b) \cong [0, 1)$

$\langle 2 \rangle 1$ . ASSUME:  $[x_i, x_{i+1}) \cong [0, 1)$  for all  $i$

$\langle 2 \rangle 2$ . PICK an order isomorphism  $f_i : [x_i, x_{i+1}) \cong [1/2^i, 2/2^{i+1})$  for each  $i$ .

PROOF: By Lemma 13

$\langle 2 \rangle 3$ . The union of the  $f_i$ s is an order isomorphism  $[x_0, b) \cong [0, 1)$

□



### 3 Real Analysis

**Definition 16.** Let  $\mathbb{R}^\infty$  be the subset of  $\mathbb{R}^\omega$  consisting of all sequences  $(a_n)$  such that  $a_n = 0$  for all but finitely many  $n$ .

## 4 Group Theory

**Definition 17.** Given a group  $G$  and sets  $A, B \subseteq G$ , let  $AB = \{ab \mid a \in A, b \in B\}$ .

**Definition 18.** Given a group  $G$  and a set  $A \subseteq G$ , let  $A^{-1} = \{a^{-1} \mid a \in A\}$ .

## 5 Topological Spaces

**Definition 19** (Topology). A *topology* on a set  $X$  is a set  $\mathcal{T} \subseteq \mathcal{P}X$  such that:

- $X \in \mathcal{T}$ .
- For all  $\mathcal{U} \subseteq \mathcal{T}$  we have  $\bigcup \mathcal{U} \in \mathcal{T}$ .
- For all  $U, V \in \mathcal{T}$  we have  $U \cap V \in \mathcal{T}$ .

We call the elements of  $X$  *points* and the elements of  $\mathcal{T}$  *open sets*.

**Definition 20** (Topological Space). A *topological space*  $X$  consists of a set  $X$  and a topology on  $X$ .

**Definition 21** (Discrete Space). For any set  $X$ , the *discrete* topology on  $X$  is  $\mathcal{P}X$ .

**Definition 22** (Indiscrete Space). For any set  $X$ , the *indiscrete* or *trivial* topology on  $X$  is  $\{\emptyset, X\}$ .

**Definition 23** (Finite Complement Topology). For any set  $X$ , the *finite complement topology* on  $X$  is  $\{U \in \mathcal{P}X \mid X \setminus U \text{ is finite}\} \cup \{\emptyset\}$ .

**Definition 24** (Countable Complement Topology). For any set  $X$ , the *countable complement topology* on  $X$  is  $\{U \in \mathcal{P}X \mid X \setminus U \text{ is countable}\} \cup \{\emptyset\}$ .

**Definition 25** (Finer, Coarser). Suppose that  $\mathcal{T}$  and  $\mathcal{T}'$  are two topologies on a given set  $X$ . If  $\mathcal{T}' \supseteq \mathcal{T}$ , we say that  $\mathcal{T}'$  is *finer* than  $\mathcal{T}$ ; if  $\mathcal{T}'$  *properly* contains  $\mathcal{T}$ , we say that  $\mathcal{T}'$  is *strictly finer* than  $\mathcal{T}$ . We also say that  $\mathcal{T}$  is *coarser* than  $\mathcal{T}'$ , or *strictly coarser*, in these two respective situations. We say  $\mathcal{T}$  is *comparable* with  $\mathcal{T}'$  if either  $\mathcal{T}' \supseteq \mathcal{T}$  or  $\mathcal{T} \supseteq \mathcal{T}'$ .

**Lemma 26.** Let  $X$  be a topological space and  $U \subseteq X$ . Then  $U$  is open if and only if, for all  $x \in U$ , there exists an open set  $V$  such that  $x \in V \subseteq U$ .

PROOF:

$\langle 1 \rangle 1. \Rightarrow$

PROOF: Take  $V = U$

$\langle 1 \rangle 2. \Leftarrow$

PROOF: We have  $U = \bigcup \{V \text{ open in } X \mid V \subseteq U\}$ .

□

**Lemma 27.** Let  $X$  be a set and  $\mathcal{T}$  a nonempty set of topologies on  $X$ . Then  $\bigcap \mathcal{T}$  is a topology on  $X$ , and is the finest topology that is coarser than every member of  $\mathcal{T}$ .

PROOF:

$\langle 1 \rangle 1. X \in \bigcap \mathcal{T}$

PROOF: Since  $X$  is in every member of  $\mathcal{T}$ .

$\langle 1 \rangle 2. \bigcap \mathcal{T}$  is closed under union.

- ⟨2⟩1. LET:  $\mathcal{U} \subseteq \bigcap \mathcal{T}$
- ⟨2⟩2. For all  $T \in \mathcal{T}$  we have  $\mathcal{U} \subseteq T$
- ⟨2⟩3. For all  $T \in \mathcal{T}$  we have  $\bigcup \mathcal{U} \in T$
- ⟨2⟩4.  $\bigcup \mathcal{U} \in \bigcap \mathcal{T}$
- ⟨1⟩3.  $\bigcap \mathcal{T}$  is closed under binary intersection.
- ⟨2⟩1. LET:  $U, V \in \bigcap \mathcal{T}$
- ⟨2⟩2. For all  $T \in \mathcal{T}$  we have  $U, V \in T$
- ⟨2⟩3. For all  $T \in \mathcal{T}$  we have  $U \cap V \in T$
- ⟨2⟩4.  $U \cap V \in \bigcap \mathcal{T}$

□

**Lemma 28.** *Let  $X$  be a set and  $\mathcal{T}$  a set of topologies on  $X$ . Then there exists a unique coarsest topology that is finer than every member of  $\mathcal{T}$ .*

PROOF: The required topology is given by

$$\bigcap \{T \in \mathcal{P}X \mid T \text{ is a topology on } X \text{ that is finer than every member of } \mathcal{T}\},$$

The set is nonempty since it contains the discrete topology. □

**Definition 29** (Neighbourhood). A *neighbourhood* of a point  $x$  is an open set that contains  $x$ .

## 6 Closed Set

**Definition 30** (Closed Set). Let  $X$  be a topological space and  $A \subseteq X$ . Then  $A$  is *closed* if and only if  $X \setminus A$  is open.

**Lemma 31.** *The empty set is closed.*

PROOF: Since the whole space  $X$  is always open. □

**Lemma 32.** *The topological space  $X$  is closed.*

PROOF: Since  $\emptyset$  is open. □

**Lemma 33.** *The intersection of a nonempty set of closed sets is closed.*

PROOF: Let  $\mathcal{C}$  be a nonempty set of closed sets. Then  $X \setminus \bigcap \mathcal{C} = \bigcup \{X \setminus C \mid C \in \mathcal{C}\}$  is open. □

**Lemma 34.** *The union of two closed sets is closed.*

PROOF: Let  $C$  and  $D$  be closed. Then  $X \setminus (C \cup D) = (X \setminus C) \cap (X \setminus D)$  is open. □

**Proposition 35.** *Let  $X$  be a set and  $\mathcal{C} \subseteq \mathcal{P}X$  a set such that:*

1.  $\emptyset \in \mathcal{C}$
2.  $X \in \mathcal{C}$
3. For all  $\mathcal{A} \subseteq \mathcal{C}$  nonempty we have  $\bigcap \mathcal{A} \in \mathcal{C}$

4. For all  $C, D \in \mathcal{C}$  we have  $C \cup D \in \mathcal{C}$ .

Then there exists a unique topology  $\mathcal{T}$  such that  $\mathcal{C}$  is the set of closed sets, namely

$$\mathcal{T} = \{X \setminus C \mid C \in \mathcal{C}\}$$

PROOF:

$\langle 1 \rangle 1$ . LET:  $\mathcal{T} = \{X \setminus C \mid C \in \mathcal{C}\}$

$\langle 1 \rangle 2$ .  $\mathcal{T}$  is a topology

$\langle 2 \rangle 1$ .  $X \in \mathcal{T}$

PROOF: Since  $\emptyset \in \mathcal{C}$

$\langle 2 \rangle 2$ . For all  $\mathcal{U} \subseteq \mathcal{T}$  we have  $\bigcup \mathcal{U} \in \mathcal{T}$

$\langle 3 \rangle 1$ . LET:  $\mathcal{U} \subseteq \mathcal{T}$

$\langle 3 \rangle 2$ . CASE:  $\mathcal{U} = \emptyset$

PROOF: In this case  $\bigcup \mathcal{U} = \emptyset \in \mathcal{T}$  since  $X \in \mathcal{C}$

$\langle 3 \rangle 3$ . CASE:  $\mathcal{U} \neq \emptyset$

PROOF: In this case  $X \setminus \bigcup \mathcal{U} = \bigcap \{X \setminus U \mid U \in \mathcal{U}\} \in \mathcal{C}$ .

$\langle 2 \rangle 3$ . For all  $U, V \in \mathcal{T}$  we have  $U \cap V \in \mathcal{T}$

PROOF: Since  $X \setminus (U \cap V) = (X \setminus U) \cup (X \setminus V) \in \mathcal{C}$ .

$\langle 1 \rangle 3$ .  $\mathcal{C}$  is the set of all closed sets in  $\mathcal{T}$

PROOF:

$C$  is closed in  $\mathcal{T}$

$$\Leftrightarrow X \setminus C \in \mathcal{T}$$

$$\Leftrightarrow C \in \mathcal{C}$$

$\langle 1 \rangle 4$ . If  $\mathcal{T}'$  is a topology and  $\mathcal{C}$  is the set of closed sets in  $\mathcal{T}'$  then  $\mathcal{T}' = \mathcal{T}$

PROOF: We have

$$U \in \mathcal{T}$$

$$\Leftrightarrow X \setminus U \in \mathcal{C}$$

$$\Leftrightarrow X \setminus U \text{ is closed in } \mathcal{T}'$$

$$\Leftrightarrow U \in \mathcal{T}'$$

□

**Proposition 36.** If  $U$  is open and  $A$  is closed then  $U \setminus A$  is open.

PROOF:  $U \setminus A = U \cap (X \setminus A)$  is the intersection of two open sets. □

**Proposition 37.** If  $U$  is open and  $A$  is closed then  $A \setminus U$  is closed.

PROOF:  $A \setminus U = A \cap (X \setminus U)$  is the intersection of two closed sets. □

## 7 Interior

**Definition 38** (Interior). Let  $X$  be a topological space and  $A \subseteq X$ . The *interior* of  $A$ ,  $\text{Int } A$ , is the union of all the open subsets of  $A$ .

**Lemma 39.** The interior of a set is open.

PROOF: It is a union of open sets.  $\square$

**Lemma 40.**

$$\text{Int } A \subseteq A$$

PROOF: Immediate from definition.  $\square$

**Lemma 41.** *If  $U$  is open and  $U \subseteq A$  then  $U \subseteq \text{Int } A$*

PROOF: Immediate from definition.  $\square$

**Lemma 42.** *A set  $A$  is open if and only if  $A = \text{Int } A$ .*

PROOF: If  $A = \text{Int } A$  then  $A$  is open by Lemma 39. Conversely if  $A$  is open then  $A \subseteq \text{Int } A$  by the definition of interior and so  $A = \text{Int } A$ .

## 8 Closure

**Definition 43** (Closure). Let  $X$  be a topological space and  $A \subseteq X$ . The *closure* of  $A$ ,  $\overline{A}$ , is the intersection of all the closed sets that include  $A$ .

This intersection exists since  $X$  is a closed set that includes  $A$  (Lemma 32).

**Lemma 44.** *The closure of a set is closed.*

PROOF: Dual to Lemma 39.  $\square$

**Lemma 45.**

$$A \subseteq \overline{A}$$

PROOF: Immediate from definition.  $\square$

**Lemma 46.** *If  $C$  is closed and  $A \subseteq C$  then  $\overline{A} \subseteq C$ .*

PROOF: Immediate from definition.  $\square$

**Lemma 47.** *A set  $A$  is closed if and only if  $A = \overline{A}$ .*

PROOF: Dual to Lemma 42.  $\square$

**Theorem 48.** *Let  $X$  be a topological space,  $A \subseteq X$  and  $x \in X$ . Then  $x \in \overline{A}$  if and only if every neighbourhood of  $x$  intersects  $A$ .*

PROOF: We have

$$\begin{aligned} x \in \overline{A} \\ \Leftrightarrow \forall C. C \text{ closed} \wedge A \subseteq C \Rightarrow x \in C \\ \Leftrightarrow \forall U. U \text{ open} \wedge A \cap U = \emptyset \Rightarrow x \notin U \\ \Leftrightarrow \forall U. U \text{ open} \wedge x \in U \Rightarrow U \text{ intersects } A \end{aligned} \quad \square$$

**Proposition 49.** *If  $A \subseteq B$  then  $\overline{A} \subseteq \overline{B}$ .*

PROOF: This holds because  $\overline{B}$  is a closed set that includes  $A$ .  $\square$

**Proposition 50.**

$$\overline{A \cup B} = \overline{A} \cup \overline{B}$$

PROOF:

$\langle 1 \rangle 1. \overline{A} \subseteq \overline{A \cup B}$

PROOF: By Proposition 49.

$\langle 1 \rangle 2. \overline{B} \subseteq \overline{A \cup B}$

PROOF: By Proposition 49.

$\langle 1 \rangle 3. \overline{A \cup B} \subseteq \overline{A \cup B}$

$\langle 2 \rangle 1. \text{ LET: } x \in \overline{A \cup B}$

$\langle 2 \rangle 2. \text{ ASSUME: } x \notin \overline{A}$

PROVE:  $x \in \overline{B}$

$\langle 2 \rangle 3. \text{ PICK a neighbourhood } U \text{ of } x \text{ that does not intersect } A$

$\langle 2 \rangle 4. \text{ LET: } V \text{ be any neighbourhood of } x$

$\langle 2 \rangle 5. U \cap V \text{ is a neighbourhood of } x$

$\langle 2 \rangle 6. U \cap V \text{ intersects } A \cup B$

PROOF: From  $\langle 2 \rangle 1$  and Theorem 48.

$\langle 2 \rangle 7. U \cap V \text{ intersects } B$

PROOF: From  $\langle 2 \rangle 3$

$\langle 2 \rangle 8. V \text{ intersects } B$

$\langle 2 \rangle 9. \text{ Q.E.D.}$

PROOF: We have  $x \in \overline{B}$  from Theorem 48.

$\square$

## 9 Boundary

**Definition 51** (Boundary). The *boundary* of a set  $A$  is the set  $\partial A = \overline{A} \cap \overline{X \setminus A}$ .

**Proposition 52.**

$$\text{Int } A \cap \partial A = \emptyset$$

PROOF: Since  $\overline{X \setminus A} = X \setminus \text{Int } A$ .  $\square$

**Proposition 53.**

$$\overline{A} = \text{Int } A \cup \partial A$$

PROOF:

$$\begin{aligned} \text{Int } A \cup \partial A &= \text{Int } A \cup (\overline{A} \cap \overline{X \setminus A}) \\ &= (\text{Int } A \cup \overline{A}) \cap (\text{Int } A \cup \overline{X \setminus A}) \\ &= \overline{A} \cap X \\ &= \overline{A} \end{aligned}$$

**Proposition 54.**  $\partial A = \emptyset$  if and only if  $A$  is open and closed.

PROOF: If  $\partial A = \emptyset$  then  $\overline{A} = \text{Int } A$  by Proposition 53.

**Proposition 55.** *A set  $U$  is open if and only if  $\partial U = \overline{U} \setminus U$ .*

PROOF:

$$\begin{aligned} \partial U &= \overline{U} \setminus U \\ \Leftrightarrow \overline{U} \setminus \text{Int } U &= \overline{U} \setminus U && (\text{Propositions 52, 53}) \\ \Leftrightarrow \text{Int } U &= U && \square \end{aligned}$$

## 10 Limit Points

**Definition 56** (Limit Point). Let  $X$  be a topological space,  $a \in X$  and  $A \subseteq X$ . Then  $a$  is a *limit point*, *cluster point* or *point of accumulation* for  $A$  if and only if every neighbourhood of  $a$  intersects  $A$  at a point other than  $a$ .

**Lemma 57.** *The point  $a$  is an accumulation point for  $A$  if and only if  $a \in \overline{A \setminus \{a\}}$ .*

PROOF: From Theorem 48.  $\square$

**Theorem 58.** *Let  $X$  be a topological space and  $A \subseteq X$ . Let  $A'$  be the set of all limit points of  $A$ . Then  $\overline{A} = A \cup A'$ .*

PROOF:

$\langle 1 \rangle 1$ . For all  $x \in \overline{A}$ , if  $x \notin A$  then  $x \in A'$

PROOF: From Theorem 48.

$\langle 1 \rangle 2$ .  $A \subseteq \overline{A}$

PROOF: Lemma 45.

$\langle 1 \rangle 3$ .  $A' \subseteq \overline{A}$

PROOF: From Theorem 48.

$\square$

**Corollary 58.1.** *A set is closed if and only if it contains all its limit points.*

**Proposition 59.** *In an indiscrete topology, every point is a limit point of any set that has more than one point.*

PROOF: Let  $X$  be an indiscrete space. Let  $A$  be a set with more than one point and  $x$  be a point. The only neighbourhood of  $x$  is  $X$ , which must intersect  $A$  at a point other than  $x$ .  $\square$

**Lemma 60.** *Let  $X$  be a topological space and  $A \subseteq B \subseteq X$ . Then every limit point of  $A$  is a limit point of  $B$ .*

PROOF: Immediate from definitions.  $\square$

## 11 Basis for a Topology

**Definition 61** (Basis). If  $X$  is a set, a *basis* for a topology on  $X$  is a set  $\mathcal{B} \subseteq \mathcal{P}X$  called *basis elements* such that



1. For all  $x \in X$ , there exists  $B \in \mathcal{B}$  such that  $x \in B$ .
2. For all  $B_1, B_2 \in \mathcal{B}$  and  $x \in B_1 \cap B_2$ , there exists  $B_3 \in \mathcal{B}$  such that  $x \in B_3 \subseteq B_1 \cap B_2$ .

If  $\mathcal{B}$  satisfies these two conditions, then we define the topology *generated* by  $\mathcal{B}$  to be  $\mathcal{T} = \{U \in \mathcal{P}X \mid \forall x \in U. \exists B \in \mathcal{B}. x \in B \subseteq U\}$ .

We prove this is a topology.

PROOF:

$\langle 1 \rangle 1.$   $X \in \mathcal{T}$

PROOF: For all  $x \in X$  there exists  $B \in \mathcal{B}$  such that  $x \in B \subseteq X$  by condition 1.

$\langle 1 \rangle 2.$  For all  $\mathcal{U} \subseteq \mathcal{T}$  we have  $\bigcup \mathcal{U} \in \mathcal{T}$

$\langle 2 \rangle 1.$  LET:  $\mathcal{U} \subseteq \mathcal{T}$

$\langle 2 \rangle 2.$  LET:  $x \in \bigcup \mathcal{U}$

$\langle 2 \rangle 3.$  PICK  $U \in \mathcal{U}$  such that  $x \in U$

$\langle 2 \rangle 4.$  PICK  $B \in \mathcal{B}$  such that  $x \in B \subseteq U$

PROOF: Since  $U \in \mathcal{T}$  by  $\langle 2 \rangle 1$  and  $\langle 2 \rangle 3$ .

$\langle 2 \rangle 5.$   $x \in B \subseteq \bigcup \mathcal{U}$

$\langle 1 \rangle 3.$  For all  $U, V \in \mathcal{T}$  we have  $U \cap V \in \mathcal{T}$

$\langle 2 \rangle 1.$  LET:  $U, V \in \mathcal{T}$

$\langle 2 \rangle 2.$  LET:  $x \in U \cap V$

$\langle 2 \rangle 3.$  PICK  $B_1 \in \mathcal{B}$  such that  $x \in B_1 \subseteq U$

$\langle 2 \rangle 4.$  PICK  $B_2 \in \mathcal{B}$  such that  $x \in B_2 \subseteq V$

$\langle 2 \rangle 5.$  PICK  $B_3 \in \mathcal{B}$  such that  $x \in B_3 \subseteq B_1 \cap B_2$

PROOF: By condition 2.

$\langle 2 \rangle 6.$   $x \in B_3 \subseteq U \cap V$

□

**Lemma 62.** *Let  $X$  be a set. Let  $\mathcal{B}$  be a basis for a topology  $\mathcal{T}$  on  $X$ . Then  $\mathcal{T}$  is the set of all unions of subsets of  $\mathcal{B}$ .*

PROOF:

$\langle 1 \rangle 1.$  For all  $U \in \mathcal{T}$ , there exists  $\mathcal{A} \subseteq \mathcal{B}$  such that  $U = \bigcup \mathcal{A}$

$\langle 2 \rangle 1.$  LET:  $U \in \mathcal{T}$

$\langle 2 \rangle 2.$  LET:  $\mathcal{A} = \{B \in \mathcal{B} \mid B \subseteq U\}$

$\langle 2 \rangle 3.$   $U \subseteq \bigcup \mathcal{A}$

$\langle 3 \rangle 1.$  LET:  $x \in U$

$\langle 3 \rangle 2.$  PICK  $B \in \mathcal{B}$  such that  $x \in B \subseteq U$

PROOF: Since  $\mathcal{B}$  is a basis for  $\mathcal{T}$ .

$\langle 3 \rangle 3.$   $x \in B \in \mathcal{A}$

$\langle 2 \rangle 4.$   $\bigcup \mathcal{A} \subseteq U$

PROOF: From the definition of  $\mathcal{A}$  ( $\langle 2 \rangle 2$ ).

$\langle 1 \rangle 2.$  For all  $\mathcal{A} \subseteq \mathcal{B}$  we have  $\bigcup \mathcal{A} \in \mathcal{T}$

$\langle 2 \rangle 1.$   $\bigcup \mathcal{A} \in \mathcal{T}$

PROOF: If  $B \in \mathcal{B}$  and  $x \in B$ , then there exists  $B' \in \mathcal{B}$  such that  $x \in B' \subseteq B$ , namely  $B' = B$ .

⟨2⟩2. Q.E.D.

PROOF: Since  $\mathcal{T}$  is closed under union.

□

**Corollary 62.1.** *Let  $X$  be a set. Let  $\mathcal{B}$  be a basis for a topology  $\mathcal{T}$  on  $X$ . Then  $\mathcal{T}$  is the coarsest topology that includes  $\mathcal{B}$ .*

PROOF: Since every topology that includes  $\mathcal{B}$  includes all unions of subsets of  $\mathcal{B}$ . □

**Lemma 63.** *Let  $X$  be a topological space. Suppose that  $\mathcal{C}$  is a set of open sets such that, for every open set  $U$  and every point  $x \in U$ , there exists  $C \in \mathcal{C}$  such that  $x \in C \subseteq U$ . Then  $\mathcal{C}$  is a basis for the topology on  $X$ .*

PROOF:

⟨1⟩1. For all  $x \in X$ , there exists  $C \in \mathcal{C}$  such that  $x \in C$

PROOF: Immediate from hypothesis.

⟨1⟩2. For all  $C_1, C_2 \in \mathcal{C}$  and  $x \in C_1 \cap C_2$ , there exists  $C_3 \in \mathcal{C}$  such that  $x \in C_3 \subseteq C_1 \cap C_2$

PROOF: Since  $C_1 \cap C_2$  is open.

⟨1⟩3. Every open set is open in the topology generated by  $\mathcal{C}$

PROOF: Immediate from hypothesis.

⟨1⟩4. Every union of a subset of  $\mathcal{C}$  is open.

PROOF: Since every member of  $\mathcal{C}$  is open.

□

**Lemma 64.** *Let  $\mathcal{B}$  and  $\mathcal{B}'$  be bases for the topologies  $\mathcal{T}$  and  $\mathcal{T}'$  respectively on the set  $X$ . Then the following are equivalent.*

1.  $\mathcal{T} \subseteq \mathcal{T}'$

2. For all  $B \in \mathcal{B}$  and  $x \in B$ , there exists  $B' \in \mathcal{B}'$  such that  $x \in B' \subseteq B$ .

PROOF:

⟨1⟩1.  $1 \Rightarrow 2$

⟨2⟩1. ASSUME:  $\mathcal{T} \subseteq \mathcal{T}'$

⟨2⟩2. LET:  $B \in \mathcal{B}$  and  $x \in B$

⟨2⟩3.  $B \in \mathcal{T}$

PROOF: Corollary 62.1.

⟨2⟩4.  $B \in \mathcal{T}'$

PROOF: By ⟨2⟩1

⟨2⟩5. There exists  $B' \in \mathcal{B}'$  such that  $x \in B' \subseteq B$

PROOF: Since  $\mathcal{B}'$  is a basis for  $\mathcal{T}'$ .

⟨1⟩2.  $2 \Rightarrow 1$

⟨2⟩1. ASSUME: 2

⟨2⟩2. LET:  $U \in \mathcal{T}$

PROVE:  $U \in \mathcal{T}'$

⟨2⟩3. LET:  $x \in U$

PROVE: There exists  $B' \in \mathcal{B}'$  such that  $x \in B' \subseteq U$

- $\langle 2 \rangle 4$ . PICK  $B \in \mathcal{B}$  such that  $x \in B \subseteq U$   
 PROOF: Since  $\mathcal{B}$  is a basis for  $\mathcal{T}$ .  
 $\langle 2 \rangle 5$ . PICK  $B' \in \mathcal{B}'$  such that  $x \in B' \subseteq B$   
 PROOF: By  $\langle 2 \rangle 1$ .  
 $\langle 2 \rangle 6$ .  $x \in B' \subseteq U$

□

**Theorem 65.** *Let  $X$  be a topological space,  $A \subseteq X$  and  $x \in X$ . Let  $\mathcal{B}$  be a basis for  $X$ . Then  $x \in \overline{A}$  if and only if, for all  $B \in \mathcal{B}$ , if  $x \in B$  then  $B$  intersects  $A$ .*

PROOF:

- $\langle 1 \rangle 1$ . If  $x \in \overline{A}$  then, for all  $B \in \mathcal{B}$ , if  $x \in B$  then  $B$  intersects  $A$ .  
 PROOF: This follows from Theorem 48 since every element of  $\mathcal{B}$  is open (Corollary 62.1).  
 $\langle 1 \rangle 2$ . Suppose that, for all  $B \in \mathcal{B}$ , if  $x \in B$  then  $B$  intersects  $A$ . Then  $x \in \overline{A}$ .  
 $\langle 2 \rangle 1$ . ASSUME: For all  $B \in \mathcal{B}$ , if  $x \in B$  then  $B$  intersects  $A$ .  
 $\langle 2 \rangle 2$ . LET:  $U$  be an open set that contains  $x$   
 PROVE:  $U$  intersects  $A$ .  
 $\langle 2 \rangle 3$ . PICK  $B \in \mathcal{B}$  such that  $x \in B \subseteq U$ .  
 $\langle 2 \rangle 4$ .  $B$  intersects  $A$ .  
 PROOF: From  $\langle 2 \rangle 1$ .  
 $\langle 2 \rangle 5$ .  $U$  intersects  $A$ .  
 $\langle 2 \rangle 6$ . Q.E.D.  
 PROOF: By Theorem 48.

□

**Definition 66** (Lower Limit Topology on the Real Line). The *lower limit topology on the real line* is the topology on  $\mathbb{R}$  generated by the basis consisting of all half-open intervals of the form  $[a, b)$ .

We write  $\mathbb{R}_l$  for the topological space  $\mathbb{R}$  under the lower limit topology.

We prove this is a basis for a topology.

PROOF:

- $\langle 1 \rangle 1$ . For all  $x \in \mathbb{R}$  there exists an interval  $[a, b)$  such that  $x \in [a, b)$ .  
 PROOF: Take  $[a, b) = [x, x + 1)$ .  
 $\langle 1 \rangle 2$ . For any open intervals  $[a, b)$ ,  $[c, d)$  if  $x \in [a, b) \cap [c, d)$ , then there exists an interval  $[e, f)$  such that  $x \in [e, f) \subseteq [a, b) \cap [c, d)$   
 PROOF: Take  $[e, f) = [\max(a, c), \min(b, d))$ .

□

**Definition 67** ( $K$ -topology on the Real Line). Let  $K = \{1/n \mid n \in \mathbb{Z}^+\}$ .

The  *$K$ -topology on the real line* is the topology on  $\mathbb{R}$  generated by the basis consisting of all open intervals  $(a, b)$  and all sets of the form  $(a, b) \setminus K$ .

We write  $\mathbb{R}_K$  for the topological space  $\mathbb{R}$  under the  $K$ -topology.

We prove this is a basis for a topology.

PROOF:

- ⟨1⟩1. For all  $x \in \mathbb{R}$  there exists an open interval  $(a, b)$  such that  $x \in (a, b)$ .  
PROOF: Take  $(a, b) = (x - 1, x + 1)$ .
- ⟨1⟩2. For any basic open sets  $B_1, B_2$  if  $x \in B_1 \cap B_2$ , then there exists a basic open set  $B_3$  such that  $x \in B_3 \subseteq B_1 \cap B_2$
- ⟨2⟩1. CASE:  $B_1 = (a, b), B_2 = (c, d)$   
PROOF: Take  $B_3 = (\max(a, c), \min(b, d))$ .
- ⟨2⟩2. CASE:  $B_1 = (a, b)$  or  $(a, b) \setminus K, B_2 = (c, d)$  or  $(c, d) \setminus K$ , and they are not both open intervals.  
PROOF: Take  $B_3 = (\max(a, c), \min(b, d)) \setminus K$ .

□

**Lemma 68.** *The lower limit topology and the  $K$ -topology are incomparable.*

PROOF:

- ⟨1⟩1. The interval  $[10, 11)$  is not open in the  $K$ -topology.  
PROOF: There is no open interval  $(a, b)$  such that  $10 \in (a, b) \subseteq [10, 11)$  or  $10 \in (a, b) \setminus K \subseteq [10, 11)$ .
- ⟨1⟩2. The set  $(-1, 1) \setminus K$  is not open in the lower limit topology.  
PROOF: There is no half-open interval  $[a, b)$  such that  $0 \in [a, b) \subseteq (-1, 1) \setminus K$ , since there must be a positive integer  $n$  with  $1/n \in [a, b)$ .

□

**Definition 69** (Subbasis). A *subbasis*  $\mathcal{S}$  for a topology on  $X$  is a set  $\mathcal{S} \subseteq \mathcal{P}X$  such that  $\bigcup \mathcal{S} = X$ .

The topology *generated* by the subbasis  $\mathcal{S}$  is the set of all unions of finite intersections of elements of  $\mathcal{S}$ .

We prove this is a topology.

PROOF:

- ⟨1⟩1. The set  $\mathcal{B}$  of all finite intersections of elements of  $\mathcal{S}$  forms a basis for a topology on  $X$ .  
⟨2⟩1.  $\bigcup \mathcal{B} = X$   
PROOF: Since  $\mathcal{S} \subseteq \mathcal{B}$ .
- ⟨2⟩2.  $\mathcal{B}$  is closed under binary intersection.  
PROOF: By definition.
- ⟨1⟩2. Q.E.D.  
PROOF: By Lemma 62.

□

We have simultaneously proved:

**Proposition 70.** *Let  $\mathcal{S}$  be a subbasis for the topology on  $X$ . Then the set of all finite intersections of elements of  $\mathcal{S}$  is a basis for the topology on  $X$ .*

**Proposition 71.** *Let  $X$  be a set. Let  $\mathcal{S}$  be a subbasis for a topology  $\mathcal{T}$  on  $X$ . Then  $\mathcal{T}$  is the coarsest topology that includes  $\mathcal{S}$ .*

PROOF: Since every topology that includes  $\mathcal{S}$  includes every union of finite intersections of elements of  $\mathcal{S}$ . □

## 12 Local Basis at a Point

**Definition 72** (Local Basis). Let  $X$  be a topological space and  $a \in X$ . A *(local) basis at  $a$*  is a set  $\mathcal{B}$  of neighbourhoods of  $a$  such that every neighbourhood of  $a$  includes some member of  $\mathcal{B}$ .

**Lemma 73.** *If there exists a countable local basis at a point  $a$ , then there exists a countable local basis  $\{B_n \mid n \geq 1\}$  such that  $B_1 \supseteq B_2 \supseteq \dots$ .*

PROOF: Given a countable local basis  $\{C_n \mid n \geq 1\}$ , take  $B_n = C_1 \cap \dots \cap C_n$ .  $\square$

## 13 Convergence

**Definition 74** (Convergence). Let  $X$  be a topological space. Let  $(a_n)_{n \in \mathbb{N}}$  be a sequence of points in  $X$  and  $l \in X$ . Then the sequence  $(a_n)_{n \in \mathbb{N}}$  *converges* to the *limit*  $l$ ,  $a_n \rightarrow l$  as  $n \rightarrow \infty$ , if and only if, for every neighbourhood  $U$  of  $l$ , there exists  $N$  such that, for all  $n \geq N$ , we have  $a_n \in U$ .

**Lemma 75.** *Let  $X$  be a topological space. Let  $A \subseteq X$  and  $l \in X$ . If there is a sequence of points in  $A$  that converges to  $l$  then  $l \in \overline{A}$ .*

PROOF:

- $\langle 1 \rangle 1$ . LET:  $(a_n)$  be a sequence of points in  $A$  that converges to  $l$ .
- $\langle 1 \rangle 2$ . LET:  $U$  be a neighbourhood of  $l$ .
- $\langle 1 \rangle 3$ . PICK  $N$  such that, for all  $n \geq N$ , we have  $a_n \in U$ .
- $\langle 1 \rangle 4$ .  $a_N \in U \cap A$
- $\langle 1 \rangle 5$ . Q.E.D.

PROOF: Theorem 48.

$\square$

**Proposition 76.** *Let  $X$  be a topological space. Let  $\mathcal{B}$  be a basis for the topology on  $X$ . Let  $(a_n)$  be a sequence in  $X$  and  $l \in X$ . Then  $a_n \rightarrow l$  as  $n \rightarrow \infty$  if and only if, for every  $B \in \mathcal{B}$  with  $l \in B$ , there exists  $N$  such that, for all  $n \geq N$ , we have  $a_n \in B$ .*

PROOF:

- $\langle 1 \rangle 1$ . If  $a_n \rightarrow l$  as  $n \rightarrow \infty$  then, for every  $B \in \mathcal{B}$  with  $l \in B$ , there exists  $N$  such that, for all  $n \geq N$ , we have  $a_n \in B$ .

PROOF: Since every element of  $\mathcal{B}$  is open (Corollary 62.1).

- $\langle 1 \rangle 2$ . If, for every  $B \in \mathcal{B}$  with  $l \in B$ , there exists  $N$  such that, for all  $n \geq N$ , we have  $a_n \in B$ , then  $a_n \rightarrow l$  as  $n \rightarrow \infty$ .

- $\langle 2 \rangle 1$ . ASSUME: for every  $B \in \mathcal{B}$  with  $l \in B$ , there exists  $N$  such that, for all  $n \geq N$ , we have  $a_n \in B$

- $\langle 2 \rangle 2$ . LET:  $U$  be a neighbourhood of  $l$ .

- $\langle 2 \rangle 3$ . PICK  $B \in \mathcal{B}$  such that  $l \in B \subseteq U$

- $\langle 2 \rangle 4$ . PICK  $N$  such that, for all  $n \geq N$ , we have  $a_n \in B$

PROOF: From  $\langle 2 \rangle 1$ .

- $\langle 2 \rangle 5$ . For all  $n \geq N$  we have  $a_n \in U$

□

**Lemma 77.** *If a sequence  $(a_n)$  is constant with  $a_n = l$  for all  $n$ , then  $a_n \rightarrow l$  as  $n \rightarrow \infty$ .*

PROOF: Immediate from definitions. □

**Theorem 78.** *Let  $X$  be a linearly ordered set. Let  $(s_n)$  be a monotone increasing sequence in  $X$  with a supremum  $s$ . Then  $s_n \rightarrow s$  as  $n \rightarrow \infty$ .*

PROOF:

⟨1⟩1. ASSUME:  $s$  is not least in  $X$ .

PROOF: Otherwise  $(s_n)$  is the constant sequence  $s$  and the result follows from Lemma 77.

⟨1⟩2. LET:  $U$  be a neighbourhood of  $s$ .

⟨1⟩3. PICK  $a < s$  such that  $(a, s] \subseteq U$

⟨1⟩4. PICK  $N$  such that  $a < a_N$ .

⟨1⟩5. For all  $n \geq N$  we have  $a_n \in (a, s]$

⟨1⟩6. For all  $n \geq N$  we have  $a_n \in U$ .

□

**Theorem 79.** *If  $\sum_{i=0}^{\infty} a_i = s$  and  $\sum_{i=0}^{\infty} b_i = t$  then  $\sum_{i=0}^{\infty} (ca_i + b_i) = cs + t$ .*

PROOF:  $\sum_{i=0}^N (ca_i + b_i) = c \sum_{i=0}^N a_i + \sum_{i=0}^N b_i \rightarrow cs + t$  as  $n \rightarrow \infty$ . □

**Theorem 80** (Comparison Test). *If  $|a_i| \leq b_i$  for all  $i$  and  $\sum_{i=0}^{\infty} b_i$  converges then  $\sum_{i=0}^{\infty} a_i$  converges.*

PROOF:

⟨1⟩1.  $\sum_{i=0}^{\infty} |a_i|$  converges

PROOF: The partial sums  $\sum_{i=0}^N |a_i|$  form an increasing sequence bounded above by  $\sum_{i=0}^{\infty} b_i$ .

⟨1⟩2. LET:  $c_i = |a_i| + a_i$  for all  $i$

⟨1⟩3.  $\sum_{i=0}^{\infty} c_i$  converges

PROOF: Each  $c_i$  is either  $2|a_i|$  or 0. So the partial sums  $\sum_{i=0}^N c_i$  form an increasing sequence bounded above by  $2 \sum_{i=0}^{\infty} b_i$ .

⟨1⟩4. Q.E.D.

PROOF: Since  $a_i = c_i - |a_i|$ .

□

**Corollary 80.1.** *If  $\sum_{i=0}^{\infty} |a_i|$  converges then  $\sum_{i=0}^{\infty} a_i$  converges.*

**Theorem 81** (Weierstrass M-test). *Let  $X$  be a set and  $(f_n : X \rightarrow \mathbb{R})$  be a sequence of functions. Let*

$$s_n(x) = \sum_{i=0}^n f_i(x)$$

*for all  $n, x$ . Suppose  $|f_i(x)| \leq M_i$  for all  $i \geq 0$  and  $x \in X$ . If the series  $\sum_{i=0}^{\infty} M_i$  converges, then the sequence  $(s_n)$  converges uniformly to*

$$s(x) = \sum_{i=0}^{\infty} f_i(x) .$$

PROOF:

- ⟨1⟩1. LET:  $r_n = \sum_{i=n+1}^{\infty} M_i$  for all  $n$   
 ⟨1⟩2. Given  $0 \leq n < k$ , we have  $|s_k(x) - s_n(x)| \leq r_n$

PROOF:

$$\begin{aligned} |s_k(x) - s_n(x)| &= \left| \sum_{i=n+1}^k f_i(x) \right| \\ &\leq \sum_{i=n+1}^k |f_i(x)| \\ &\leq \sum_{i=n+1}^k M_i \\ &\leq r_n \end{aligned}$$

- ⟨1⟩3. Given  $n \geq 0$  we have  $|s(x) - s_n(x)| \leq r_n$

PROOF: By taking the limit  $k \rightarrow \infty$  in ⟨1⟩2.

- ⟨1⟩4. Q.E.D.

PROOF: Since  $r_n \rightarrow 0$  as  $n \rightarrow \infty$ .

□

## 14 Locally Finite Sets

**Definition 82** (Locally Finite). Let  $X$  be a topological space and  $\{A_\alpha\}$  a family of subsets of  $X$ . Then  $\mathcal{A}$  is *locally finite* if and only if every point in  $X$  has a neighbourhood that intersects  $A_\alpha$  for only finitely many  $\alpha$ .

**Theorem 83** (Pasting Lemma). Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Let  $\{A_\alpha\}$  be a locally finite family of closed subsets of  $X$  that cover  $X$ . Suppose  $f \upharpoonright A_\alpha$  is continuous for all  $\alpha$ . Then  $f$  is continuous.

PROOF:

- ⟨1⟩1. Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Let  $A$  and  $B$  be closed subsets of  $X$  such that  $X = A \cup B$ . Suppose  $f \upharpoonright A$  and  $f \upharpoonright B$  are continuous. Then  $f$  is continuous.

- ⟨2⟩1. LET:  $C \subseteq Y$  be closed.

- ⟨2⟩2.  $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$

- ⟨2⟩3.  $f^{-1}(C)$  and  $g^{-1}(C)$  are closed in  $X$ .

PROOF: Theorems 93 and 143.

- ⟨2⟩4.  $h^{-1}(C)$  is closed in  $X$ .

PROOF: Lemma 34.

- ⟨2⟩5. Q.E.D.

PROOF: Theorem 93.

- ⟨1⟩2. Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Let  $\{A_\alpha\}$  be a finite family of closed subsets of  $X$  that cover  $X$ . Suppose  $f \upharpoonright A_\alpha$  is continuous for all  $\alpha$ . Then  $f$  is continuous.

PROOF: From ⟨1⟩1 by induction.

- ⟨1⟩3. Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Let  $\{A_\alpha\}$  be a locally finite family of closed subsets of  $X$  that cover  $X$ . Suppose  $f \upharpoonright A_\alpha$  is continuous for all  $\alpha$ . Then  $f$  is continuous.
- ⟨2⟩1. LET:  $x \in X$   
PROVE:  $f$  is continuous at  $x$
- ⟨2⟩2. PICK a neighbourhood  $U$  of  $x$  that intersects  $A_\alpha$  for only finitely many  $\alpha$ .
- ⟨2⟩3.  $f \upharpoonright U$  is continuous  
PROOF: By ⟨1⟩2.
- ⟨2⟩4. Q.E.D.  
PROOF: Lemma 103.

□

The following example shows that we cannot remove the assumption of local finiteness.

**Example 84.** Define  $f : [-1, 1] \rightarrow \mathbb{R}$  by:  $f(x) = 1$  if  $x < -1$ ,  $f(x) = 0$  if  $x > 1$ . Let  $C_n = [-1, -1/n]$  for  $n \geq 1$ , and  $D = [0, 1]$ . Then  $[-1, 1] = \bigcup_{n=1}^{\infty} C_n \cup D$  and  $f$  is continuous on each  $C_n$  and each  $D$ , but  $f$  is not continuous on  $[-1, 1]$ .

## 15 Open Maps

**Definition 85** (Open Map). Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Then  $f$  is an *open map* if and only if, for every open set  $U$  in  $X$ , the set  $f(U)$  is open in  $Y$ .

**Lemma 86.** Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Let  $\mathcal{B}$  be a basis for the topology on  $X$ . If  $f(B)$  is open in  $Y$  for all  $B \in \mathcal{B}$ , then  $f$  is an open map.

PROOF: From Lemma 62. □

**Proposition 87.** Let  $X$  and  $Y$  be topological spaces. Let  $\mathcal{B}$  be a basis for the topology on  $X$ . Let  $f : X \rightarrow Y$ . Suppose that, for all  $B \in \mathcal{B}$ , we have  $f(B)$  is open in  $Y$ . Then  $f$  is an open map.

PROOF: For any  $\mathcal{A} \subseteq \mathcal{B}$ , we have  $f(\bigcup \mathcal{A}) = \bigcup_{B \in \mathcal{A}} f(B)$  is open in  $Y$ . The result follows from Lemma 62. □

## 16 Continuous Functions

**Definition 88** (Continuous). Let  $X$  and  $Y$  be topological spaces. A function  $f : X \rightarrow Y$  is *continuous* if and only if, for every open set  $V$  in  $Y$ , the set  $f^{-1}(V)$  is open in  $X$ .

**Proposition 89.** Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Let  $\mathcal{B}$  be a basis for  $Y$ . Then  $f$  is continuous if and only if, for all  $B \in \mathcal{B}$ , we have  $f^{-1}(B)$  is open in  $X$ .



PROOF:

⟨1⟩1. If  $f$  is continuous then, for all  $B \in \mathcal{B}$ , we have  $f^{-1}(B)$  is open in  $X$ .

PROOF: Since every element of  $\mathcal{B}$  is open (Lemma 62).

⟨1⟩2. Suppose that, for all  $B \in \mathcal{B}$ , we have  $f^{-1}(B)$  is open in  $X$ . Then  $f$  is continuous.

⟨2⟩1. ASSUME: For all  $B \in \mathcal{B}$ , we have  $f^{-1}(B)$  is open in  $X$ .

⟨2⟩2. LET:  $V$  be open in  $Y$ .

⟨2⟩3. PICK  $\mathcal{A} \subseteq \mathcal{B}$  such that  $V = \bigcup \mathcal{A}$

PROOF: By Lemma 62.

⟨2⟩4.  $f^{-1}(V)$  is open in  $X$ .

PROOF:

$$\begin{aligned} f^{-1}(V) &= f^{-1}\left(\bigcup \mathcal{A}\right) \\ &= \bigcup_{B \in \mathcal{A}} f^{-1}(B) \end{aligned}$$

□

**Proposition 90.** *Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Let  $\mathcal{S}$  be a subbasis for  $Y$ . Then  $f$  is continuous if and only if, for all  $S \in \mathcal{S}$ , we have  $f^{-1}(S)$  is open in  $X$ .*

PROOF:

⟨1⟩1. If  $f$  is continuous then, for all  $S \in \mathcal{S}$ , we have  $f^{-1}(S)$  is open in  $X$ .

PROOF: Since every element of  $\mathcal{S}$  is open.

⟨1⟩2. Suppose that, for all  $S \in \mathcal{S}$ , we have  $f^{-1}(S)$  is open in  $X$ . Then  $f$  is continuous.

⟨2⟩1. ASSUME: For all  $S \in \mathcal{S}$ , we have  $f^{-1}(S)$  is open in  $X$ .

⟨2⟩2. LET:  $S_1, \dots, S_n \in \mathcal{S}$

⟨2⟩3.  $f^{-1}(S_1 \cap \dots \cap S_n)$  is open in  $X$

PROOF: Since  $f^{-1}(S_1 \cap \dots \cap S_n) = f^{-1}(S_1) \cap \dots \cap f^{-1}(S_n)$ .

⟨2⟩4. Q.E.D.

PROOF: By Propositions 89 and 70.

□

**Proposition 91.** *Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Let  $\mathcal{S}$  be a basis for  $Y$ . Then  $f$  is continuous if and only if, for all  $V \in \mathcal{S}$ , we have  $f^{-1}(V)$  is open in  $X$ .*

PROOF:

⟨1⟩1. If  $f$  is continuous then, for all  $V \in \mathcal{S}$ , we have  $f^{-1}(V)$  is open in  $X$ .

PROOF: Since every element of  $\mathcal{S}$  is open.

⟨1⟩2. Suppose that, for all  $V \in \mathcal{S}$ , we have  $f^{-1}(V)$  is open in  $X$ . Then  $f$  is continuous.

⟨2⟩1. ASSUME: For all  $V \in \mathcal{S}$ , we have  $f^{-1}(V)$  is open in  $X$ .

⟨2⟩2. For every set  $B$  that is the finite intersection of elements of  $\mathcal{S}$ , we have  $f^{-1}(B)$  is open in  $X$ .

PROOF: Because  $f^{-1}(V_1 \cap \dots \cap V_n) = f^{-1}(V_1) \cap \dots \cap f^{-1}(V_n)$ .

⟨2⟩3. Q.E.D.

PROOF: From Propositions 70 and 89.

□

**Definition 92** (Continuous at a Point). Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$  and  $x \in X$ . Then  $f$  is *continuous at  $x$*  if and only if, for every neighbourhood  $V$  of  $f(x)$ , there exists a neighbourhood  $U$  of  $x$  such that  $f(U) \subseteq V$ .

**Theorem 93.** Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Then the following are equivalent:

1.  $f$  is continuous.
2. For all  $A \subseteq X$ , we have  $f(\overline{A}) \subseteq \overline{f(A)}$
3. For all  $B \subseteq Y$  closed, we have  $f^{-1}(B)$  is closed in  $X$ .
4.  $f$  is continuous at every point of  $X$ .

PROOF:

⟨1⟩1.  $1 \Rightarrow 2$

⟨2⟩1. ASSUME:  $f$  is continuous.

⟨2⟩2. LET:  $A \subseteq X$

⟨2⟩3. LET:  $x \in \overline{A}$

PROVE:  $f(x) \in \overline{f(A)}$

⟨2⟩4. LET:  $V$  be a neighbourhood of  $f(x)$

⟨2⟩5.  $f^{-1}(V)$  is a neighbourhood of  $x$

⟨2⟩6. PICK  $y \in A \cap f^{-1}(V)$

PROOF: By Theorem 48.

⟨2⟩7.  $f(y) \in V \cap f(A)$

⟨2⟩8. Q.E.D.

PROOF: By Theorem 48.

⟨1⟩2.  $2 \Rightarrow 3$

⟨2⟩1. ASSUME: 2

⟨2⟩2. LET:  $B$  be closed in  $Y$

⟨2⟩3. LET:  $x \in \overline{f^{-1}(B)}$

PROVE:  $x \in f^{-1}(B)$

⟨2⟩4.  $f(x) \in B$

PROOF:

$$f(x) \in \overline{f(f^{-1}(B))}$$

$$\subseteq \overline{f(f^{-1}(B))}$$

(⟨2⟩1)

$$\subseteq \overline{B}$$

(Proposition 49)

$$= B$$

⟨1⟩3.  $3 \Rightarrow 1$

⟨2⟩1. ASSUME: 3

⟨2⟩2. LET:  $V$  be open in  $Y$

⟨2⟩3.  $Y \setminus V$  is closed in  $Y$

$\langle 2 \rangle 4$ .  $f^{-1}(Y \setminus V)$  is closed in  $X$   
 $\langle 2 \rangle 5$ .  $X \setminus f^{-1}(V)$  is closed in  $X$   
 $\langle 2 \rangle 6$ .  $f^{-1}(V)$  is open in  $X$   
 $\langle 1 \rangle 4$ .  $1 \Rightarrow 4$   
 PROOF: For any neighbourhood  $V$  of  $f(x)$ , the set  $U = f^{-1}(V)$  is a neighbourhood of  $x$  such that  $f(U) \subseteq V$ .  
 $\langle 1 \rangle 5$ .  $4 \Rightarrow 1$   
 $\langle 2 \rangle 1$ . ASSUME: 4  
 $\langle 2 \rangle 2$ . LET:  $V$  be open in  $Y$   
 $\langle 2 \rangle 3$ . LET:  $x \in f^{-1}(V)$   
 $\langle 2 \rangle 4$ .  $V$  is a neighbourhood of  $f(x)$   
 $\langle 2 \rangle 5$ . PICK a neighbourhood  $U$  of  $x$  such that  $f(U) \subseteq V$   
 $\langle 2 \rangle 6$ .  $x \in U \subseteq f^{-1}(V)$   
 $\langle 2 \rangle 7$ . Q.E.D.  
 PROOF: By Lemma 26.

□

**Theorem 94.** *A constant function is continuous.*

PROOF: Let  $X$  and  $Y$  be topological spaces. Let  $b \in Y$ , and let  $f : X \rightarrow Y$  be the constant function with value  $b$ . For any open  $V \subseteq Y$ , the set  $f^{-1}(V)$  is either  $X$  (if  $b \in V$ ) or  $\emptyset$  (if  $b \notin V$ ). □

**Theorem 95.** *If  $A$  is a subspace of  $X$  then the inclusion  $j : A \rightarrow X$  is continuous.*

PROOF: For any  $V$  open in  $X$ , we have  $j^{-1}(V) = V \cap A$  is open in  $A$ . □

**Theorem 96.** *The composite of two continuous functions is continuous.*

PROOF: Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  be continuous. For any  $V$  open in  $Z$ , we have  $(g \circ f)^{-1}(V) = f^{-1}(g^{-1}(V))$  is open in  $X$ . □

**Theorem 97.** *Let  $f : X \rightarrow Y$  be a continuous function and  $A$  be a subspace of  $X$ . Then the restriction  $f \upharpoonright A : A \rightarrow Y$  is continuous.*

PROOF: Let  $V$  be open in  $Y$ . Then  $(f \upharpoonright A)^{-1}(V) = f^{-1}(V) \cap A$  is open in  $A$ . □

**Theorem 98.** *Let  $f : X \rightarrow Y$  be continuous. Let  $Z$  be a subspace of  $Y$  such that  $f(X) \subseteq Z$ . Then the corestriction  $f : X \rightarrow Z$  is continuous.*

PROOF:

$\langle 1 \rangle 1$ . LET:  $V$  be open in  $Z$ .  
 $\langle 1 \rangle 2$ . PICK  $U$  open in  $Y$  such that  $V = U \cap Z$ .  
 $\langle 1 \rangle 3$ .  $f^{-1}(V) = f^{-1}(U)$   
 $\langle 1 \rangle 4$ .  $f^{-1}(V)$  is open in  $X$ .  
 □

**Theorem 99.** *Let  $f : X \rightarrow Y$  be continuous. Let  $Z$  be a space such that  $Y$  is a subspace of  $Z$ . Then the expansion  $f : X \rightarrow Z$  is continuous.*

PROOF: Let  $V$  be open in  $Z$ . Then  $f^{-1}(V) = f^{-1}(V \cap Y)$  is open in  $X$ .  $\square$

**Theorem 100.** *Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Suppose  $\mathcal{U}$  is a set of open sets in  $X$  such that  $X = \bigcup \mathcal{U}$  and, for all  $U \in \mathcal{U}$ , we have  $f \upharpoonright U : U \rightarrow Y$  is continuous. Then  $f$  is continuous.*

PROOF:

- $\langle 1 \rangle 1$ . LET:  $V$  be open in  $Y$
- $\langle 1 \rangle 2$ .  $f^{-1}(V) = \bigcup_{U \in \mathcal{U}} (f \upharpoonright U)^{-1}(V)$
- $\langle 1 \rangle 3$ . For all  $U \in \mathcal{U}$ , we have  $(f \upharpoonright U)^{-1}(V)$  is open in  $U$ .
- $\langle 1 \rangle 4$ . For all  $U \in \mathcal{U}$ , we have  $(f \upharpoonright U)^{-1}(V)$  is open in  $X$ .

PROOF: Lemma 142.

$\square$

**Proposition 101.** *Let  $X$  and  $X'$  be the same set  $X$  under two topologies  $\mathcal{T}$  and  $\mathcal{T}'$ . Let  $i : X \rightarrow X'$  be the identity function. Then  $i$  is continuous if and only if  $\mathcal{T}' \subseteq \mathcal{T}$ .*

PROOF: Immediate from definitions.  $\square$

**Proposition 102.** *Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $a \in \mathbb{R}$ . Then  $f$  is continuous on the right at  $a$  if and only if  $f$  is continuous at  $a$  as a function  $\mathbb{R}_l \rightarrow \mathbb{R}$ .*

PROOF:

- $\langle 1 \rangle 1$ . If  $f$  is continuous on the right at  $a$  then  $f$  is continuous at  $a$  as a function  $\mathbb{R}_l \rightarrow \mathbb{R}$ .
  - $\langle 2 \rangle 1$ . ASSUME:  $f$  is continuous on the right at  $a$ .
  - $\langle 2 \rangle 2$ . LET:  $V$  be a neighbourhood of  $f(a)$
  - $\langle 2 \rangle 3$ . PICK  $b, c$  such that  $f(a) \in (b, c) \subseteq V$ .
  - $\langle 2 \rangle 4$ . LET:  $\epsilon = \min(c - f(a), f(a) - b)$
  - $\langle 2 \rangle 5$ . PICK  $\delta > 0$  such that, for all  $x$ , if  $a < x < a + \delta$  then  $|f(x) - f(a)| < \epsilon$
  - $\langle 2 \rangle 6$ . LET:  $U = [a, a + \delta)$
  - $\langle 2 \rangle 7$ .  $f(U) \subseteq V$
- $\langle 1 \rangle 2$ . If  $f$  is continuous at  $a$  as a function  $\mathbb{R}_l \rightarrow \mathbb{R}$  then  $f$  is continuous on the right at  $a$ .
  - $\langle 2 \rangle 1$ . ASSUME:  $f$  is continuous at  $a$  as a function  $\mathbb{R}_l \rightarrow \mathbb{R}$
  - $\langle 2 \rangle 2$ . LET:  $\epsilon > 0$
  - $\langle 2 \rangle 3$ . PICK a neighbourhood  $U$  of  $a$  such that  $f(U) \subseteq (f(a) - \epsilon, f(a) + \epsilon)$
  - $\langle 2 \rangle 4$ . PICK  $b, c$  such that  $a \in [b, c) \subset U$
  - $\langle 2 \rangle 5$ . LET:  $\delta = c - a$
  - $\langle 2 \rangle 6$ . For all  $x$ , if  $a < x < a + \delta$  then  $|f(x) - f(a)| < \epsilon$

$\square$

**Lemma 103.** *Let  $f : X \rightarrow Y$ . Let  $Z$  be an open subspace of  $X$  and  $a \in Z$ . If  $f \upharpoonright Z$  is continuous at  $a$  then  $f$  is continuous at  $a$ .*

PROOF:

- $\langle 1 \rangle 1$ . LET:  $V$  be a neighbourhood of  $f(a)$

⟨1⟩2. PICK a neighbourhood  $W$  of  $x$  in  $Z$  such that  $f(W) \subseteq V$

⟨1⟩3.  $W$  is a neighbourhood of  $x$  in  $X$  such that  $f(W) \subseteq V$

PROOF: Lemma 142.

□

**Proposition 104.** *Let  $f : A \rightarrow B$  and  $g : C \rightarrow D$  be continuous. Define  $f \times g : A \times C \rightarrow B \times D$  by*

$$(f \times g)(a, c) = (f(a), g(c)) .$$

*Then  $f \times g$  is continuous.*

PROOF:  $\pi_1 \circ (f \times g) = f \circ \pi_1$  and  $\pi_2 \circ (f \times g) = g \circ \pi_2$  are continuous by Theorem 96. The result follows by Theorem 132.

**Proposition 105.** *Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$  be continuous. If  $a_n \rightarrow l$  as  $n \rightarrow \infty$  in  $X$  then  $f(a_n) \rightarrow f(l)$  as  $n \rightarrow \infty$ .*

PROOF:

⟨1⟩1. LET:  $V$  be a neighbourhood of  $f(l)$

⟨1⟩2. PICK a neighbourhood  $U$  of  $l$  such that  $f(U) \subseteq V$

⟨1⟩3. PICK  $N$  such that, for all  $n \geq N$ , we have  $a_n \in U$

⟨1⟩4. For all  $n \geq N$  we have  $f(a_n) \in V$

□

## 17 Homeomorphisms

**Definition 106** (Homeomorphism). Let  $X$  and  $Y$  be topological spaces. A *Homeomorphism*  $f$  between  $X$  and  $Y$ ,  $f : X \cong Y$ , is a bijection  $f : X \rightarrow Y$  such that both  $f$  and  $f^{-1}$  are continuous.

**Lemma 107.** *Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$  a bijection. Then the following are equivalent:*

1.  $f$  is a homeomorphism.
2.  $f$  is continuous and an open map.
3.  $f$  is continuous and a closed map.
4. For any  $U \subseteq X$ , we have  $U$  is open if and only if  $f(U)$  is open.

PROOF: Immediate from definitions. □

**Proposition 108.** *Let  $X$  and  $X'$  be the same set  $X$  under two topologies  $\mathcal{T}$  and  $\mathcal{T}'$ . Let  $i : X \rightarrow X'$  be the identity function. Then  $i$  is a homeomorphism if and only if  $\mathcal{T} = \mathcal{T}'$ .*

PROOF: Immediate from definitions. □

**Definition 109** (Topological Property). Let  $P$  be a property of topological spaces. Then  $P$  is a *topological* property if and only if, for any spaces  $X$  and  $Y$ , if  $P$  holds of  $X$  and  $X \cong Y$  then  $P$  holds of  $Y$ .

**Definition 110** (Topological Imbedding). Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Then  $f$  is a *topological imbedding* if and only if the corestriction  $f : X \rightarrow f(X)$  is a homeomorphism.

**Proposition 111.** *Let  $X$  and  $Y$  be topological spaces and  $a \in X$ . The function  $i : Y \rightarrow X \times Y$  that maps  $y$  to  $(a, y)$  is an imbedding.*

PROOF:

$\langle 1 \rangle 1$ .  $i$  is injective

$\langle 1 \rangle 2$ .  $i$  is continuous.

PROOF: For  $U$  open in  $X$  and  $V$  open in  $Y$ , we have  $i^{-1}(U \times V)$  is  $V$  if  $a \in U$ , and  $\emptyset$  if  $a \notin U$ .

$\langle 1 \rangle 3$ .  $i : Y \rightarrow i(Y)$  is an open map.

PROOF: For  $V$  open in  $Y$  we have  $i(V) = (X \times V) \cap i(Y)$ .

□

## 18 The Order Topology

**Definition 112** (Order Topology). Let  $X$  be a linearly ordered set with at least two points. The *order topology* on  $X$  is the topology generated by the basis  $\mathcal{B}$  consisting of:

- all open intervals  $(a, b)$ ;
- all intervals of the form  $[\perp, b)$  where  $\perp$  is least in  $X$ ;
- all intervals of the form  $(a, \top]$  where  $\top$  is greatest in  $X$ .

We prove this is a basis for a topology.

PROOF:

$\langle 1 \rangle 1$ . For all  $x \in X$  there exists  $B \in \mathcal{B}$  such that  $x \in B$ .

$\langle 2 \rangle 1$ . LET:  $x \in X$

$\langle 2 \rangle 2$ . CASE:  $x$  is greatest in  $X$ .

$\langle 3 \rangle 1$ . PICK  $y \in X$  with  $y \neq x$

$\langle 3 \rangle 2$ .  $x \in (y, x] \in \mathcal{B}$

$\langle 2 \rangle 3$ . CASE:  $x$  is least in  $X$ .

$\langle 3 \rangle 1$ . PICK  $y \in X$  with  $y \neq x$

$\langle 3 \rangle 2$ .  $x \in [x, y) \in \mathcal{B}$

$\langle 2 \rangle 4$ . CASE:  $x$  is neither greatest nor least in  $X$ .

$\langle 3 \rangle 1$ . PICK  $a, b \in X$  with  $a < x$  and  $x < b$

$\langle 3 \rangle 2$ .  $x \in (a, b) \in \mathcal{B}$

$\langle 1 \rangle 2$ . For all  $B_1, B_2 \in \mathcal{B}$  and  $x \in B_1 \cap B_2$ , there exists  $B_3 \in \mathcal{B}$  such that  $x \in B_3 \subseteq B_1 \cap B_2$

- ⟨2⟩1. LET:  $B_1, B_2 \in \mathcal{B}$  and  $x \in B_1 \cap B_2$
- ⟨2⟩2. CASE:  $B_1 = (a, b)$ ,  $B_2 = (c, d)$   
PROOF: Take  $B_3 = (\max(a, c), \min(b, d))$ .
- ⟨2⟩3. CASE:  $B_1 = (a, b)$ ,  $B_2 = [\perp, d)$   
PROOF: Take  $B_3 = (a, \min(b, d))$ .
- ⟨2⟩4. CASE:  $B_1 = (a, b)$ ,  $B_2 = (c, \top]$   
PROOF: Take  $B_3 = (\max(a, c), b)$ .
- ⟨2⟩5. CASE:  $B_1 = [\perp, b)$ ,  $B_2 = [\perp, d)$   
PROOF: Take  $B_3 = [\perp, \min(b, d))$ .
- ⟨2⟩6. CASE:  $B_1 = [\perp, b)$ ,  $B_2 = (c, \top]$   
PROOF: Take  $B_3 = (c, b)$ .

□

**Lemma 113.** *Let  $X$  be a linearly ordered set. Then the open rays form a subbasis for the order topology on  $X$ .*

PROOF:

- ⟨1⟩1. Every open ray is open.
- ⟨2⟩1. For all  $a \in X$ , the ray  $(-\infty, a)$  is open.
  - ⟨3⟩1. LET:  $x \in (-\infty, a)$
  - ⟨3⟩2. CASE:  $x$  is least in  $X$   
PROOF:  $x \in [x, a) = (-\infty, a)$ .
  - ⟨3⟩3. CASE:  $x$  is not least in  $X$ 
    - ⟨4⟩1. PICK  $y < x$
    - ⟨4⟩2.  $x \in (y, a) \subseteq (-\infty, a)$
- ⟨2⟩2. For all  $a \in X$ , the ray  $(a, +\infty)$  is open.  
PROOF: Similar.
- ⟨1⟩2. Every basic open set is a finite intersection of open rays.  
PROOF: We have  $(a, b) = (a, +\infty) \cap (-\infty, b)$ ,  $[\perp, b) = (-\infty, b)$  and  $(a, \top] = (a, +\infty)$ .

□

**Definition 114** (Standard Topology on the Real Line). The *standard topology on the real line* is the order topology on  $\mathbb{R}$  generated by the standard order.

**Lemma 115.** *The lower limit topology is strictly finer than the standard topology on  $\mathbb{R}$ .*

PROOF:

- ⟨1⟩1. Every open interval is open in the lower limit topology.  
PROOF: If  $x \in (a, b)$  then  $x \in [x, b) \subseteq (a, b)$ .
- ⟨1⟩2. The half-open interval  $[0, 1)$  is not open in the standard topology.  
PROOF: There is no open interval  $(a, b)$  such that  $0 \in (a, b) \subseteq [0, 1)$ .

□

**Lemma 116.** *The  $K$ -topology is strictly finer than the standard topology on  $\mathbb{R}$ .*

PROOF:

⟨1⟩1. Every open interval is open in the  $K$ -topology.

PROOF: Corollary 62.1.

⟨1⟩2. The set  $(-1, 1) \setminus K$  is not open in the standard topology.

PROOF: There is no open interval  $(a, b)$  such that  $0 \in (a, b) \subseteq (-1, 1) \setminus K$ , since there must be a positive integer  $n$  with  $1/n \in (a, b)$ .

□

**Lemma 117.** *Let  $X$  be a topological space. Let  $Y$  be a linearly ordered set in the order topology. Let  $f, g : X \rightarrow Y$  be continuous. Then  $C = \{x \in X \mid f(x) \leq g(x)\}$  is closed.*

PROOF:

⟨1⟩1. LET:  $x \in X \setminus C$

⟨1⟩2.  $f(x) > g(x)$

PROVE: There exists a neighbourhood  $U$  of  $x$  such that  $U \subseteq X \setminus C$

⟨1⟩3. CASE: There exists  $y$  such that  $g(x) < y < f(x)$

PROOF: Take  $U = g^{-1}((-\infty, y)) \cup f^{-1}(y, +\infty)$ .

⟨1⟩4. CASE: There is no  $y$  such that  $g(x) < y < f(x)$

PROOF: Take  $U = g^{-1}((-\infty, f(x))) \cup f^{-1}(g(x), +\infty)$ .

□

**Proposition 118.** *Let  $X$  be a topological space. Let  $Y$  be a linearly ordered set in the order topology. Let  $f, g : X \rightarrow Y$  be continuous. Define  $h : X \rightarrow Y$  by  $h(x) = \min(f(x), g(x))$ . Then  $h$  is continuous.*

PROOF: By the Pasting Lemma applied to  $\{x \in X \mid f(x) \leq g(x)\}$  and  $\{x \in X \mid g(x) \leq f(x)\}$ , which are closed by Lemma 117.

**Proposition 119.** *Let  $X$  and  $Y$  be linearly ordered sets in the order topology. Let  $f : X \rightarrow Y$  be strictly monotone and surjective. Then  $f$  is a homeomorphism.*

PROOF:

⟨1⟩1.  $f$  is bijective.

PROOF: Proposition 6.

⟨1⟩2.  $f$  is continuous.

⟨2⟩1. For all  $y \in Y$  we have  $f^{-1}((y, +\infty))$  is open.

⟨3⟩1. LET:  $y \in Y$

⟨3⟩2. PICK  $x \in X$  such that  $f(x) = y$

PROOF: Since  $f$  is surjective.

⟨3⟩3.  $f^{-1}((y, +\infty)) = (x, +\infty)$

PROOF: By strict monotonicity.

⟨2⟩2. For all  $y \in Y$  we have  $f^{-1}((-\infty, y))$  is open.

PROOF: Similar.

⟨1⟩3.  $f^{-1}$  is continuous.

⟨2⟩1. For all  $x \in X$  we have  $f((x, +\infty))$  is open.

PROOF:  $f((x, +\infty)) = (f(x), +\infty)$ .

⟨2⟩2. For all  $x \in X$  we have  $f((-\infty, x))$  is open.

PROOF:  $f((-\infty, x)) = (-\infty, f(x))$ .

□



## 19 The $n$ th Root Function

**Proposition 120.** For all  $n \geq 1$ , the function  $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  defined by  $f(x) = x^n$  is a homeomorphism.

PROOF:

$\langle 1 \rangle 1$ .  $f$  is strictly monotone.

$\langle 2 \rangle 1$ . LET:  $x, y \in \mathbb{R}$  with  $0 \leq x < y$

$\langle 2 \rangle 2$ .  $x^n < y^n$

$$y^n - x^n = (y - x)(y^{n-1} + y^{n-2}x + y^{n-3}x^2 + \cdots + x^{n-1}) > 0$$

$\langle 1 \rangle 2$ .  $f$  is surjective.

$\langle 2 \rangle 1$ . LET:  $y \in \mathbb{R}_{\geq 0}$

$\langle 2 \rangle 2$ . PICK  $x \in \mathbb{R}$  such that  $y \leq x^n$

PROOF: If  $y \leq 1$  take  $x = 1$ , otherwise take  $x = y$ .

$\langle 2 \rangle 3$ . There exists  $x' \in [0, x]$  such that  $(x')^n = y$

PROOF: By the Intermediate Value Theorem.

$\langle 1 \rangle 3$ . Q.E.D.

PROOF: Proposition 119.

□

**Definition 121.** For  $n \geq 1$ , the  $n$ th root function is the function  $\mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  that is the inverse of  $\lambda x.x^n$ .

## 20 The Product Topology

**Definition 122** (Product Topology). Let  $\{A_i\}_{i \in I}$  be a family of topological spaces. The *product topology* on  $\prod_{i \in I} A_i$  is the topology generated by the subbasis consisting of the sets of the form  $\pi_i^{-1}(U)$  where  $i \in I$  and  $U$  is open in  $A_i$ .

**Proposition 123.** The product topology on  $\prod_{i \in I} A_i$  is generated by the basis consisting of all sets of the form  $\prod_{i \in I} U_i$  where  $\{U_i\}_{i \in I}$  is a family such that each  $U_i$  is an open set in  $A_i$  and  $U_i = A_i$  for all but finitely many  $i$ .

PROOF: From Proposition 70. □

**Proposition 124.** If  $A_i$  is closed in  $X_i$  for all  $i \in I$  then  $\prod_{i \in I} A_i$  is closed in  $\prod_{i \in I} X_i$ .

PROOF:

$$\left(\prod_{i \in I} X_i\right) \setminus \left(\prod_{i \in I} A_i\right) = \bigcup_{j \in I} \left(\prod_{i \in I} X_i \setminus \pi_j^{-1}(A_j)\right) \square$$

**Proposition 125.** Let  $\{A_i\}_{i \in I}$  be a family of topological spaces. For  $i \in I$ , let  $\mathcal{B}_i$  be a basis for the topology on  $A_i$ . Then  $\mathcal{B} = \{\prod_{i \in I} B_i \mid \forall i \in I. B_i \in \mathcal{B}_i, B_i = A_i \text{ for all but finitely many } i\}$  is a basis for the box topology on  $\prod_{i \in I} A_i$ .

PROOF:

- ⟨1⟩1. Every set in  $\mathcal{B}$  is open.
- ⟨1⟩2. For every point  $a \in \prod_{i \in I} A_i$  and every open set  $U$  with  $a \in U$ , there exists  $B \in \mathcal{B}$  such that  $a \in B \subseteq U$ .
  - ⟨2⟩1. LET:  $U$  be open and  $a \in U$
  - ⟨2⟩2. PICK a family  $\{U_i\}_{i \in I}$  such that each  $U_i$  is open in  $A_i$ , such that  $U_i = A_i$  except for  $i = i_1, \dots, i_n$ , and such that  $a \in \prod_{i \in I} U_i \subseteq U$ .
  - ⟨2⟩3. For  $j = 1, \dots, n$ , PICK  $B_{i_j} \in \mathcal{B}_{i_j}$  such that  $a_{i_j} \in B_{i_j} \subseteq U_{i_j}$
  - ⟨2⟩4. LET:  $B = \prod_{i \in I} B_i$  where  $B_i = A_i$  for  $i \neq i_1, \dots, i_n$
  - ⟨2⟩5.  $B \in \mathcal{B}$
  - ⟨2⟩6.  $a \in B \subseteq U$
- ⟨1⟩3. Q.E.D.

PROOF: Lemma 63.

□

**Proposition 126.** *Let  $\{A_i\}_{i \in I}$  be a family of topological spaces. Then the projections  $\pi_i : \prod_{i \in I} A_i \rightarrow A_i$  are open maps.*

PROOF: From Lemma 86. □

**Example 127.** The projections are not always closed maps. For example,  $\pi_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$  maps the closed set  $\{(x, 1/x) \mid x > 0\}$  to  $(0, +\infty)$ .

**Proposition 128.** *Let  $\{X_i\}_{i \in I}$  be a family of sets. For  $i \in I$ , let  $\mathcal{T}_i$  and  $\mathcal{U}_i$  be topologies on  $X_i$ . Let  $\mathcal{P}$  be the product topology on  $\prod_{i \in I} X_i$  generated by the topologies  $\mathcal{T}_i$ , and  $\mathcal{Q}$  the product topology on the same set generated by the topologies  $\mathcal{U}_i$ . Then  $\mathcal{P} \subseteq \mathcal{Q}$  if and only if  $\mathcal{T}_i \subseteq \mathcal{U}_i$  for all  $i$ .*

PROOF:

- ⟨1⟩1. If  $\mathcal{T}_i \subseteq \mathcal{U}_i$  for all  $i$  then  $\mathcal{P} \subseteq \mathcal{Q}$ 

PROOF: By Corollary 62.1.
- ⟨1⟩2. If  $\mathcal{P} \subseteq \mathcal{Q}$  then  $\mathcal{T}_i \subseteq \mathcal{U}_i$  for all  $i$ 
  - ⟨2⟩1. ASSUME:  $\mathcal{P} \subseteq \mathcal{Q}$
  - ⟨2⟩2. LET:  $i \in I$
  - ⟨2⟩3. LET:  $U \in \mathcal{T}_i$
  - ⟨2⟩4. LET:  $U_i = U$  and  $U_j = X_j$  for  $j \neq i$
  - ⟨2⟩5.  $\prod_{i \in I} U_i \in \mathcal{P}$
  - ⟨2⟩6.  $\prod_{i \in I} U_i \in \mathcal{Q}$
  - ⟨2⟩7.  $U \in \mathcal{U}_i$

PROOF: From Proposition 126.

□

**Proposition 129 (AC).** *Let  $\{X_i\}_{i \in I}$  be a family of topological spaces. Let  $A_i \subseteq X_i$  for all  $i \in I$ . Then*

$$\prod_{i \in I} \overline{A_i} = \overline{\prod_{i \in I} A_i} .$$

PROOF:

- (1)1.  $\overline{\prod_{i \in I} A_i} \subseteq \prod_{i \in I} \overline{A_i}$   
 (2)1. For all  $i \in I$  we have  $A_i \subseteq \overline{A_i}$   
 PROOF: Lemma 45.  
 (2)2.  $\prod_{i \in I} A_i \subseteq \prod_{i \in I} \overline{A_i}$   
 (2)3. Q.E.D.  
 PROOF: Since  $\prod_{i \in I} A_i$  is closed by Proposition 124.  
 (1)2.  $\prod_{i \in I} \overline{A_i} \subseteq \overline{\prod_{i \in I} A_i}$   
 (2)1. LET:  $x \in \prod_{i \in I} \overline{A_i}$   
 (2)2. LET:  $U$  be a neighbourhood of  $x$   
 (2)3. PICK  $V_i$  open in  $X_i$  such that  $x \in \prod_{i \in I} V_i \subseteq U$  with  $V_i = X_i$  except for  
 $i = i_1, \dots, i_n$   
 (2)4. For  $i \in I$ , pick  $a_i \in V_i \cap A_i$   
 PROOF: By Theorem 48 and (2)1 using the Axiom of Choice.  
 (2)5.  $U$  intersects  $\prod_{i \in I} A_i$   
 (2)6. Q.E.D.  
 PROOF:  $a \in U \cap \prod_{i \in I} A_i$

□

**Example 130.** The closure of  $\mathbb{R}^\infty$  in  $\mathbb{R}^\omega$  is  $\mathbb{R}^\omega$

PROOF:

- (1)1. LET:  $a \in \mathbb{R}^\omega$   
 (1)2. LET:  $U$  be any neighbourhoods of  $a$ .  
 (1)3. PICK  $U_n$  open in  $\mathbb{R}$  for all  $n$  such that  $a \in \prod_{n \geq 0} U_n \subseteq U$  and  $U_n = \mathbb{R}$  for  
 all  $n$  except  $n_1, \dots, n_k$   
 (1)4. LET:  $b_n = a_n$  for  $n = n_1, \dots, n_k$  and  $b_n = 0$  for all other  $n$   
 (1)5.  $b \in \mathbb{R}^\infty \cap U$   
 (1)6. Q.E.D.

PROOF: From Theorem 48.

□

**Proposition 131.** Let  $\{X_i\}_{i \in I}$  be a family of topological spaces. Let  $(a_n)$  be a sequence in  $\prod_{i \in I} X_i$  and  $l \in \prod_{i \in I} X_i$ . Then  $a_n \rightarrow l$  as  $n \rightarrow \infty$  if and only if, for all  $i \in I$ , we have  $\pi_i(a_n) \rightarrow \pi_i(l)$  as  $n \rightarrow \infty$ .

PROOF:

- (1)1. If  $a_n \rightarrow l$  as  $n \rightarrow \infty$  then, for all  $i \in I$ , we have  $\pi_i(a_n) \rightarrow \pi_i(l)$  as  $n \rightarrow \infty$   
 PROOF: Proposition 105.  
 (1)2. If, for all  $i \in I$ , we have  $\pi_i(a_n) \rightarrow \pi_i(l)$  as  $n \rightarrow \infty$ , then  $a_n \rightarrow l$  as  $n \rightarrow \infty$   
 (2)1. ASSUME: For all  $i \in I$ , we have  $\pi_i(a_n) \rightarrow \pi_i(l)$  as  $n \rightarrow \infty$   
 (2)2. LET:  $V$  be a neighbourhood of  $l$   
 (2)3. PICK open sets  $U_i$  in  $X_i$  such that  $l \in \prod_{i \in I} U_i \subseteq V$  and  $U_i = X_i$  for all  
 $i$  except  $i = i_1, \dots, i_k$   
 (2)4. For  $j = 1, \dots, k$ , PICK  $N_j$  such that, for all  $n \geq N_j$ , we have  $\pi_{i_j}(a_n) \in$   
 $U_{i_j}$   
 (2)5. LET:  $N = \max(N_1, \dots, N_k)$   
 (2)6. For all  $n \geq N$  we have  $a_n \in V$

□

**Theorem 132.** *Let  $A$  be a topological space and  $\{X_i\}_{i \in I}$  be a family of topological spaces. Let  $f : A \rightarrow \prod_{i \in I} X_i$  be a function. If  $\pi_i \circ f$  is continuous for all  $i \in I$  then  $f$  is continuous.*

PROOF:

⟨1⟩1. LET:  $i \in I$  and  $U$  be open in  $X_i$

⟨1⟩2.  $f^{-1}(\pi_i^{-1}(U))$  is open in  $A$

⟨1⟩3. Q.E.D.

PROOF: Proposition 90.

□

## 20.1 Continuous in Each Variable Separately

**Definition 133** (Continuous in Each Variable Separately). Let  $F : X \times Y \rightarrow Z$ . Then  $F$  is *continuous in each variable separately* if and only if:

- for every  $a \in X$  the function  $\lambda y \in Y. F(a, y)$  is continuous;
- for every  $b \in Y$  the function  $\lambda x \in X. F(x, b)$  is continuous.

**Proposition 134.** *Let  $F : X \times Y \rightarrow Z$ . If  $F$  is continuous then  $F$  is continuous in each variable separately.*

PROOF: For  $a \in X$ , the function  $\lambda y \in Y. F(a, y)$  is  $F \circ i$  where  $i : Y \rightarrow X \times Y$  maps  $y$  to  $(a, y)$ . We have  $i$  is continuous by Proposition 111, hence  $F \circ i$  is continuous by Theorem 96.

Similarly for  $\lambda x \in X. F(x, b)$  for  $b \in Y$ . □

**Example 135.** Define  $F : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  by

$$F(x, y) = \begin{cases} xy/(x^2 + y^2) & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

Then  $F$  is continuous in each variable separately but not continuous.

**Proposition 136.** *Let  $f : A \rightarrow C$  and  $g : B \rightarrow D$  be open maps. Then  $f \times g : A \times B \rightarrow C \times D$  is an open map.*

PROOF: Given  $U$  open in  $A$  and  $V$  open in  $B$ . Then  $(f \times g)(U \times V) = f(U) \times g(V)$  is open in  $C \times D$ . The result follows from Proposition 87. □

## 21 The Subspace Topology

**Definition 137** (Subspace Topology). Let  $X$  be a topological space and  $Y \subseteq X$ . The *subspace topology* on  $Y$  is  $\mathcal{T} = \{U \cap Y \mid U \text{ is open in } X\}$ .

We prove this is a topology.

PROOF:

⟨1⟩1.  $Y \in \mathcal{T}$

PROOF: Since  $Y = X \cap Y$

⟨1⟩2. For all  $\mathcal{U} \subseteq \mathcal{T}$ , we have  $\bigcup \mathcal{U} \in \mathcal{T}$

⟨2⟩1. LET:  $\mathcal{U} \subseteq \mathcal{T}$

⟨2⟩2. LET:  $\mathcal{V} = \{V \text{ open in } X \mid V \cap Y \in \mathcal{U}\}$

⟨2⟩3.  $\bigcup \mathcal{U} = (\bigcup \mathcal{V}) \cap Y$

⟨1⟩3. For all  $U, V \in \mathcal{T}$ , we have  $U \cap V \in \mathcal{T}$

⟨2⟩1. LET:  $U, V \in \mathcal{T}$

⟨2⟩2. PICK  $U', V'$  open in  $X$  such that  $U = U' \cap Y$  and  $V = V' \cap Y$

⟨2⟩3.  $(U \cap V) = (U' \cap V') \cap Y$

□

**Theorem 138.** *Let  $X$  be a topological space and  $Y$  a subspace of  $X$ . Let  $A \subseteq Y$ . Then  $A$  is closed in  $Y$  if and only if there exists a closed set  $C$  in  $X$  such that  $A = C \cap Y$ .*

PROOF: We have

$$\begin{aligned} & A \text{ is closed in } Y \\ \Leftrightarrow & Y \setminus A \text{ is open in } Y \\ \Leftrightarrow & \exists U \text{ open in } X. Y \setminus A = Y \cap U \\ \Leftrightarrow & \exists C \text{ closed in } X. Y \setminus A = Y \cap (X \setminus U) \\ \Leftrightarrow & \exists C \text{ closed in } X. A = Y \cap U \end{aligned} \quad \square$$

**Theorem 139.** *Let  $Y$  be a subspace of  $X$ . Let  $A \subseteq Y$ . Let  $\bar{A}$  be the closure of  $A$  in  $X$ . Then the closure of  $A$  in  $Y$  is  $\bar{A} \cap Y$ .*

PROOF: The closure of  $A$  in  $Y$  is

$$\begin{aligned} & \bigcap \{C \text{ closed in } Y \mid A \subseteq C\} \\ &= \bigcap \{D \cap Y \mid D \text{ closed in } X, A \subseteq D \cap Y\} \quad (\text{Theorem 138}) \\ &= \bigcap \{D \mid D \text{ closed in } X, A \subseteq D\} \cap Y \\ &= \bar{A} \cap Y \end{aligned} \quad \square$$

**Lemma 140.** *Let  $X$  be a topological space and  $Y \subseteq X$ . Let  $\mathcal{B}$  be a basis for the topology on  $X$ . Then  $\mathcal{B}' = \{B \cap Y \mid B \in \mathcal{B}\}$  is a basis for the subspace topology on  $Y$ .*

PROOF:

⟨1⟩1. Every element in  $\mathcal{B}'$  is open in  $Y$

⟨1⟩2. For every open set  $U$  in  $Y$  and point  $y \in U$ , there exists  $B' \in \mathcal{B}'$  such that  $y \in B' \subseteq U$

⟨2⟩1. LET:  $U$  be open in  $Y$  and  $y \in U$

⟨2⟩2. PICK  $V$  open in  $X$  such that  $U = V \cap Y$

⟨2⟩3. PICK  $B \in \mathcal{B}$  such that  $y \in B \subseteq V$

⟨2⟩4. LET:  $B' = B \cap Y$

⟨2⟩5.  $B' \in \mathcal{B}'$

⟨2⟩6.  $y \in B' \subseteq U$

⟨1⟩3. Q.E.D.

PROOF: By Lemma 63.

□

**Lemma 141.** *Let  $X$  be a topological space and  $Y \subseteq X$ . Let  $\mathcal{S}$  be a basis for the topology on  $X$ . Then  $\mathcal{S}' = \{S \cap Y \mid S \in \mathcal{S}\}$  is a subbasis for the subspace topology on  $Y$ .*

PROOF: The set  $\{B \cap Y \mid B \text{ is a finite intersection of elements of } \mathcal{S}\}$  is a basis for the subspace topology by Lemma 140, and this is the set of all finite intersections of elements of  $\mathcal{S}'$ . □

**Lemma 142.** *Let  $Y$  be a subspace of  $X$ . If  $U$  is open in  $Y$  and  $Y$  is open in  $X$  then  $U$  is open in  $X$ .*

PROOF:

⟨1⟩1. PICK  $V$  open in  $X$  such that  $U = V \cap Y$

⟨1⟩2.  $U$  is open in  $X$

PROOF: Since it is the intersection of two open sets  $V$  and  $Y$ .

□

**Theorem 143.** *Let  $Y$  be a subspace of  $X$  and  $A \subseteq Y$ . If  $A$  is closed in  $Y$  and  $Y$  is closed in  $X$  then  $A$  is closed in  $X$ .*

PROOF: Pick a closed set  $C$  in  $X$  such that  $A = C \cap Y$  (Theorem 138). Then  $A$  is the intersection of two sets closed in  $X$ , hence  $A$  is closed in  $X$  (Lemma 33).

□

**Theorem 144.** *Let  $\{X_i\}_{i \in I}$  be a family of topological spaces. Let  $A_i$  be a subspace of  $X_i$  for all  $i \in I$ . Then the product topology on  $\prod_{i \in I} A_i$  is the same as the topology it inherits as a subspace of  $\prod_{i \in I} X_i$ .*

PROOF: The product topology is generated by the subbasis

$$\begin{aligned} & \{\pi_i^{-1}(U) \mid i \in I, U \text{ open in } A_i\} \\ &= \{\pi_i^{-1}(V) \cap A_i \mid i \in I, V \text{ open in } X_i\} \\ &= \{\pi_i^{-1}(V) \mid i \in I, V \text{ open in } X_i\} \cap \prod_{i \in I} A_i \end{aligned}$$

and this is a subbasis for the subspace topology by Lemma 141. □

**Theorem 145.** *Let  $X$  be an ordered set in the order topology. Let  $Y \subseteq X$  be an interval. Then the order topology on  $Y$  is the same as the subspace topology on  $Y$ .*

PROOF:

⟨1⟩1. The order topology is finer than the subspace topology.

⟨2⟩1. For every open ray  $R$  in  $X$ , the set  $R \cap Y$  is open in the order topology.

⟨3⟩1. For all  $a \in X$ , we have  $(-\infty, a) \cap Y$  is open in the order topology.

⟨4⟩1. CASE: For all  $y \in Y$  we have  $y < a$   
 PROOF: In this case  $(-\infty, a) \cap Y = Y$ .  
 ⟨4⟩2. CASE: For all  $y \in Y$  we have  $a < y$   
 PROOF: In this case  $(-\infty, a) \cap Y = \emptyset$ .  
 ⟨4⟩3. CASE: There exists  $y \in Y$  such that  $y \leq a$  and  $y \in Y$  such that  
 $a \leq y$   
 ⟨5⟩1.  $a \in Y$   
 PROOF: Because  $Y$  is an interval.  
 ⟨5⟩2.  $(-\infty, a) \cap Y = \{y \in Y \mid y < a\}$   
 ⟨3⟩2. For all  $a \in X$ , we have  $(a, +\infty) \cap Y$  is open in the order topology.  
 PROOF: Similar.  
 ⟨2⟩2. Q.E.D.  
 PROOF: By Lemmas 113 and 141 and Proposition 71.  
 ⟨1⟩2. The subspace topology is finer than the order topology.  
 ⟨2⟩1. Every open ray in  $Y$  is open in the subspace topology.  
 PROOF: For any  $a \in Y$  we have  $(-\infty, a)_Y = (-\infty, a)_X \cap Y$  and  $(a, +\infty)_Y = (a, +\infty)_X \cap Y$ .  
 ⟨2⟩2. Q.E.D.  
 PROOF: By Lemma 113 and Proposition 71  
 □

This example shows that we cannot remove the hypothesis that  $Y$  is an interval:

**Example 146.** The order topology on  $I_o^2$  is different from the subspace topology as a subspace of  $\mathbb{R}^2$  under the dictionary order topology.

PROOF: The set  $\{1/2\} \times (1/2, 1)$  is open in the subspace topology but not in the order topology. □

**Proposition 147.** Let  $X$  be a topological space,  $Y$  a subspace of  $X$ , and  $Z$  a subspace of  $Y$ . Then the subspace topology on  $Z$  inherited from  $X$  is the same as the subspace topology on  $Z$  inherited from  $Y$ .

PROOF: The subspace topology inherited from  $Y$  is

$$\begin{aligned}
 & \{V \cap Z \mid V \text{ open in } Y\} \\
 &= \{U \cap Y \cap Z \mid U \text{ open in } X\} \\
 &= \{U \cap Z \mid U \text{ open in } X\}
 \end{aligned}$$

which is the subspace topology inherited from  $X$ . □

**Definition 148** (Unit Circle). The *unit circle*  $S^1$  is  $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$  as a subspace of  $\mathbb{R}^2$ .

**Definition 149** (Unit 2-sphere). The *unit 2-sphere* is  $S^2 = \{(x, y, z) \mid x^2 + y^2 + z^2 \leq 1\}$  as a subspace of  $\mathbb{R}^3$ .

**Proposition 150.** Let  $f : X \rightarrow Y$  be an open map and  $A \subseteq X$  be open. Then the restriction  $f \upharpoonright A : A \rightarrow f(A)$  is an open map.

PROOF:

$\langle 1 \rangle 1$ . LET:  $U$  be open in  $A$

$\langle 1 \rangle 2$ .  $U$  is open in  $X$

PROOF: Lemma 142.

$\langle 1 \rangle 3$ .  $f(U)$  is open in  $Y$

$\langle 1 \rangle 4$ .  $f(U)$  is open in  $f(A)$

PROOF: Since  $f(U) = f(U) \cap f(A)$ .

□

**Example 151.** This example shows that we cannot remove the hypothesis that  $A$  is open.

Let  $A = \{(x, y) \in \mathbb{R}^2 \mid (x > 0 \text{ and } y = 1/x) \text{ or } x = y = 0\}$ . Then  $\pi_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$  is an open map, but  $\pi_1 \upharpoonright A : A \rightarrow [0, +\infty)$  is not, because it maps the set  $\{0, 0\}$  which is open in  $A$  to  $\{0\}$  which is not open in  $[0, +\infty)$ .

**Proposition 152.** *Let  $Y$  be a subspace of  $X$ . Let  $A \subseteq Y$  and  $l \in Y$ . Then  $l$  is a limit point of  $A$  in  $Y$  if and only if  $l$  is a limit point of  $A$  in  $X$ .*

PROOF: Both are equivalent to the condition that any neighbourhood of  $l$  in  $X$  intersects  $A$  in a point other than  $l$ . □

## 22 The Box Topology

**Definition 153** (Box Topology). Let  $\{A_i\}_{i \in I}$  be a family of topological spaces. The *box topology* on  $\prod_{i \in I} A_i$  is the topology generated by the set of all sets of the form  $\prod_{i \in I} U_i$  where  $\{U_i\}_{i \in I}$  is a family such that each  $U_i$  is open in  $A_i$ .

This is a basis since it covers  $\prod_{i \in I} A_i$  and is closed under intersection.

**Proposition 154.** *The box topology is finer than the product topology.*

PROOF: From Proposition 123. □

**Corollary 154.1.** *If  $A_i$  is closed in  $X_i$  for all  $i \in I$  then  $\prod_{i \in I} A_i$  is closed in  $\prod_{i \in I} X_i$  under the box topology.*

PROOF: From Proposition 124.

**Proposition 155** (AC). *Let  $\{A_i\}_{i \in I}$  be a family of topological spaces. For  $i \in I$ , let  $\mathcal{B}_i$  be a basis for the topology on  $A_i$ . Then  $\mathcal{B} = \{\prod_{i \in I} B_i \mid \forall i \in I, B_i \in \mathcal{B}_i\}$  is a basis for the box topology on  $\prod_{i \in I} A_i$ .*

PROOF:

$\langle 1 \rangle 1$ . Every set of the form  $\prod_{i \in I} B_i$  is open.

$\langle 1 \rangle 2$ . For every point  $a \in \prod_{i \in I} A_i$  and every open set  $U$  with  $a \in U$ , there exists  $B \in \mathcal{B}$  such that  $a \in B \subseteq U$ .

$\langle 2 \rangle 1$ . LET:  $U$  be open and  $a \in U$

$\langle 2 \rangle 2$ . PICK a family  $\{U_i\}_{i \in I}$  such that each  $U_i$  is open in  $A_i$  and  $a \in \prod_{i \in I} U_i \subseteq U$ .



⟨2⟩3. For  $i \in I$ , PICK  $B_i \in \mathcal{B}_i$  such that  $a_i \in B_i \subseteq U_i$

PROOF: Using the Axiom of Choice.

⟨2⟩4.  $a \in \prod_{i \in I} B_i \subseteq U$

⟨1⟩3. Q.E.D.

PROOF: Lemma 63.

□

**Theorem 156.** Let  $\{X_i\}_{i \in I}$  be a family of topological spaces. Let  $A_i$  be a subspace of  $X_i$  for all  $i \in I$ . Give  $\prod_{i \in I} X_i$  the box topology. Then the box topology on  $\prod_{i \in I} A_i$  is the same as the topology it inherits as a subspace of  $\prod_{i \in I} X_i$ .

PROOF: The box topology is generated by the basis

$$\begin{aligned} & \left\{ \prod_{i \in I} U_i \mid \forall i \in I, U_i \text{ open in } A_i \right\} \\ &= \left\{ \prod_{i \in I} (V_i \cap A_i) \mid \forall i \in I, V_i \text{ open in } X_i \right\} \\ &= \left\{ \prod_{i \in I} V_i \mid \forall i \in I, V_i \text{ open in } X_i \right\} \cap \prod_{i \in I} A_i \end{aligned}$$

and this is a basis for the subspace topology by Lemma 140. □

**Proposition 157 (AC).** Let  $\{X_i\}_{i \in I}$  be a family of topological spaces. Give  $\prod_{i \in I} X_i$  the box topology. Let  $A_i \subseteq X_i$  for all  $i \in I$ . Then

$$\prod_{i \in I} \overline{A_i} = \overline{\prod_{i \in I} A_i}.$$

PROOF:

⟨1⟩1.  $\overline{\prod_{i \in I} A_i} \subseteq \prod_{i \in I} \overline{A_i}$

⟨2⟩1. For all  $i \in I$  we have  $A_i \subseteq \overline{A_i}$

PROOF: Lemma 45.

⟨2⟩2.  $\prod_{i \in I} A_i \subseteq \prod_{i \in I} \overline{A_i}$

⟨2⟩3. Q.E.D.

PROOF: Since  $\prod_{i \in I} A_i$  is closed by Corollary 154.1.

⟨1⟩2.  $\prod_{i \in I} \overline{A_i} \subseteq \overline{\prod_{i \in I} A_i}$

⟨2⟩1. LET:  $x \in \prod_{i \in I} \overline{A_i}$

⟨2⟩2. LET:  $U$  be a neighbourhood of  $x$

⟨2⟩3. PICK  $V_i$  open in  $X_i$  such that  $x \in \prod_{i \in I} V_i \subseteq U$

⟨2⟩4. For  $i \in I$ , pick  $a_i \in V_i \cap A_i$

PROOF: By Theorem 48 and ⟨2⟩1 using the Axiom of Choice.

⟨2⟩5.  $U$  intersects  $\prod_{i \in I} A_i$

⟨2⟩6. Q.E.D.

PROOF:  $a \in U \cap \prod_{i \in I} A_i$ .

□

The following example shows that Theorem 132 fails in the box topology.

**Example 158.** Define  $f : \mathbb{R} \rightarrow \mathbb{R}^\omega$  by  $f(t) = (t, t, \dots)$ . Then  $\pi_n \circ f = \text{id}_{\mathbb{R}}$  is continuous for all  $n$ . But  $f$  is not continuous when  $\mathbb{R}^\omega$  is given the box topology because the inverse image of

$$(-1, 1) \times (-1/2, 1/2) \times (-1/3, 1/3) \times \dots$$

is  $\{0\}$  which is not open.

The following example shows that Proposition 131 fails in the box topology.

**Example 159.** Give  $\mathbb{R}^\omega$  the box topology. Let  $a_n = (1/n, 1/n, \dots)$  for  $n \geq 1$  and  $l = (0, 0, \dots)$ . Then  $\pi_i(a_n) \rightarrow \pi_i(l)$  as  $n \rightarrow \infty$  for all  $i$ , but  $a_n \not\rightarrow l$  as  $n \rightarrow \infty$  since the open set

$$(-1, 1) \times (-1/2, 1/2) \times (-1/3, 1/3) \times \dots$$

contains  $l$  but does not contain any  $a_n$ .

**Example 160.** The set  $\mathbb{R}^\infty$  is closed in  $\mathbb{R}^\omega$  under the box topology. For let  $(a_n)$  be any sequence not in  $\mathbb{R}^\infty$ . Let  $U_n$  be an open interval around  $a_n$  that does not contain 0 if  $a_n \neq 0$ , and  $U_n = \mathbb{R}$  if  $a_n = 0$ . Then  $\prod_{n \geq 0} U_n$  is a neighbourhood of  $(a_n)$  that does not intersect  $\mathbb{R}^\infty$ .

## 23 $T_1$ Spaces

**Definition 161** ( $T_1$  Space). A topological space is  $T_1$  if and only if every singleton is closed.

**Lemma 162.** A space is  $T_1$  if and only if every finite set is closed.

PROOF: From Lemma 34.  $\square$

**Theorem 163.** In a  $T_1$  space, a point  $a$  is a limit point of a set  $A$  if and only if every neighbourhood of  $a$  contains infinitely many points of  $A$ .

PROOF:

$\langle 1 \rangle 1$ . If  $a$  is a limit point of  $A$  then every neighbourhood of  $a$  contains infinitely many points of  $A$ .

$\langle 2 \rangle 1$ . ASSUME:  $a$  is a limit point of  $A$ .

$\langle 2 \rangle 2$ . LET:  $U$  be a neighbourhood of  $a$ .

$\langle 2 \rangle 3$ . ASSUME: for a contradiction  $U$  contains only finitely many points of  $A$ .

$\langle 2 \rangle 4$ .  $(U \cap A) \setminus \{a\}$  is closed.

PROOF: By the  $T_1$  axiom.

$\langle 2 \rangle 5$ .  $(U \setminus A) \cup \{a\}$  is open.

PROOF: It is  $U \setminus ((U \cap A) \setminus \{a\})$ .

$\langle 2 \rangle 6$ .  $(U \setminus A) \cup \{a\}$  intersects  $A$  in a point other than  $a$ .

PROOF: From  $\langle 2 \rangle 1$ .

$\langle 2 \rangle 7$ . Q.E.D.

$\square$

⟨1⟩2. If every neighbourhood of  $a$  contains infinitely many points of  $A$  then  $a$  is a limit point of  $A$ .

PROOF: Immediate from definitions.

□

(To see this does not hold in every space, see Proposition 59.)

**Proposition 164.** *A space is  $T_1$  if and only if, for any two distinct points  $x$  and  $y$ , there exist neighbourhoods  $U$  of  $x$  and  $V$  of  $y$  such that  $x \notin V$  and  $y \notin U$ .*

PROOF:

⟨1⟩1. LET:  $X$  be a topological space.

⟨1⟩2. If  $X$  is  $T_1$  then, for any two distinct points  $x$  and  $y$ , there exist neighbourhoods  $U$  of  $x$  and  $V$  of  $y$  such that  $x \notin V$  and  $y \notin U$ .

PROOF: This holds because  $\{x\}$  and  $\{y\}$  are closed.

⟨1⟩3. Suppose, for any two distinct points  $x$  and  $y$ , there exist neighbourhoods  $U$  of  $x$  and  $V$  of  $y$  such that  $x \notin V$  and  $y \notin U$ . Then  $X$  is  $T_1$ .

⟨2⟩1. ASSUME: For any two distinct points  $x$  and  $y$ , there exist neighbourhoods  $U$  of  $x$  and  $V$  of  $y$  such that  $x \notin V$  and  $y \notin U$ .

⟨2⟩2. LET:  $a \in X$

⟨2⟩3.  $\{a\}$  is closed.

PROOF: For all  $b \neq a$  there exists a neighbourhood  $U$  of  $b$  such that  $U \subseteq X \setminus \{a\}$ .

□

**Proposition 165.** *A subspace of a  $T_1$  space is  $T_1$ .*

PROOF: From Proposition 143.

## 24 Hausdorff Spaces

**Definition 166** (Hausdorff Space). A topological space is *Hausdorff* if and only if, for any points  $x, y$  with  $x \neq y$ , there exist disjoint open sets  $U$  and  $V$  such that  $x \in U$  and  $y \in V$ .

**Theorem 167.** *Every Hausdorff space is  $T_1$ .*

PROOF:

⟨1⟩1. LET:  $X$  be a Hausdorff space.

⟨1⟩2. LET:  $b \in X$

PROVE:  $\overline{\{b\}} = \{b\}$

⟨1⟩3. ASSUME:  $a \in \overline{\{b\}}$  and  $a \neq b$

⟨1⟩4. PICK disjoint neighbourhoods  $U$  of  $a$  and  $V$  of  $b$ .

⟨1⟩5.  $U$  intersects  $\{b\}$

PROOF: Theorem 48.

⟨1⟩6.  $b \in U$

⟨1⟩7. Q.E.D.

PROOF: This contradicts the fact that  $U$  and  $V$  are disjoint (⟨1⟩4).

□

**Proposition 168.** *An infinite set under the finite complement topology is  $T_1$  but not Hausdorff.*

PROOF:

⟨1⟩1. LET:  $X$  be an infinite set under the finite complement topology.

⟨1⟩2. Every singleton is closed.

PROOF: By definition.

⟨1⟩3. PICK  $a, b \in X$  with  $a \neq b$

⟨1⟩4. There are no disjoint neighbourhoods  $U$  of  $a$  and  $V$  of  $b$ .

⟨2⟩1. LET:  $U$  be a neighbourhood of  $a$  and  $V$  a neighbourhood of  $b$ .

⟨2⟩2.  $X \setminus U$  and  $X \setminus V$  are finite.

⟨2⟩3. PICK  $c \in X$  that is not in  $X \setminus U$  or  $X \setminus V$ .

⟨2⟩4.  $c \in U \cap V$

□

**Proposition 169.** *The product of a family of Hausdorff spaces is Hausdorff.*

PROOF:

⟨1⟩1. LET:  $\{X_i\}_{i \in I}$  be a family of Hausdorff spaces.

⟨1⟩2. LET:  $a, b \in \prod_{i \in I} X_i$  with  $a \neq b$

⟨1⟩3. PICK  $i \in I$  such that  $a_i \neq b_i$

⟨1⟩4. PICK  $U, V$  disjoint open sets in  $X_i$  with  $a_i \in U$  and  $b_i \in V$

⟨1⟩5.  $\pi_i^{-1}(U)$  and  $\pi_i^{-1}(V)$  are disjoint open sets in  $\prod_{i \in I} X_i$  with  $a \in \pi_i^{-1}(U)$  and  $b \in \pi_i^{-1}(V)$

□

**Theorem 170.** *Every linearly ordered set under the order topology is Hausdorff.*

PROOF:

⟨1⟩1. LET:  $X$  be a linearly ordered set under the order topology.

⟨1⟩2. LET:  $a, b \in X$  with  $a \neq b$

⟨1⟩3. ASSUME: w.l.o.g.  $a < b$

⟨1⟩4. CASE: There exists  $c$  such that  $a < c < b$

PROOF: The sets  $(-\infty, c)$  and  $(c, +\infty)$  are disjoint neighbourhoods of  $a$  and  $b$  respectively.

⟨1⟩5. CASE: There is no  $c$  such that  $a < c < b$

PROOF: The sets  $(-\infty, b)$  and  $(a, +\infty)$  are disjoint neighbourhoods of  $a$  and  $b$  respectively.

□

**Theorem 171.** *A subspace of a Hausdorff space is Hausdorff.*

PROOF:

⟨1⟩1. LET:  $X$  be a Hausdorff space and  $Y$  a subspace of  $X$ .

⟨1⟩2. LET:  $x, y \in Y$  with  $x \neq y$

⟨1⟩3. PICK disjoint neighbourhoods  $U$  of  $x$  and  $V$  of  $y$  in  $X$ .

⟨1⟩4.  $U \cap Y$  and  $V \cap Y$  are disjoint neighbourhoods of  $x$  and  $y$  respectively in  $Y$ .

□

**Proposition 172.** *A space  $X$  is Hausdorff if and only if the diagonal  $\Delta = \{(x, x) \mid x \in X\}$  is closed in  $X^2$ .*

PROOF:

$X$  is Hausdorff

$$\Leftrightarrow \forall x, y \in X. x \neq y \Rightarrow \exists V, W \text{ open. } x \in V \wedge y \in W \wedge V \cap W = \emptyset$$

$$\Leftrightarrow \forall (x, y) \in X^2 \setminus \Delta. \exists V, W \text{ open. } (x, y) \subseteq V \times W \subseteq X^2 \setminus \Delta$$

$$\Leftrightarrow \Delta \text{ is closed}$$

□

**Theorem 173.** *In a Hausdorff space, a sequence has at most one limit.*

PROOF:

⟨1⟩1. LET:  $X$  be a Hausdorff space.

⟨1⟩2. ASSUME: for a contradiction  $a_n \rightarrow l$  as  $n \rightarrow \infty$ ,  $a_n \rightarrow m$  as  $n \rightarrow \infty$ , and  $l \neq m$

⟨1⟩3. PICK disjoint neighbourhoods  $U$  of  $l$  and  $V$  of  $m$

PROOF: By the Hausdorff axiom.

⟨1⟩4. PICK  $M$  and  $N$  such that  $a_n \in U$  for  $n \geq M$  and  $a_n \in V$  for  $n \geq N$

⟨1⟩5.  $a_{\max(M, N)} \in U \cap V$

⟨1⟩6. Q.E.D.

PROOF: This contradicts the fact that  $U$  and  $V$  are disjoint (⟨1⟩3).

□

To see this is not always true in spaces that are  $T_1$  but not Hausdorff:

**Proposition 174.** *Let  $X$  be an infinite set under the finite complement topology. Let  $(a_n)_{n \in \mathbb{N}}$  be a sequence with all points distinct. Then for every  $l \in X$  we have  $a_n \rightarrow l$  as  $n \rightarrow \infty$ .*

PROOF: Let  $U$  be any neighbourhood of  $l$ . Since  $X \setminus U$  is finite, there must exist  $N$  such that, for all  $n \geq N$ , we have  $a_n \in U$ . □

**Proposition 175.** *Let  $X$  be a topological space. Let  $Y$  a Hausdorff space. Let  $A \subseteq X$ . Let  $f, g : \overline{A} \rightarrow Y$  be continuous. If  $f$  and  $g$  agree on  $A$  then  $f = g$ .*

PROOF:

⟨1⟩1. LET:  $x \in \overline{A}$

⟨1⟩2. ASSUME:  $f(x) \neq g(x)$

⟨1⟩3. PICK disjoint neighbourhoods  $V$  of  $f(x)$  and  $W$  of  $g(x)$ .

⟨1⟩4. PICK  $y \in f^{-1}(V) \cap g^{-1}(W) \cap A$

PROOF: Since  $f^{-1}(V) \cap g^{-1}(W)$  is a neighbourhood of  $x$  and hence intersects  $A$ .

⟨1⟩5.  $f(y) = g(y) \in V \cap W$

⟨1⟩6. Q.E.D.

PROOF: This contradicts the fact that  $V$  and  $W$  are disjoint ( $\langle 1 \rangle 3$ ).

□

**Proposition 176.** *Let  $\{X_i\}_{i \in I}$  be a family of Hausdorff spaces. Then  $\prod_{i \in I} X_i$  under the box topology is Hausdorff.*

PROOF:

- $\langle 1 \rangle 1$ . LET:  $\{X_i\}_{i \in I}$  be a family of Hausdorff spaces.
- $\langle 1 \rangle 2$ . LET:  $a, b \in \prod_{i \in I} X_i$  with  $a \neq b$
- $\langle 1 \rangle 3$ . PICK  $i \in I$  such that  $a_i \neq b_i$
- $\langle 1 \rangle 4$ . PICK  $U, V$  disjoint open sets in  $X_i$  with  $a_i \in U$  and  $b_i \in V$
- $\langle 1 \rangle 5$ .  $\pi_i^{-1}(U)$  and  $\pi_i^{-1}(V)$  are disjoint open sets in  $\prod_{i \in I} X_i$  with  $a \in \pi_i^{-1}(U)$  and  $b \in \pi_i^{-1}(V)$

□

**Proposition 177.** *Let  $\mathcal{T}$  and  $\mathcal{T}'$  be two topologies on the same set  $X$  with  $\mathcal{T} \subseteq \mathcal{T}'$ . If  $\mathcal{T}$  is Hausdorff then  $\mathcal{T}'$  is Hausdorff.*

PROOF: Immediate from definitions.

## 25 The First Countability Axiom

**Definition 178** (First Countability Axiom). A topological space  $X$  satisfies the *first countability axiom*, or is *first countable*, if and only if every point has a countable local basis.

**Lemma 179** (Sequence Lemma (CC)). *Let  $X$  be a first countable space. Let  $A \subseteq X$  and  $l \in \bar{A}$ . Then there exists a sequence in  $A$  that converges to  $l$ .*

PROOF:

- $\langle 1 \rangle 1$ . PICK a countable local basis  $\{B_n \mid n \in \mathbb{Z}^+\}$  at  $l$  such that  $B_1 \supseteq B_2 \supseteq \dots$ .

PROOF: Lemma 73.

- $\langle 1 \rangle 2$ . For all  $n \geq 1$ , PICK  $a_n \in A \cap B_n$ .

PROVE:  $a_n \rightarrow l$  as  $n \rightarrow \infty$

- $\langle 1 \rangle 3$ . LET:  $U$  be a neighbourhood of  $l$

- $\langle 1 \rangle 4$ . PICK  $N$  such that  $B_N \subseteq U$

- $\langle 1 \rangle 5$ . For  $n \geq N$  we have  $a_n \in U$

PROOF:  $a_n \in B_n \subseteq B_N \subseteq U$

□

**Theorem 180** (CC). *Let  $X$  be a first countable space and  $Y$  a topological space. Let  $f : X \rightarrow Y$ . Suppose that, for every sequence  $(x_n)$  in  $X$  and  $l \in X$ , if  $x_n \rightarrow l$  as  $n \rightarrow \infty$ , then  $f(x_n) \rightarrow f(l)$  as  $n \rightarrow \infty$ . Then  $f$  is continuous.*

PROOF:

- $\langle 1 \rangle 1$ . LET:  $A \subseteq X$

- $\langle 1 \rangle 2$ . LET:  $a \in A$

PROVE:  $f(a) \in \overline{f(A)}$

⟨1⟩3. PICK a sequence  $(x_n)$  in  $A$  that converges to  $a$ .

PROOF: By the Sequence Lemma.

⟨1⟩4.  $f(x_n) \rightarrow f(a)$

⟨1⟩5.  $f(a) \in f(A)$

PROOF: By Lemma 75.

⟨1⟩6. Q.E.D.

PROOF: By Theorem 93.

□

**Example 181 (CC).** The space  $\mathbb{R}^\omega$  under the box product is not first countable.

PROOF: Let  $\{B_n \mid n \geq 0\}$  be a countable set of neighbourhoods of  $\vec{0}$ . We will construct a neighbourhood of  $\vec{0}$  that does not include any of these.

For  $n \geq 0$ , pick a neighbourhood  $U_n$  of 0 such that  $U_n \subset \pi_n(B_p)$ . Then  $\prod_{n=0}^\infty U_n$  is a neighbourhood of  $\vec{0}$  that does not include any  $B_n$ . □

**Example 182.** If  $J$  is an uncountable set then  $\mathbb{R}^J$  is not first countable.

PROOF:

⟨1⟩1. LET:  $\{B_n \mid n \geq 0\}$  be any countable set of neighbourhoods of  $\vec{0}$ .

⟨1⟩2. For  $n \geq 0$ , PICK a basis element  $\prod_{\alpha \in J} U_{n\alpha}$  that contains  $\vec{0}$  and is included in  $B_n$ .

PROOF: Using the Axiom of Countable Choice.

⟨1⟩3. For  $n \geq 0$ ,

LET:  $J_n = \{\alpha \in J \mid U_{n\alpha} \neq \mathbb{R}\}$

⟨1⟩4. PICK  $\beta \in J$  such that  $\beta \notin J_n$  for any  $n$ .

PROOF: Since each  $J_n$  is finite so  $\bigcup_n J_n$  is countable.

⟨1⟩5.  $\pi_\beta((-1, 1))$  is a neighbourhood of  $\vec{0}$  that does not include any  $B_n$ .

□

**Example 183.** The space  $\mathbb{R}_l$  is first countable.

PROOF: For any  $a \in \mathbb{R}$ , the set  $\{[a, a + 1/n) \mid n \geq 1\}$  is a countable local basis.

**Example 184.** The ordered square is first countable.

PROOF: For any  $(a, b) \in I_o^2$  with  $b \neq 0, 1$ , the set  $\{(\{a\} \times (b - 1/n, b + 1/n)) \cap I_o^2 \mid n \geq 1\}$  is a countable local basis.

## 26 Strong Continuity

**Definition 185** (Strongly Continuous). Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$  be a function. Then  $f$  is *strongly continuous* if and only if, for every subset  $U \subseteq Y$ , we have  $U$  is open in  $Y$  if and only if  $f^{-1}(U)$  is open in  $X$ .

**Proposition 186.** Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$  be a function. Then  $f$  is strongly continuous if and only if, for every subset  $C \subseteq Y$ , we have  $C$  is closed in  $Y$  if and only if  $f^{-1}(C)$  is closed in  $X$ .

PROOF: Since  $X \setminus f^{-1}(C) = f^{-1}(Y \setminus C)$ .  $\square$

**Proposition 187.** *Let  $X, Y$  and  $Z$  be topological spaces. Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ . If  $f$  and  $g$  are strongly continuous then so is  $g \circ f$ .*

PROOF: Since  $(g \circ f)^{-1}(U) = f^{-1}(g^{-1}(U))$ .  $\square$

**Proposition 188.** *Let  $X, Y$  and  $Z$  be topological spaces. Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ . If  $g \circ f$  is continuous and  $f$  is strongly continuous then  $g$  is continuous.*

PROOF:

$\langle 1 \rangle 1$ . LET:  $V \subseteq Z$  be open.

$\langle 1 \rangle 2$ .  $f^{-1}(g^{-1}(V))$  is open in  $X$ .

PROOF: Since  $g \circ f$  is continuous.

$\langle 1 \rangle 3$ .  $f^{-1}(V)$  is open in  $Y$ .

PROOF: Since  $g$  is strongly continuous.

$\square$

**Proposition 189.** *Let  $X, Y$  and  $Z$  be topological spaces. Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ . If  $g \circ f$  is strongly continuous and  $f$  is strongly continuous then  $g$  is strongly continuous.*

PROOF: For  $V \subseteq Z$ , we have  $V$  is open iff  $f^{-1}(g^{-1}(V))$  is open iff  $g^{-1}(V)$  is open.

## 27 Saturated Sets

**Definition 190.** Let  $X$  and  $Y$  be sets and  $p : X \rightarrow Y$  a surjective function. Let  $C \subseteq X$ . Then  $C$  is *saturated* with respect to  $p$  if and only if, for all  $x, y \in X$ , if  $x \in C$  and  $p(x) = p(y)$  then  $y \in C$ .

**Proposition 191.** *Let  $X$  and  $Y$  be sets and  $p : X \rightarrow Y$  a surjective function. Let  $C \subseteq X$ . Then the following are equivalent:*

1.  $C$  is saturated with respect to  $p$ .
2. There exists  $D \subseteq Y$  such that  $C = p^{-1}(D)$
3.  $C = p^{-1}(p(C))$ .

PROOF:

$\langle 1 \rangle 1$ .  $1 \Rightarrow 3$

$\langle 2 \rangle 1$ . ASSUME:  $C$  is saturated with respect to  $p$ .

$\langle 2 \rangle 2$ .  $C \subseteq p^{-1}(p(C))$

PROOF: Trivial.

$\langle 2 \rangle 3$ .  $p^{-1}(p(C)) \subseteq C$

$\langle 3 \rangle 1$ . LET:  $x \in p^{-1}(p(C))$

$\langle 3 \rangle 2$ .  $p(x) \in p(C)$



$\langle 3 \rangle 3$ . There exists  $y \in C$  such that  $p(x) = p(y)$

$\langle 3 \rangle 4$ .  $x \in C$

PROOF: From  $\langle 2 \rangle 1$ .

$\langle 1 \rangle 2$ .  $3 \Rightarrow 2$

PROOF: Trivial.

$\langle 1 \rangle 3$ .  $2 \Rightarrow 1$

PROOF: This follows because if  $p(x) \in D$  and  $p(x) = p(y)$  then  $p(y) \in D$ .

□

## 28 Quotient Maps

**Definition 192** (Quotient Map). Let  $X$  and  $Y$  be topological spaces and  $p : X \rightarrow Y$ . Then  $p$  is a *quotient map* if and only if  $p$  is surjective and strongly continuous.

**Proposition 193.** Let  $X$  and  $Y$  be topological spaces and  $p : X \rightarrow Y$  be a surjective function. Then the following are equivalent.

1.  $p$  is a quotient map.
2.  $p$  is continuous and maps saturated open sets to open sets.
3.  $p$  is continuous and maps saturated closed sets to closed sets.

PROOF:

$\langle 1 \rangle 1$ .  $1 \Rightarrow 2$

$\langle 2 \rangle 1$ . ASSUME:  $p$  is a quotient map.

$\langle 2 \rangle 2$ . LET:  $U$  be a saturated open set in  $X$ .

$\langle 2 \rangle 3$ .  $p^{-1}(p(U))$  is open in  $X$ .

PROOF: Since  $U = p^{-1}(p(U))$  by Proposition 191.

$\langle 2 \rangle 4$ .  $p(U)$  is open in  $Y$ .

PROOF: From  $\langle 2 \rangle 1$ .

$\langle 1 \rangle 2$ .  $1 \Rightarrow 3$

PROOF: Similar.

$\langle 1 \rangle 3$ .  $2 \Rightarrow 1$

$\langle 2 \rangle 1$ . ASSUME:  $p$  is continuous and maps saturated open sets to open sets.

$\langle 2 \rangle 2$ . LET:  $U \subseteq Y$

$\langle 2 \rangle 3$ . ASSUME:  $p^{-1}(U)$  is open in  $X$

$\langle 2 \rangle 4$ .  $p^{-1}(U)$  is saturated.

PROOF: Proposition 191.

$\langle 2 \rangle 5$ .  $U$  is open in  $Y$ .

$\langle 1 \rangle 4$ .  $3 \Rightarrow 1$

PROOF: Similar.

□

**Corollary 193.1.** Every surjective continuous open map is a quotient map.

**Corollary 193.2.** Every surjective continuous closed map is a quotient map.

**Example 194.** The converses of these corollaries do not hold.

Let  $A = \{(x, y) \mid x \geq 0\} \cup \{(x, y) \mid y = 0\}$ . Then  $\pi_1 : A \rightarrow \mathbb{R}$  is a quotient map, but not an open map or a closed map.

We prove that  $\pi_1$  maps saturated open sets to open sets:

- $\langle 1 \rangle 1$ . LET:  $\pi_1^{-1}(U)$  be a saturated open set in  $A$   
PROVE:  $U$  is open in  $\mathbb{R}$
- $\langle 1 \rangle 2$ . LET:  $x \in U$
- $\langle 1 \rangle 3$ .  $(x, 0) \in \pi_1(U)^{-1}$
- $\langle 1 \rangle 4$ . PICK  $W, V$  open in  $\mathbb{R}$  such that  $(x, 0) \subseteq W \times V \subseteq \pi_1(U)^{-1}$
- $\langle 1 \rangle 5$ .  $x \in W \subseteq U$

It is not an open map because it maps  $((-1, 1) \times (1, 2)) \cap A$  to  $[0, 1)$ .

It is not a closed map because it maps  $\{(x, 1/x) \mid x > 0\}$  to  $(0, +\infty)$ .

**Proposition 195.** Let  $p : X \twoheadrightarrow Y$  be a quotient map. Let  $A \subseteq X$  be saturated with respect to  $p$ . Let  $q : A \twoheadrightarrow p(A)$  be the restriction of  $p$ .

- 1. If  $A$  is either open or closed in  $X$  then  $q$  is a quotient map.
- 2. If  $p$  is either an open map or a closed map then  $q$  is a quotient map.

PROOF:

- $\langle 1 \rangle 1$ . LET:  $p : X \twoheadrightarrow Y$  be a quotient map.
- $\langle 1 \rangle 2$ . LET:  $A \subseteq X$  be saturated with respect to  $p$ .
- $\langle 1 \rangle 3$ . LET:  $q : A \twoheadrightarrow p(A)$  be the restriction of  $p$ .
- $\langle 1 \rangle 4$ .  $q$  is continuous.  
PROOF: Theorem 97.
- $\langle 1 \rangle 5$ . If  $A$  is open in  $X$  then  $q$  is a quotient map.
  - $\langle 2 \rangle 1$ . ASSUME:  $A$  is open in  $X$ .
  - $\langle 2 \rangle 2$ .  $q$  maps saturated open sets to open sets.
    - $\langle 3 \rangle 1$ . LET:  $U \subseteq A$  be saturated with respect to  $q$  and open in  $A$
    - $\langle 3 \rangle 2$ .  $U$  is saturated with respect to  $p$ 
      - $\langle 4 \rangle 1$ . LET:  $x, y \in X$
      - $\langle 4 \rangle 2$ . ASSUME:  $x \in U$
      - $\langle 4 \rangle 3$ . ASSUME:  $p(x) = p(y)$
      - $\langle 4 \rangle 4$ .  $x \in A$   
PROOF: From  $\langle 3 \rangle 1$  and  $\langle 4 \rangle 2$ .
      - $\langle 4 \rangle 5$ .  $y \in A$   
PROOF: From  $\langle 1 \rangle 2$  and  $\langle 4 \rangle 3$
      - $\langle 4 \rangle 6$ .  $q(x) = x(y)$   
PROOF: From  $\langle 1 \rangle 3$ ,  $\langle 4 \rangle 3$ ,  $\langle 4 \rangle 4$ ,  $\langle 4 \rangle 5$ .
      - $\langle 4 \rangle 7$ .  $y \in U$   
PROOF: From  $\langle 3 \rangle 1$ ,  $\langle 4 \rangle 2$ ,  $\langle 4 \rangle 6$
  - $\langle 3 \rangle 3$ .  $U$  is open in  $X$   
PROOF: Lemma 142,  $\langle 2 \rangle 1$ ,  $\langle 3 \rangle 1$ .
  - $\langle 3 \rangle 4$ .  $p(U)$  is open in  $Y$   
PROOF: Proposition 193,  $\langle 1 \rangle 1$ ,  $\langle 3 \rangle 2$ ,  $\langle 3 \rangle 3$

⟨3⟩5.  $q(U)$  is open in  $p(A)$

PROOF: Since  $q(U) = p(U) = p(U) \cap p(A)$ .

⟨2⟩3. Q.E.D.

PROOF: By Proposition 193.

⟨1⟩6. If  $A$  is closed in  $X$  then  $q$  is a quotient map.

PROOF: Similar.

⟨1⟩7. If  $p$  is an open map then  $q$  is a quotient map.

⟨2⟩1. ASSUME:  $p$  is an open map

⟨2⟩2.  $q$  maps saturated open sets to open sets.

⟨3⟩1. LET:  $U$  be open in  $A$  and saturated with respect to  $q$

⟨3⟩2. PICK  $V$  open in  $X$  such that  $U = A \cap V$

⟨3⟩3.  $p(V)$  is open in  $Y$

⟨3⟩4.  $q(U) = p(V) \cap p(A)$

⟨4⟩1.  $q(U) \subseteq p(V) \cap p(A)$

PROOF: From ⟨3⟩2.

⟨4⟩2.  $p(V) \cap p(A) \subseteq q(U)$

⟨5⟩1. LET:  $y \in p(V) \cap p(A)$

⟨5⟩2. PICK  $x \in V$  and  $x' \in A$  such that  $p(x) = p(x') = y$

⟨5⟩3.  $x \in A$

PROOF: By ⟨1⟩2.

⟨5⟩4.  $x \in U$

PROOF: From ⟨3⟩2

⟨2⟩3. Q.E.D.

PROOF: By Proposition 193.

⟨1⟩8. If  $p$  is a closed map then  $q$  is a quotient map.

PROOF: Similar.

□

**Example 196.** This example shows we cannot remove the hypotheses on  $A$  and  $p$ .

Define  $f : [0, 1] \rightarrow [2, 3] \rightarrow [0, 2]$  by  $f(x) = x$  if  $x \leq 1$ ,  $f(x) = x - 1$  if  $x \geq 2$ . Then  $f$  is a quotient map but its restriction  $f'$  to  $[0, 1] \cup [2, 3]$  is not, because  $f'^{-1}([1, 2])$  is open but  $[1, 2]$  is not.

For a counterexample where  $A$  is saturated, see Example 202.

**Proposition 197.** Let  $p : A \rightarrow C$  and  $q : B \rightarrow D$  be open quotient maps. Then  $p \times q : A \times B \rightarrow C \times D$  is an open quotient map.

PROOF: From Corollary 193.1, Proposition 136 and Theorem 132. □

**Theorem 198.** Let  $p : X \rightarrow Y$  be a quotient map. Let  $Z$  be a topological space and  $f : Y \rightarrow Z$  be a function. Then

1.  $f \circ p$  is continuous if and only if  $f$  is continuous.

2.  $f \circ p$  is a quotient map if and only if  $f$  is a quotient map.

PROOF:

- $\langle 1 \rangle 1$ . If  $f \circ p$  is continuous then  $f$  is continuous.  
 PROOF: Proposition 188.  
 $\langle 1 \rangle 2$ . If  $f$  is continuous then  $f \circ p$  is continuous.  
 PROOF: Theorem 96.  
 $\langle 1 \rangle 3$ . If  $f \circ p$  is a quotient map then  $f$  is a quotient map.  
 PROOF: Proposition 189.  
 $\langle 1 \rangle 4$ . If  $f$  is a quotient map then  $f \circ p$  is a quotient map.  
 PROOF: From Proposition 187.

□

**Proposition 199.** *Let  $X$  and  $Y$  be topological spaces. Let  $p : X \rightarrow Y$  and  $f : Y \rightarrow X$  be continuous maps such that  $p \circ f = \text{id}_Y$ . Then  $p$  is a quotient map.*

PROOF:

- $\langle 1 \rangle 1$ . LET:  $V \subseteq Y$   
 $\langle 1 \rangle 2$ . ASSUME:  $p^{-1}(V)$  is open in  $X$ .  
 $\langle 1 \rangle 3$ .  $f^{-1}(p^{-1}(V))$  is open in  $Y$ .  
 PROOF: Because  $f$  is continuous.  
 $\langle 1 \rangle 4$ .  $V$  is open in  $Y$ .  
 PROOF: Because  $f^{-1}(p^{-1}(V)) = V$ .

□

## 29 Quotient Topology

**Definition 200** (Quotient Topology). Let  $X$  be a topological space,  $Y$  a set and  $p : X \rightarrow Y$  be a surjective function. Then the *quotient topology* on  $Y$  is the unique topology on  $Y$  with respect to which  $p$  is a quotient map, namely  $\mathcal{T} = \{U \in \mathcal{P}Y \mid p^{-1}(U) \text{ is open in } X\}$ .

We prove this is a topology.

PROOF:

- $\langle 1 \rangle 1$ .  $Y \in \mathcal{T}$   
 PROOF: Since  $p^{-1}(Y) = X$  by surjectivity.  
 $\langle 1 \rangle 2$ . For all  $\mathcal{A} \subseteq \mathcal{T}$  we have  $\bigcup \mathcal{A} \in \mathcal{T}$   
 PROOF: Since  $p^{-1}(\bigcup \mathcal{A}) = \bigcup_{U \in \mathcal{A}} p^{-1}(U)$   
 $\langle 1 \rangle 3$ . For all  $U, V \in \mathcal{T}$  we have  $U \cap V \in \mathcal{T}$   
 PROOF: Since  $p^{-1}(U \cap V) = p^{-1}(U) \cap p^{-1}(V)$ .

□

**Definition 201** (Quotient Space). Let  $X$  be a topological space and  $\sim$  an equivalence relation on  $X$ . Let  $p : X \rightarrow X/\sim$  be the canonical surjection. Then  $X/\sim$  under the quotient topology is called a *quotient space*, *identification space* or *decomposition space* of  $X$ .

Here is a counterexample showing we cannot remove all the hypotheses of Proposition 195 except that  $A$  is saturated.

**Example 202.** Let  $X = (0, 1/2] \cup \{1\} \cup \{1 + 1/n : n \geq 2\}$  as a subspace of  $\mathbb{R}$ . Define  $R$  to be the equivalence relation on  $X$  where  $xRy$  iff  $(x = y \text{ or } |x - y| = 1)$ , so we identify  $1/n$  with  $1 + 1/n$  for all  $n \geq 2$ . Let  $Y$  be the resulting quotient space  $X/R$  in the quotient topology and  $p : X \twoheadrightarrow Y$  the canonical surjection.

Let  $A = \{1\} \cup (0, 1/2] \setminus \{1/n : n \geq 2\} \subseteq X$ . Then  $A$  is saturated under  $p$  but the restriction  $q$  of  $p$  to  $A$  is not a quotient map because it maps the saturated open set  $\{1\}$  to  $\{1\}$  which is not open in  $p(A)$ .

**Proposition 203.** Let  $X, Y$  and  $Z$  be topological spaces. Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ . If  $f$  and  $g$  are quotient maps then so is  $g \circ f$ .

PROOF: From Proposition 187.  $\square$

**Example 204.** The product of two quotient maps is not necessarily a quotient map.

Let  $X = \mathbb{R}$  and  $X^*$  the quotient space formed by identifying all positive integers. Let  $p : X \twoheadrightarrow X^*$  be the canonical surjection.

We prove  $p \times \text{id}_{\mathbb{Q}} : X \times \mathbb{Q} \rightarrow X^* \times \mathbb{Q}$  is not a quotient map.

PROOF:

$\langle 1 \rangle 1$ . For  $n \geq 1$ ,

LET:  $c_n = \sqrt{2}/n$

$\langle 1 \rangle 2$ . For  $n \geq 1$ ,

LET:  $U_n = \{(x, y) \in X \times \mathbb{Q} \mid n - 1/4 < x < n + 1/4, (y + n > x + c_n \text{ and } y + n > -x + c_n) \text{ or } (y + n < x + c_n \text{ and } y + n < -x + c_n)\}$

$\langle 1 \rangle 3$ . For  $n \geq 1$ , we have  $U_n$  is open in  $X \times \mathbb{Q}$

$\langle 1 \rangle 4$ . For  $n \geq 1$ , we have  $\{n\} \times \mathbb{Q} \subseteq U_n$

$\langle 1 \rangle 5$ . LET:  $U = \bigcup_{n=1}^{\infty} U_n$

$\langle 1 \rangle 6$ .  $U$  is open in  $X \times \mathbb{Q}$

$\langle 1 \rangle 7$ .  $U$  is saturated with respect to  $p \times \text{id}_{\mathbb{Q}}$

$\langle 1 \rangle 8$ . LET:  $U' = (p \times \text{id}_{\mathbb{Q}})(U)$

$\langle 1 \rangle 9$ . ASSUME: for a contradiction  $U'$  is open in  $X^* \times \mathbb{Q}$

$\langle 1 \rangle 10$ .  $(1, 0) \in U'$

$\langle 1 \rangle 11$ . PICK a neighbourhood  $W$  of 1 in  $X^*$  and  $\delta > 0$  such that  $W \times (-\delta, \delta) \subseteq U'$

$\langle 1 \rangle 12$ .  $p^{-1}(W) \times (-\delta, \delta) \subseteq U$

$\langle 1 \rangle 13$ . PICK  $n$  such that  $c_n < \delta$

$\langle 1 \rangle 14$ .  $n \in p^{-1}(W)$

$\langle 1 \rangle 15$ . PICK  $\epsilon > 0$  such that  $\epsilon < \delta - c_n$  and  $\epsilon < 1/4$  and  $(n - \epsilon, n + \epsilon) \subseteq p^{-1}(W)$

$\langle 1 \rangle 16$ .  $(n - \epsilon, n + \epsilon) \times (-\delta, \delta) \subseteq U$

$\langle 1 \rangle 17$ . PICK a rational  $y$  such that  $c_n - \epsilon/2 < y < c_n + \epsilon/2$

$\langle 1 \rangle 18$ .  $(n + \epsilon/2, y) \notin U$

$\langle 1 \rangle 19$ . Q.E.D.

PROOF: This contradicts  $\langle 1 \rangle 16$ .

$\square$

**Proposition 205.** Let  $X$  be a topological space and  $\sim$  an equivalence relation on  $X$ . Then  $X/\sim$  is  $T_1$  if and only if every equivalence class is closed in  $X$ .

PROOF: Immediate from definitions.  $\square$

## 30 Retractions

**Definition 206** (Retraction). Let  $X$  be a topological space and  $A \subseteq X$ . A *retraction* of  $X$  onto  $A$  is a continuous map  $r : X \rightarrow A$  such that, for all  $a \in A$ , we have  $r(a) = a$ .

**Proposition 207.** *Every retraction is a quotient map.*

PROOF: Proposition 199 with  $f$  the inclusion  $A \hookrightarrow X$ .  $\square$

## 31 Homogeneous Spaces

**Definition 208** (Homogeneous). A topological space  $X$  is *homogeneous* if and only if, for any points  $a, b \in X$ , there exists a homeomorphism  $\phi : X \cong X$  such that  $\phi(a) = b$ .

## 32 Regular Spaces

**Definition 209** (Regular Space). A topological space  $X$  is *regular* if and only if, for any closed set  $A$  and point  $a \notin A$ , there exist disjoint open sets  $U, V$  such that  $A \subseteq U$  and  $a \in V$ .

## 33 Connected Spaces

**Definition 210** (Separation). A *separation* of a topological space  $X$  is a pair of disjoint open sets  $U, V$  such that  $U \cup V = \emptyset$ .

**Definition 211** (Connected). A topological space is *connected* if and only if it has no separation; otherwise it is *disconnected*.

**Proposition 212.** *A topological space  $X$  is connected if and only if the only sets that are both open and closed are  $X$  and  $\emptyset$ .*

Immediate from definitions.

**Lemma 213.** *If  $Y$  is a subspace of  $X$ , a separation of  $Y$  is a pair of disjoint nonempty sets  $A$  and  $B$  whose union is  $Y$ , neither of which contains a limit point of the other.*

PROOF:

$\langle 1 \rangle 1$ . LET:  $A, B \subseteq Y$

$\langle 1 \rangle 2$ . If  $A$  and  $B$  form a separation of  $Y$  then  $A$  and  $B$  are disjoint and nonempty,  $A \cup B = Y$ , and neither of  $A$  and  $B$  contains a limit point of the other.

$\langle 2 \rangle 1$ . ASSUME:  $A$  and  $B$  form a separation of  $Y$

$\langle 2 \rangle 2$ .  $A$  and  $B$  are disjoint and nonempty and  $A \cup B = Y$

PROOF: From  $\langle 2 \rangle 1$  and the definition of separation.

$\langle 2 \rangle 3$ .  $A$  does not contain a limit point of  $B$

⟨3⟩1. ASSUME: for a contradiction  $l \in A$  and  $l$  is a limit point of  $B$  in  $X$ .  
 ⟨3⟩2.  $l$  is a limit point of  $B$  in  $Y$   
 PROOF: Proposition 152.  
 ⟨3⟩3.  $l \in B$   
 ⟨4⟩1.  $B$  is closed in  $Y$   
 PROOF: Since  $A$  is open in  $Y$  and  $B = Y \setminus A$  from ⟨2⟩1.  
 ⟨4⟩2. Q.E.D.  
 PROOF: Corollary 58.1.  
 ⟨3⟩4. Q.E.D.  
 PROOF: This contradicts the fact that  $A \cap B = \emptyset$  (⟨2⟩1).  
 ⟨2⟩4.  $B$  does not contain a limit point of  $A$   
 PROOF: Similar.  
 ⟨1⟩3. If  $A$  and  $B$  are disjoint and nonempty,  $A \cup B = Y$ , and neither of  $A$  and  $B$  contains a limit point of the other, then  $A$  and  $B$  form a separation of  $Y$ .  
 ⟨2⟩1. ASSUME:  $A$  and  $B$  are disjoint and nonempty,  $A \cup B = Y$ , and neither of  $A$  and  $B$  contains a limit point of the other.  
 ⟨2⟩2.  $A$  is open in  $Y$   
 ⟨3⟩1.  $B$  is closed in  $Y$   
 ⟨4⟩1. LET:  $l$  be a limit point of  $B$  in  $Y$   
 ⟨4⟩2.  $l$  is a limit point of  $B$  in  $X$   
 PROOF: Proposition 152.  
 ⟨4⟩3.  $l \notin A$   
 PROOF: By ⟨2⟩1  
 ⟨4⟩4.  $l \in B$   
 PROOF: By ⟨2⟩1 since  $A \cup B = Y$   
 ⟨4⟩5. Q.E.D.  
 PROOF: Corollary 58.1.  
 ⟨3⟩2. Q.E.D.  
 PROOF: Since  $A = Y \setminus B$ .  
 ⟨2⟩3.  $B$  is open in  $Y$   
 PROOF: Similar.

□

**Example 214.** Every set under the indiscrete topology is connected.

**Example 215.** The discrete topology on a set  $X$  is connected if and only if  $|X| \leq 1$ .

**Example 216.** The finite complement topology on a set  $X$  is connected if and only if either  $|X| \leq 1$  or  $X$  is infinite.

**Example 217.** The countable complement topology on a set  $X$  is connected if and only if either  $|X| \leq 1$  or  $X$  is uncountable.

**Example 218.** The rationals  $\mathbb{Q}$  are disconnected. For any irrational  $a$ , the sets  $(-\infty, a) \cap \mathbb{Q}$  and  $(a, +\infty) \cap \mathbb{Q}$  form a separation of  $\mathbb{Q}$ .

**Lemma 219.** *Let  $X$  be a topological space. If  $C$  and  $D$  form a separation of  $X$ , and  $Y$  is a connected subspace of  $X$ , then either  $Y \subseteq C$  or  $Y \subseteq D$ .*

PROOF: Otherwise  $Y \cap C$  and  $Y \cap D$  would form a separation of  $Y$ .  $\square$

**Theorem 220.** *The union of a set of connected subspaces of a space  $X$  that have a point in common is connected.*

PROOF:

$\langle 1 \rangle 1$ . LET:  $\mathcal{A}$  be a set of connected subspaces of the space  $X$  that have the point  $a$  in common.

$\langle 1 \rangle 2$ . ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $\bigcup \mathcal{A}$

$\langle 1 \rangle 3$ . ASSUME: without loss of generality  $a \in C$

$\langle 1 \rangle 4$ . For all  $A \in \mathcal{A}$  we have  $A \subseteq C$

PROOF: Lemma 219.

$\langle 1 \rangle 5$ .  $D = \emptyset$

$\langle 1 \rangle 6$ . Q.E.D.

PROOF: This contradicts  $\langle 1 \rangle 2$ .

$\square$

**Theorem 221.** *Let  $X$  be a topological space and  $A$  a connected subspace of  $X$ . If  $A \subseteq B \subseteq \overline{A}$  then  $B$  is connected.*

PROOF:

$\langle 1 \rangle 1$ . ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $B$ .

$\langle 1 \rangle 2$ . ASSUME: without loss of generality  $A \subseteq C$

PROOF: Lemma 219.

$\langle 1 \rangle 3$ .  $B \subseteq C$

$\langle 2 \rangle 1$ . LET:  $x \in B$

$\langle 2 \rangle 2$ .  $x \in \overline{A}$

$\langle 2 \rangle 3$ . Either  $x \in A$  or  $x$  is a limit point of  $A$ .

PROOF: Theorem 58.

$\langle 2 \rangle 4$ . Either  $x \in A$  or  $x$  is a limit point of  $C$ .

PROOF: Lemma 60,  $\langle 1 \rangle 2$ .

$\langle 2 \rangle 5$ .  $x \in C$

PROOF: Lemma 213.

$\langle 1 \rangle 4$ .  $D = \emptyset$

$\langle 1 \rangle 5$ . Q.E.D.

PROOF: This contradicts  $\langle 1 \rangle 1$ .

$\square$

**Theorem 222.** *The image of a connected space under a continuous map is connected.*

PROOF:

$\langle 1 \rangle 1$ . LET:  $f : X \rightarrow Y$  be a surjective continuous map where  $X$  is connected.

$\langle 1 \rangle 2$ . ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $Y$ .

$\langle 1 \rangle 3$ .  $f^{-1}(C)$  and  $f^{-1}(D)$  form a separation of  $X$ .



□

**Theorem 223.** *The product of a family of connected spaces is connected.*

PROOF:

⟨1⟩1. The product of two connected spaces is connected.

⟨2⟩1. LET:  $X$  and  $Y$  be connected spaces.

⟨2⟩2. PICK  $a \in X$  and  $b \in Y$

PROOF: We may assume  $X$  and  $Y$  are nonempty since otherwise  $X \times Y = \emptyset$  which is connected.

⟨2⟩3.  $X \times \{b\}$  is connected.

PROOF: It is homeomorphic to  $X$ .

⟨2⟩4. For all  $x \in X$  we have  $\{x\} \times Y$  is connected.

PROOF: It is homeomorphic to  $Y$ .

⟨2⟩5. For any  $x \in X$

LET:  $T_x = (X \times \{b\}) \cup (\{x\} \times Y)$

⟨2⟩6. For all  $x \in X$ ,  $T_x$  is connected.

PROOF: Theorem 220 since  $(x, b) \in (X \times \{b\}) \cap (\{x\} \times Y)$ .

⟨2⟩7.  $X \times Y$  is connected.

PROOF: Theorem 220 since  $X \times Y = \bigcup_{x \in X} T_x$  and  $(a, b)$  is a point in every  $T_x$ .

⟨1⟩2. The product of a finite family of connected spaces is connected.

PROOF: From ⟨1⟩1 by induction.

⟨1⟩3. The product of any family of connected spaces is connected.

⟨2⟩1. LET:  $\{X_\alpha\}_{\alpha \in J}$  be a family of connected spaces.

⟨2⟩2. LET:  $X = \prod_{\alpha \in J} X_\alpha$

⟨2⟩3. PICK  $a \in X$

PROOF: We may assume  $X \neq \emptyset$  as the empty space is connected.

⟨2⟩4. For every finite subset  $K$  of  $J$ ,

LET:  $X_K = \{x \in X \mid \forall \alpha \in J \setminus K, x_\alpha = a_\alpha\}$

⟨2⟩5. For every finite  $K \subseteq J$ , we have  $X_K$  is connected.

PROOF: From ⟨1⟩2 since  $X_K \cong \prod_{\alpha \in K} X_\alpha$ .

⟨2⟩6. LET:  $Y = \bigcup_K X_K$

⟨2⟩7.  $Y$  is connected

PROOF: Theorem 220 since  $a$  is a common point.

⟨2⟩8.  $X = \overline{Y}$

⟨3⟩1. LET:  $x \in X$

⟨3⟩2. LET:  $U = \prod_{\alpha \in J} U_\alpha$  be a basic neighbourhood of  $x$  where  $U_\alpha = X_\alpha$  for all  $\alpha$  except  $\alpha \in K$  for some finite  $K \subseteq J$

⟨3⟩3. LET:  $y \in X$  be the point with  $y_\alpha = x_\alpha$  for  $\alpha \in K$  and  $y_\alpha = a_\alpha$  for all other  $\alpha$

⟨3⟩4.  $y \in U \cap X_K$

⟨3⟩5.  $y \in U \cap Y$

⟨2⟩9.  $X$  is connected.

PROOF: Theorem 221.

□

**Example 224.** The set  $\mathbb{R}^\omega$  is disconnected under the box topology. The set of bounded sequences and the set of unbounded sequences form a separation.

**Proposition 225.** Let  $\mathcal{T}$  and  $\mathcal{T}'$  be two topologies on the same set  $X$ . If  $\mathcal{T} \subseteq \mathcal{T}'$  and  $(X, \mathcal{T}')$  is connected then  $(X, \mathcal{T})$  is connected.

PROOF: If  $U$  and  $V$  form a separation of  $(X, \mathcal{T})$  then they form a separation of  $(X, \mathcal{T}')$ .  $\square$

**Proposition 226.** Let  $X$  be a topological space and  $(A_n)$  a sequence of connected subspaces of  $X$ . If  $A_n \cap A_{n+1} \neq \emptyset$  for all  $n$  then  $\bigcup_n A_n$  is connected.

PROOF:

$\langle 1 \rangle 1$ . ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $\bigcup_n A_n$

$\langle 1 \rangle 2$ . ASSUME: without loss of generality  $A_0 \subseteq C$

PROOF: Lemma 219.

$\langle 1 \rangle 3$ . For all  $n$  we have  $A_n \subseteq C$

PROOF:

$\langle 2 \rangle 1$ . ASSUME:  $A_n \subseteq C$

$\langle 2 \rangle 2$ . PICK  $x \in A_n \cap A_{n+1}$

$\langle 2 \rangle 3$ .  $x \in C$

$\langle 2 \rangle 4$ .  $A_{n+1} \subseteq C$

PROOF: Lemma 219.

$\langle 2 \rangle 5$ . Q.E.D.

PROOF: The result follows by induction.

$\langle 1 \rangle 4$ .  $D = \emptyset$

$\langle 1 \rangle 5$ . Q.E.D.

PROOF: This contradicts  $\langle 1 \rangle 1$ .

$\square$

**Proposition 227.** Let  $X$  be a topological space. Let  $A, C \subseteq X$ . If  $C$  is connected and intersects both  $A$  and  $X \setminus A$  then  $C$  intersects  $\partial A$ .

PROOF: Otherwise  $C \cap A^\circ$  and  $C \setminus \overline{A}$  would form a separation of  $C$ .  $\square$

**Example 228.** The space  $\mathbb{R}_l$  is disconnected. For any real  $x$ , the sets  $(-\infty, x)$  and  $[x, +\infty)$  form a separation.

**Proposition 229.** Let  $X$  and  $Y$  be connected spaces. Let  $A$  be a proper subset of  $X$  and  $B$  a proper subset of  $Y$ . Then  $(X \times Y) \setminus (A \times B)$  is connected.

PROOF:

$\langle 1 \rangle 1$ . PICK  $a \in X \setminus A$  and  $b \in Y \setminus B$

$\langle 1 \rangle 2$ . For  $x \in X \setminus A$  we have  $(X \times \{b\}) \cup (\{x\} \times Y)$  is connected.

PROOF: Theorem 220 since  $(x, b)$  is a common point.

$\langle 1 \rangle 3$ . For  $y \in Y \setminus B$  we have  $(X \times \{y\}) \cup (\{a\} \times Y)$  is connected.

PROOF: Theorem 220 since  $(a, y)$  is a common point.

$\langle 1 \rangle 4$ .  $(X \times Y) \setminus (A \times B)$  is connected.

PROOF: Theorem 220 since it is the union of the sets in  $\langle 1 \rangle 2$  and  $\langle 1 \rangle 3$  with  $(a, b)$  as a common point.

□

**Proposition 230.** *Let  $p : X \rightarrow Y$  be a quotient map. If  $Y$  is connected and  $p^{-1}(y)$  is connected for all  $y \in Y$ , then  $X$  is connected.*

PROOF:

⟨1⟩1. ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $X$ .

⟨1⟩2.  $C$  is saturated.

⟨2⟩1. LET:  $x \in C, y \in X$  with  $p(x) = p(y) = a$ , say

⟨2⟩2.  $y \notin D$

PROOF: Otherwise  $p^{-1}(a) \cap C$  and  $p^{-1}(a) \cap D$  form a separation of  $p^{-1}(a)$ .

⟨2⟩3.  $y \in C$

⟨1⟩3.  $D$  is saturated.

PROOF: Similar.

⟨1⟩4.  $p(C)$  and  $p(D)$  form a separation of  $Y$ .

□

**Proposition 231.** *Let  $X$  be a connected space and  $Y$  a connected subspace of  $X$ . Suppose  $A$  and  $B$  form a separation of  $X \setminus Y$ . Then  $Y \cup A$  and  $Y \cup B$  are both connected.*

PROOF:

⟨1⟩1.  $Y \cup A$  is connected.

⟨2⟩1. ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $Y \cup A$

⟨2⟩2. ASSUME: without loss of generality  $Y \subseteq C$

⟨2⟩3. PICK open sets  $A_1, B_1, C_1, D_1$  in  $X$  with

$$A = A_1 \setminus Y$$

$$B = B_1 \setminus Y$$

$$C = C_1 \cap (Y \cup A)$$

$$D = D_1 \cap (Y \cup A)$$

⟨2⟩4.  $B_1 \cup C_1$  and  $A_1 \cap D_1$  form a separation of  $X$

⟨1⟩2.  $Y \cup B$  is connected.

PROOF: Similar.

□

**Theorem 232.** *Let  $L$  be a linearly ordered set under the order topology. Then  $L$  is connected if and only if  $L$  is a linear continuum.*

PROOF:

⟨1⟩1. If  $L$  is a linear continuum then  $L$  is connected.

⟨2⟩1. LET:  $L$  be a linear continuum under the order topology.

⟨2⟩2. ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $L$ .

⟨2⟩3. PICK  $a \in C$  and  $b \in D$ .

⟨2⟩4. ASSUME: without loss of generality  $a < b$ .

⟨2⟩5. LET:  $S = \{x \in L \mid a < x \text{ and } [a, x) \subseteq C\}$

⟨2⟩6.  $S$  is nonempty.

PROOF: Since  $a \in C$  and  $C$  is open.

(2)7.  $S$  is bounded above by  $b$ .  
 PROOF: Since  $b \notin C$ .  
 (2)8. LET:  $s = \sup S$   
 (2)9.  $s \in S$   
 (3)1. LET:  $y \in [a, s)$   
 PROVE:  $y \in C$   
 (3)2. PICK  $z$  with  $y < z \in S$   
 PROOF: By minimality of  $s$ .  
 (3)3.  $y \in [a, z) \subseteq C$   
 (2)10. CASE:  $s \in C$   
 (3)1. PICK  $x$  such that  $s < x$  and  $[s, x) \subseteq C$   
 PROOF: Since  $C$  is open and  $s$  is not greatest in  $L$  because  $s < b$ .  
 (3)2.  $x \in S$   
 PROOF: Since  $[a, x) = [a, s) \cup [s, x) \subseteq C$ .  
 (3)3. Q.E.D.  
 PROOF: This contradicts the fact that  $s$  is an upper bound for  $S$ .  
 (2)11. CASE:  $s \in D$   
 (3)1. PICK  $x < s$  such that  $(x, s] \subseteq D$   
 (3)2. PICK  $y$  with  $x < y < s$   
 PROOF: Since  $L$  is dense.  
 (3)3.  $y \in C$   
 PROOF: From (2)9.  
 (3)4.  $y \in D$   
 PROOF: From (3)1.  
 (3)5. Q.E.D.  
 (3)6. LET:  $L$  be a linear continuum under the order topology.  
 (3)7. ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $L$ .  
 (3)8. PICK  $a \in C$  and  $b \in D$ .  
 (3)9. ASSUME: without loss of generality  $a < b$ .  
 (3)10. LET:  $S = \{x \in L \mid a < x \text{ and } [a, x) \subseteq C\}$   
 (3)11.  $S$  is nonempty.  
 PROOF: Since  $a \in C$  and  $C$  is open.  
 (3)12.  $S$  is bounded above by  $b$ .  
 PROOF: Since  $b \notin C$ .  
 (3)13. LET:  $s = \sup S$   
 (3)14.  $s \in S$   
 (4)1. LET:  $y \in [a, s)$   
 PROVE:  $y \in C$   
 (4)2. PICK  $z$  with  $y < z \in S$   
 PROOF: By minimality of  $s$ .  
 (4)3.  $y \in [a, z) \subseteq C$   
 (3)15. CASE:  $s \in C$   
 (4)1. PICK  $x$  such that  $s < x$  and  $[s, x) \subseteq C$   
 PROOF: Since  $C$  is open and  $s$  is not greatest in  $L$  because  $s < b$ .  
 (4)2.  $x \in S$   
 PROOF: Since  $[a, x) = [a, s) \cup [s, x) \subseteq C$ .

⟨4⟩3. Q.E.D.

PROOF: This contradicts the fact that  $s$  is an upper bound for  $S$ .

$\langle 3 \rangle 16$ . CASE:  $s \in D$

⟨4⟩1. PICK  $x < s$  such that  $(x, s] \subseteq D$

⟨4⟩2. PICK  $y$  with  $x < y < s$

PROOF: Since  $L$  is dense.

⟨4⟩3.  $y \in C$

PROOF: From  $\langle 2 \rangle 9$ .

$\langle 4 \rangle 4. \ y \in D$

PROOF: From  $\langle 3 \rangle 1$ .

⟨4⟩5. Q.E.D.

PROOF: This contradicts  $\langle 2 \rangle 2$ .

$\langle 1 \rangle 2$ . If  $L$  is connected then  $L$  is a linear continuum.

$\langle 2 \rangle 1$ . ASSUME:  $L$  is connected.

(2) Every nonempty subset of  $L$  that is bounded above has a supremum.

**(3)1. LET:**  $X$  be a nonempty subset of  $L$  bounded above by  $b$ .

⟨3⟩2. ASSUME: for a contradiction  $X$  has no supremum.

**3. LET:**  $U$  be the set of upper bounds of  $X$ ,

$\langle 3 \rangle 4$ .  $U$  is nonempty.

PROOF: Since  $b \in U$ .

$\langle 3 \rangle 5$ .  $U$  is open.

$\langle 4 \rangle 1$ . LET:  $x \in U$

(4)2. PICK an upper bound  $y$  for  $X$  such that  $y < x$

⟨4⟩3. Either  $x$  is greatest in  $L$  and  $(y, x] \subseteq U$ , or there exists  $z > x$  such that  $(y, z) \subseteq U$

**⟨3⟩6.** LET:  $V$  be the set of lower bounds of  $U$ .

$\langle 3 \rangle 7$ .  $V$  is nonempty.

PROOF: Since  $X \subset V$

$\langle 3 \rangle 8$ .  $V$  is open.

$\langle 4 \rangle 1$ . LET:  $x \in V$

⟨4⟩2. PICK  $y \in X$  with  $x < y$

PROOF:  $x$  cannot be an upper bound for  $X$ , because it would be the supremum of  $X$ .

(4)3. Either  $x$  least in  $L$  and  $[x, y] \subseteq V$ , or there exists  $z < x$  such that  $(z, y) \subseteq V$

⟨3⟩9.  $L = U \cup V$

$\langle 4 \rangle 1$ . LET:  $x \in L \setminus U$

⟨4⟩2. PICK  $y \in X$  such that  $x < y$

⟨4⟩3. For all  $u \in U$  we have  $x < y \leq u$

$$\langle 4 \rangle 4. \quad x \in V$$

⟨3⟩10.  $U \cap V = \emptyset$

PROOF: Any element of  $U \cap V$  would be a supremum of  $X$ .

⟨3⟩11.  $U$  and  $V$  form a separation of  $L$ .

$\langle 3 \rangle 12$ . Q.E.D.

PROOF: This contradicts  $\langle 2 \rangle 1$ .

$\langle 2 \rangle 3$ .  $L$  is dense.

⟨3⟩1. LET:  $x, y \in L$  with  $x < y$

⟨3⟩2. There exists  $z \in L$  such that  $x < z < y$

PROOF: Otherwise  $(-\infty, y)$  and  $(x, +\infty)$  would form a separation of  $L$ .

□

**Corollary 232.1.** *The real line  $\mathbb{R}$  is connected.*

**Corollary 232.2.** *Every interval in  $\mathbb{R}$  is connected.*

**Corollary 232.3.** *The ordered square is connected.*

**Theorem 233** (Intermediate Value Theorem). *Let  $X$  be a connected space. Let  $Y$  be a linearly ordered set under the order topology. Let  $f : X \rightarrow Y$  be continuous. Let  $a, b \in X$  and  $r \in Y$ . Suppose  $f(a) < r < f(b)$ . Then there exists  $c \in X$  such that  $f(c) = r$ .*

PROOF: Otherwise  $f^{-1}((-\infty, r))$  and  $f^{-1}((r, +\infty))$  would form a separation of  $X$ . □

**Proposition 234.** *Every function  $f : [0, 1] \rightarrow [0, 1]$  has a fixed point.*

PROOF:

⟨1⟩1. LET:  $g : [0, 1] \rightarrow [-1, 1]$  be the function  $g(x) = f(x) - x$

PROVE: there exists  $x \in [0, 1]$  such that  $g(x) = 0$

⟨1⟩2. ASSUME: without loss of generality  $g(0) \neq 0$  and  $g(1) \neq 0$

⟨1⟩3.  $g(0) > 0$

⟨1⟩4.  $g(1) < 0$

⟨1⟩5. There exists  $x \in (0, 1)$  such that  $g(x) = 0$

PROOF: By the Intermediate Value Theorem.

**Proposition 235.** *Give  $\mathbb{R}^\omega$  the box topology. Let  $x, y \in \mathbb{R}^\omega$ . Then  $x$  and  $y$  lie in the same component if and only if  $x - y$  is eventually zero, i.e. there exists  $N$  such that, for all  $n \geq N$ , we have  $x_n = y_n$ .*

PROOF:

⟨1⟩1. The component containing 0 is the set of sequences that are eventually zero.

⟨2⟩1. LET:  $B$  be the set of sequences that are eventually zero.

⟨2⟩2.  $B$  is path-connected.

⟨3⟩1. LET:  $x, y \in B$

⟨3⟩2. PICK  $N$  such that, for all  $n \geq N$ , we have  $x_n = y_n = 0$

⟨3⟩3. LET:  $p : [0, 1] \rightarrow \mathbb{R}^\omega$ ,  $p(t) = (1 - t)x + ty$

PROVE:  $p$  is continuous.

⟨3⟩4. LET:  $t \in [0, 1]$  and  $\prod_j U_j$  be a basic open neighbourhood of  $p(t)$ , where each  $U_j$  is open in  $\mathbb{R}$

⟨3⟩5. PICK  $\delta$  such that, for all  $n < N$  and all  $s \in [0, 1]$ , if  $|s - t| < \delta$  then  $p(s)_n \in U_n$

⟨3⟩6. For all  $s \in [0, 1]$ , if  $|s - t| < \delta$  then  $p(s) \in \prod_j U_j$

⟨2⟩3.  $B$  is connected.

PROOF: Proposition 241.

⟨2⟩4. If  $C$  is connected and  $B \subseteq C$  then  $B = C$ .

⟨3⟩1. ASSUME:  $C$  is connected and  $B \subseteq C$

⟨3⟩2. ASSUME: for a contradiction  $x \in C \setminus B$

⟨3⟩3. For  $n \geq 1$ ,

LET:  $c_n = 1$  if  $x_n = 0$ ,  $c_n = n/x_n$  otherwise

⟨3⟩4. LET:  $h : \mathbb{R}^\omega \rightarrow \mathbb{R}^\omega$  be the function  $h(x) = (c_n x_n)_{n \geq 1}$

⟨3⟩5.  $h$  is a homeomorphism of  $\mathbb{R}^\omega$  with itself.

⟨3⟩6.  $h(x)$  is unbounded.

PROOF: For any  $b > 0$ , pick  $N > b$  such that  $x_N \neq 0$ . Then  $h(x)_N > b$ .

⟨3⟩7.  $h^{-1}(\{\text{bounded sequences}\}) \cap C$  and  $h^{-1}(\{\text{unbounded sequences}\}) \cap C$  form a separation of  $C$

⟨3⟩8. Q.E.D.

PROOF: This contradicts ⟨3⟩1.

⟨1⟩2. Q.E.D.

PROOF: Since  $\lambda x.x - y$  is a homeomorphism of  $\mathbb{R}^\omega$  with itself.

□

## 34 Totally Disconnected Spaces

**Definition 236** (Totally Disconnected). A topological space  $X$  is *totally disconnected* if and only if the only connected subspaces are the singletons.

**Example 237.** Every discrete space is totally disconnected.

**Example 238.** The rationals  $\mathbb{Q}$  are totally disconnected.

## 35 Paths and Path Connectedness

**Definition 239** (Path). Let  $X$  be a topological space and  $a, b \in X$ . A *path* from  $a$  to  $b$  is a continuous function  $p : [0, 1] \rightarrow X$  such that  $p(0) = a$  and  $p(1) = b$ .

**Definition 240** (Path Connected). A topological space is *path connected* if and only if there exists a path between any two points.

**Proposition 241.** *Every path connected space is connected.*

PROOF:

⟨1⟩1. LET:  $X$  be a path connected space.

⟨1⟩2. ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $X$ .

⟨1⟩3. PICK  $a \in C$  and  $b \in D$ .

⟨1⟩4. PICK a path  $p : [0, 1] \rightarrow X$  from  $a$  to  $b$ .

⟨1⟩5.  $p^{-1}(C)$  and  $p^{-1}(D)$  form a separation of  $[0, 1]$ .

⟨1⟩6. Q.E.D.

PROOF: This contradicts Corollary 232.2.

□

An example that shows the converse does not hold:

**Example 242.** The ordered square is not path connected.

PROOF:

⟨1⟩1. ASSUME: for a contradiction  $p : [0, 1] \rightarrow I_o^2$  is a path from  $(0, 0)$  to  $(1, 1)$ .

⟨1⟩2.  $p$  is surjective.

PROOF: By the Intermediate Value Theorem.

⟨1⟩3. For  $x \in [0, 1]$ , PICK a rational  $q_x \in p^{-1}((x, 0), (x, 1))$

PROOF: Since  $p^{-1}((x, 0), (x, 1))$  is open and nonempty by ⟨1⟩2.

⟨1⟩4. For  $x, y \in [0, 1]$ , if  $x \neq y$  then  $q_x \neq q_y$

PROOF: We have  $p(q_x) \neq p(q_y)$  because  $((x, 0), (x, 1))$  and  $((y, 0), (y, 1))$  are disjoint.

⟨1⟩5.  $\{q_x \mid x \in [0, 1]\}$  is an uncountable set of rationals.

⟨1⟩6. Q.E.D.

PROOF: This contradicts the fact that the rationals are countable.

□

**Proposition 243.** *The continuous image of a path connected space is path connected.*

PROOF:

⟨1⟩1. LET:  $X$  be a path connected space,  $Y$  a topological space, and  $f : X \rightarrow Y$  be continuous and surjective.

⟨1⟩2. LET:  $a, b \in Y$

⟨1⟩3. PICK  $c, d \in X$  with  $f(c) = a$  and  $f(d) = b$

⟨1⟩4. PICK a path  $p : [0, 1] \rightarrow X$  from  $c$  to  $d$ .

⟨1⟩5.  $f \circ p$  is a path from  $a$  to  $b$  in  $Y$ .

□

**Proposition 244 (AC).** *The product of a family of path-connected spaces is path-connected.*

PROOF:

⟨1⟩1. LET:  $\{X_\alpha\}_{\alpha \in J}$  be a family of path-connected spaces.

⟨1⟩2. LET:  $a, b \in \prod_{\alpha \in J} X_\alpha$

⟨1⟩3. For  $\alpha \in J$ , PICK a path  $p_\alpha : [0, 1] \rightarrow X_\alpha$  from  $a_\alpha$  to  $b_\alpha$

PROOF: Using the Axiom of Choice.

⟨1⟩4. Define  $p : [0, 1] \rightarrow \prod_{\alpha \in J} X_\alpha$  by  $p(t)_\alpha = p_\alpha(t)$

⟨1⟩5.  $p$  is a path from  $a$  to  $b$ .

PROOF: Theorem 132.

□

**Proposition 245.** *The continuous image of a path-connected space is path-connected.*

PROOF:

⟨1⟩1. LET:  $f : X \rightarrow Y$  be continuous and surjective where  $X$  is path-connected.



- ⟨1⟩2. LET:  $a, b \in Y$
- ⟨1⟩3. PICK  $a', b' \in X$  with  $f(a') = a$  and  $f(b') = b$ .
- ⟨1⟩4. PICK a path  $p : [0, 1] \rightarrow X$  from  $a'$  to  $b'$ .
- ⟨1⟩5.  $f \circ p$  is a path from  $a$  to  $b$ .

□

**Proposition 246.** *Let  $X$  be a topological space. The union of a set of path-connected subspaces of  $X$  that have a point in common is path-connected.*

PROOF:

- ⟨1⟩1. LET:  $\mathcal{A}$  be a set of path-connected subspaces of  $X$  with the point  $a$  in common.
- ⟨1⟩2. LET:  $b, c \in \bigcup \mathcal{A}$
- ⟨1⟩3. PICK  $B, C \in \mathcal{A}$  with  $b \in B$  and  $c \in C$ .
- ⟨1⟩4. PICK a path  $p$  in  $B$  from  $b$  to  $a$ .
- ⟨1⟩5. PICK a path  $q$  in  $C$  from  $a$  to  $c$ .
- ⟨1⟩6. The concatenation of  $p$  and  $q$  is a path from  $b$  to  $c$  in  $\bigcup \mathcal{A}$ .

□

**Proposition 247.** *Let  $A \subseteq \mathbb{R}^2$  be countable. Then  $\mathbb{R}^2 \setminus A$  is path-connected.*

PROOF:

- ⟨1⟩1. LET:  $a, b \in \mathbb{R}^2 \setminus A$
- ⟨1⟩2. PICK a line  $l$  in  $\mathbb{R}^2$  with  $a$  on one side and  $b$  on the other.
- ⟨1⟩3. For every point  $x$  on  $l$ ,  
LET:  $p_x$  be the path in  $\mathbb{R}^2$  consisting of a line from  $a$  to  $x$  then a line from  $x$  to  $b$
- ⟨1⟩4. For  $x \neq y$  we have  $p_x$  and  $p_y$  have no points in common except  $a$  and  $b$
- ⟨1⟩5. There are only countably many  $x$  such that a point of  $A$  lies on  $p_x$ .
- ⟨1⟩6. There exists  $x$  such that  $p_x$  is a path from  $a$  to  $b$  in  $\mathbb{R}^2 \setminus A$ .

□

**Proposition 248.** *Every open connected subspace of  $\mathbb{R}^2$  is path-connected.*

PROOF:

- ⟨1⟩1. LET:  $U$  be an open connected subspace of  $\mathbb{R}^2$ .
- ⟨1⟩2. For all  $x_0 \in U$ ,  
LET:  $PC(x_0) = \{y \in U \mid \text{there exists a path from } x \text{ to } y\}$
- ⟨1⟩3. For all  $x_0 \in U$ , the set  $PC(x_0)$  is open and closed in  $U$ .
  - ⟨2⟩1. LET:  $x_0 \in U$
  - ⟨2⟩2.  $PC(x_0)$  is open in  $U$ 
    - ⟨3⟩1. LET:  $y \in PC(x_0)$
    - ⟨3⟩2. PICK  $\epsilon > 0$  such that  $B(y, \epsilon) \subseteq U$   
PROOF: Since  $U$  is open.
    - ⟨3⟩3.  $B(y, \epsilon) \subseteq PC(x_0)$   
PROOF: For all  $z \in B(y, \epsilon)$ , pick a path from  $x_0$  to  $y$  then concatenate the straight line from  $y$  to  $z$ .
  - ⟨2⟩3.  $PC(x_0)$  is closed in  $U$

$\langle 3 \rangle 1$ . LET:  $y \in U$  be a limit point of  $PC(x_0)$

$\langle 3 \rangle 2$ . PICK  $\epsilon > 0$  such that  $B(y, \epsilon) \subseteq U$

$\langle 3 \rangle 3$ . PICK  $z \in PC(x_0) \cap B(y, \epsilon)$

$\langle 3 \rangle 4$ .  $y \in PC(x_0)$

PROOF: Pick a path from  $x_0$  to  $z$  then concatenate the straight line from  $z$  to  $y$ .

$\langle 1 \rangle 4$ .  $PC(x_0) = U$

PROOF: Proposition 212.

□

**Example 249.** If  $A$  is a connected subspace of  $X$ , then  $A^\circ$  is not necessarily connected.

Take two closed circles in  $\mathbb{R}^2$  that touch at one point. The interior of this space is two open circles, and these two circles form a separation.

**Example 250.** If  $A$  is a connected subspace of  $X$  then  $\partial A$  is not necessarily connected.

We have  $[0, 1]$  is connected but  $\partial[0, 1] = \{0, 1\}$  is not.

**Example 251.** If  $A$  is a subspace of  $X$  and  $A^\circ$  and  $\partial A$  are connected, then  $A$  is not necessarily connected.

We have  $\mathbb{Q}^\circ = \emptyset$  and  $\partial\mathbb{Q} = \mathbb{R}$  are connected but  $\mathbb{Q}$  is not connected.

## 36 The Topologist's Sine Curve

**Definition 252** (The Topologist's Sine Curve). Let  $S = \{(x, \sin 1/x) \mid 0 < x \leq 1\}$ , The *topologist's sine curve* is the closure  $\bar{S}$  of  $S$  in  $\mathbb{R}^2$ .

**Proposition 253.** The topologist's sine curve is connected.

PROOF:

$\langle 1 \rangle 1$ . LET:  $S = \{(x, \sin 1/x) \mid 0 < x \leq 1\}$

$\langle 1 \rangle 2$ .  $S$  is connected.

PROOF: Theorem 222.

$\langle 1 \rangle 3$ .  $\bar{S}$  is connected.

PROOF: Theorem 221.

□

**Proposition 254.** The topologist's sine curve is  $\{(x, \sin 1/x) \mid 0 < x \leq 1\} \cup (\{0\} \times [-1, 1])$ .

PROOF: Sketch proof: Given a point  $(0, y)$  with  $-1 \leq y \leq 1$ , pick  $a$  such that  $\sin a = y$ . Then  $((1/a, y), (1/(a + 2\pi), y), (1/(a + 4\pi), y), \dots)$  is a sequence in  $S$  that converges to  $(0, y)$ .

Conversely, let  $(x, y)$  be any point not in  $S \cup (\{0\} \times [-1, 1])$ . If  $x < 0$  or  $y > 1$  or  $y < -1$  then we can easily find a neighbourhood that does not intersect  $S \cup (\{0\} \times [-1, 1])$ . If  $x > 0$  and  $-1 \leq y \leq 1$ , then we have  $y \neq \sin 1/x$ . Hence pick a neighbourhood that does not intersect  $S$ .

**Proposition 255.** *Every closed subset of  $\mathbb{R}$  that is bounded above has a greatest element.*

PROOF: It has a supremum, which is a limit point of the set and hence an element.  $\square$

**Proposition 256 (CC).** *The topologist's sine curve is not path connected.*

PROOF:

$\langle 1 \rangle 1$ . ASSUME: For a contradiction  $p : [0, 1] \rightarrow \bar{S}$  is a path from  $(0, 0)$  to  $(1, \sin 1)$ .

$\langle 1 \rangle 2$ .  $\{t \in [0, 1] \mid p(t) \in \{0\} \times [-1, 1]\}$  is closed.

PROOF: Since  $p$  is continuous and  $\{0\} \times [-1, 1]$  is closed.

$\langle 1 \rangle 3$ . LET:  $b$  be the largest number in  $[0, 1]$  such that  $p(b) \in \{0\} \times [-1, 1]$ .

PROOF: Proposition 255.

$\langle 1 \rangle 4$ . LET:  $x : [b, 1] \rightarrow \bar{S}$  be the function  $\pi_1 \circ p$

$\langle 1 \rangle 5$ . LET:  $y : [b, 1] \rightarrow \bar{S}$  be the function  $\pi_2 \circ p$

$\langle 1 \rangle 6$ . PICK a sequence  $t_n$  in  $(b, 1]$  such that  $t_n \rightarrow b$  and  $y(t_n) = (-1)^n$  for all  $n$

$\langle 2 \rangle 1$ . LET:  $n \geq 1$

$\langle 2 \rangle 2$ . PICK  $u$  with  $0 < u < x(1/n)$  and  $\sin(1/u) = (-1)^n$

$\langle 2 \rangle 3$ . PICK  $t_n$  with  $b < t_n < 1/n$  and  $x(t_n) = u$

PROOF: By the Intermediate Value Theorem

$\langle 1 \rangle 7$ . Q.E.D.

PROOF: This contradicts Proposition 105 since  $y$  is continuous and  $y(t_n)$  does not converge.

$\square$

**Corollary 256.1.** *The closure of a path-connected subspace of a space is not necessarily path-connected.*

## 37 The Long Line

**Definition 257** (The Long Line). The *long line* is the space  $\omega_1 \times [0, 1]$  in the dictionary order under the order topology, where  $\omega_1$  is the first uncountable ordinal.

**Lemma 258.** *For any ordinal  $\alpha$  with  $0 < \alpha < \omega_1$  we have  $[(0, 0), (\alpha, 0)) \cong [0, 1]$*

$\langle 1 \rangle 1$ .  $[(0, 0), (1, 0)) \cong [0, 1]$

PROOF: The map  $\pi_2$  is a homeomorphism.

$\langle 1 \rangle 2$ . If  $[(0, 0), (\alpha, 0)) \cong [0, 1]$  then  $[(0, 0), (\alpha + 1, 0)) \cong [0, 1]$

PROOF: Proposition 14.

$\langle 1 \rangle 3$ . If  $\lambda$  is a limit ordinal with  $\lambda < \omega_1$  and  $[(0, 0), (\alpha, 0)) \cong [0, 1]$  for all  $\alpha$  with  $0 < \alpha < \lambda$  then  $[(0, 0), (\lambda, 0)) \cong [0, 1]$

$\langle 2 \rangle 1$ . LET:  $\lambda$  be a limit ordinal  $< \omega_1$

$\langle 2 \rangle 2$ . ASSUME:  $[(0, 0), (\alpha, 0)) \cong [0, 1]$  for all  $\alpha$  with  $0 < \alpha < \lambda$

$\langle 2 \rangle 3$ . PICK a sequence of ordinals  $\alpha_0 < \alpha_1 < \dots$  with limit  $\lambda$

PROOF: Since  $\lambda$  is countable.

$\langle 2 \rangle 4.$   $[(\alpha_i, 0), (\alpha_{i+1}, 0)) \cong [0, 1]$  for all  $i$

PROOF: Lemma 13.

$\langle 2 \rangle 5.$  Q.E.D.

PROOF: By Proposition 15.

$\langle 1 \rangle 4.$  Q.E.D.

PROOF: By transfinite induction.

**Proposition 259 (CC).** *The long line is path-connected.*

PROOF:

$\langle 1 \rangle 1.$  LET:  $(\alpha, i), (\beta, j) \in \omega_1 \times [0, 1]$

$\langle 1 \rangle 2.$  ASSUME: without loss of generality  $(\alpha, i) < (\beta, j)$

$\langle 1 \rangle 3.$   $[(0, 0), (\beta + 1, 0)) \cong [0, 1]$

PROOF: By Lemma 258

$\langle 1 \rangle 4.$   $[(\alpha, i), (\beta, j)) \cong [0, 1]$

PROOF: Lemma 13.

$\langle 1 \rangle 5.$  PICK a homeomorphism  $q : [0, 1] \rightarrow [(\alpha, i), (\beta, j))$

$\langle 1 \rangle 6.$   $q \cup \{(1, (\beta, j))\}$  is a path from  $(\alpha, i)$  to  $(\beta, j)$

□

**Proposition 260.** *Every point in the long line has a neighbourhood homeomorphic to an interval in  $\mathbb{R}$ .*

PROOF: For any  $(\alpha, i)$  in the long line, the neighbourhood  $[(0, 0), (\alpha + 1, 0))$  satisfies the condition by Lemma 258.

**Proposition 261.** *The long line  $L$  is not second countable.*

PROOF:

$\langle 1 \rangle 1.$  LET:  $\mathcal{B}$  be a basis for  $L$ .

$\langle 1 \rangle 2.$  For  $\alpha < \omega_1$ , PICK  $B_\alpha \in \mathcal{B}$  such that  $(\alpha, 1/2) \in B_\alpha \subseteq ((\alpha, 0), (\alpha + 1, 0))$

$\langle 1 \rangle 3.$   $\mathcal{B}$  is uncountable.

PROOF: The mapping  $\alpha \mapsto B_\alpha$  is an injection  $\omega_1 \rightarrow \mathcal{B}$ .

**Corollary 261.1.** *The long line cannot be imbedded into  $\mathbb{R}^n$  for any  $n$ .*

## 38 Components

**Proposition 262.** *Let  $X$  be a topological space. Define the relation  $\sim$  on  $X$  by  $a \sim b$  if and only if there exists a connected subspace  $A \subseteq X$  with  $a, b \in A$ . Then  $\sim$  is an equivalence relation on  $X$ .*

PROOF:

$\langle 1 \rangle 1.$   $\sim$  is reflexive.

PROOF: For any  $a \in X$  we have  $\{a\}$  is a connected subspace that contains  $a$ .

$\langle 1 \rangle 2.$   $\sim$  is symmetric.

PROOF: Trivial.

$\langle 1 \rangle 3.$   $\sim$  is transitive.

- ⟨2⟩1. LET:  $a, b, c \in X$
  - ⟨2⟩2. ASSUME:  $a \sim b$  and  $b \sim c$
  - ⟨2⟩3. PICK connected subspaces  $A$  and  $B$  with  $a, b \in A$  and  $b, c \in B$
  - ⟨2⟩4.  $A \cup B$  is a connected subspace that contains  $a$  and  $c$
- PROOF: Theorem 220.

□

**Definition 263** ((Connected) Component). Let  $X$  be a topological space. The *(connected) components* of  $X$  are the equivalence classes under the above  $\sim$ .

**Lemma 264.** *Let  $X$  be a topological space. If  $A \subseteq X$  is connected and nonempty then there exists a unique component  $C$  of  $X$  such that  $A \subseteq C$ .*

PROOF:

- ⟨1⟩1. PICK  $a \in A$
  - ⟨1⟩2. LET:  $C$  be the  $\sim$ -equivalence class of  $a$ .
  - ⟨1⟩3.  $A \subseteq C$
- PROOF: For all  $x \in A$  we have  $x \sim a$ .
- ⟨1⟩4. If  $C'$  is a component and  $A \subseteq C'$  then  $C = C'$
- PROOF: Since we have  $a \in C'$ .

□

**Theorem 265.** *Let  $X$  be a topological space. The components of  $X$  are connected disjoint subspaces of  $X$  whose union is  $X$  such that each nonempty connected subspace of  $X$  intersects only one of them.*

PROOF:

- ⟨1⟩1. Every component of  $X$  is connected.
- PROOF: For  $a \in X$ , the  $\sim$ -equivalence class of  $a$  is  $\bigcup\{A \subseteq X \mid A \text{ is connected, } a \in A\}$  which is connected by Theorem 220.
- ⟨1⟩2. The components form a partition of  $X$ .
- PROOF: Immediate from the definition.
- ⟨1⟩3. Every nonempty connected subspace of  $X$  intersects a unique component of  $X$ .
- ⟨2⟩1. LET:  $A \subseteq X$  be connected and nonempty.
  - ⟨2⟩2. LET:  $C$  be the component such that  $A \subseteq C$
- PROOF: Lemma 264.
- ⟨2⟩3.  $A$  intersects  $C$
  - ⟨2⟩4. If  $A$  intersects the component  $C'$  then  $C' = C$
- ⟨3⟩1. LET:  $C'$  be a component that intersects  $A$
  - ⟨3⟩2. PICK  $b \in A \cap C'$
  - ⟨3⟩3.  $A \subseteq C'$
- PROOF: For all  $x \in A$  we have  $x \sim b$ .
- ⟨3⟩4.  $C = C'$
- PROOF: By uniqueness in ⟨2⟩2.

□

**Proposition 266.** *Every component of a space is closed.*

PROOF:

$\langle 1 \rangle 1$ . LET:  $X$  be a topological space and  $C$  a component of  $X$ .

$\langle 1 \rangle 2$ .  $\overline{C}$  is connected.

PROOF: Theorem 221.

$\langle 1 \rangle 3$ .  $C = \overline{C}$

PROOF: Lemma 219.

$\langle 1 \rangle 4$ .  $C$  is closed.

PROOF: Lemma 47.

□

**Proposition 267.** *If a topological space has finitely many components then every component is open.*

PROOF: Each component is the complement of a finite union of closed sets. □

## 39 Path Components

**Proposition 268.** *Let  $X$  be a topological space. Define the relation  $\sim$  on  $X$  by:  $a \sim b$  if and only if there exists a path in  $X$  from  $a$  to  $b$ . Then  $\sim$  is an equivalence relation on  $X$ .*

PROOF:

$\langle 1 \rangle 1$ .  $\sim$  is reflexive.

PROOF: For  $a \in X$ , the constant function  $[0, 1] \rightarrow X$  with value  $a$  is a path from  $a$  to  $a$ .

$\langle 1 \rangle 2$ .  $\sim$  is symmetric.

PROOF: If  $p : [0, 1] \rightarrow X$  is a path from  $a$  to  $b$ , then  $\lambda t.p(1-t)$  is a path from  $b$  to  $a$ .

$\langle 1 \rangle 3$ .  $\sim$  is transitive.

PROOF: Concatenate paths.

□

**Definition 269** (Path Component). Let  $X$  be a topological space. The *path components* of  $X$  are the equivalence relations under  $\sim$ .

**Theorem 270.** *The path components of  $X$  are path-connected disjoint subspaces of  $X$  whose union is  $X$  such that every nonempty path-connected subspace of  $X$  intersects exactly one path component.*

PROOF:

$\langle 1 \rangle 1$ . Every path component is path-connected.

PROOF: If  $a$  and  $b$  are in the same path component then  $a \sim b$ , i.e. there exists a path from  $a$  to  $b$ .

$\langle 1 \rangle 2$ . The path components are disjoint and their union is  $X$ .

PROOF: Immediate from the definition.

- ⟨1⟩3. Every non-empty path-connected subspace of  $X$  intersects exactly one path component.
- ⟨2⟩1. LET:  $A$  be a nonempty path-connected subspace of  $X$ .
- ⟨2⟩2. PICK  $a \in A$
- ⟨2⟩3.  $A$  intersects the  $\sim$ -equivalence class of  $a$ .
- ⟨2⟩4. LET:  $C$  be any path component that intersects  $A$ .
- ⟨2⟩5. PICK  $b \in A \cap C$
- ⟨2⟩6.  $a \sim b$
- PROOF: Since  $A$  is path-connected.
- ⟨2⟩7.  $C$  is the  $\sim$ -equivalence class of  $a$ .

□

**Proposition 271.** *Every path component is included in a component.*

PROOF:

- ⟨1⟩1. LET:  $X$  be a topological space and  $C$  a path component of  $X$ .
- ⟨1⟩2.  $C$  is path-connected.
- PROOF: Theorem 270.
- ⟨1⟩3.  $C$  is connected.
- PROOF: Proposition 241.
- ⟨1⟩4.  $C$  is included in a component.
- PROOF: Lemma 264.

□

## 40 Local Connectedness

**Definition 272** (Locally Connected). Let  $X$  be a topological space and  $a \in X$ . Then  $X$  is *locally connected* at  $a$  if and only if every neighbourhood of  $a$  includes a connected neighbourhood of  $a$ .

The space  $X$  is *locally connected* if and only if it is locally connected at every point.

**Example 273.** The real line is both connected and locally connected.

**Example 274.** The space  $\mathbb{R} \setminus \{0\}$  is disconnected but locally connected.

**Example 275.** The topologist's sine curve is connected but not locally connected.

**Example 276.** The rationals  $\mathbb{Q}$  are neither connected nor locally connected.

**Theorem 277.** *A topological space  $X$  is locally connected if and only if, for every open set  $U$  in  $X$ , every component of  $U$  is open in  $X$ .*

PROOF:

- ⟨1⟩1. If  $X$  is locally connected then, for every open set  $U$  in  $X$ , every component of  $U$  is open in  $X$ .
- ⟨2⟩1. ASSUME:  $X$  is locally connected.

- ⟨2⟩2. LET:  $U$  be open in  $X$ .
- ⟨2⟩3. LET:  $C$  be a component of  $U$ .
- ⟨2⟩4. LET:  $a \in C$
- ⟨2⟩5. LET:  $V$  be a connected neighbourhood of  $a$  such that  $V \subseteq U$
- ⟨2⟩6.  $V \subseteq C$

PROOF: Lemma 264.

- ⟨2⟩7. Q.E.D.

PROOF: Lemma 26.

- ⟨1⟩2. If, for every open set  $U$  in  $X$ , every component of  $U$  is open in  $X$ , then  $X$  is locally connected.
- ⟨2⟩1. ASSUME: for every open set  $U$  in  $X$ , every component of  $U$  is open in  $X$ .
- ⟨2⟩2. LET:  $a \in X$
- ⟨2⟩3. LET:  $U$  be a neighbourhood of  $a$
- ⟨2⟩4. The component of  $U$  that contains  $a$  is a connected neighbourhood of  $a$  included in  $U$ .

□

**Example 278.** The ordered square is locally connected.

PROOF: Every neighbourhood of a point includes an interval around that point, which is connected by Theorem 232.

**Example 279.** Let  $X$  be the set of all rational points on the line segment  $[0, 1] \times \{0\}$ , and  $Y$  the set of all rational points on the line segment  $[0, 1] \times \{1\}$ . Let  $A$  be the space consisting of all line segments joining the point  $(0, 1)$  to a point of  $X$ , and all line segments joining the point  $(1, 0)$  to a point of  $Y$ . Then  $A$  is path-connected but is not locally connected at any point,

**Proposition 280.** Let  $X$  and  $Y$  be topological spaces and  $p : X \twoheadrightarrow Y$  be a quotient map. If  $X$  is locally connected then so is  $Y$ .

PROOF:

- ⟨1⟩1. LET:  $U$  be an open set in  $Y$ .
- ⟨1⟩2. LET:  $C$  be a component of  $U$ .
- ⟨1⟩3.  $p^{-1}(C)$  is a union of components of  $p^{-1}(U)$ 
  - ⟨2⟩1. LET:  $x \in p^{-1}(C)$
  - ⟨2⟩2. LET:  $D$  be the component of  $p^{-1}(U)$  that contains  $x$ .
  - ⟨2⟩3.  $p(D)$  is connected.

PROOF: Theorem 222.

- ⟨2⟩4.  $p(D) \subseteq C$ .

PROOF: From ⟨1⟩2 since  $p(x) \in p(D) \cap C$  (⟨2⟩1, ⟨2⟩2).

- ⟨2⟩5.  $D \subseteq p^{-1}(C)$

- ⟨1⟩4.  $p^{-1}(C)$  is open in  $p^{-1}(U)$

PROOF: Theorem 277.

- ⟨1⟩5.  $C$  is open in  $U$

PROOF: Since the restriction of  $p$  to  $p : p^{-1}(U) \twoheadrightarrow U$  is a quotient map by Proposition 195.



⟨1⟩6. Q.E.D.

PROOF: Theorem 277.

□

## 41 Local Path Connectedness

**Definition 281** (Locally Path-Connected). Let  $X$  be a topological space and  $a \in X$ . Then  $X$  is *locally path-connected* at  $a$  if and only if every neighbourhood of  $a$  includes a path-connected neighbourhood of  $a$ .

The space  $X$  is *locally path-connected* if and only if it is locally path-connected at every point.

**Theorem 282.** *A topological space  $X$  is locally path-connected if and only if, for every open set  $U$  in  $X$ , every path component of  $U$  is open in  $X$ .*

PROOF:

⟨1⟩1. If  $X$  is locally path-connected then, for every open set  $U$  in  $X$ , every path component of  $U$  is open in  $X$ .

⟨2⟩1. ASSUME:  $X$  is locally path-connected.

⟨2⟩2. LET:  $U$  be open in  $X$ .

⟨2⟩3. LET:  $C$  be a path component of  $U$ .

⟨2⟩4. LET:  $a \in C$

⟨2⟩5. LET:  $V$  be a path-connected neighbourhood of  $a$  such that  $V \subseteq U$

⟨2⟩6.  $V \subseteq C$

PROOF: Lemma 264.

⟨2⟩7. Q.E.D.

PROOF: Lemma 26.

⟨1⟩2. If, for every open set  $U$  in  $X$ , every component of  $U$  is open in  $X$ , then  $X$  is locally connected.

⟨2⟩1. ASSUME: for every open set  $U$  in  $X$ , every component of  $U$  is open in  $X$ .

⟨2⟩2. LET:  $a \in X$

⟨2⟩3. LET:  $U$  be a neighbourhood of  $a$

⟨2⟩4. The component of  $U$  that contains  $a$  is a connected neighbourhood of  $a$  included in  $U$ .

□

**Theorem 283.** *If a space is locally path connected then its components and its path components are the same.*

PROOF:

⟨1⟩1. LET:  $X$  be a locally path connected space.

⟨1⟩2. LET:  $C$  be a component of  $X$ .

⟨1⟩3. LET:  $x \in C$

⟨1⟩4. LET:  $P$  be the path component of  $x$

PROVE:  $P = C$

⟨1⟩5.  $P \subseteq C$

PROOF: Proposition 271.

⟨1⟩6. LET:  $Q$  be the union of the other path components included in  $C$

⟨1⟩7.  $C = P \cup Q$

PROOF: Proposition 271.

⟨1⟩8.  $P$  and  $Q$  are open in  $C$

⟨2⟩1.  $C$  is open.

PROOF: Theorem 277.

⟨2⟩2. Q.E.D.

PROOF: Theorem 282.

⟨1⟩9.  $Q = \emptyset$

PROOF: Otherwise  $P$  and  $Q$  would form a separation of  $C$ .

□

**Example 284.** The ordered square is not locally path connected, since it is connected but not path connected.

**Proposition 285.** *Let  $X$  be a locally path-connected space. Then every connected open subspace of  $X$  is path-connected.*

PROOF:

⟨1⟩1. LET:  $U$  be a connected open subspace of  $X$ .

⟨1⟩2. LET:  $P$  be a path component of  $U$ .

⟨1⟩3. LET:  $Q$  be the union of the other path components of  $U$ .

⟨1⟩4.  $P$  and  $Q$  are open in  $U$ .

PROOF: Theorem 282.

⟨1⟩5.  $Q = \emptyset$

PROOF: Otherwise  $P$  and  $Q$  form a separation of  $U$ .

□

## 42 Weak Local Connectedness

**Definition 286** (Weakly Locally Connected). Let  $X$  be a topological space and  $a \in X$ . Then  $X$  is *weakly locally connected* at  $a$  if and only if every neighbourhood of  $a$  includes a connected subspace that includes a neighbourhood of  $a$ .

**Proposition 287.** *Let  $X$  be a topological space. If  $X$  is weakly locally connected at every point then  $X$  is locally connected.*

PROOF:

⟨1⟩1. ASSUME:  $X$  is weakly locally connected at every point.

⟨1⟩2. LET:  $U$  be open in  $X$ .

⟨1⟩3. LET:  $C$  be a component of  $U$ .

⟨1⟩4.  $C$  is open in  $X$ .

⟨2⟩1. LET:  $x \in C$

⟨2⟩2. PICK a connected subspace  $D$  of  $U$  that includes a neighbourhood  $V$  of  $x$ .

⟨2⟩3.  $D \subseteq C$

PROOF: Lemma 264.

⟨2⟩4.  $x \in V \subseteq C$

⟨2⟩5. Q.E.D.

PROOF: Lemma 26.

⟨1⟩5. Q.E.D.

PROOF: Theorem 277.

□

**Example 288.** The 'infinite broom' (Munkres p.163) is an example of a space that is weakly locally connected at a point  $p$  but not locally connected at  $p$ .

## 43 Quasicomponents

**Proposition 289.** Let  $X$  be a topological space. Define  $\sim$  on  $X$  by  $x \sim y$  if and only if there exists no separation  $U$  and  $V$  of  $X$  such that  $x \in U$  and  $y \in V$ . Then  $\sim$  is an equivalence relation on  $X$ .

PROOF:

⟨1⟩1.  $\sim$  is reflexive.

PROOF: Since the two sets that make up a separation are disjoint.

⟨1⟩2.  $\sim$  is symmetric.

PROOF: Immediate from the definition.

⟨1⟩3.  $\sim$  is transitive.

⟨2⟩1. ASSUME:  $x \sim y$  and  $y \sim z$

⟨2⟩2. ASSUME: for a contradiction there is a separation  $U$  and  $V$  of  $X$  with  
 $x \in U$  and  $z \in V$

⟨2⟩3.  $y \in U$  or  $y \in V$

⟨2⟩4. Q.E.D.

PROOF: Either case contradicts ⟨2⟩1.

□

**Definition 290** (Quasicomponents). For  $X$  a topological space, the *quasicomponents* of  $X$  are the equivalence classes under  $\sim$ .

**Proposition 291.** Let  $X$  be a topological space. Then every component of  $X$  is included in a quasicomponent of  $X$ .

PROOF:

⟨1⟩1. LET:  $C$  be a component of  $X$ .

⟨1⟩2. LET:  $x, y \in C$

PROVE:  $x \sim y$

⟨1⟩3. ASSUME: for a contradiction there exists a separation  $U$  and  $V$  of  $X$  with  
 $x \in U$  and  $y \in V$

⟨1⟩4.  $C \cap U$  and  $C \cap V$  form a separation of  $C$ .

⟨1⟩5. Q.E.D.

PROOF: This contradicts ⟨1⟩1.

**Proposition 292.** *In a locally connected space, the components and the quasi-components are the same.*

PROOF:

⟨1⟩1. LET:  $X$  be a locally connected space and  $Q$  a quasicomponent of  $X$ .

⟨1⟩2. PICK a component  $C$  of  $X$  such that  $C \subseteq Q$

⟨1⟩3. LET:  $D$  be the union of the components of  $X$

⟨1⟩4.  $C$  and  $D$  are open in  $X$ .

PROOF: Theorem 277.

⟨1⟩5.  $D$  cannot contain any points of  $Q$ .

PROOF: If it did, then  $C$  and  $D$  would form a separation of  $X$  and there would be points  $x, y \in Q$  with  $x \in C$  and  $y \in D$ .

⟨1⟩6.  $C = Q$

□

## 44 Open Coverings

**Definition 293** (Open Covering). Let  $X$  be a topological space. An *open covering* of  $X$  is a covering of  $X$  whose elements are all open sets.

## 45 Compact Spaces

**Definition 294** (Compact). A topological space is *compact* if and only if every open covering has a finite subcovering.

**Lemma 295.** *Let  $X$  be a topological space and  $Y$  a subspace of  $X$ . Then  $Y$  is compact if and only if every covering of  $Y$  by sets open in  $X$  has a finite subcovering.*

PROOF:

⟨1⟩1. If  $Y$  is compact then every covering of  $Y$  by sets open in  $X$  has a finite subcovering.

⟨2⟩1. ASSUME:  $Y$  is compact.

⟨2⟩2. LET:  $\mathcal{U}$  be a covering of  $Y$  by sets open in  $X$ .

⟨2⟩3.  $\{U \cap Y \mid U \in \mathcal{U}\}$  is an open covering of  $Y$ .

⟨2⟩4. PICK a finite subcovering  $\{U_1 \cap Y, \dots, U_n \cap Y\}$

⟨2⟩5.  $\{U_1, \dots, U_n\}$  is a finite subcovering of  $\mathcal{U}$ .

⟨1⟩2. If every covering of  $Y$  by sets open in  $X$  has a finite subcovering then  $Y$  is compact.

⟨2⟩1. LET:  $\mathcal{U}$  be an open covering of  $Y$ .

⟨2⟩2. LET:  $\mathcal{V} = \{V \text{ open in } X \mid V \cap Y \in \mathcal{U}\}$ .

⟨2⟩3.  $\mathcal{V}$  is a covering of  $Y$  by sets open in  $X$ .

⟨2⟩4. PICK a finite subcovering  $\{V_1, \dots, V_n\}$

⟨2⟩5.  $\{V_1 \cap Y, \dots, V_n \cap Y\}$  is a finite subcovering of  $\mathcal{U}$ .

□

**Proposition 296.** *Every closed subspace of a compact space is compact.*

PROOF:

- <1>1. LET:  $X$  be a compact space and  $Y \subseteq X$  be closed.
- <1>2. LET:  $\mathcal{U}$  be a covering of  $Y$  by sets open in  $X$ .
- <1>3.  $\mathcal{U} \cup \{X \setminus Y\}$  is an open covering of  $X$ .
- <1>4. PICK a finite subcovering  $\mathcal{U}_0$
- <1>5.  $\mathcal{U}_0 \cap \mathcal{U}$  is a finite subset of  $\mathcal{U}$  that covers  $Y$ .

□

**Theorem 297.** *The continuous image of a compact space is compact.*

PROOF:

- <1>1. LET:  $f : X \rightarrow Y$  be continuous and surjective.
- <1>2. LET:  $\mathcal{V}$  be an open covering of  $Y$
- <1>3.  $\{f^{-1}(V) \mid V \in \mathcal{V}\}$  is an open covering of  $X$ .
- <1>4. PICK a finite subcovering  $\{f^{-1}(V_1), \dots, f^{-1}(V_n)\}$
- <1>5.  $\{V_1, \dots, V_n\}$  covers  $Y$ .

□

**Theorem 298.** *Let  $A$  and  $B$  be compact subspaces of  $X$  and  $Y$  respectively. Let  $N$  be an open set in  $X \times Y$  that includes  $A \times B$ . Then there exist open sets  $U$  and  $V$  in  $X$  and  $Y$  respectively such that  $A \subseteq U$ ,  $B \subseteq V$  and  $U \times V \subseteq N$ .*

PROOF:

- <1>1. For all  $x \in A$ , there exist neighbourhoods  $U$  of  $x$  and  $V$  of  $B$  such that  $U \times V \subseteq N$ .
- <2>1. LET:  $x \in A$
- <2>2. For all  $y \in B$ , there exist neighbourhoods  $U$  of  $x$  and  $V$  of  $y$  such that  $U \times V \subseteq N$
- <2>3.  $\{V \text{ open in } Y \mid \exists \text{ neighbourhood } U \text{ of } x, U \times V \subseteq N\}$  covers  $B$ .
- <2>4. PICK a finite subcover  $\{V_1, \dots, V_n\}$
- <2>5. For  $i = 1, \dots, n$ , PICK a neighbourhood  $U_i$  of  $x$  such that  $U_i \times V_i \subseteq N$
- <2>6. LET:  $U = U_1 \cap \dots \cap U_n$
- <2>7. LET:  $V = V_1 \cup \dots \cup V_n$
- <2>8.  $U$  is a neighbourhood of  $x$ .
- <2>9.  $V$  is a neighbourhood of  $B$ .
- <2>10.  $U \times V \subseteq N$
- <1>2.  $\{U \text{ open in } X \mid \exists \text{ neighbourhood } V \text{ of } B, U \times V \subseteq N\}$  covers  $A$ .
- <1>3. PICK a finite subcover  $\{U_1, \dots, U_n\}$
- <1>4. For  $i = 1, \dots, n$ , PICK a neighbourhood  $V_i$  of  $B$  such that  $U_i \times V_i \subseteq N$
- <1>5. LET:  $U = U_1 \cup \dots \cup U_n$
- <1>6. LET:  $V = V_1 \cap \dots \cap V_n$
- <1>7.  $U$  and  $V$  are open.
- <1>8.  $A \subseteq U$
- <1>9.  $B \subseteq V$
- <1>10.  $U \times V \subseteq N$

□

**Corollary 298.1** (Tube Lemma). *Let  $X$  and  $Y$  be topological spaces with  $Y$  compact. Let  $a \in X$  and  $N$  be an open set in  $X \times Y$  that includes  $\{a\} \times Y$ . Then there exists a neighbourhood  $W$  of  $a$  such that  $N$  includes the tube  $W \times Y$ .*

**Theorem 299.** *Let  $X$  be a topological space. Then  $X$  is compact if and only if every set of closed sets that has the finite intersection property has nonempty intersection.*

PROOF: The following are equivalent.

1.  $X$  is compact.
2. Every open covering of  $X$  has a finite subcovering.
3. For any set  $\mathcal{C}$  of closed sets, if  $\{X \setminus C \mid C \in \mathcal{C}\}$  covers  $X$  then there is a finite subset  $\mathcal{C}_0$  such that  $\{X \setminus C \mid C \in \mathcal{C}_0\}$  covers  $X$
4. For any set  $\mathcal{C}$  of closed sets, if  $\bigcap \mathcal{C} = \emptyset$  then there is a finite subset  $\mathcal{C}_0$  with empty intersection.
5. Any set of closed sets with the finite intersection property has nonempty intersection.

□

**Corollary 299.1.** *Let  $X$  be a topological space and  $C_1 \supseteq C_2 \supseteq \dots$  a nested sequence of nonempty closed sets. Then  $\bigcap_n C_n$  is nonempty.*

**Proposition 300.** *Let  $\mathcal{T}$  and  $\mathcal{T}'$  be two topologies on the same set  $X$  with  $\mathcal{T} \subseteq \mathcal{T}'$ . If  $\mathcal{T}'$  is compact then  $\mathcal{T}$  is compact.*

PROOF:

- ⟨1⟩1. LET:  $\mathcal{U} \subseteq \mathcal{T}$  cover  $X$   
 ⟨1⟩2.  $\mathcal{U} \subseteq \mathcal{T}'$   
 ⟨1⟩3. A finite subset of  $\mathcal{U}$  covers  $X$ .

□

**Corollary 300.1.** *If  $\mathcal{T}$  and  $\mathcal{T}'$  are two compact Hausdorff topologies on the same set  $X$ , then either  $\mathcal{T} = \mathcal{T}'$  or  $\mathcal{T}$  and  $\mathcal{T}'$  are incomparable.*

PROOF: From the Proposition and Proposition 177. □

**Example 301.** Any set under the finite complement topology is compact.

**Proposition 302.** *Let  $X$  be a topological space. A finite union of compact subspaces of  $X$  is compact.*

PROOF:

- ⟨1⟩1. LET:  $A$  and  $B$  be compact subspaces of  $X$ .  
 ⟨1⟩2. LET:  $\mathcal{U}$  be a set of open sets in  $X$  that covers  $A \cup B$   
 ⟨1⟩3. PICK a finite subset  $\mathcal{U}_1$  that covers  $A$ .

PROOF: Lemma 295.

⟨1⟩4. PICK a finite subset  $\mathcal{U}_2$  that covers  $B$ .

PROOF: Lemma 295.

⟨1⟩5.  $\mathcal{U}_1 \cup \mathcal{U}_2$  is a finite subset that covers  $A \cup B$ .

⟨1⟩6. Q.E.D.

PROOF: Lemma 295.

□

**Proposition 303.** *Let  $A$  and  $B$  be disjoint compact subspaces of the Hausdorff space  $X$ . Then there exist disjoint open sets  $U$  and  $V$  that include  $A$  and  $B$  respectively.*

PROOF: From Theorem 298 with  $N = X^2 \setminus \{(x, x) \mid x \in X\}$ . □

**Corollary 303.1.** *Every compact subspace of a Hausdorff space is closed.*

**Theorem 304.** *Let  $f : X \rightarrow Y$  be a bijective continuous function. If  $X$  is compact and  $Y$  is Hausdorff then  $f$  is a homeomorphism.*

PROOF:

⟨1⟩1. LET:  $C \subseteq X$  be closed.

⟨1⟩2.  $C$  is compact.

PROOF: Proposition 296.

⟨1⟩3.  $f(C)$  is compact.

PROOF: Theorem 297.

⟨1⟩4.  $f(C)$  is closed.

PROOF: Corollary 303.1.

⟨1⟩5. Q.E.D.

PROOF: Lemma 107.

□

**Proposition 305.** *Let  $X$  be a compact space,  $Y$  a Hausdorff space, and  $f : X \rightarrow Y$  a continuous map. Then  $f$  is a closed map.*

PROOF:

⟨1⟩1. LET:  $C \subseteq X$  be closed.

⟨1⟩2.  $C$  is compact.

PROOF: Proposition 296.

⟨1⟩3.  $f(C)$  is compact.

PROOF: Theorem 297.

⟨1⟩4.  $f(C)$  is closed.

PROOF: Corollary 303.1.

□

**Proposition 306.** *If  $Y$  is compact then the projection  $\pi_1 : X \times Y \rightarrow X$  is a closed map.*

PROOF:

⟨1⟩1. LET:  $A \subseteq X \times Y$  be closed.

⟨1⟩2. LET:  $x \in X \setminus \pi_1(A)$

⟨1⟩3. PICK a neighbourhood  $U$  of  $x$  such that  $U \times Y \subseteq (X \times Y) \setminus A$

PROOF: By the Tube Lemma.

⟨1⟩4.  $x \in U \subseteq X \setminus \pi_1(A)$

⟨1⟩5. Q.E.D.

PROOF: So  $X \setminus \pi_1(A)$  is open by Lemma 26.

□

**Theorem 307.** *Let  $X$  be a topological space and  $Y$  a compact Hausdorff space. Let  $f : X \rightarrow Y$  be a function. Then  $f$  is continuous if and only if the graph of  $f$  is closed in  $X \times Y$ .*

PROOF:

⟨1⟩1. LET:  $G_f$  be the graph of  $f$ .

⟨1⟩2. If  $f$  is continuous then  $G_f$  is closed.

⟨2⟩1. ASSUME:  $f$  is continuous.

⟨2⟩2. LET:  $(x, y) \in (X \times Y) \setminus G_f$

⟨2⟩3. PICK disjoint neighbourhoods  $U$  and  $V$  of  $y$  and  $f(x)$  respectively.

⟨2⟩4.  $f^{-1}(V) \times U$  is a neighbourhood of  $(x, y)$  disjoint from  $G_f$ .

⟨1⟩3. If  $G_f$  is closed then  $f$  is continuous.

⟨2⟩1. ASSUME:  $G_f$  is closed.

⟨2⟩2. LET:  $x \in X$  and  $V$  be a neighbourhood of  $f(x)$ .

⟨2⟩3.  $G_f \cap (X \times (Y \setminus V))$  is closed.

⟨2⟩4.  $\pi_1(G_f \cap (X \times (Y \setminus V)))$  is closed.

PROOF: Proposition 306.

⟨2⟩5. LET:  $U = X \setminus \pi_1(G_f \cap (X \times (Y \setminus V)))$

⟨2⟩6.  $U$  is a neighbourhood of  $x$

⟨2⟩7.  $f(U) \subseteq V$

□

**Theorem 308.** *Let  $X$  be a compact topological space. Let  $(f_n : X \rightarrow \mathbb{R})$  be a monotone increasing sequence of continuous functions and  $f : X \rightarrow \mathbb{R}$  a continuous function. If  $(f_n)$  converges pointwise to  $f$ , then  $(f_n)$  converges uniformly to  $f$ .*

PROOF:

⟨1⟩1. LET:  $\epsilon > 0$

⟨1⟩2. For all  $x \in X$ , there exists  $N$  such that, for all  $n \geq N$ , we have  $|f_n(x) - f(x)| < \epsilon$

⟨1⟩3. For  $n \geq 1$ ,

LET:  $U_n = \{x \in X \mid |f_n(x) - f(x)| < \epsilon\}$

⟨1⟩4. For  $n \geq 1$ , we have  $U_n$  is open in  $X$ .

⟨2⟩1. LET:  $x \in X$

⟨2⟩2. LET:  $\delta = \epsilon - |f_n(x) - f(x)|$

⟨2⟩3. PICK a neighbourhood  $U$  of  $x$  such that  $f(U) \subseteq B(f(x), \delta/2)$

⟨2⟩4. PICK a neighbourhood  $V$  of  $x$  such that  $f_n(V) \subseteq B(f_n(x), \delta/2)$

⟨2⟩5.  $f(U \cap V) \subseteq U_n$



PROOF: For  $y \in U \cap V$  we have

$$\begin{aligned} |f_n(y) - f(y)| &\leq |f_n(y) - f_n(x)| + |f_n(x) - f(x)| + |f(x) - y(y)| \\ &< \delta/2 + |f_n(x) - f(x)| + \delta/2 \\ &= \epsilon \end{aligned}$$

$\langle 1 \rangle 5$ .  $\{U_n \mid n \geq 1\}$  covers  $X$

PROOF: From  $\langle 1 \rangle 2$

$\langle 1 \rangle 6$ . PICK  $N$  such that  $X = U_N$

$\langle 2 \rangle 1$ . PICK  $n_1, \dots, n_k$  such that  $U_{n_1}, \dots, U_{n_k}$  cover  $X$ .

$\langle 2 \rangle 2$ . LET:  $N = \max(n_1, \dots, n_k)$

$\langle 2 \rangle 3$ . For all  $i$  we have  $U_{n_i} \subseteq U_N$

PROOF: Since  $(f_n)$  is monotone increasing.

$\langle 2 \rangle 4$ .  $X = U_N$

$\langle 1 \rangle 7$ . For all  $x \in X$  and  $n \geq N$  we have  $|f_n(x) - f(x)| < \epsilon$

□

An example to show that we cannot remove the hypothesis that  $X$  is compact:

**Example 309.** Let  $X = (0, 1)$ ,  $f_n(x) = -x^n$  and  $f(x) = 0$  for  $x \in X$  and  $n \geq 1$ . Then  $f_n \rightarrow f$  pointwise and  $(f_n)$  is monotone increasing but the convergence is not uniform since, for all  $N \geq 1$ , there exists  $x \in (0, 1)$  such that  $-x^N < -1/2$ .

An example to show that we cannot remove the hypothesis that  $(f_n)$  is monotone increasing:

**Example 310.** Let  $X = [0, 1]$ ,  $f_n(x) = 1/(n^3(x - 1/n)^2 + 1)$  and  $f(x) = 0$  for  $x \in X$  and  $n \geq 1$ . Then  $X$  is compact and  $f_n \rightarrow f$  pointwise but the convergence is not uniform since, for all  $N \geq 1$ , there exists  $x \in [0, 1]$  such that  $f_N(x) = 1$ , namely  $x = 1/N$ .

**Theorem 311.** Let  $X$  be a compact Hausdorff space. Let  $\mathcal{A}$  be a chain of closed connected subsets of  $X$ . Then  $\bigcap \mathcal{A}$  is connected.

PROOF:

$\langle 1 \rangle 1$ . ASSUME: for a contradiction  $C$  and  $D$  form a separation of  $\bigcap \mathcal{A}$ .

$\langle 1 \rangle 2$ . PICK disjoint open sets  $U$  and  $V$  that include  $C$  and  $D$  respectively.

PROOF: Proposition 303.

$\langle 1 \rangle 3$ .  $\{A \setminus (U \cup V) \mid A \in \mathcal{A}\}$  is a set of closed sets with the finite intersection property.

$\langle 2 \rangle 1$ . For all  $A \in \mathcal{A}$  we have  $A \setminus (U \cup V)$  is closed.

$\langle 2 \rangle 2$ . For all  $A_1, \dots, A_n \in \mathcal{A}$  we have  $(A_1 \cap \dots \cap A_n) \setminus (U \cup V)$  is nonempty.

PROOF:

$\langle 3 \rangle 1$ . LET:  $A_1, \dots, A_n \in \mathcal{A}$

$\langle 3 \rangle 2$ . ASSUME: without loss of generality  $A_1 \subseteq A_2, \dots, A_n$

PROOF: Since  $\mathcal{A}$  is a chain.

$\langle 3 \rangle 3$ .  $A_1 \setminus (U \cup V)$  is nonempty

PROOF: Otherwise  $(A_1 \cap \dots \cap A_n \cap U)$  and  $(A_1 \cap \dots \cap A_n \cap V)$  would form a separation of  $A_n$ .

⟨1⟩4.  $\bigcap \mathcal{A} \setminus (U \cup V)$  is nonempty.

PROOF: Theorem 299.

⟨1⟩5. Q.E.D.

PROOF: This contradicts ⟨1⟩1 since  $\bigcap \mathcal{A} \setminus (U \cup V) = \bigcap \mathcal{A} \setminus (C \cup D)$ .

□

## 46 Perfect Maps

**Definition 312** (Perfect Map). Let  $X$  and  $Y$  be topological spaces and  $f : X \rightarrow Y$ . Then  $f$  is a *perfect map* if and only if  $f$  is a closed map, continuous, surjective and  $p^{-1}(y)$  is compact for all  $y \in Y$ .

**Proposition 313.** Let  $X$  be a topological space,  $Y$  a compact space, and  $p : X \rightarrow Y$  a closed map such that, for all  $y \in Y$ , we have  $p^{-1}(y)$  is compact. Then  $X$  is compact.

PROOF:

⟨1⟩1. LET:  $\mathcal{A}$  be a set of closed sets in  $X$  with the finite intersection property.

⟨1⟩2.  $\mathcal{B} = \{p(A_1 \cap \cdots \cap A_n) \mid A_1, \dots, A_n \in \mathcal{A}\}$  is a set of closed sets in  $Y$  with the finite intersection property.

PROOF: Since  $p$  is a closed map.

⟨1⟩3. PICK  $y \in \bigcap \mathcal{B}$

PROOF: Theorem 299 since  $Y$  is compact.

⟨1⟩4.  $\{A \cap p^{-1}(y) \mid A \in \mathcal{A}\}$  is a set of closed sets in  $p^{-1}(y)$  with the finite intersection property.

⟨1⟩5. PICK  $x \in \bigcap_{A \in \mathcal{A}} (A \cap p^{-1}(y))$

PROOF: Theorem 299 since  $p^{-1}(y)$  is compact.

⟨1⟩6.  $x \in \bigcap \mathcal{A}$

⟨1⟩7. Q.E.D.

PROOF: Theorem 299.

□

## 47 Topological Groups

**Definition 314** (Topological Group). A *topological group*  $G$  consists of a  $T_1$  space  $G$  and continuous maps  $\cdot : G^2 \rightarrow G$  and  $(\ )^{-1} : G \rightarrow G$  such that  $(G, \cdot, (\ )^{-1})$  is a group.

**Example 315.** 1. The integers  $\mathbb{Z}$  under addition are a topological group.

2. The real numbers  $\mathbb{R}$  under addition are a topological group.

3. The positive reals under multiplication are a topological group.

4. The set  $\{z \in \mathbb{C} \mid |z| = 1\}$  under multiplication and given the topology of  $S^1$  is a topological group.

5. For any  $n \geq 0$ , the general linear group  $GL_n(\mathbb{R})$  is a topological group under matrix multiplication, considered as a subspace of  $\mathbb{R}^{n^2}$ .

**Lemma 316.** *Let  $G$  be a  $T_1$  space and  $\cdot : G^2 \rightarrow G$ ,  $(\ )^{-1} : G \rightarrow G$  be functions such that  $(G, \cdot, (\ )^{-1})$  is a group. Then  $G$  is a topological group if and only if the function  $f : G^2 \rightarrow G$  that maps  $(x, y)$  to  $xy^{-1}$  is continuous.*

PROOF:

- $\langle 1 \rangle 1$ . If  $G$  is a topological group then  $f$  is continuous.

PROOF: From Theorem 96.

- $\langle 1 \rangle 2$ . If  $f$  is continuous then  $G$  is a topological group.

- $\langle 2 \rangle 1$ . ASSUME:  $f$  is continuous.

- $\langle 2 \rangle 2$ .  $(\ )^{-1}$  is continuous.

PROOF: Since  $x^{-1} = f(e, x)$ .

- $\langle 2 \rangle 3$ .  $\cdot$  is continuous.

PROOF: Since  $xy = f(x, y^{-1})$ .

□

**Lemma 317.** *Let  $G$  be a topological group and  $H$  a subgroup of  $G$ . Then  $H$  is a topological group under the subspace topology.*

PROOF:

- $\langle 1 \rangle 1$ .  $H$  is  $T_1$ .

PROOF: From Proposition 165.

- $\langle 1 \rangle 2$ . multiplication and inverse on  $H$  are continuous.

PROOF: From Theorem 97.

□

**Lemma 318.** *Let  $G$  be a topological group and  $H$  a subgroup of  $G$ . Then  $\overline{H}$  is a subgroup of  $G$ .*

PROOF:

- $\langle 1 \rangle 1$ . LET:  $x, y \in \overline{H}$

PROVE:  $xy^{-1} \in \overline{H}$

- $\langle 1 \rangle 2$ . LET:  $U$  be any neighbourhood of  $xy^{-1}$

- $\langle 1 \rangle 3$ . LET:  $f : G^2 \rightarrow G$ ,  $f(a, b) = ab^{-1}$

- $\langle 1 \rangle 4$ .  $f^{-1}(U)$  is a neighbourhood of  $(x, y)$

- $\langle 1 \rangle 5$ . PICK neighbourhoods  $V, W$  of  $x$  and  $y$  respectively such that  $f(V \times W) \subseteq U$ .

- $\langle 1 \rangle 6$ . PICK  $a \in V \cap H$  and  $b \in W \cap H$

PROOF: Theorem 48.

- $\langle 1 \rangle 7$ .  $ab^{-1} \in U \cap H$

- $\langle 1 \rangle 8$ . Q.E.D.

PROOF: By Theorem 48.

□

**Proposition 319.** *Let  $G$  be a topological group and  $\alpha \in G$ . Then the maps  $l_\alpha, r_\alpha : G \rightarrow G$  defined by  $l_\alpha(x) = \alpha x$ ,  $r_\alpha(x) = x\alpha$  are homeomorphisms of  $G$  with itself.*

PROOF: They are continuous with continuous inverses  $l_{\alpha^{-1}}$  and  $r_{\alpha^{-1}}$ .  $\square$

**Corollary 319.1.** *Every topological group is homogeneous.*

PROOF: Given a topological group  $G$  and  $a, b \in G$ , we have  $l_{ba^{-1}}$  is a homeomorphism that maps  $a$  to  $b$ .  $\square$

**Proposition 320.** *Let  $G$  be a topological group and  $H$  a subgroup of  $G$ . Give  $G/H$  the quotient topology. For all  $\alpha \in G$ , the map  $\overline{f_\alpha}$  that sends  $xH$  to  $\alpha xH$  is a homeomorphism  $G/H \cong G/H$ .*

PROOF:

$\langle 1 \rangle 1.$   $\overline{f_\alpha}$  is well-defined.

PROOF: If  $xy^{-1} \in H$  then  $(\alpha x)(\alpha y)^{-1} = xy^{-1} \in H$ .

$\langle 1 \rangle 2.$   $\overline{f_\alpha}$  is continuous.

PROOF: Theorem 198 since  $\overline{f_\alpha} \circ p = p \circ f_\alpha$  is continuous, where  $p : G \twoheadrightarrow G/H$  is the canonical surjection.

$\langle 1 \rangle 3.$   $\overline{f_\alpha}^{-1}$  is continuous.

PROOF: Similar since  $\overline{f_\alpha}^{-1} = \overline{f_{\alpha^{-1}}}$ .

$\square$

**Corollary 320.1.** *Let  $G$  be a topological group and  $H$  a subgroup of  $G$ . Give  $G/H$  the quotient topology. Then  $G/H$  is homogeneous.*

**Proposition 321.** *Let  $G$  be a topological group and  $H$  a subgroup of  $G$ . Give  $G/H$  the quotient topology. If  $H$  is closed in  $G$  then  $G/H$  is  $T_1$ .*

PROOF:

$\langle 1 \rangle 1.$  LET:  $p : G \twoheadrightarrow G/H$  be the canonical surjection

$\langle 1 \rangle 2.$  LET:  $x \in G$

$\langle 1 \rangle 3.$   $p^{-1}(xH) = f_x(H)$

$\langle 1 \rangle 4.$   $p^{-1}(xH)$  is closed in  $G$

PROOF: Since  $H$  is closed and  $f_x$  is a homeomorphism of  $G$  with itself.

$\langle 1 \rangle 5.$   $\{xH\}$  is closed in  $G/H$

$\square$

**Proposition 322.** *Let  $G$  be a topological group and  $H$  a subgroup of  $G$ . Give  $G/H$  the quotient topology. Then the canonical surjection  $p : G \twoheadrightarrow G/H$  is an open map.*

PROOF:

$\langle 1 \rangle 1.$  LET:  $U \subseteq G$  be open.

$\langle 1 \rangle 2.$   $p^{-1}(p(U)) = \bigcup_{h \in H} r_h(U)$

$\langle 1 \rangle 3.$   $p^{-1}(p(U))$  is open.

$\langle 1 \rangle 4.$   $p(U)$  is open.

$\square$

**Proposition 323.** *Let  $G$  be a topological group and  $H$  a closed normal subgroup of  $G$ . Then  $G/H$  is a topological group under the quotient topology.*

PROOF:

$\langle 1 \rangle 1$ .  $G/H$  is  $T_1$

PROOF: Proposition 321.

$\langle 1 \rangle 2$ . The map  $\bar{m} : (xH, yH) \mapsto xy^{-1}H$  is continuous.

$\langle 2 \rangle 1$ .  $p^2 : G^2 \rightarrow (G/H)^2$  is a quotient map.

PROOF: Propositions 197, 322.

$\langle 2 \rangle 2$ .  $\bar{m} \circ p^2$  is continuous.

PROOF: As it is  $p^2 \circ m$  where  $m : G^2 \rightarrow G$  with  $m(x, y) = xy^{-1}$

□

**Lemma 324.** *Let  $G$  be a topological group and  $A, B \subseteq G$ . If either  $A$  or  $B$  is open then  $AB$  is open.*

PROOF: If  $A$  is open we have  $AB = \bigcup_{b \in B} r_b(A)$ . Similarly if  $B$  is open. □

**Definition 325** (Symmetric Neighbourhood). Let  $G$  be a topological group. A neighbourhood  $V$  of  $e$  is *symmetric* if and only if  $V = V^{-1}$ .

**Lemma 326.** *Let  $G$  be a topological group. Let  $V$  be a neighbourhood of  $e$ . Then  $V$  is symmetric if and only if, for all  $x \in V$ , we have  $x^{-1} \in V$ .*

PROOF:

$\langle 1 \rangle 1$ . If  $V$  is symmetric then, for all  $x \in V$ , we have  $x^{-1} \in V$

PROOF: Immediate from definitions.

$\langle 1 \rangle 2$ . If, for all  $x \in V$ , we have  $x^{-1} \in V$ , then  $V$  is symmetric.

$\langle 2 \rangle 1$ . ASSUME: for all  $x \in V$  we have  $x^{-1} \in V$

$\langle 2 \rangle 2$ .  $V \subseteq V^{-1}$

PROOF: If  $x \in V$  then there exists  $y \in V$  such that  $x = y^{-1}$ , namely  $y = x^{-1}$

$\langle 2 \rangle 3$ .  $V^{-1} \subseteq V$

PROOF: Immediate from  $\langle 2 \rangle 1$ .

□

**Lemma 327.** *Let  $G$  be a topological group. For every neighbourhood  $U$  of  $e$ , there exists a symmetric neighbourhood  $V$  of  $e$  such that  $V^2 \subseteq U$ .*

PROOF:

$\langle 1 \rangle 1$ . LET:  $U$  be a neighbourhood of  $e$ .

$\langle 1 \rangle 2$ . PICK a neighbourhood  $V'$  of  $e$  such that  $V'V' \subseteq U$

PROOF: Such a neighbourhood exists because multiplication in  $G$  is continuous.

$\langle 1 \rangle 3$ . PICK a neighbourhood  $W$  of  $e$  such that  $WW^{-1} \subseteq V'$

PROOF: Such a neighbourhood exists because the function that maps  $(x, y)$  to  $xy^{-1}$  is continuous.

$\langle 1 \rangle 4$ . LET:  $V = WW^{-1}$

$\langle 1 \rangle 5$ .  $V$  is a neighbourhood of  $e$

$\langle 2 \rangle 1$ .  $e \in V$

PROOF: Since  $e \in W$  so  $e = ee^{-1} \in V$ .

$\langle 2 \rangle 2$ .  $V$  is open  
 PROOF: Lemma 324.  
 $\langle 1 \rangle 6$ .  $V$  is symmetric  
 $\langle 2 \rangle 1$ . For all  $x \in V$  we have  $x^{-1} \in V$   
 $\langle 3 \rangle 1$ . LET:  $x \in V$   
 $\langle 3 \rangle 2$ . PICK  $y, z \in W$  such that  $x = yz^{-1}$   
 $\langle 3 \rangle 3$ .  $x^{-1} = zy^{-1}$   
 $\langle 3 \rangle 4$ .  $x^{-1} \in V$   
 $\langle 3 \rangle 5$ .  $x \in V^{-1}$   
 $\langle 2 \rangle 2$ . Q.E.D.  
 PROOF: Lemma 326  
 $\langle 1 \rangle 7$ .  $V^2 \subseteq U$   
 PROOF: We have  $V^2 \subseteq (V')^2 \subseteq U$   
 $\square$

**Proposition 328.** *Every topological group is Hausdorff.*

PROOF:  
 $\langle 1 \rangle 1$ . LET:  $G$  be a topological group.  
 $\langle 1 \rangle 2$ . LET:  $x, y \in G$  with  $x \neq y$   
 $\langle 1 \rangle 3$ . LET:  $U = G \setminus \{x^{-1}y\}$   
 $\langle 1 \rangle 4$ . PICK a symmetric neighbourhood  $V$  of  $e$  such that  $VV \subseteq U$   
 $\langle 2 \rangle 1$ .  $U$  is open  
 PROOF: Since  $G$  is  $T_1$ .  
 $\langle 2 \rangle 2$ .  $e \in U$   
 PROOF: Since  $x \neq y$   
 $\langle 2 \rangle 3$ . Q.E.D.  
 PROOF: Lemma 327.  
 $\langle 1 \rangle 5$ .  $Vx$  and  $Vy$  are disjoint neighbourhoods of  $x$  and  $y$  respectively.  
 $\langle 2 \rangle 1$ .  $Vx$  is open  
 PROOF: Since  $Vx = r_x(V)$   
 $\langle 2 \rangle 2$ .  $Vy$  is open  
 PROOF: Similar.  
 $\langle 2 \rangle 3$ .  $Vx \cap Vy = \emptyset$   
 $\langle 3 \rangle 1$ . ASSUME: for a contradiction  $z \in Vx \cap Vy$   
 $\langle 3 \rangle 2$ . PICK  $a, b \in V$  such that  $z = ax = by$   
 $\langle 3 \rangle 3$ .  $xy^{-1} \in VV$   
 PROOF: Since  $xy^{-1} = a^{-1}b$   
 $\langle 3 \rangle 4$ .  $xy^{-1} \in U$   
 $\langle 3 \rangle 5$ . Q.E.D.  
 PROOF: From  $\langle 1 \rangle 3$ .  
 $\square$

**Proposition 329.** *Every topological group is regular.*

PROOF:  
 $\langle 1 \rangle 1$ . LET:  $G$  be a topological group.

- ⟨1⟩2. LET:  $A \subseteq G$  be a closed set and  $a \notin A$ .
- ⟨1⟩3. LET:  $U = G \setminus Aa^{-1}$
- ⟨1⟩4. PICK a symmetric neighbourhood  $V$  of  $e$  such that  $VV \subseteq U$ 
  - ⟨2⟩1.  $U$  is open  
PROOF: Since  $Aa^{-1} = r_{a^{-1}}(A)$  is closed.
  - ⟨2⟩2.  $e \in U$   
PROOF: Since  $a \notin A$ .
  - ⟨2⟩3. Q.E.D.  
PROOF: Lemma 327.
- ⟨1⟩5.  $VA$  and  $Va$  are disjoint open sets with  $A \subseteq VA$  and  $a \in Va$ 
  - ⟨2⟩1.  $VA$  is open  
PROOF: Lemma 324
  - ⟨2⟩2.  $Va$  is open  
PROOF: Lemma 324
  - ⟨2⟩3.  $VA \cap Va = \emptyset$ 
    - ⟨3⟩1. ASSUME: for a contradiction  $z \in VA \cap Va$
    - ⟨3⟩2. PICK  $b, c \in V$  and  $d \in A$  with  $z = bd = ca$
    - ⟨3⟩3.  $da^{-1} \in U$   
PROOF: Since  $da^{-1} = b^{-1}c \in VV \subseteq U$
    - ⟨3⟩4. Q.E.D.  
PROOF: This contradicts ⟨1⟩3

□

**Proposition 330.** *Let  $G$  be a topological group and  $H$  a subgroup of  $G$ . Give  $G/H$  the quotient topology. If  $H$  is closed in  $G$  then  $G/H$  is regular.*

PROOF:

- ⟨1⟩1. LET:  $p : G \rightarrow G/H$  be the canonical surjection.
- ⟨1⟩2. LET:  $A$  be a closed set in  $G/H$  and  $aH \in (G/H) \setminus A$ .
- ⟨1⟩3. LET:  $B = p^{-1}(A)$
- ⟨1⟩4.  $B$  is a closed saturated set in  $G$ .
- ⟨1⟩5.  $B \cap aH = \emptyset$
- ⟨1⟩6.  $B = BH$
- ⟨1⟩7. PICK a symmetric neighbourhood  $V$  of  $e$  such that  $VB$  does not intersect  $Va$ 
  - ⟨2⟩1. LET:  $U = G \setminus Ba^{-1}$
  - ⟨2⟩2. PICK a symmetric neighbourhood  $V$  of  $e$  such that  $VV \subseteq U$ 
    - ⟨3⟩1.  $U$  is open  
PROOF: Since  $Ba^{-1} = r_{a^{-1}}(B)$  is closed.
    - ⟨3⟩2.  $e \in U$   
PROOF: If  $e \in Ba^{-1}$  then  $a \in B$
    - ⟨3⟩3. Q.E.D.  
PROOF: Lemma 327
  - ⟨2⟩3.  $VB \cap Va = \emptyset$   
PROOF: If  $vb = v'a$  for  $v, v' \in V$  and  $b \in B$  then we have  $ba^{-1} = v^{-1}v' \in VV \subseteq U$   
 $Ba \cap VV \subseteq Ba \cap U$ .
- ⟨1⟩8.  $p(VB)$  and  $p(Va)$  are disjoint open sets

$\langle 2 \rangle 1.$   $p(VB)$  and  $p(Va)$  are open.

PROOF: Proposition 322.

$\langle 2 \rangle 2.$   $p(VB) \cap p(Va) = \emptyset$

PROOF: If  $vbH = v'aH$  for  $v, v' \in V, b \in B$  then  $v'a = vbh$  for some  $h \in H$ .

Hence  $v'a \in Va \cap VBH = Va \cap VB$ .

$\langle 1 \rangle 9.$   $A \subseteq p(VB)$

$\langle 1 \rangle 10.$   $aH \in p(Va)$

□

**Proposition 331.** *Let  $G$  be a topological group. The component of  $G$  that contains  $e$  is a normal subgroup of  $G$ .*

PROOF:

$\langle 1 \rangle 1.$  LET:  $C$  be the component of  $G$  that contains  $e$ .

$\langle 1 \rangle 2.$  For all  $x \in G$ ,  $xC$  is the component of  $G$  that contains  $x$ .

$\langle 2 \rangle 1.$  LET:  $x \in G$

$\langle 2 \rangle 2.$  LET:  $D$  be the component of  $G$  that contains  $x$ .

$\langle 2 \rangle 3.$   $xC \subseteq D$

PROOF: Since  $xC$  is connected by Theorem 222.

$\langle 2 \rangle 4.$   $D \subseteq xC$

PROOF: Since  $x^{-1}D \subseteq C$  similarly.

$\langle 1 \rangle 3.$  For all  $x \in G$ ,  $Cx$  is the component of  $G$  that contains  $x$ .

PROOF: Similar.

$\langle 1 \rangle 4.$  For all  $x \in C$  we have  $xC = Cx = C$

$\langle 1 \rangle 5.$  For all  $x \in C$  we have  $x^{-1}C = C$

$\langle 1 \rangle 6.$  For all  $x \in C$  we have  $x^{-1} \in C$

$\langle 1 \rangle 7.$  For all  $x, y \in C$  we have  $xy \in C$

PROOF: Since  $xyC = xC = x$ .

$\langle 1 \rangle 8.$  For all  $x \in G$  we have  $xC = Cx$ .

PROOF: From  $\langle 1 \rangle 2$  and  $\langle 1 \rangle 3$ .

□

**Lemma 332.** *Let  $G$  be a topological group. Let  $A$  be a closed set in  $G$  and  $B$  a compact subspace of  $G$  such that  $A \cap B = \emptyset$ . Then there exists a symmetric neighbourhood  $U$  of  $e$  such that  $AU \cap BU = \emptyset$ .*

PROOF:

$\langle 1 \rangle 1.$  For all  $b \in B$  there exists a symmetric neighbourhood  $V$  of  $e$  such that  $bV^2 \cap A = \emptyset$

$\langle 2 \rangle 1.$  LET:  $b \in B$

$\langle 2 \rangle 2.$  LET:  $W = b^{-1}(G \setminus A)$

$\langle 2 \rangle 3.$   $W$  is a neighbourhood of  $e$  and  $bW \cap A = \emptyset$

$\langle 2 \rangle 4.$  PICK a symmetric neighbourhood  $V$  of  $e$  such that  $V^2 \subseteq W$

$\langle 1 \rangle 2.$   $\{bV^2 \mid b \in B, V \text{ is a symmetric neighbourhood of } e, bV^2 \cap A = \emptyset\}$  is an open cover of  $B$

$\langle 1 \rangle 3.$  PICK a finite subcover  $b_1V_1^2, \dots, b_nV_n^2$ , say.

$\langle 1 \rangle 4.$  LET:  $U = V_1 \cap \dots \cap V_n$



$\langle 1 \rangle 5. BU^2 \cap A = \emptyset$

$\langle 1 \rangle 6. AU \cap BU = \emptyset$

PROOF: If  $av \in BU$  where  $a \in A$  and  $v \in V$  then  $a = avv^{-1} \in BU^2 \cap A$ .

□

**Proposition 333 (AC).** *Let  $G$  be a topological group. Let  $A$  be a closed set in  $G$ , and  $B$  a compact subspace of  $G$ . Then  $AB$  is closed.*

PROOF:

$\langle 1 \rangle 1.$  LET:  $x \in G \setminus AB$

$\langle 1 \rangle 2. A^{-1}x \cap B = \emptyset$

$\langle 1 \rangle 3. A^{-1}x$  is closed.

$\langle 1 \rangle 4.$  PICK a symmetric neighbourhood  $U$  of  $e$  such that  $A^{-1}xU \cap BU = \emptyset$

$\langle 1 \rangle 5. xU^2$  is open

PROOF: Lemma 324.

$\langle 1 \rangle 6. x \in xU^2 \subseteq G \setminus AB$

□

**Corollary 333.1.** *Let  $G$  be a topological group and  $H \leq G$ . Let  $p : G \twoheadrightarrow G/H$  be the quotient map. If  $H$  is compact then  $p$  is a closed map.*

PROOF: For  $A$  closed in  $G$ , we have  $p^{-1}(p(A)) = AH$  is closed, and so  $p(A)$  is closed. □

**Corollary 333.2.** *Let  $G$  be a topological group and  $H \leq G$ . If  $H$  and  $G/H$  are compact then  $G$  is compact.*

PROOF: From Proposition 313 since, for all  $aH \in G/H$ , we have  $p^{-1}(aH) = aH$  is compact because it is homomorphic to  $H$ . □

## 48 The Metric Topology

**Definition 334** (Metric). Let  $X$  be a set. A *metric* on  $X$  is a function  $d : X^2 \rightarrow \mathbb{R}$  such that:

1. For all  $x, y \in X$ ,  $d(x, y) \geq 0$
2. For all  $x, y \in X$ ,  $d(x, y) = 0$  if and only if  $x = y$
3. For all  $x, y \in X$ ,  $d(x, y) = d(y, x)$
4. (*Triangle Inequality*) For all  $x, y, z \in X$ ,  $d(x, z) \leq d(x, y) + d(y, z)$

We call  $d(x, y)$  the *distance* between  $x$  and  $y$ .

**Definition 335** (Open Ball). Let  $X$  be a metric space. Let  $a \in X$  and  $\epsilon > 0$ . The *open ball* with *centre*  $a$  and *radius*  $\epsilon$  is

$$B(a, \epsilon) = \{x \in X \mid d(a, x) < \epsilon\} .$$

**Definition 336** (Metric Topology). Let  $X$  be a metric space. The *metric topology* on  $X$  is the topology generated by the basis consisting of all the open balls.

We prove this is a basis for a topology.

PROOF:

$\langle 1 \rangle 1$ . For every point  $a$ , there exists a ball  $B$  such that  $a \in B$

PROOF: We have  $a \in B(a, 1)$ .

$\langle 1 \rangle 2$ . For any balls  $B_1, B_2$  and point  $a \in B_1 \cap B_2$ , there exists a ball  $B_3$  such that  $a \in B_3 \subseteq B_1 \cap B_2$

$\langle 2 \rangle 1$ . LET:  $B_1 = B(c_1, \epsilon_1)$  and  $B_2 = B(c_2, \epsilon_2)$

$\langle 2 \rangle 2$ . LET:  $\delta = \min(\epsilon_1 - d(c_1, a), \epsilon_2 - d(c_2, a))$

PROVE:  $B(a, \delta) \subseteq B_1 \cap B_2$

$\langle 2 \rangle 3$ . LET:  $x \in B(a, \delta)$

$\langle 2 \rangle 4$ .  $x \in B_1$

PROOF:

$$\begin{aligned} d(x, c_1) &= d(x, a) + d(a, c_1) \\ &< \delta + d(a, c_1) \\ &\leq \epsilon_1 \end{aligned}$$

$\langle 2 \rangle 5$ .  $x \in B_2$

PROOF: Similar.

□

**Proposition 337.** Let  $X$  be a metric space and  $U \subseteq X$ . Then  $U$  is open if and only if, for every  $x \in U$ , there exists  $\epsilon > 0$  such that  $B(x, \epsilon) \subseteq U$ .

PROOF:

$\langle 1 \rangle 1$ . If  $U$  is open then, for all  $x \in U$ , there exists  $\epsilon > 0$  such that  $B(x, \epsilon) \subseteq U$ .

$\langle 2 \rangle 1$ . ASSUME:  $U$  is open.

- ⟨2⟩2. LET:  $x \in U$   
 ⟨2⟩3. PICK  $a \in X$  and  $\delta > 0$  such that  $x \in B(a, \delta) \subseteq U$   
 ⟨2⟩4. LET:  $\epsilon = \delta - d(a, x)$   
         PROVE:  $B(x, \epsilon) \subseteq U$   
 ⟨2⟩5. LET:  $y \in B(x, \epsilon)$   
 ⟨2⟩6.  $d(y, a) < \delta$   
         PROOF:

$$\begin{aligned}
 d(y, a) &\leq d(a, x) + d(x, y) \\
 &< \delta + d(x, y) \\
 &= \epsilon
 \end{aligned}$$

- ⟨2⟩7.  $y \in U$   
 ⟨1⟩2. If, for all  $x \in U$ , there exists  $\epsilon > 0$  such that  $B(x, \epsilon) \subseteq U$ , then  $U$  is open.  
 PROOF: Immediate from definitions.

□

**Definition 338** (Discrete Metric). Let  $X$  be a set. The *discrete metric* on  $X$  is defined by

$$d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

**Proposition 339.** *The discrete metric induces the discrete topology.*

PROOF: For any (open) set  $U$  and point  $a \in U$ , we have  $a \in B(a, 1) \subseteq U$ . □

**Definition 340** (Standard Metric on  $\mathbb{R}$ ). The *standard metric* on  $\mathbb{R}$  is defined by  $d(x, y) = |x - y|$ .

**Proposition 341.** *The standard metric on  $\mathbb{R}$  induces the standard topology on  $\mathbb{R}$ .*

PROOF:

- ⟨1⟩1. Every open ball is open in the standard topology on  $\mathbb{R}$ .  
         PROOF:  $B(a, \epsilon) = (a - \epsilon, a + \epsilon)$   
 ⟨1⟩2. For every open set  $U$  and point  $a \in U$ , there exists  $\epsilon > 0$  such that  
          $B(a, \epsilon) \subseteq U$   
 ⟨2⟩1. LET:  $U$  be an open set and  $a \in U$   
 ⟨2⟩2. PICK an open interval  $b, c$  such that  $a \in (b, c) \subseteq U$   
 ⟨2⟩3. LET:  $\epsilon = \min(a - b, c - a)$   
 ⟨2⟩4.  $B(a, \epsilon) \subseteq U$

□

**Definition 342** (Metriizable). A topological space  $X$  is *metrizable* if and only if there exists a metric on  $X$  that induces the topology.

**Definition 343** (Bounded). Let  $X$  be a metric space and  $A \subseteq X$ . Then  $A$  is *bounded* if and only if there exists  $M$  such that, for all  $x, y \in A$ , we have  $d(x, y) \leq M$ .

**Definition 344** (Diameter). Let  $X$  be a metric space and  $A \subseteq X$ . The *diameter* of  $A$  is

$$\text{diam } A = \sup_{x,y \in A} d(x,y) .$$

**Definition 345** (Standard Bounded Metric). Let  $d$  be a metric on  $X$ . The *standard bounded metric* corresponding to  $d$  is the metric  $\bar{d}$  defined by

$$\bar{d}(x,y) = \min(d(x,y), 1) .$$

We prove this is a metric.

PROOF:

$\langle 1 \rangle 1. \bar{d}(x,y) \geq 0$

PROOF: Since  $d(x,y) \geq 0$

$\langle 1 \rangle 2. \bar{d}(x,y) = 0$  if and only if  $x = y$

PROOF:  $\bar{d}(x,y) = 0$  if and only if  $d(x,y) = 0$  if and only if  $x = y$

$\langle 1 \rangle 3. \bar{d}(x,y) = \bar{d}(y,x)$

PROOF: Since  $d(x,y) = d(y,x)$

$\langle 1 \rangle 4. \bar{d}(x,z) \leq \bar{d}(x,y) + \bar{d}(y,z)$

PROOF:

$$\begin{aligned} \bar{d}(x,y) + \bar{d}(y,z) &= \min(d(x,y), 1) + \min(d(y,z), 1) \\ &= \min(d(x,y) + d(y,z), d(x,y) + 1, d(y,z) + 1, 2) \\ &\geq \min(d(x,z), 1) \\ &= \bar{d}(x,z) \end{aligned}$$

□

**Lemma 346.** In any metric space  $X$ , the set  $\mathcal{B} = \{B(a, \epsilon) \mid a \in X, \epsilon < 1\}$  is a basis for the metric topology.

PROOF:

$\langle 1 \rangle 1.$  Every element of  $\mathcal{B}$  is open.

PROOF: From Lemma 62.

$\langle 1 \rangle 2.$  For every open set  $U$  and point  $a \in U$ , there exists  $B \in \mathcal{B}$  such that  $a \in B \subseteq U$

$\langle 2 \rangle 1.$  LET:  $U$  be an open set and  $a \in U$

$\langle 2 \rangle 2.$  PICK  $\epsilon > 0$  such that  $B(a, \epsilon) \subseteq U$

$\langle 2 \rangle 3.$   $B(a, \min(\epsilon, 1/2)) \subseteq U$

$\langle 1 \rangle 3.$  Q.E.D.

PROOF: Lemma 63.

□

**Proposition 347.** Let  $d$  be a metric on the set  $X$ . Then the standard bounded metric  $\bar{d}$  induces the same metric as  $d$ .

PROOF: This follows from Lemma 346 since the open balls with radius  $< 1$  are the same under both metrics. □

**Lemma 348.** *Let  $d$  and  $d'$  be two metrics on the same set  $X$ . Let  $\mathcal{T}$  and  $\mathcal{T}'$  be the topologies they induce. Then  $\mathcal{T} \subseteq \mathcal{T}'$  if and only if, for all  $x \in X$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that*

$$B_{d'}(x, \delta) \subseteq B_d(x, \epsilon) .$$

PROOF:

$\langle 1 \rangle 1$ . If  $\mathcal{T} \subseteq \mathcal{T}'$  then, for all  $x \in X$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$

PROOF: From Proposition 337 since  $x \in B_d(x, \epsilon) \in \mathcal{T}'$ .

$\langle 1 \rangle 2$ . If, for all  $x \in X$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$ , then  $\mathcal{T} \subseteq \mathcal{T}'$

$\langle 2 \rangle 1$ . ASSUME: For all  $x \in X$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$

$\langle 2 \rangle 2$ . LET:  $U \in \mathcal{T}$

$\langle 2 \rangle 3$ . For all  $x \in U$  there exists  $\delta > 0$  such that  $B_{d'}(x, \delta) \subseteq U$ .

$\langle 3 \rangle 1$ . LET:  $x \in U$

$\langle 3 \rangle 2$ . PICK  $\epsilon > 0$  such that  $B_d(x, \epsilon) \subseteq U$

PROOF: Proposition 337

$\langle 3 \rangle 3$ . PICK  $\delta > 0$  such that  $B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$

PROOF: By  $\langle 2 \rangle 1$

$\langle 3 \rangle 4$ .  $B_{d'}(x, \delta) \subseteq U$

$\langle 2 \rangle 4$ .  $U \in \mathcal{T}'$

PROOF: Proposition 337.

□

**Proposition 349.**  $\mathbb{R}^2$  under the dictionary order topology is metrizable.

PROOF: Define  $d : \mathbb{R}^2 \rightarrow \mathbb{R}$  by

$$d((x, y), (x, z)) = \max(|y - z|, 1)$$

$$d((x, y), (x', y')) = 1$$

$$\text{if } x \neq x' \square$$

$\langle 1 \rangle 1$ .  $x \in \bigcap_{i=1}^N \pi_i^{-1}() \subseteq B_D(a, \epsilon)$

**Proposition 350.** *Let  $d : X^2 \rightarrow \mathbb{R}$  be a metric on  $X$ . Then the metric topology on  $X$  is the coarsest topology such that  $d$  is continuous.*

PROOF:

$\langle 1 \rangle 1$ .  $d$  is continuous.

$\langle 2 \rangle 1$ . LET:  $a, b \in X$

$\langle 2 \rangle 2$ . LET:  $\epsilon > 0$

$\langle 2 \rangle 3$ . LET:  $\delta = \epsilon/2$

$\langle 2 \rangle 4$ . LET:  $x, y \in X$

$\langle 2 \rangle 5$ . ASSUME:  $\rho((a, b), (x, y)) < \delta$

$\langle 2 \rangle 6$ .  $|d(a, b) - d(x, y)| < \epsilon$

$\langle 3 \rangle 1$ .  $d(a, b) - d(x, y) < \epsilon$

PROOF:

$$\begin{aligned}
d(a, b) &\leq d(a, x) + d(x, y) + d(y, b) \\
&\leq d(x, y) + 2\rho((a, b), (x, y)) \\
&< d(x, y) + 2\delta \\
&= d(x, y) + \epsilon
\end{aligned}$$

$$\langle 3 \rangle 2. \ d(a, b) - d(x, y) > -\epsilon$$

PROOF: Similar.

$\langle 2 \rangle 7.$  Q.E.D.

$\langle 1 \rangle 2.$  If  $\mathcal{T}$  is any topology under which  $d$  is continuous then  $\mathcal{T}$  is finer than the metric topology.

PROOF: Since  $B(a, \epsilon) = d_a^{-1}((-\infty, \epsilon))$

□

**Proposition 351.** *Let  $X$  be a metric space with metric  $d$  and  $A \subseteq X$ . The restriction of  $d$  to  $A$  is a metric on  $A$  that induces the subspace topology.*

PROOF:

$\langle 1 \rangle 1.$  The restriction of  $d$  to  $A$  is a metric on  $A$ .

$\langle 1 \rangle 2.$  Every open ball under  $d \upharpoonright A$  is open under the subspace topology.

PROOF:  $B_{d \upharpoonright A}(a, \epsilon) = B_d(a, \epsilon) \cap A$ .

$\langle 1 \rangle 3.$  If  $U$  is open in the subspace topology and  $x \in U$ , then there exists a  $d \upharpoonright A$ -ball  $B$  such that  $x \in B \subseteq U$ .

$\langle 2 \rangle 1.$  PICK  $V$  open in  $X$  such that  $U = V \cap A$

$\langle 2 \rangle 2.$  PICK  $\epsilon > 0$  such that  $B_d(x, \epsilon) \subseteq V$

$\langle 2 \rangle 3.$  Take  $B = B_{d \upharpoonright A}(x, \epsilon)$

□

**Corollary 351.1.** *A subspace of a metrizable space is metrizable.*

**Proposition 352.** *Every metrizable space is Hausdorff.*

PROOF:

$\langle 1 \rangle 1.$  LET:  $X$  be a metric space

$\langle 1 \rangle 2.$  LET:  $a, b \in X$  with  $a \neq b$

$\langle 1 \rangle 3.$  LET:  $\epsilon = d(a, b)/2$

$\langle 1 \rangle 4.$  LET:  $U = B(a, \epsilon)$  and  $V = B(b, \epsilon)$

$\langle 1 \rangle 5.$   $U$  and  $V$  are disjoint neighbourhoods of  $a$  and  $b$  respectively.

□

**Proposition 353 (CC).** *The product of a countable family of metrizable spaces is metrizable.*

PROOF:

$\langle 1 \rangle 1.$  LET:  $(X_n, d_n)$  be a sequence of metric spaces.

$\langle 1 \rangle 2.$  ASSUME: w.l.o.g. each  $d_n$  is bounded above by 1.

PROOF: By Proposition 347.

$\langle 1 \rangle 3.$  LET:  $D$  be the metric on  $\mathbb{R}^\omega$  defined by  $D(x, y) = \sup_i (d_i(x_i, y_i)/i)$ .

- ⟨2⟩1.  $D(x, y) \geq 0$
- ⟨2⟩2.  $D(x, y) = 0$  if and only if  $x = y$
- ⟨2⟩3.  $D(x, y) = D(y, x)$
- ⟨2⟩4.  $D(x, z) \leq D(x, y) + D(y, z)$

PROOF:

$$\begin{aligned}
D(x, z) &= \sup_i \frac{d_i(x_i, z_i)}{i} \\
&\leq \sup_i \frac{d_i(x_i, y_i) + d_i(y_i, z_i)}{i} \\
&\leq \sup_i \frac{d_i(x_i, y_i)}{i} + \sup_i \frac{d_i(y_i, z_i)}{i} \\
&= D(x, y) + D(y, z)
\end{aligned}$$

- ⟨1⟩4. Every open ball  $B_D(a, \epsilon)$  is open in the product topology.
- ⟨2⟩1. PICK  $N$  such that  $1/\epsilon < N$
- ⟨2⟩2.  $B_D(a, \epsilon) = \prod_{i=1}^{\infty} U_i$  where  $U_i = B(a_i, i\epsilon)$  if  $i \leq N$ , and  $U_i = X_i$  if  $i > N$
- ⟨1⟩5. For any open set  $U$  and  $a \in U$ , there exists  $\epsilon > 0$  such that  $B_D(a, \epsilon) \subseteq U$ .
- ⟨2⟩1. LET:  $n \geq 1$ ,  $V$  be an open set in  $\mathbb{R}$  and  $a \in \pi_n^{-1}(V)$
- ⟨2⟩2. PICK  $\epsilon > 0$  such that  $B_{d_n}(a, \epsilon) \subseteq V$
- ⟨2⟩3.  $B_D(a, \epsilon/n) \subseteq \pi_n^{-1}(V)$

□

**Theorem 354.** *Let  $X$  and  $Y$  be metric spaces and  $f : X \rightarrow Y$ . Then  $f$  is continuous if and only if, for all  $x \in X$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that, for all  $y \in X$ , if  $d(x, y) < \delta$  then  $d(f(x), f(y)) < \epsilon$ .*

PROOF:

- ⟨1⟩1. If  $f$  is continuous then, for all  $x \in X$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that, for all  $y \in X$ , if  $d(x, y) < \delta$  then  $d(f(x), f(y)) < \epsilon$
- ⟨2⟩1. ASSUME:  $f$  is continuous.
- ⟨2⟩2. LET:  $x \in X$  and  $\epsilon > 0$
- ⟨2⟩3. PICK a neighbourhood  $U$  of  $x$  such that  $f(U) \subseteq B(f(x), \epsilon)$
- PROOF: Theorem 93.
- ⟨2⟩4. PICK  $\delta > 0$  such that  $B(x, \delta) \subseteq U$
- PROOF: Proposition 337.
- ⟨2⟩5. For all  $y \in X$ , if  $d(x, y) < \delta$  then  $d(f(x), f(y)) < \epsilon$
- ⟨1⟩2. If for all  $x \in X$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that, for all  $y \in X$ , if  $d(x, y) < \delta$  then  $d(f(x), f(y)) < \epsilon$ , then  $f$  is continuous.
- ⟨2⟩1. ASSUME: for all  $x \in X$  and  $\epsilon > 0$ , there exists  $\delta > 0$  such that, for all  $y \in X$ , if  $d(x, y) < \delta$  then  $d(f(x), f(y)) < \epsilon$
- ⟨2⟩2. LET:  $x \in X$  and  $V$  be a neighbourhood of  $f(x)$
- ⟨2⟩3. PICK  $\epsilon > 0$  such that  $B(f(x), \epsilon) \subseteq V$
- PROOF: Proposition 337.
- ⟨2⟩4. PICK  $\delta > 0$  such that, for all  $y \in X$ , if  $d(x, y) < \delta$  then  $d(f(x), f(y)) < \epsilon$
- PROOF: By ⟨2⟩1
- ⟨2⟩5. LET:  $U = B(x, \delta)$
- ⟨2⟩6.  $U$  is a neighbourhood of  $x$  with  $f(U) \subseteq V$

⟨2⟩7. Q.E.D.

PROOF: Theorem 93.

□

**Proposition 355.** *Let  $X$  be a metric space. Let  $(a_n)$  be a sequence in  $X$  and  $l \in X$ . Then  $a_n \rightarrow l$  as  $n \rightarrow \infty$  if and only if, for all  $\epsilon > 0$ , there exists  $N$  such that, for all  $n \geq N$ , we have  $d(a_n, l) < \epsilon$ .*

PROOF: From Proposition 76. □

**Proposition 356.** *Every metrizable space is first countable.*

PROOF: In any metric space  $X$ , the open balls  $B(a, 1/n)$  for  $n \geq 1$  form a local basis at  $a$ .

**Example 357.**  $\mathbb{R}^\omega$  under the box topology is not metrizable.

**Example 358.** If  $J$  is uncountable then  $\mathbb{R}^J$  under the product topology is not metrizable.

**Proposition 359.** *A compact subspace of a metric space is bounded.*

PROOF:

⟨1⟩1. LET:  $X$  be a metric space and  $A \subseteq X$  be compact.

⟨1⟩2. PICK  $a \in A$

⟨1⟩3.  $\{B(a, n) \mid n \in \mathbb{Z}^+\}$  covers  $A$

⟨1⟩4. PICK a finite subcover  $\{B(a, n_1), \dots, B(a, n_k)\}$

⟨1⟩5. LET:  $N = \max(n_1, \dots, n_k)$

⟨1⟩6. For all  $x, y \in A$  we have  $d(x, y) < 2N$

PROOF:

$$\begin{aligned} d(x, y) &\leq d(x, a) + d(a, y) \\ &< N + N \end{aligned}$$

□

This example shows the converse does not hold:

**Example 360.** The space  $\mathbb{R}$  under the standard bounded metric is bounded but not compact.

## 49 Real Linear Algebra

**Definition 361** (Square Metric). The *square metric*  $\rho$  on  $\mathbb{R}^n$  is defined by

$$\rho(\vec{x}, \vec{y}) = \max(|x_1 - y_1|, \dots, |x_n - y_n|)$$

We prove this is a metric.

PROOF:

⟨1⟩1.  $\rho(\vec{x}, \vec{y}) \geq 0$

PROOF: Immediate from definition.



- $\langle 1 \rangle 2.$   $\rho(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} = \vec{y}$   
 PROOF: Immediate from definition.  
 $\langle 1 \rangle 3.$   $\rho(\vec{x}, \vec{y}) = \rho(\vec{y}, \vec{x})$   
 PROOF: Immediate from definition.  
 $\langle 1 \rangle 4.$   $\rho(\vec{x}, \vec{z}) \leq \rho(\vec{x}, \vec{y}) + \rho(\vec{y}, \vec{z})$   
 PROOF: Since  $|x_i - z_i| \leq |x_i - y_i| + |y_i - z_i|$ .  
 $\square$

**Proposition 362.** *The square metric induces the standard topology on  $\mathbb{R}^n$ .*

PROOF:

- $\langle 1 \rangle 1.$  For every  $a \in X$  and  $\epsilon > 0$ , we have  $B_\rho(a, \epsilon)$  is open in the standard product topology.

PROOF:

$$B_\rho(a, \epsilon) = (a_1 - \epsilon, a_1 + \epsilon) \times \cdots \times (a_n - \epsilon, a_n + \epsilon)$$

- $\langle 1 \rangle 2.$  For any open sets  $U_1, \dots, U_n$  in  $\mathbb{R}$ , we have  $U_1 \times \cdots \times U_n$  is open in the square metric topology.  
 $\langle 2 \rangle 1.$  LET:  $\vec{a} \in U_1 \times \cdots \times U_n$   
 $\langle 2 \rangle 2.$  For  $i = 1, \dots, n$ , PICK  $\epsilon_i > 0$  such that  $(a_i - \epsilon_i, a_i + \epsilon_i) \subseteq U_i$   
 $\langle 2 \rangle 3.$  LET:  $\epsilon = \min(\epsilon_1, \dots, \epsilon_n)$   
 $\langle 2 \rangle 4.$   $B_\rho(\vec{a}, \epsilon) \subseteq U$   
 $\square$

**Definition 363.** Given  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , define the *sum*  $\vec{x} + \vec{y}$  by

$$(x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n) .$$

**Definition 364.** Given  $\lambda \in \mathbb{R}$  and  $\vec{x} \in \mathbb{R}^n$ , define the *scalar product*  $\lambda \vec{x} \in \mathbb{R}^n$  by

$$\lambda(x_1, \dots, x_n) = (\lambda x_1, \dots, \lambda x_n)$$

**Definition 365** (Inner Product). Given  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , define the *inner product*  $\vec{x} \cdot \vec{y} \in \mathbb{R}$  by

$$(x_1, \dots, x_n) \cdot (y_1, \dots, y_n) = x_1 y_1 + \cdots + x_n y_n .$$

We write  $\vec{x}^2$  for  $\vec{x} \cdot \vec{x}$ .

**Definition 366** (Norm). Let  $n \geq 1$ . The *norm* on  $\mathbb{R}^n$  is the function  $\| \cdot \| : \mathbb{R}^n \rightarrow \mathbb{R}$  defined by

$$\|(x_1, \dots, x_n)\| = \sqrt{x_1^2 + \cdots + x_n^2}$$

**Lemma 367.**

$$\|\vec{x}\|^2 = \vec{x}^2$$

PROOF: Immediate from definitions.  $\square$

**Lemma 368.**

$$\vec{x} \cdot (\vec{y} + \vec{z}) = \vec{x} \cdot \vec{y} + \vec{x} \cdot \vec{z}$$

PROOF: Each is equal to  $(x_1y_1 + x_1z_1, \dots, x_ny_n + x_nz_n)$ .  $\square$

**Lemma 369.**

$$|\vec{x} \cdot \vec{y}| \leq \|\vec{x}\| \|\vec{y}\|$$

PROOF:

$\langle 1 \rangle 1$ . ASSUME:  $\vec{x} \neq \vec{0} \neq \vec{y}$

PROOF: Otherwise both sides are 0.

$\langle 1 \rangle 2$ . LET:  $a = 1/\|\vec{x}\|$

$\langle 1 \rangle 3$ . LET:  $b = 1/\|\vec{y}\|$

$\langle 1 \rangle 4$ .  $(a\vec{x} + b\vec{y})^2 \geq 0$  and  $(a\vec{x} - b\vec{y})^2 \geq 0$

$\langle 1 \rangle 5$ .  $a^2\|\vec{x}\|^2 + 2ab\vec{x} \cdot \vec{y} + b^2\|\vec{y}\|^2 \geq 0$  and  $a^2\|\vec{x}\|^2 - 2ab\vec{x} \cdot \vec{y} + b^2\|\vec{y}\|^2 \geq 0$

$\langle 1 \rangle 6$ .  $2ab\vec{x} \cdot \vec{y} + 2 \geq 0$  and  $-2ab\vec{x} \cdot \vec{y} + 2 \geq 0$

$\langle 1 \rangle 7$ .  $\vec{x} \cdot \vec{y} \geq -1/ab$  and  $\vec{x} \cdot \vec{y} \leq 1/ab$

$\langle 1 \rangle 8$ .  $\vec{x} \cdot \vec{y} \geq -\|\vec{x}\| \|\vec{y}\|$  and  $\vec{x} \cdot \vec{y} \leq \|\vec{x}\| \|\vec{y}\|$

$\square$

**Lemma 370** (Triangle Inequality).

$$\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$$

PROOF:

$$\begin{aligned} \|\vec{x} + \vec{y}\|^2 &= \|\vec{x}\|^2 + 2\vec{x} \cdot \vec{y} + \|\vec{y}\|^2 \\ &\leq \|\vec{x}\|^2 + 2\|\vec{x}\| \|\vec{y}\| + \|\vec{y}\|^2 && \text{(Lemma 369)} \\ &= (\|\vec{x}\| + \|\vec{y}\|)^2 && \square \end{aligned}$$

**Definition 371** (Euclidean Metric). Let  $n \geq 1$ . The *Euclidean metric* on  $\mathbb{R}^n$  is defined by

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

We prove this is a metric.

$\langle 1 \rangle 1$ .  $d(\vec{x}, \vec{y}) \geq 0$

PROOF: Immediate from definition.

$\langle 1 \rangle 2$ .  $d(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} = \vec{y}$

PROOF:  $d(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} - \vec{y} = \vec{0}$ .

$\langle 1 \rangle 3$ .  $d(\vec{x}, \vec{y}) = d(\vec{y}, \vec{x})$

PROOF: Immediate from definition.

$\langle 1 \rangle 4$ .  $d(\vec{x}, \vec{z}) \leq d(\vec{x}, \vec{y}) + d(\vec{y}, \vec{z})$

PROOF:

$$\begin{aligned} \|\vec{x} - \vec{z}\| &= \|(\vec{x} - \vec{y}) + (\vec{y} - \vec{z})\| \\ &\leq \|\vec{x} - \vec{y}\| + \|\vec{y} - \vec{z}\| && \text{(Lemma 370)} \end{aligned}$$

$\square$

**Proposition 372.** The Euclidean metric induces the standard topology on  $\mathbb{R}^n$ .

PROOF:

$\langle 1 \rangle 1$ . LET:  $\rho$  be the square metric.

- (1)2. For all  $\vec{a} \in \mathbb{R}^n$  and  $\epsilon > 0$ , we have  $B_d(\vec{a}, \epsilon) \subseteq B_\rho(\vec{a}, \epsilon)$   
 (2)1. LET:  $\vec{x} \in B_d(\vec{a}, \epsilon)$   
 (2)2.  $\sqrt{(x_1 - a_1)^2 + \cdots + (x_n - a_n)^2} < \epsilon$   
 (2)3.  $(x_1 - a_1)^2 + \cdots + (x_n - a_n)^2 < \epsilon^2$   
 (2)4. For all  $i$  we have  $(x_i - a_i)^2 < \epsilon^2$   
 (2)5. For all  $i$  we have  $|x_i - a_i| < \epsilon$   
 (2)6.  $\rho(\vec{x}, \vec{a}) < \epsilon$   
 (1)3. For all  $\vec{a} \in \mathbb{R}^n$  and  $\epsilon > 0$ , we have  $B_\rho(\vec{a}, \epsilon/\sqrt{n}) \subseteq B_d(\vec{a}, \epsilon)$   
 (2)1. LET:  $\vec{x} \in B_\rho(\vec{a}, \epsilon/\sqrt{n})$   
 (2)2.  $\rho(\vec{x}, \vec{a}) < \epsilon/\sqrt{n}$   
 (2)3. For all  $i$  we have  $|x_i - a_i| < \epsilon/\sqrt{n}$   
 (2)4. For all  $i$  we have  $(x_i - a_i)^2 < \epsilon^2/n$   
 (2)5.  $d(\vec{x}, \vec{a}) < \epsilon$   
 (1)4. Q.E.D.  
 PROOF: By Lemma 348.

□

**Proposition 373.** *Let  $n \geq 0$ . For all  $c \in \mathbb{R}^n$  and  $\epsilon > 0$ , the open ball  $B(c, \epsilon)$  is path connected.*

PROOF:

- (1)1. LET:  $a, b \in B(c, \epsilon)$   
 (1)2. LET:  $p : [0, 1] \rightarrow B(c, \epsilon)$  be the function  $p(t) = (1 - t)a + tb$   
 PROOF: We have  $p(t) \in B(c, \epsilon)$  for all  $t$  because
 
$$\begin{aligned}
 d(p(t), c) &= \|(1 - t)a + tb - c\| \\
 &= \|(1 - t)(a - c) + t(b - c)\| \\
 &\leq (1 - t)\|a - c\| + t\|b - c\| \\
 &< (1 - t)\epsilon + t\epsilon \\
 &= \epsilon
 \end{aligned}$$

- (1)3.  $p$  is a path from  $a$  to  $b$ .

□

**Proposition 374.** *Let  $n \geq 0$ . For all  $c \in \mathbb{R}^n$  and  $\epsilon > 0$ , the closed ball  $\overline{B(c, \epsilon)}$  is path connected.*

PROOF:

- (1)1. LET:  $a, b \in \overline{B(c, \epsilon)}$   
 (1)2. LET:  $p : [0, 1] \rightarrow \overline{B(c, \epsilon)}$  be the function  $p(t) = (1 - t)a + tb$   
 PROOF: We have  $p(t) \in \overline{B(c, \epsilon)}$  for all  $t$  because
 
$$\begin{aligned}
 d(p(t), c) &= \|(1 - t)a + tb - c\| \\
 &= \|(1 - t)(a - c) + t(b - c)\| \\
 &\leq (1 - t)\|a - c\| + t\|b - c\| \\
 &\leq (1 - t)\epsilon + t\epsilon \\
 &= \epsilon
 \end{aligned}$$

- (1)3.  $p$  is a path from  $a$  to  $b$ .

□

**Lemma 375.** *If  $\sum_{i=0}^{\infty} x_i^2$  and  $\sum_{i=0}^{\infty} y_i^2$  converge then  $\sum_{i=0}^{\infty} |x_i y_i|$  converges.*

PROOF:

⟨1⟩1. For all  $N \geq 0$  we have  $\sum_{i=0}^N |x_i y_i| \leq \sqrt{\sum_{i=0}^N |x_i|^2} \sqrt{\sum_{i=0}^N |y_i|^2}$

PROOF: By the Cauchy-Schwarz inequality

⟨1⟩2. Q.E.D.

PROOF: Since  $\sum_{i=0}^N |x_i y_i|$  is an increasing sequence bounded above by  $(\sum_{i=0}^{\infty} x_i^2)(\sum_{i=0}^{\infty} y_i^2)$ .

□

**Corollary 375.1.** *If  $\sum_{i=0}^{\infty} x_i^2$  and  $\sum_{i=0}^{\infty} y_i^2$  converge then  $\sum_{i=0}^{\infty} (x_i + y_i)^2$  converges.*

PROOF: Since  $\sum_{i=0}^{\infty} x_i^2$ ,  $\sum_{i=0}^{\infty} y_i^2$  and  $2 \sum_{i=0}^{\infty} x_i y_i$  all converge.

**Definition 376** ( $l^2$ -metric). The  $l^2$ -metric on

$$\left\{ (x_n) \in \mathbb{R}^{\omega} \mid \sum_{n=0}^{\infty} x_n^2 \text{ converges} \right\}$$

is defined by

$$d(x, y) = \left( \sum_{n=0}^{\infty} (x_n - y_n)^2 \right)^{1/2}$$

We prove this is a metric.

PROOF:

⟨1⟩1.  $d$  is well-defined.

PROOF: By Corollary 375.1.

⟨1⟩2.  $d(x, y) \geq 0$

⟨1⟩3.  $d(x, y) = 0$  if and only if  $x = y$

⟨1⟩4.  $d(x, y) = d(y, x)$

⟨1⟩5.  $d(x, z) \leq d(x, y) + d(y, z)$

PROOF: By Lemma 370.

□

**Theorem 377.** *Addition is a continuous function  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .*

PROOF:

⟨1⟩1. LET:  $a, b \in \mathbb{R}$

⟨1⟩2. LET:  $\epsilon > 0$

⟨1⟩3. LET:  $\delta = \epsilon/2$

⟨1⟩4. LET:  $x, y \in \mathbb{R}$

⟨1⟩5. ASSUME:  $\rho((a, b), (x, y)) < \delta$

⟨1⟩6.  $|(a + b) - (x + y)| < \epsilon$

PROOF:

$$\begin{aligned}
|(a+b) - (x+y)| &= |a-x| + |b-y| \\
&\leq 2\rho((a,b), (x,y)) \\
&< 2\delta \\
&= \epsilon
\end{aligned}$$

⟨1⟩7. Q.E.D.

PROOF: Theorem 354

□

**Theorem 378.** *Multiplication is a continuous function  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .*

PROOF:

⟨1⟩1. LET:  $a, b \in \mathbb{R}$

⟨1⟩2. LET:  $\epsilon > 0$

⟨1⟩3. LET:  $\delta = \min(\epsilon/(|a| + |b| + 1), 1)$

⟨1⟩4. LET:  $x, y \in \mathbb{R}$

⟨1⟩5. ASSUME:  $\rho((a,b), (x,y)) < \delta$

⟨1⟩6.  $|ab - xy| < \epsilon$

PROOF:

$$\begin{aligned}
|ab - xy| &= |a(b-y) + (a-x)b - (a-x)(b-y)| \\
&\leq |a||b-y| + |b||a-x| + |a-x||b-y| \\
&< |a|\delta + |b|\delta + \delta^2 & (\langle 1 \rangle 5) \\
&\leq |a|\delta + |b|\delta + \delta & (\langle 1 \rangle 3) \\
&\leq \epsilon & (\langle 1 \rangle 3)
\end{aligned}$$

⟨1⟩7. Q.E.D.

PROOF: Theorem 354

□

**Theorem 379.** *The function  $f : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$  defined by  $f(x) = x^{-1}$  is continuous.*

PROOF:

⟨1⟩1. For all  $a \in \mathbb{R}$  we have  $f^{-1}((a, +\infty))$  is open.

PROOF: The set is

$$(a^{-1}, +\infty) \text{ if } a > 0$$

$$(0, +\infty) \text{ if } a = 0$$

$$(-\infty, a^{-1}) \cup (0, +\infty) \text{ if } a < 0$$

⟨1⟩2. For all  $a \in \mathbb{R}$  we have  $f^{-1}((-\infty, a))$  is open.

PROOF: Similar.

⟨1⟩3. Q.E.D.

PROOF: By Proposition 90 and Lemma 113.

□

**Definition 380.** For  $n \geq 0$ , the *unit ball*  $B^n$  is the space  $\{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$ .

**Proposition 381.** *For all  $n \geq 0$ , the unit ball  $B^n$  is path connected.*

PROOF:

⟨1⟩1. LET:  $a, b \in B^n$

⟨1⟩2. LET:  $p : [0, 1] \rightarrow B^n$  be the function  $p(t) = (1 - t)a + tb$

PROOF: We have  $p(t) \in B^n$  for all  $t$  because

$$\begin{aligned}\|(1 - t)a + tb\| &\leq (1 - t)\|a\| + t\|b\| \\ &\leq (1 - t) + t \\ &= 1\end{aligned}$$

⟨1⟩3.  $p$  is a path from  $a$  to  $b$ .

□

**Definition 382** (Punctured Euclidean Space). For  $n \geq 0$ , defined *punctured Euclidean space* to be  $\mathbb{R}^n \setminus \{0\}$ .

**Proposition 383.** For  $n > 1$ , *punctured Euclidean space*  $\mathbb{R}^n \setminus \{0\}$  is path connected.

PROOF:

⟨1⟩1. LET:  $a, b \in \mathbb{R}^n \setminus \{0\}$

⟨1⟩2. CASE: 0 is on the line from  $a$  to  $b$

⟨2⟩1. PICK a point  $c$  not on the line from  $a$  to  $b$

⟨2⟩2. The path consisting of a straight line from  $a$  to  $c$  followed by a straight line from  $c$  to  $b$  is a path from  $a$  to  $b$ .

⟨1⟩3. CASE: 0 is not on the line from  $a$  to  $b$

PROOF: The straight line from  $a$  to  $b$  is a path from  $a$  to  $b$ .

**Corollary 383.1.** For  $n > 1$ , the spaces  $\mathbb{R}$  and  $\mathbb{R}^n$  are not homeomorphic.

PROOF: For any point  $a$ , the space  $\mathbb{R} \setminus \{a\}$  is disconnected.

**Definition 384** (Unit Sphere). For  $n \geq 1$ , the *unit sphere*  $S^{n-1}$  is the space

$$S^{n-1} = \{x \in \mathbb{R}^n \mid \|x\| = 1\}.$$

**Proposition 385.** For  $n > 1$ , the unit sphere  $S^{n-1}$  is path connected.

PROOF: The map  $g : \mathbb{R}^n \setminus \{0\} \rightarrow S^{n-1}$  defined by  $g(x) = x/\|x\|$  is continuous and surjective. The result follows by Proposition 243. □

**Proposition 386.** Let  $f : S^1 \rightarrow \mathbb{R}$  be continuous. Then there exists  $x \in S^1$  such that  $f(x) = f(-x)$ .

PROOF:

⟨1⟩1. LET:  $g : S^1 \rightarrow \mathbb{R}$  be the function  $g(x) = f(x) - f(-x)$

PROVE: There exists  $x \in S^1$  such that  $g(x) = 0$

⟨1⟩2. ASSUME: without loss of generality  $g((1, 0)) > 0$

⟨1⟩3.  $g((-1, 0)) < 0$

⟨1⟩4. There exists  $x$  such that  $g(x) = 0$

PROOF: By the Intermediate Value Theorem.

□

**Definition 387** (Topologist's Sine Curve). Let  $S = \{(x, \sin 1/x) \mid 0 < x \leq 1\}$ . The *topologist's sine curve* is the closure  $\bar{S}$  of  $S$ .

**Proposition 388.**

$$\bar{S} = S \cup (\{0\} \times [-1, 1])$$

**Proposition 389.** *The topologist's sine curve is connected.*

PROOF:

$\langle 1 \rangle 1$ . LET:  $S = \{(x, \sin 1/x) \mid 0 < x \leq 1\}$

$\langle 1 \rangle 2$ .  $S$  is connected.

PROOF: Theorem 222.

$\langle 1 \rangle 3$ .  $\bar{S}$  is connected.

PROOF: Theorem 221.

□

**Proposition 390** (CC). *The topologist's sine curve is not path connected.*

PROOF:

$\langle 1 \rangle 1$ . ASSUME: for a contradiction  $p : [0, 1] \rightarrow \bar{S}$  is a path from  $(0, 0)$  to  $(1, \sin 1)$ .

$\langle 1 \rangle 2$ .  $p^{-1}(\{0\} \times [0, 1])$  is closed.

$\langle 1 \rangle 3$ . LET:  $b$  be the greatest element of  $p^{-1}(\{0\} \times [0, 1])$ .

$\langle 1 \rangle 4$ .  $b < 1$

PROOF: Since  $p(1) = (1, \sin 1)$ .

$\langle 1 \rangle 5$ . PICK a sequence  $(t_n)_{n \geq 1}$  in  $(b, 1]$  such that  $t_n \rightarrow b$  and  $\pi_2(p(t_n)) = (-1)^n$

$\langle 2 \rangle 1$ . LET:  $n \geq 1$

$\langle 2 \rangle 2$ . PICK  $u$  with  $0 < u < \pi_1(p(1/n))$  such that  $\sin(1/u) = (-1)^n$

$\langle 2 \rangle 3$ . PICK  $t_n$  such that  $0 < t_n < 1/n$  and  $\pi_1(p(t_n)) = u$

PROOF: One exists by the Intermediate Value Theorem.

$\langle 1 \rangle 6$ . Q.E.D.

PROOF: This contradicts 105.

□

## 50 The Uniform Topology

**Definition 391** (Uniform Metric). Let  $J$  be a set. The *uniform metric*  $\bar{\rho}$  on  $\mathbb{R}^J$  is defined by

$$\bar{\rho}(a, b) = \sup_{j \in J} \bar{d}(a_j, b_j)$$

where  $\bar{d}$  is the standard bounded metric on  $\mathbb{R}$ .

The *uniform topology* on  $\mathbb{R}^J$  is the topology induced by the uniform metric.

We prove this is a metric.

PROOF:

$\langle 1 \rangle 1$ .  $\bar{\rho}(a, b) \geq 0$

PROOF: Immediate from definitions.

$\langle 1 \rangle 2$ .  $\bar{\rho}(a, b) = 0$  if and only if  $a = b$

PROOF: Immediate from definitions.

⟨1⟩3.  $\bar{\rho}(a, b) = \bar{\rho}(b, a)$

PROOF: Immediate from definitions.

⟨1⟩4.  $\bar{\rho}(a, c) \leq \bar{\rho}(a, b) + \bar{\rho}(b, c)$

PROOF:

$$\begin{aligned}\bar{\rho}(a, c) &= \sup_{j \in J} \bar{d}(a_j, c_j) \\ &\leq \sup_{j \in J} (\bar{d}(a_j, b_j) + \bar{d}(b_j, c_j)) \\ &\leq \sup_{j \in J} \bar{d}(a_j, b_j) + \sup_{j \in J} \bar{d}(b_j, c_j) \\ &= \bar{\rho}(a, b) + \bar{\rho}(b, c)\end{aligned}$$

□

**Proposition 392.** *The uniform topology on  $\mathbb{R}^J$  is finer than the product topology.*

PROOF:

⟨1⟩1. LET:  $j \in J$  and  $U$  be open in  $\mathbb{R}$

PROVE:  $\pi_j^{-1}(U)$  is open in the uniform topology.

⟨1⟩2. LET:  $a \in \pi_j^{-1}(U)$

⟨1⟩3. PICK  $\epsilon > 0$  such that  $(a_j - \epsilon, a_j + \epsilon) \subseteq U$

⟨1⟩4.  $B_{\bar{\rho}}(a, \epsilon) \subseteq \pi_j^{-1}(U)$

□

**Proposition 393.** *The uniform topology on  $\mathbb{R}^J$  is coarser than the box topology.*

PROOF:

⟨1⟩1. LET:  $a \in \mathbb{R}^J$  and  $\epsilon > 0$

PROVE:  $B(a, \epsilon)$  is open in the box topology.

⟨1⟩2. LET:  $b \in B(a, \epsilon)$

⟨1⟩3. For  $j \in J$  we have  $|a_j - b_j| < \epsilon$

⟨1⟩4. For  $j \in J$ ,

LET:  $\delta_j = (\epsilon - |a_j - b_j|)/2$

⟨1⟩5.  $\prod_{j \in J} (b_j - \delta_j, b_j + \delta_j) \subseteq B(a, \epsilon)$

□

**Proposition 394.** *The uniform topology on  $\mathbb{R}^J$  is strictly finer than the product topology if and only if  $J$  is infinite.*

PROOF:

⟨1⟩1. If  $J$  is finite then the uniform and product topologies coincide.

PROOF: The uniform, box and product topologies are all the same.

⟨1⟩2. If  $J$  is infinite then the uniform and product topologies are different.

PROOF: The set  $B(\vec{0}, 1)$  is open in the uniform topology but not the product topology.

□



**Proposition 395 (DC).** *The uniform topology on  $\mathbb{R}^J$  is strictly coarser than the box topology if and only if  $J$  is infinite.*

PROOF:

$\langle 1 \rangle 1$ . If  $J$  is finite then the uniform and box topologies coincide.

PROOF: The uniform, box and product topologies are all the same.

$\langle 1 \rangle 2$ . If  $J$  is infinite then the uniform and box topologies are different.

PROOF: Pick an  $\omega$ -sequence  $(j_1, j_2, \dots)$  in  $J$ . Let  $U = \prod_{j \in J} U_j$  where  $U_{j_i} = (-1/i, 1/i)$  and  $U_j = (-1, 1)$  for all other  $j$ . Then  $\vec{0} \in U$  but there is no  $\epsilon > 0$  such that  $B(\vec{0}, \epsilon) \subseteq U$ .

□

**Proposition 396.** *The closure of  $\mathbb{R}^\infty$  in  $\mathbb{R}^\omega$  under the uniform topology is  $\mathbb{R}^\omega$ .*

PROOF: Given any open ball  $B(a, \epsilon)$ , pick an integer  $N$  such that  $1/\epsilon < N$ . Then  $B(a, \epsilon)$  includes sequences whose  $n$ th entry is 0 for all  $n \geq N$ . □

**Example 397.** The space  $\mathbb{R}^\omega$  is disconnected under the uniform topology. The set of bounded sequences and the set of unbounded sequences form a separation.

**Proposition 398.** *Give  $\mathbb{R}^\omega$  the uniform topology. Let  $x, y \in \mathbb{R}^\omega$ . Then  $x$  and  $y$  are in the same component if and only if  $x - y$  is bounded.*

PROOF:

$\langle 1 \rangle 1$ . The component containing 0 is the set of bounded sequences.

$\langle 2 \rangle 1$ . LET:  $B$  be the set of bounded sequences.

$\langle 2 \rangle 2$ .  $B$  is path-connected.

$\langle 3 \rangle 1$ . LET:  $x, y \in B$

$\langle 3 \rangle 2$ . PICK  $b > 0$  such that  $|x_j|, |y_j| \leq b$  for all  $j$

$\langle 3 \rangle 3$ . LET:  $p : [0, 1] \rightarrow B$  be the function  $p(t) = (1 - t)x + ty$

PROVE:  $p$  is continuous.

$\langle 3 \rangle 4$ . LET:  $t \in [0, 1]$  and  $\epsilon > 0$

$\langle 3 \rangle 5$ . LET:  $\delta = \epsilon/2b$

$\langle 3 \rangle 6$ . LET:  $s \in [0, 1]$  with  $|s - t| < \delta$

$\langle 3 \rangle 7$ .  $\bar{p}(p(s), p(t)) < \epsilon$

PROOF:

$$\begin{aligned} \bar{p}(p(s), p(t)) &= \sup_j \bar{d}((1 - s)x_j + sy_j, (1 - t)x_j + ty_j) \\ &\leq |(s - t)x_j + (t - s)y_j| \\ &\leq |s - t||x_j - y_j| \\ &< 2b\delta \\ &= \epsilon \end{aligned}$$

$\langle 2 \rangle 3$ .  $B$  is connected.

PROOF: Proposition 241.

$\langle 2 \rangle 4$ . If  $C$  is connected and  $B \subseteq C$  then  $B = C$ .

PROOF: Otherwise  $B \cap C$  and  $C \setminus B$  form a separation of  $C$ .

$\langle 1 \rangle 2$ . Q.E.D.

PROOF: Since  $\lambda x. x - y$  is a Homeomorphism of  $\mathbb{R}^\omega$  with itself.

□

## 51 Uniform Convergence

**Definition 399** (Uniform Convergence). Let  $X$  be a set and  $Y$  a metric space. Let  $(f_n : X \rightarrow Y)$  be a sequence of functions and  $f : X \rightarrow Y$  be a function. Then  $f_n$  converges uniformly to  $f$  as  $n \rightarrow \infty$  if and only if, for all  $\epsilon > 0$ , there exists  $N$  such that, for all  $n \geq N$  and  $x \in X$ , we have  $d(f_n(x), f(x)) < \epsilon$ .

**Example 400.** Define  $f_n : [0, 1] \rightarrow \mathbb{R}$  by  $f_n(x) = x^n$  for  $n \geq 1$ , and  $f : [0, 1] \rightarrow \mathbb{R}$  by  $f(x) = 0$  if  $x < 1$ ,  $f(1) = 1$ . Then  $f_n$  converges to  $f$  pointwise but not uniformly.

**Theorem 401** (Uniform Limit Theorem). Let  $X$  be a topological space and  $Y$  a metric space. Let  $(f_n : X \rightarrow Y)$  be a sequence of continuous functions and  $f : X \rightarrow Y$  be a function. If  $f_n$  converges uniformly to  $f$  as  $n \rightarrow \infty$ , then  $f$  is continuous.

PROOF:

$\langle 1 \rangle 1$ . LET:  $x \in X$  and  $\epsilon > 0$

$\langle 1 \rangle 2$ . PICK  $N$  such that, for all  $n \geq N$  and  $y \in X$ , we have  $d(f_n(y), f(y)) < \epsilon/3$

$\langle 1 \rangle 3$ . PICK a neighbourhood  $U$  of  $x$  such that  $f_N(U) \subseteq B(f_N(x), \epsilon/3)$

PROVE:  $f(U) \subseteq B(f(x), \epsilon)$

$\langle 1 \rangle 4$ . LET:  $y \in U$

$\langle 1 \rangle 5$ .  $d(f(y), f(x)) < \epsilon$

PROOF:

$$\begin{aligned} d(f(y), f(x)) &\leq d(f(y), f_N(y)) + d(f_N(y), f_N(x)) + d(f_N(x), f(x)) \quad (\text{Triangle Inequality}) \\ &< \epsilon/3 + \epsilon/3 + \epsilon/3 \quad (\langle 1 \rangle 2, \langle 1 \rangle 3) \\ &= \epsilon \end{aligned}$$

□

**Proposition 402.** Let  $X$  be a topological space and  $Y$  a metric space. Let  $(f_n : X \rightarrow Y)$  be a sequence of continuous functions and  $f : X \rightarrow Y$  be a function. Let  $(a_n)$  be a sequence of points in  $X$  and  $a \in X$ . If  $f_n$  converges uniformly to  $f$  and  $a_n$  converges to  $a$  in  $X$  then  $f_n(a_n)$  converges to  $f(a)$  uniformly in  $Y$ .

PROOF:

$\langle 1 \rangle 1$ . LET:  $\epsilon > 0$

$\langle 1 \rangle 2$ . PICK  $N_1$  such that, for all  $n \geq N_1$  and  $x \in X$ , we have  $d(f_n(x), f(x)) < \epsilon/2$

$\langle 1 \rangle 3$ . PICK  $N_2$  such that, for all  $n \geq N_2$ , we have  $a_n \in f^{-1}(B(a, \epsilon/2))$

PROOF: Using the fact that  $f$  is continuous from the Uniform Limit Theorem.

$\langle 1 \rangle 4$ . LET:  $N = \max(N_1, N_2)$

$\langle 1 \rangle 5$ . LET:  $n \geq N$

$\langle 1 \rangle 6$ .  $d(f_n(a_n), f(a)) < \epsilon$

PROOF:

$$\begin{aligned} d(f_n(a_n), f(a)) &\leq d(f_n(a_n), f(a_n)) + d(f(a_n), f(a)) \quad (\text{Triangle Inequality}) \\ &< \epsilon/2 + \epsilon/2 \quad (\langle 1 \rangle 2, \langle 1 \rangle 3) \\ &= \epsilon \end{aligned}$$

□

**Proposition 403.** *Let  $X$  be a set. Let  $(f_n : X \rightarrow \mathbb{R})$  be a sequence of functions and  $f : X \rightarrow \mathbb{R}$  be a function. Then  $f_n$  converges uniformly to  $f$  as  $n \rightarrow \infty$  if and only if  $f_n \rightarrow f$  as  $n \rightarrow \infty$  in  $\mathbb{R}^X$  under the uniform topology.*

PROOF:

- ⟨1⟩1. If  $f_n$  converges uniformly to  $f$  then  $f_n$  converges to  $f$  under the uniform topology.
- ⟨2⟩1. ASSUME:  $f_n$  converges uniformly to  $f$
- ⟨2⟩2. LET:  $\epsilon > 0$
- ⟨2⟩3. PICK  $N$  such that, for all  $n \geq N$  and  $x \in X$ , we have  $d(f_n(x), f(x)) < \epsilon/2$
- ⟨2⟩4. For all  $n \geq N$  we have  $\bar{\rho}(f_n, f) \leq \epsilon/2$
- ⟨2⟩5. For all  $n \geq N$  we have  $\bar{\rho}(f_n, f) < \epsilon$
- ⟨1⟩2. If  $f_n$  converges to  $f$  under the uniform topology then  $f_n$  converges uniformly to  $f$ .
- ⟨2⟩1. ASSUME:  $f_n$  converges to  $f$  under the uniform topology.
- ⟨2⟩2. LET:  $\epsilon > 0$
- ⟨2⟩3. PICK  $N$  such that, for all  $n \geq N$ , we have  $\bar{\rho}(f_n, f) < \min(\epsilon, 1/2)$
- ⟨2⟩4. LET:  $n \geq N$
- ⟨2⟩5. LET:  $x \in X$
- ⟨2⟩6.  $\bar{\rho}(f_n, f) < \min(\epsilon, 1/2)$
- PROOF: From ⟨2⟩3.
- ⟨2⟩7.  $d(f_n(x), f(x)) < \min(\epsilon, 1/2)$
- ⟨2⟩8.  $d(f_n(x), f(x)) < \epsilon$

□

## 52 Isometric Imbeddings

**Definition 404.** Let  $X$  and  $Y$  be metric spaces. An *isometric imbedding*  $f : X \rightarrow Y$  is a function such that, for all  $x, y \in X$ , we have  $d(f(x), f(y)) = d(x, y)$ .

**Proposition 405.** *Every isometric imbedding is an imbedding.*

PROOF:

- ⟨1⟩1. LET:  $f : X \rightarrow Y$  be an isometric imbedding.
- ⟨1⟩2.  $f$  is injective.
- PROOF: If  $f(x) = f(y)$  then  $d(f(x), f(y)) = 0$  hence  $d(x, y) = 0$  hence  $x = y$ .
- ⟨1⟩3.  $f$  is continuous.
- PROOF: For all  $\epsilon > 0$ , if  $d(x, y) < \epsilon$  then  $d(f(x), f(y)) < \epsilon$ .
- ⟨1⟩4.  $f : X \rightarrow f(X)$  is an open map.
- PROOF:  $f(B(a, \epsilon)) = B(f(a), \epsilon) \cap f(X)$ .

□