

## 1. A short sketch of laser history

1917: Einstein – stimulated absorption and emission of light

1954: Charles Townes and Schawlow – maser, prediction of the optical laser - Nobel Prize (1964)

1960: Maimann – first demonstration of a laser: Ruby laser

Rapid progress in the 1960s:

1961: first gas laser, first Nd laser

1962: first semiconductor laser

1963: CO<sub>2</sub> laser (IR)

## 2. Introduction

- ❖ A laser is a device that generates light by a process called **STIMULATED EMISSION**.
- ❖ The acronym LASER stands for Light Amplification by Stimulated Emission of Radiation

## 3. Characteristic of laser

The laser light exhibits some peculiar properties than compare with the convectional light. Those are


- ➡ Highly directionality
- ➡ Highly monochromatic
- ➡ Highly intense
- ➡ Highly coherence

### Highly directionality

The light ray coming ordinary light source travels in all directions, but laser light travels in single direction.

For example the light emitted from torch light travels 1km distance it spreads around 1 km wide.

But the laser light spreads a few centimetres distance even it travels lacks of kilometre distance.



The directionality of laser beam is expressed in terms of divergence  $\phi$

$$\phi = \frac{\text{arc}}{\text{radius}} = \frac{d_2 - d_1}{s_2 - s_1}$$

Where  $d_2$  and  $d_1$  are the diameters of laser spots at distances of  $s_2$  and  $s_1$  respectively from laser source.

For laser light divergence  $\phi = 10^{-3} \text{ radians}$ .

Since the divergence of light is very low, so we say that the laser light having highly directional.

### Highly monochromatic

In laser radiation, all the photons emitted between discrete energy levels will have same wavelength.

As a result the radiation is monochromatic in nature.

Due to the stimulated characteristic of laser light, the laser light is more monochromatic than that of a convectional light.

laser radiation -the wavelength spread = 0.001 nm

So it is clear that the laser radiation is highly monochromatic



### Highly intense


- ✚ Laser light is highly intense than the conventional light.
- ✚ one mill watt He-Ne laser is highly intense than the sun intensity
- ✚ when two photons each of amplitude  $a$  are in phase with other; the resultant amplitude of two photons is  $2a$  and the intensity is  $4a^2$
- ✚ in laser much number of photons are in phase with each other, the amplitude of the resulting wave becomes  $na$  and hence the intensity of laser is proportional to  $n^2a^2$
- ✚ So 1mW He-Ne laser is highly intense than the sun

### Highly coherence

- ✚ Coherence is the property of the wave being in phase with itself and also with another wave over a period of time, and space or distance. There are two types of coherence
- ✚ Temporal coherence
- ✚ Spatial coherence.
- ✚ For laser radiation all the emitted photons are in phase, the resultant radiation obeys spatial and temporal coherence.


#### Temporal coherence (or longitudinal coherence):-

The predictable correlation of amplitude and phase at one point on the wave train w.r.t another point on the same wave train, then the wave is said to be temporal coherence.



#### spatial coherence (or transverse coherence).

The predictable correlation of amplitude and phase at one point on the wave train w.r.t another point on a second wave, then the waves are said to be spatially coherence (or transverse coherence).



### 4. Stimulated absorption

Let  $E_1$  and  $E_2$  are the energies and  $N_1$  and  $N_2$  are the number of atoms per unit volume of ground and excited states and  $\rho(\nu)$  be the density of photon density.

Suppose, if a photon of energy  $E_2 - E_1 = h\nu$  interacts with an atom present in the ground state, the atom gets excitation from ground state to excited state by absorbing the photon energy.

It is the **process of excitation of atom into excited state from ground state by absorbing the incident photon.**



Stimulated absorption rate depends upon the number of atoms available in the lowest energy state as well as the energy density photons.

Stimulated absorption rate  $\propto N_1$

$$\propto \rho(\nu)$$

$$\propto \rho(\nu) N_1$$


$$= B_{12} \rho(\nu) N_1$$

Where  $B_{12}$  is known as Einstein coefficient of stimulated absorption



## 5. Spontaneous emission

- Spontaneous emission was postulated by Bohr.
- The excited atom does not stay in a long time in the excited state.
- The lifetime of excited atom in higher state  $E_2$  is up to  $10^{-8}$  seconds.
- After the life time of the excited atom, gets de-excited into ground by emitting a photon of energy  $E_2 - E_1 = h\nu$ .



It is the process of de-excitation of atom itself into ground state after its life time from excited state by emitting a photon

$$\begin{aligned}\text{Spontaneous emission rate} &\propto N_2 \\ &= A_{21} N_2\end{aligned}$$

Where  $A_{21}$  is known as Einstein coefficient of spontaneous emission


Characteristics of spontaneous emitted radiation

- The emitted radiation is poly-monochromatic
- The emitted radiation is Incoherent
- The emitted radiation is less intense
- The emitted radiation has less directionality
- Example: light from sodium or mercury lamp

## 6. Stimulated emission

Stimulated emission was postulated by Einstein.

Let, a photon of energy  $E_2 - E_1 = h\nu$  interacts with the excited atom within their life time; the atom gets de-excitation into ground state by emitting of additional photon.



It is the process of de-excitation of atom into ground state from excited state by interacting with an additional photon within its life time by emitting of an additional photon.

Stimulated emission rate depends upon the number of atoms available in the excited state as well as the energy density of photons.

$$\begin{aligned}\text{Stimulated emission rate} &\propto N_2 \\ &\propto \rho \\ &\propto N_2 \rho \\ &= B_{21} N_2 \rho\end{aligned}$$

Where  $B_{21}$  is known as Einstein coefficient of stimulated emission

### Characteristics of stimulated emitted radiation

- The emitted radiation is monochromatic
- The emitted radiation is Coherent
- The emitted radiation is high intense
- The emitted radiation has high directionality
- Example: light from laser source

### 7. Spontaneous and Stimulated emission

Spontaneous emission	Stimulated emission
1. The spontaneous emission was postulated by Bohr	1. The stimulated emission was postulated by Einstein
2. Additional photons are not required in spontaneous emission	2. Additional photons are required in stimulated emission
3. One photon is emitted in spontaneous emission	3. Two photons are emitted in stimulated emission
4. The emitted radiation is poly-monochromatic	4. The emitted radiation is monochromatic
5. The emitted radiation is Incoherent	5. The emitted radiation is Coherent
6. The emitted radiation is less intense	6. The emitted radiation is high intense
7. The emitted radiation have less directionality	7. The emitted radiation have high directionality
8. Example: : light from sodium or mercury lamp	8. Example: light from laser source


### 9. Population inversion

Let us consider two level energy system of energies  $E_1$  and  $E_2$  as shown in figure. Let  $N_1$  and  $N_2$  be the populations of energy levels  $E_1$  and  $E_2$ . The number of atoms present in an energy level is known as population of that energy level. At ordinary conditions, i.e., the population in the ground or lower state is always greater than the population in the excited or higher states. The stage of making, population of higher energy level is greater than the population of lower energy level is called population inversion.

According to Boltzmann's distribution the population of an energy level  $E_i$  at temperature  $T$  is given by

$$N_i = N_0 e^{\left(\frac{-E_i}{KT}\right)}$$

Where  $N_0$  is the population of the lower level or ground state and  $k$  is the Boltzmann's constant.




*The number of atoms present in the excited state ( $N_2$ ) is greater than the number of atoms present in the ground state ( $N_1$ ) is called population inversion.*



The process of raising the particles from ground state to excited state to achieve population inversion is called pumping. (Or the process of achieving of population inversion is called pumping)

To understand the concept of laser emission (stimulated emission) let us consider a three energy level system with energies  $E_1$ ,  $E_2$  and  $E_3$  of populations,  $N_1$ ,  $N_2$  and  $N_3$ . At normal conditions,  $E_1 < E_2 < E_3$  and  $N_1 > N_2 > N_3$ . In the ground state the life time of atom is more and the life time of atom in the excited state is  $10^{-8}$  sec. But in the intermediate state the atom has more life time. So it is called metastable state.



When a suitable energy is supplied to the system, atoms get excited into  $E_3$ . After their lifetime the atoms are transit into  $E_2$ . Due to more lifetime of an atom in state  $E_2$ , the atoms stay for longer time than compare with the state  $E_3$ . Due to the accumulation of atoms in state  $E_2$ , the population inversion is established in between the  $E_1$  and  $E_2$  states.

## 10. Types of lasers


On the basis of active medium used in the laser systems, lasers are classified into several types

- |                         |  |
|-------------------------|--|
| I. Solid lasers         | : Ruby laser, Nd:YAG laser, Nd:Glass       |
| II. Liquid lasers       | : Europium Chelate laser, $\text{SeOCl}_2$ |
| III. Gas lasers         | : $\text{CO}_2$ , He-Ne, Argon-Ion Laser   |
| IV. Dye lasers          | : Rhodamine 6G                             |
| V. Semiconductor lasers | : InP, GaAs.                               |
| VI. Chemical lasers     | : HF, DF.                                  |

## 11. Construction and components of laser

Generally, every laser system consists of three components. They are

- Energy source
- Active medium
- Optical cavity or resonator



## Energy source

- ✦ To get laser emission, first we must have population inversion in the active medium.
- ✦ The energy source supplies the energy to the active medium.
- ✦ By absorbing that energy, the atoms or molecules or ions can be excited into higher levels.
- ✦ As a result we get population inversion in the active medium.



## Active medium

**Definition:** - *In which medium we are creating population inversion to get stimulated emission of radiation is called active medium.*

- ✚ After receiving the energy from the source, the atoms or molecules or ions get excited into higher energy levels.
- ✚ While de-excitation to lower energy level, the emitted photons start stimulated emission which results in laser emission.
- ✚ Depending upon the active medium, lasers are classified as solid state, liquid state, gaseous state and semiconductor lasers.

## Optical cavity or resonator

- ✚ *The active medium is enclosed between a fully reflective mirror and partially reflective mirror. These mirrors constitute the optical cavity or resonator.*
- ✚ The reflectors enhance the stimulated emission process by reflecting the photons into the active medium.
- ✚ As a result, we get high-intensity monochromatic and coherent laser light through the partially reflecting portion of the mirror.

## 12. Excitation mechanisms

Excitation of an atom can be done by number of ways. The most commonly used excitation methods are

- ✚ Optical pumping
- ✚ Electrical discharge pumping
- ✚ Chemical pumping
- ✚ Injection current pumping

## Optical pumping

- ✚ Optical pumping is a process in which light is used to raise the atoms from a lower energy level to a higher level to create population inversion.
- ✚ Optical pumping is used in solid lasers.
- ✚ The solid materials have a very broad absorption band, so a sufficient amount of energy is absorbed from the emission band of a flash lamp to create population inversion.
- ✚ Xenon flash tubes are used for optical pumping.
- ✚ Examples: - Ruby laser, Nd:YAG Laser (Neodymium: Yttrium Aluminum Garnet), Nd: Glass Laser

## Electrical discharge pumping

- ✚ In electric discharge pumping, atoms are excited into an excited state by collisions with fast moving electrons in an electric discharge tube.
- ✚ Electrical discharge pumping is used in gas lasers.
- ✚ Since gas lasers have a very narrow absorption band, so optical pumping is not suitable for gas lasers.
- ✚ Examples:- He-Ne laser, CO<sub>2</sub> laser, argon-ion laser, etc

## Chemical pumping

- ✚ In this method, the chemical energy released during the chemical process, that energy will excite the atoms to a higher level and create population inversion.
- ✚ Whenever hydrogen reacts with fluorine, it liberates a lot of heat energy. By utilizing this heat energy, the atoms get excited into higher states and create population inversion.
- ✚ Examples:- HF and DF lasers.

## Injection current pumping

- ✚ This pumping mechanism is used in semiconductor lasers.
- ✚ In semiconductor lasers, by passing high currents across the junction, the population inversion will be created.
- ✚ In semiconductor lasers, the population inversion always occurs among majority and minority charge carriers.
- ✚ Examples:- InP and GaAs lasers




## 6. Ruby laser

Ruby laser is a three level solid state laser and was constructed by Mainmann in 1960. Ruby ( $\text{Al}_2\text{O}_3 + \text{Cr}_2\text{O}_3$ ) is a crystal of Aluminium oxide, in which 0.05% of  $\text{Al}^{+3}$  ions are replaced by the  $\text{Cr}^{+3}$  ions. The colour of the rod is pink. The active medium in the ruby rod is  $\text{Cr}^{+3}$  ions.

### Construction

In ruby laser 4cm length and 5mm diameter rod is generally used. Both the ends of the rods are highly polished and made strictly parallel. The ends are silvered in such a way, one becomes partially reflected and the other end fully reflected. The ruby rod is surrounded by xenon flash tube, which provides the pumping light to excite the chromium ions in to upper energy levels.




Xenon flash tube emits thousands joules of energy in few milli seconds, but only a part of that energy is utilized by the chromium ions while the rest energy heats up the apparatus. A cooling arrangement is provided to keep the experimental set up at normal temperatures

### Working

The energy level diagram of chromium ions is shown in figure.

The chromium ions get excitation into higher energy levels by absorbing of  $5600\text{\AA}$  of wave length radiation. The excited chromium ions stay in the level H for short interval of time ( $10^{-8}$  Sec). After their life time most of the chromium ions are de-excited from H to G and a few chromium ions are de-excited from H to M.



The transition between H and M is non-radioactive transition i.e. the chromium ions give their energy to the lattice in the form of heat. In the Metastable state the life time of chromium ions is  $10^{-3}$  sec. The life time of chromium ions in the Metastable state is  $10^5$  times greater than the life time of chromium ions in higher state.

Due to the continuous working of flash lamp, the chromium ions are excited to higher state H and returned to M level. After few milliseconds the level M is more populated than the level G and hence the desired population inversion is achieved. The state of population inversion is not a stable one. The process of spontaneous transition is very high. When the excited chromium ion passes spontaneously from H to M it emits one photon of wavelength  $6943 \text{ Å}$ . The photon reflects back and forth by the silver ends and until it stimulates an excited chromium ion in M state and it to emit fresh photon in phase with the earlier photon. The process is repeated again and again until the laser beam intensity is reached to a sufficient value. When the photon beam becomes sufficient intense, it emerges through the partially silvered end of the rod. The wavelength  $6943 \text{ Å}$  is in the red region of the visible spectrum.

### Draw backs of ruby laser

- The laser requires high pumping power
- The efficiency of ruby laser is very small
- It is a pulse laser

### Uses of ruby laser


1. Ruby lasers are in optical photography
2. Ruby lasers can be used for measurement of plasma properties such as electron density and temperature.
3. Ruby lasers are used to remove the melanin of the skin.
4. Ruby laser can be used for recording of holograms.



#### 14. Semiconductor diode laser

- ✚ Laser diode is a specially fabricated  $p$ - $n$  junction device that emits coherent radiation.
- ✚ It is operated at forward biased condition.
- ✚ Direct band gap semiconductors are preferred in the fabrication of semiconductor laser diodes because they emit energy in terms of light when an electron and hole recombination takes place.
- ✚ Compound semi-conductors like GaAs and InP are examples for direct band gap semiconductors


#### Construction



- ✚ In this laser system, the active medium is a  $p$ - $n$  junction diode made from crystalline gallium arsenide.
- ✚ The  $p$ -region and  $n$ -region in the diode are obtained by heavily doping with germanium and tellurium respectively in GaAs.
- ✚ The thickness of the  $p$ - $n$  junction is very narrow so that the emitted radiation has large divergence and poor coherence.
- ✚ At the junction two sides are roughed to avoid laser emission and the remaining two faces one is partially polished and the other is fully polished.
- ✚ The laser emission takes place from the partially polished face.
- ✚ To provide bias two metal contacts are provided in the top and bottom of the diode as shown in figure.

## Working

- The semiconductor laser device is always operated in forward bias condition.
- Electrons and the holes are the minority charge carriers in n-region and p-region semiconductors.
- When a huge current ( $10^4 \text{ Amp/mm}^2$ ) is passing through the p-n junction, p-region is positively biased, holes are injected into n-region from p-region and n-region is negatively biased electrons are injected into p-region from n-region as shown in figure.



- The continuous injection of charge carriers creates the population inversion of minority carriers in n and p sides' respectively.
- The electrons and holes recombine and release of light energy takes place in or near the junction as shown in figure.
- The emitted photons increase the rate of recombination of injected electrons from the n-region and holes in p-region by inducing more recombinations.

From Planck's law  $E_g = hv = h \frac{c}{\lambda}$

$$\lambda = \frac{hc}{E_g} = \frac{6.63 \times 10^{-34} \times 3 \times 10^8}{1.4 \times 1.6 \times 10^{-19}} = 8874 \text{Å}$$

- In case of GaAs homo-junction which has an energy gap of 1.44eV gives a laser beam of wave length around 8874Å.
- The wave length of emitted radiation depends up on the concentration of donor and acceptor atoms in GaAs.
- The efficiency of the laser emission is increases when we cool the GaAs diode.



## **16. Applications of lasers**

Due to high intensity, high monochromaticity and high directionality of lasers, they are widely used in various fields like

1. communication
2. computers
3. chemistry
4. photography
5. industry
6. medicine
7. military
8. scientific research

### **1. communication**

In case of optical communication semiconductor laser diodes are used as optical sources and its bandwidth is ( $10^{14}$  Hz) is very high compared to the radio and microwave communications.

- More channels can be sent simultaneously
- Signal cannot be tapped
- As the bandwidth is large, more data can be sent.
- A laser is highly directional and less divergence, hence it has greater potential use in space crafts and submarines.

### **2. Computers**

- In LAN (local area network), data can be transferred from memory storage of one computer to other computer using laser for short time.
- Lasers are used in CD-ROMS during recording and reading the data.

### **3. Chemistry**

- Lasers are used in molecular structure identification
- Lasers are also used to accelerate some chemical reactions.
- Using lasers, new chemical compounds can be created by breaking bonds between atoms and molecules.

### **4. Photography**

- Lasers are also used in the construction of holograms.
- Lasers can be used to get 3-D lensless photography.

### **5. Industry**

- Lasers can be used to blast holes in diamonds and hard steel
- Lasers are also used as a source of intense heat
- Lasers are used to drill holes in ceramics.
- Lasers are used to cut glass and quartz.
- Lasers are used for heat treatment in the tooling and automotive industry.
- Lasers are used in electronic industry in trimming the components of ICs.
- High power lasers are used to weld or melt any material.
- Lasers are also used to cut teeth in saws and test the quality of fabric.

### **6. Medicine**

- Lasers are used for cataract removal.
- Lasers are used for eye lens curvature corrections.
- Lasers are used in bloodless surgery.
- Lasers are used in cancer diagnosis and therapy.
- Lasers are used in destroying kidney stones and gallstones.
- Argon and carbon dioxide lasers are used in the treatment of liver and lungs.
- Lasers used in endoscopy to scan the inner parts of the stomach.
- Lasers used in the elimination of moles and tumours which are developing in the skin tissue.
- Lasers are used in plastic surgery.
- Lasers are used in the treatment of mouth diseases.

### **7. Military**

- Lasers can be used as a war weapon.
- High energy lasers are used to destroy the enemy aircrafts and missiles.
- Lasers can be used in the detection and ranging like RADAR.

## 8. Scientific field

- Lasers are used for isotope preparation.
- Lasers are employed to create plasma.
- Lasers are used in air pollution, to estimate the size of the dust particles.
- Lasers are used in the field of 3D-photography
- Lasers used in Recording and reconstruction of hologram.
- Lasers used to produce certain chemical reactions.
- Lasers are used in Raman spectroscopy to identify the structure of the molecule.
- Lasers are used in the Michelson- Morley experiment.