

Report_Los_Angeles_Data

Mercat Mathilde & Radici Andrea

In this report we present preliminary exploration of the LA dataset. Highlighting the potential analysis as well as the missing data. So far we work on sepulveda and eldorado dataset.

Part 1 : Dataset description

1 - Surveillance coverage

Starting in the late 80s, the dataset includes 161 traps of 6 types (BGSENT, CO2, GRVD, NJLT, USDS, OTHER?) across 81 sites (trap location) in 2 areas (Sepulveda and Eldorado). Figure 1 shows, until 2024 only 3 traps NJLT type were set up (2 in Sepulveda and 1 in Eldorado). Then, we observe a lot of variation in the number of traps set up.

- > All traps do not have the same number of monitoring weeks.
- > Period of high monitoring after 2003.
- > Period of very intense monitoring between 2004 and 2010.

1.a. Surveillance effort over years

Surveillance efforts vary over years and areas. Higher effort of surveillance occurs around 2005 and then decreases. Early season monitoring decreases a lot after 2011 with almost no traps before mid-May.

1.b. Surveillance effort by trap type

Different types of traps are not used consistently. NJLT was the trap used at the beginning of surveillance. But since CO2 and GRVD traps are the most used.

1.c.

Trap_ID = one location and one type of trap.

Figure 4a shows that the number of traps (one location, one type of trap at least set up once a year) each year varies. Some traps have been set up for years, others as been set up sporadically (Figure 4b).

Figure 4a page 8

2 - Diversity of mosquito

19 different species have been identified. *Cx. quiquefasciatus* represents 93% of the total sample mosquitoes (Figure 5: histogram).

3 - Type of trap impact on species

Figure 6 shows that long old series NJLT traps do not capture a lot of *Cx. quiquefasciatus*.

#Figure Diapo 11

Including Plots

You can also embed plots, for example:

Note that the `echo = FALSE` parameter was added to the code chunk to prevent printing of the R code that generated the plot.