

Math 122A Homework 6

Rad Mallari

March 1, 2022

1 Problem 1

Let C be a closed, positive, and simple curve. Using Green's Theorem prove that

$$\frac{1}{2i} \int_C \bar{z} dz = \text{area enclosed by } C$$

Proof. Since $\bar{z} = x - iy$, this is equivalent to

$$\frac{1}{2i} \int_C (x - iy)(dx + idy) = \frac{1}{2i} \left[\int_C (xdx + ydy) + i \int_C (xdy - ydx) \right]$$

Where D is the area bounded by C . Now using Green's Theorem, twice we have that

$$\begin{aligned} \frac{1}{2i} \left[\iint_D (0 - 0) dx dy + i \iint_D (1 - (-1)) dx dy \right] \\ = \iint_D dx dy \end{aligned}$$

Which after evaluating the integrals is exactly the area enclosed by C . \square

2 Problem 2

Consider the function $f(z) = (z + 1)^2$ and the region R bounded by the triangle with vertices $0, 2, i$ (its boundary and interior). Find the points where $|f(z)|$ reaches its maximum and minimum value of R .

Proof. By the Maximum Modulus Theorem, we know that $|f(z)| = |z + 1|^2$. Then the minimum would be at the boundary where $|f(z)|$ is the closest to

-1 and maximum is the furthest. Then we know that since $|f(z)| = 1$ at $z = 0$, then this is the minimum, and since $|f(z)| = 9$ at $z = 2$, then this is the maximum. \square

3 Problem 3

Find the maximum of $|\sin(z)|$ on $[0, 2\pi] \times [0, 2\pi]$.

Proof. We can rewrite $\sin(z)$ as:

$$\begin{aligned}\sin(z) &= \frac{e^{iz} - e^{-iz}}{2i} = \frac{e^{x+iy} - e^{-i(x+iy)}}{2i} \\ &= \frac{e^{ix}e^{-y} - e^{-ix}e^y}{2i} \\ &= \frac{(\cos x + i \sin x)e^{-y} - (\cos x - i \sin x)e^y}{2i} \\ &= -\frac{1}{i} \cos(x) \left(\frac{e^{-y} - e^y}{2} \right) + \sin(x) \left(\frac{e^y + e^{-y}}{2} \right) \\ &= i \cos(x) \sinh(y) + \sin(x) \cosh(y)\end{aligned}\tag{1}$$

And using Maximum Modulus Theorem, and (1), we have that:

$$|\sin(z)|^2 = \cos^2(x) \sinh^2(y) + \sin^2(x) \cosh^2(y)$$

Using the identities that states $\cosh^2(y) - \sinh^2(y) = 1$ and $\cos^2(x) + \sin^2(x) = 1$, we can rewrite this as:

$$|\sin(z)|^2 = \cos^2(x) \sinh^2(y) + \sin^2(x) \cdot (1 + \sinh^2(y))$$

Leaving us with

$$|\sin(z)|^2 = \sinh^2(y) + \sin^2(x)$$

We know that that maximum of $\sin^2(x) = 1$ which is at $x = \frac{\pi}{2}$ and $x = \frac{3\pi}{2}$, meanwhile the maximum of $\sinh^2(y)$ is located at $y = 2\pi$. Therefore, our maximum is at the boundaries 2π . \square

4 Problem 4

Calculate:

(a)

$$\int_0^{2\pi} \frac{d\theta}{a + b \cos(\theta)}, \quad 0 < b < a$$

HINT: Work backwards using $\cos(\theta) = \frac{e^{i\theta} + e^{-i\theta}}{2}$ to convert the integral into a complex integral along the curve $|z| = 1$

(b)

$$\int_0^{2\pi} \frac{d\theta}{(a + b \cos(\theta))^2}$$

(c)

$$\int_0^{2\pi} \frac{\sin(\theta) d\theta}{(a + b \cos(\theta))^2}, \quad 0 < b < a$$

Proof.

(a) By the hint, we work backwards and get

$$\int_0^{2\pi} \frac{d\theta}{a + b \left(\frac{e^{i\theta} + e^{-i\theta}}{2} \right)}$$

Factoring out a 2 gives us

$$2 \int_0^{2\pi} \frac{d\theta}{2a + b(e^{i\theta} + e^{-i\theta})}$$

Letting $z(\theta) = e^{i\theta}$, where $z : [0, 2\pi] \rightarrow \mathbb{C}$, implies $d\theta = \frac{dz}{iz}$ and our line integral becomes

$$I = \frac{2}{i} \oint_{|z|=1} \frac{dz}{bz^2 + 2az + b}$$

Here our $f(z) = \frac{1}{bz^2 + 2az + b}$ is analytic except at $\frac{a \pm \sqrt{a^2 - b^2}}{b}$. These two points are:

$$z_1 = \frac{a + a\sqrt{1 - \frac{b^2}{a^2}}}{b} \quad \text{and} \quad z_2 = \frac{a - a\sqrt{1 - \frac{b^2}{a^2}}}{b}$$

By our condition that $0 < b < a$, we know z_2 must be outside our z , therefore letting $h(z) = \frac{1}{z - \left(\frac{a - a\sqrt{1 - \frac{b^2}{a^2}}}{b} \right)}$ and by Cauchy Theorem we get

$$I = \frac{2}{i} \oint_{|z|=1} \frac{h(z_2)dz}{z - \left(\frac{a + a\sqrt{1 - \frac{b^2}{a^2}}}{b} \right)} = \frac{2\pi b}{\sqrt{a^2 - b^2}}$$

(b) Similarly, letting $\cos(\theta) = \frac{e^{i\theta} + e^{-\theta}}{2}$ yields

$$\int_0^{2\pi} \frac{d\theta}{\left(a + b \left(\frac{e^{i\theta} + e^{-\theta}}{2} \right) \right)^2}$$

Then letting $z(\theta) = e^{i\theta}$, $d\theta = \frac{dz}{iz}$, we get

$$\frac{4}{i} \oint_{|z|=1} \frac{zdz}{(bz^2 + 2az + b)^2}$$

Our singularities here are the same as **(a)** which are

$$z_1 = \frac{a + a\sqrt{1 - \frac{b^2}{a^2}}}{b} \quad \text{and} \quad z_2 = \frac{a - a\sqrt{1 - \frac{b^2}{a^2}}}{b}$$

But z_1 in this case is outside of $|z| = 1$.

(c)

□

5 Problem 5

Prove that if $f : \mathbb{C} \rightarrow \mathbb{C}$ is entire such that for some $n \in \mathbb{N}$

$$\lim_{|z| \rightarrow \infty} \frac{|f(z)|}{|z|^n} = M < \infty,$$

then f is a polynomial of degree at most n .

Proof. Since f is analytic, by a theorem in **Lecture 16** which states that f has a power series expansion in the neighborhood of analyticity that is

$$f(z_0) = \sum_{n=0}^{\infty} (z_0 - z_1)^n \left(\frac{f^n(z_1)}{n!} \right)$$

In our case, $z_1 = 0$ so this becomes

$$f(z_0) = \sum_{n=0}^{\infty} (z_0)^n \left(\frac{f^n(0)}{n!} \right)$$

By **Section 49** of the book, we get

$$|f^n(0)| \leq \frac{n! M_R}{R^n}$$

Where M_R denotes the maximum value of $|f(z)|$. □

6 Problem 6

Let $A \subset \mathbb{C}$ be an open set and $f : A \rightarrow \mathbb{C}$ be an analytic function on A . Assuming that $z_0 \in A$ such that

$$\{z \in \mathbb{C} : |z - z_0| \leq R\}, \quad R > 0$$

prove that

$$f(z_0) = \frac{1}{\pi R^2} \iint_{|z-z_0| \leq R} f(x+iy) dx dy$$

Proof. □

7 Problem 7

Let $f : R \rightarrow R$ be defined as

$$f(x) = e^{\frac{-1}{x^2}} \quad \text{if } x \neq 0, \quad f(0) = 0$$

Show that f is infinitely differentiable $\forall n \in \mathbb{N}$, $f^{(n)}(0) = 0$. Verify that the power series of f at $x = 0$ does not agree with f in any neighborhood of 0.

Proof. □