

**Numerical Methods (ENUME 2019) – Project**  
**Assignment C: Solving ordinary differential equations**

1. Develop a program for solving the following differential equation:

$$9y'' + 6y' + 10y = 0 \text{ for } t \in [0, 10], \quad y(0) = 0 \text{ and } y'(0) = 2$$

by means of the implicit Lobatto IIID order 2 method defined by the following Butcher table:

$$\begin{array}{c|cc} 0 & \frac{1}{2} & \frac{1}{2} \\ 1 & -\frac{1}{2} & \frac{1}{2} \\ \hline & \frac{1}{2} & \frac{1}{2} \end{array}$$

Compare the solution, obtained by means of this program for the constant integration step  $h = 0.01$ , with the solution, obtained by means of the MATLAB operator *ode113*. Optimise the parameters *RelTol* and *AbsTol* of the latter in such a way as to get the most accurate solution  $\dot{\mathbf{y}}(t; h)$  (to be used as a reference in the following sections).

2. Carry out a systematic investigation of the dependence of the accuracy of the solution  $\hat{\mathbf{y}}(t; h)$  on the integration step  $h$ . Use the following accuracy indicators for this purpose:

$$\delta_2(h) = \frac{\|\hat{\mathbf{y}}(t; h) - \dot{\mathbf{y}}(t, h)\|_2}{\|\dot{\mathbf{y}}(t, h)\|_2} \quad (\text{the root-mean-square error})$$

$$\delta_\infty(h) = \frac{\|\hat{\mathbf{y}}(t; h) - \dot{\mathbf{y}}(t, h)\|_\infty}{\|\dot{\mathbf{y}}(t, h)\|_\infty} \quad (\text{the maximum error})$$

Make the graphs  $\delta_2(h)$  and  $\delta_\infty(h)$ .

3. Repeat the systematic investigation, defined in Section 3, for the implicit Euler method. Add the curves representative of  $\delta_2(h)$  and  $\delta_\infty(h)$  to the graph obtained in Section 2.