

Boundary currents and residue traces

Raez Lorgat

September 2026

Holomorphic BF theory and its boundary current

A current algebra can lose its central term while its bulk field theory retains a nondegenerate Batalin–Vilkovisky pairing. The mechanism is a change in the differential, with two field components kept separate. On a complex curve times a real half-line, the quadratic action density is

$$\mathcal{L}_\kappa(a, b) = b \wedge d' a + \frac{\kappa}{2} a \wedge \partial a, \quad d' = \bar{\partial} + d_t.$$

It is defined for smooth fields. On compactly supported fields the action is $S_\kappa(a, b) = \int_{\mathcal{M}}^{\text{res}} \mathcal{L}_\kappa(a, b)$. Here a is a partial connection in the antiholomorphic and real directions, and b is a holomorphic one-form. The pairing between these fields is independent of κ . At $\kappa = 0$ the action is holomorphic BF theory: the field b multiplies the partial curvature $d' a$. A compactly supported collar map identifies its quantum boundary observables with the boundary value of the bulk theory. A Cauchy contraction then extracts every current field and operator product from those supported observables. With the positive unary differential fixed below, the boundary current has level $-\kappa$. At zero its singular product vanishes, while its polynomial fields and all holomorphic jets remain.

1.1 THE FIELDS, INCLUDING THEIR ODD COMPONENTS

Let $\Sigma = \mathbb{C}$ with coordinate $z = x + iy$, and let $\mathcal{M} = \Sigma \times \mathbb{R}_{\geq 0}$ with coordinate t on the second factor. Choose the orientation whose induced boundary orientation on $\Sigma \times \{0\}$ is the complex orientation. Write $\mathcal{A}$ for smooth forms generated by $d\bar{z}, dt$. All coefficients are complex-valued, and

$$d' = d\bar{z} \wedge \partial_{\bar{z}} + dt \wedge \partial_t, \quad \partial = dz \wedge \partial_z.$$

These operators satisfy $(d')^2 = \partial^2 = d' \partial + \partial d' = 0$. The parameter $\kappa \in \mathbb{C}$ is a scalar of cohomological degree zero. Statements polynomial in κ also hold over $\mathbb{C}[\kappa]$.

The coefficient space of α is $\mathbb{C}e$, and that of β is its dual $\mathbb{C}e^\vee$. These chosen bases suppress the evaluation pairing in the scalar formulas. The fields have two summands,

$$E_\kappa = \mathcal{A}[1] \oplus (\mathcal{A} \wedge dz)[1].$$

This is the unrestricted smooth field complex. Write $E_{\kappa, c}$ for its subcomplex of compactly supported fields; compact support is required in both the curve and the real direction and may meet the boundary

of M . The notation records the following precise degrees: an ordinary p -form in either summand has cohomological degree $p - 1$. The differential is the actual exterior operator

$$Q_\kappa(\alpha, \beta) = (d'\alpha, d'\beta + \kappa d\alpha).$$

Thus the holomorphic one-form β has degree zero before adding any $d\bar{z}$ or dt . All tensor substitutions use the Koszul sign of the displayed cohomological degrees.

For a component of the field complex, the corresponding coordinate has the opposite degree. This coordinate degree is the ghost number. The complete list is

component	form degree	ghost number	role
$\alpha^{(0)}$	0	1	gauge ghost c
$\alpha^{(1)} = a$	1	0	partial connection
$\beta^{(1)} = b$	1	0	holomorphic one-form
$\alpha^{(2)}$	2	-1	antifield of b
$\beta^{(2)}$	2	-1	antifields of a
$\beta^{(3)}$	3	-2	antifield of c .

There is no $\alpha^{(3)}$, because $\mathcal{A}$ has only two exterior generators. Both summands are essential to the BF presentation.

To fix the current normalization, define

$$\int^{\text{res}} = \frac{1}{2\pi i} \int$$

on compactly supported complex-valued top forms, both on M and on Σ . The second integral is ordinary oriented integration. In contour notation $\text{Res}_{z=0} z^{-1} dz = 1$. For a compactly supported two-form θ on Σ and a compactly supported function ρ on the half-line, fiber integration has the sign

$$\int_M^{\text{res}} \theta \wedge \rho(t) dt = - \left(\int_0^\infty \rho(t) dt \right) \int_\Sigma^{\text{res}} \theta.$$

Indeed, the orientation of M is the negative of the product orientation on $\Sigma \times \mathbb{R}_{\geq 0}$. Changing the common scalar normalization rescales the action and inverse pairing together.

For homogeneous smooth fields $u = (\alpha, \beta)$ of form degree p and $v = (\alpha', \beta')$, define the local pairing density

$$\omega_{\text{loc}}(u, v) = (-1)^{p+1} (\alpha \wedge \beta' + \beta \wedge \alpha')_{[3]},$$

where $[3]$ takes the top-form component. The orientation identifies this top form with a complex-valued density. Its integral

$$\omega(u, v) = \int_M^{\text{res}} \omega_{\text{loc}}(u, v)$$

is defined whenever at least one argument has compact support. Thus the integrated pairing is defined on $E_{\kappa, c} \otimes E_\kappa$ and on $E_\kappa \otimes E_{\kappa, c}$; no integral is assigned to an arbitrary pair of unrestricted smooth fields. The local pairing has degree -1 , is independent of κ , and is fiberwise nondegenerate: the two complementary summands pair by evaluation and wedge product. Its sign gives graded skew-symmetry. Indeed, a nonzero pairing has $p + q = 3$, so interchanging the forms leaves their wedge sign unchanged and changes $(-1)^{p+1}$ to its negative.

The distinction is already necessary for fields away from the boundary. If χ is a nonzero real smooth function supported in $(1, 2)$, then $u = (\chi(t)dt, 0)$ and $v = (0, \chi(t)dz \wedge d\bar{z})$ have a well-defined local pairing density, but its integral contains the infinite area of $\mathbb{C}$. Their support is not compact in M .

1.2 THE BV IDENTITY AND THE CHIRAL BOUNDARY CONDITION

The square of the differential is zero for every κ :

$$Q_\kappa^2(\alpha, \beta) = (\circ, \kappa(d'\partial + \partial d')\alpha) = \circ.$$

The linear complex is elliptic, including at zero level. For a nonzero real covector ξ , the symbol of d' is wedge multiplication by $\xi^{\circ,1} + \xi_t dt$. This one-form is nonzero: a real covector with zero $(\circ, 1)$ part and zero t part is zero. Contraction with a dual vector contracts this symbol complex. Choose that vector with zero z component. Its contraction anticommutes with the symbol of ∂ , so it also contracts the symbol of Q_κ .

Proposition 1.1 (A nondegenerate BV family). *For every $\kappa \in \mathbb{C}$, the fields, differential and local pairing above define a free BV theory in the interior of M . The boundary condition*

$$L = \Omega^{\bullet}(\Sigma), \quad \iota^* \alpha = \circ, \quad \iota^* \beta \in L, \quad \iota : \Sigma \hookrightarrow M,$$

is a local Lagrangian boundary condition. The restricted differential is compatible with the integrated pairing whenever at least one argument has compact support.

Proof. Ellipticity and nondegeneracy were proved above. For homogeneous u of form degree p , the exterior Leibniz rule gives the local identity

$$\begin{aligned} \omega_{\text{loc}}(Q_\kappa u, v) + (-1)^{p-1} \omega_{\text{loc}}(u, Q_\kappa v) \\ = (-1)^p [d'(\alpha \wedge \beta' + \beta \wedge \alpha') + \kappa \partial(\alpha \wedge \alpha')]_{[3]}. \end{aligned}$$

Assume now that at least one of u, v has compact support. The primitives on the right then have compact support, so their Stokes integrals have no contribution from infinity. The second term integrates to zero: its possible primitive is a two-form generated by $d\bar{z}, dt$, whose boundary pullback vanishes. The first term is an ordinary total derivative, since applying ∂ to its primitive gives holomorphic degree two. Consequently

$$\begin{aligned} \omega(Q_\kappa u, v) + (-1)^{p-1} \omega(u, Q_\kappa v) \\ = (-1)^p \int_\Sigma^{\text{res}} \iota^*(\alpha \wedge \beta' + \beta \wedge \alpha'). \end{aligned}$$

Within this support domain the expression vanishes if one argument is supported away from the boundary, or if both satisfy L .

On the boundary, the two Dolbeault summands have equal rank and pair nondegenerately with one another. The second summand is isotropic, so it is Lagrangian. It is preserved by Q_κ because $\partial\beta = \circ$. Its differential is the Dolbeault operator on holomorphic one-forms, an elliptic complex on Σ . The same decomposition tensored with normal forms makes the theory topological in the normal direction. When shifts precede this tensor product, the identification of a normal one-form carries the usual suspension sign; equivalently one may use the exterior-form formula for Q_κ displayed above. Thus the boundary condition is a Lagrangian elliptic subcomplex, and the bulk complex is topological in its normal direction. $\square$

Fix the Hamiltonian convention $\iota_{Q_\kappa} \omega = -\delta S$. The full graded smooth field complex has the local quadratic density

$$S_{\text{loc}}(\Phi) = \frac{1}{2} \omega_{\text{loc}}(\Phi, Q_\kappa \Phi).$$

For $\Phi \in E_{\kappa, c}$ its integral is the well-defined functional $S(\Phi) = \int_M^{\text{res}} S_{\text{loc}}(\Phi) = \frac{1}{2} \omega(\Phi, Q_\kappa \Phi)$. Write $E_{\kappa, L, c} = \{\Phi \in E_{\kappa, c} : \iota^* \alpha = \circ\}$. With the chiral boundary condition imposed, this functional and its

Hamiltonian identity are taken on that space. For unrestricted fields, the action is interpreted as a local density, whose first variation is tested against compactly supported variations.

On ghost-number-zero components the density is

$$\frac{1}{2}(a \wedge d'b + b \wedge d'a + \kappa a \wedge \partial a) = \mathcal{L}_\kappa(a, b) - \frac{1}{2}d'(a \wedge b).$$

For compactly supported fields satisfying $i^*a = 0$, Stokes' theorem kills the total derivative, so its integral is the action stated at the start of the chapter. The local Euler–Lagrange equations, valid on the unrestricted smooth field complex, are

$$d'a = 0, \quad d'b + \kappa \partial a = 0.$$

Their local gauge symmetry, with smooth gauge parameter c vanishing on the boundary, is

$$\delta_c a = d'c, \quad \delta_c b = \kappa \partial c.$$

A compactly supported c also preserves the compactly supported field space. The second equation is invariant because $d'\partial = -\partial d'$.

For any smooth configuration satisfying L and any compactly supported variation $(\delta a, \delta b)$ satisfying the same boundary condition, the first variation of the local action integrates to

$$\begin{aligned} \int_M^{\text{res}} \delta \mathcal{L}_\kappa &= \int_M^{\text{res}} (\delta b \wedge d'a + \delta a \wedge (d'b + \kappa \partial a)) \\ &\quad - \int_\Sigma^{\text{res}} b \wedge \delta a. \end{aligned}$$

All integrands have compact support because each contains a variation or its derivative. Integration by parts is therefore legitimate, and the boundary term vanishes. This proves the variational Hamiltonian identity with its support domain. The configuration $a = t d\bar{z}$, $b = dz$ illustrates the remaining distinction: it satisfies the boundary condition but has a nonintegrable constant action density. Its local variational equations are defined even though it has no integrated action value.

The terms involving ghosts and antifields are fixed by S_{loc} on the unrestricted graded fields. On $E_{\kappa, L, c}$ the integrated functional satisfies the classical master equation: the Hamiltonian vector field of $\{S, S\}$ is $2Q_\kappa^2 = 0$, and this quadratic functional has no constant term. Equivalently, the same statement is an identity of local variational functionals when evaluated against compactly supported variations.

1.3 THE NONZERO-LEVEL CHANGE OF FIELDS AND ITS SPECIAL FIBER

For $\kappa \neq 0$, put

$$F_\kappa(\alpha, \beta) = \alpha + \kappa^{-1}\beta.$$

The target is the full de Rham field complex $\Omega^\bullet(\mathcal{M})[1]$ with differential $d = d' + \partial$ and local pairing density $\omega_{\text{CS}, \kappa, \text{loc}}(u, v) = (-1)^{p+1} \kappa(u \wedge v)_{[3]}$. Its integrated pairing is $\omega_{\text{CS}, \kappa} = \int_M^{\text{res}} \omega_{\text{CS}, \kappa, \text{loc}}$, with at least one compactly supported argument.

Proposition 1.2 (Comparison with abelian Chern–Simons theory). *For every nonzero complex κ , F_κ is an isomorphism of linear BV theories with chiral boundary condition. It identifies the local BV Hamiltonian densities and the integrated BF and level- κ abelian Chern–Simons actions on compactly supported fields satisfying that boundary condition. This field isomorphism has no specialization at $\kappa = 0$.*

Proof. The inverse separates the holomorphic degrees and multiplies the holomorphic-one-form summand by κ . Furthermore,

$$dF_\kappa(\alpha, \beta) = d'\alpha + \partial\alpha + \kappa^{-1}d'\beta = F_\kappa Q_\kappa(\alpha, \beta).$$

The product of two α summands has no three-form component. The product of two β summands has holomorphic degree two and vanishes. The two cross terms consequently give $F_\kappa^* \omega_{\text{CS}, \kappa, \text{loc}} = \omega_{\text{loc}}$ pointwise. Both F_κ and its inverse preserve compact support, so the same identity holds for the integrated pairings on their stated domains. The local Hamiltonian densities, the integrated actions on compactly supported fields, and the boundary condition are preserved. The inverse scalar κ^{-1} in F_κ excludes the zero fiber. $\square$

At zero level the field pairing remains the same nondegenerate cross pairing. In particular, b remains a field, rather than becoming κ times a fixed field. Its gauge transformation vanishes, while the gauge transformation of a remains $d'c$.

This difference can be measured before quantization. On a product of a complex disk D and an interior real interval, the local cohomology of the field complex is computed by

$$O(D) \xrightarrow{\kappa\partial} \Omega_{\text{hol}}^1(D),$$

placed in degrees $-1, 0$. The Dolbeault and real Poincaré homotopies remove the antiholomorphic and real directions. At nonzero κ , holomorphic integration on D makes the arrow surjective and its kernel consists of constants. At zero it is the zero map. Thus

	$H^{-1}(E_\kappa)$	$H^0(E_\kappa)$
$\kappa \neq 0$	$\mathbb{C}$	0
$\kappa = 0$	$O(D)$	$\Omega_{\text{hol}}^1(D)$.

These are cohomology groups of the field complex. The negative-degree summand records gauge data. The zero fiber therefore has more local field cohomology than the nonzero Chern–Simons fibers.

1.4 A GREEN OPERATOR AT ZERO LEVEL

The zero-level kinetic complex has a concrete propagator operator in the interior. Give $\mathbb{C} \times \mathbb{R}$ its flat Euclidean metric and put

$$Q^{\text{GF}} = -4\partial_{\bar{z}}\iota_{d\bar{z}} - \partial_t\iota_{dt}.$$

The contraction symbols satisfy $\iota_{d\bar{z}}(d\bar{z}) = \iota_{dt}(dt) = 1$ and kill the other basis one-forms. Then

$$Q_0 Q^{\text{GF}} + Q^{\text{GF}} Q_0 = -4\partial_z\partial_{\bar{z}} - \partial_t^2 = -\partial_x^2 - \partial_y^2 - \partial_t^2 =: \Delta.$$

The scalar Green kernel is $G(X, Y) = (4\pi|X - Y|)^{-1}$, with $\Delta_X G = \delta_Y$. Consequently $H_0 = Q^{\text{GF}} G$ satisfies

$$Q_0 H_0 + H_0 Q_0 = 1$$

on compactly supported test forms, with output interpreted as smooth forms with the corresponding decay. Its kernel is

$$H_0(X, Y) = \frac{\overline{z - w}}{2\pi|X - Y|^3} \iota_{d\bar{z}} + \frac{t - s}{4\pi|X - Y|^3} \iota_{dt}, \quad X = (z, t), \quad Y = (w, s).$$

The distribution identity follows from the outward flux of $-\nabla(4\pi r)^{-1}$ through a sphere, which is one. Differentiating G gives the two displayed coefficients. No inverse power of κ occurs. Composing with the

inverse cross pairing gives the BV contraction kernel, including the constant $2\pi i$ from the normalization of ω .

This interior operator is not a claim about a boundary Green problem. The boundary comparison below uses compactly supported normal homotopies which preserve the chiral boundary condition. Those homotopies require neither a choice of boundary Green kernel nor a limit of κ^{-1} .

1.5 THE BOUNDARY COMPLEX AND ITS QUANTUM COCYCLE

Write $\mathbf{j}(U) = \Omega_c^{\circ, \bullet}(U)$ for an open $U \subseteq \Sigma$. A generator $u \in \Omega_c^{\circ, q}(U)$ has degree $q - 1$ in the boundary observable complex. The boundary complement is the antiholomorphic summand, while the component of the boundary differential from this complement to L is $\kappa \partial$. Let $\hbar$ be a central polynomial parameter of cohomological degree zero. The boundary observables are

$$\mathcal{B}_\kappa(U) = \left(\text{Sym}(\mathbf{j}(U)[1])[\hbar], \bar{\partial} + \hbar \Delta_{\mu_\kappa} \right), \quad \mu_\kappa(u, v) = -\kappa \int_U^{\text{res}} u \wedge \partial v.$$

The pairing is zero unless the antiholomorphic degrees sum to one. As a contraction of observable generators, it has degree one. The operator Δ_{μ_κ} vanishes on constants and linear terms; on a monomial it sums contractions of unordered pairs, with the Koszul sign for bringing that pair together. This specifies all its values. The differential on a shifted generator is the displayed $\bar{\partial}$, transported without an additional minus sign.

The completion in each symmetric power has a concrete meaning. For the r th tensor power, take compactly supported smooth sections on U^r of the external tensor product of the field bundles; then pass to coinvariants for the graded action of the symmetric group. On every fixed compact support use the seminorms of all derivatives, and take the direct limit over compact supports. This is the smooth kernel, or completed bornological tensor, convention. The symmetric algebra is the direct sum of these powers. Thus polynomial degree is finite; there is no completion in the number of fields.

Lemma 1.3 (Boundary quantum differential). *The square of $\bar{\partial} + \hbar \Delta_{\mu_\kappa}$ vanishes for every κ . At $\kappa = 0$, the quantum boundary complex is exactly the commutative Dolbeault algebra $\text{Sym}(\mathbf{j}(U)[1])[\hbar]$.*

Proof. For u of antiholomorphic degree q , the compatibility equation for one contraction is

$$\mu_\kappa(\bar{\partial} u, v) + (-1)^{q-1} \mu_\kappa(u, \bar{\partial} v) = -\kappa \int_U^{\text{res}} \bar{\partial}(u \wedge \partial v) = 0.$$

Compact support removes the boundary integral. It follows that $\bar{\partial} \Delta_{\mu_\kappa} + \Delta_{\mu_\kappa} \bar{\partial} = 0$ on every symmetric power. In the square of Δ_{μ_κ} , overlapping pairs give zero because a contracted factor has disappeared. Disjoint pairs occur in both orders with opposite signs, since a contraction has odd degree. Thus $\Delta_{\mu_\kappa}^2 = 0$. Finally, $\mu_0 = 0$ by its formula. $\square$

The coefficient κ comes from the differential $\kappa \partial$, not from a degeneration of the field pairing. The next calculation derives this cocycle directly from bulk fields.

1.6 A COMPACT COLLAR MAP FOR QUANTUM OBSERVABLES

Fix a smooth compactly supported function ρ on $\mathbb{R}_{\geq 0}$ with integral one, and put $R(t) = \int_0^t \rho(s) ds$. Thus $R = 1$ for sufficiently large t . Let $E_{\kappa, L, c}(U \times \mathbb{R}_{\geq 0})$ denote compactly supported fields satisfying $\iota^* \alpha = 0$. For $u \in \Omega_c^{\circ, q}(U)$ define

$$I_\kappa(u) = u \wedge \rho(t) dt + (-1)^q (R(t) - 1) \kappa \partial u.$$

This is a degree-zero map from the unshifted complex $(\mathfrak{j}(U), \bar{\partial})$ to $(E_{\kappa, L, c}(U \times \mathbb{R}_{\geq 0}), Q_{\kappa})$. The first term is in the α summand and the second is in the β summand. Both have compact support. The first has zero boundary pullback, and the second lies in L .

The bulk observable generators are sx , for $x \in E_{\kappa, L, c}$, with $|sx| = |x| - 1$. Define their contraction by

$$B(sx, sy) = (-1)^{|x|} \omega(x, y).$$

For complementary form degrees this is the normalized integral of the two cross terms. Let Δ_B sum pair contractions as above. The bulk–boundary quantum complex is

$$\mathcal{O}_{\kappa}(U \times \mathbb{R}_{\geq 0}) = (\text{Sym}(E_{\kappa, L, c}[1])[\hbar], Q_{\kappa} + \hbar \Delta_B).$$

The shifted unary operator is $Q_{\kappa}(sx) = sQ_{\kappa}x$. The pairing identity in Proposition 1.1 and the same disjoint-pair cancellation show that this differential squares to zero.

Theorem 1.4 (The quantum boundary map at every level). *The maps I_{κ} induce quasi-isomorphisms*

$$\text{Sym}(I_{\kappa}) : \mathcal{B}_{\kappa}(U) \longrightarrow \mathcal{O}_{\kappa}(U \times \mathbb{R}_{\geq 0})$$

for every open $U \subseteq \Sigma$ and every complex κ , including zero. They commute with extension by zero and with factorization products on disjoint opens.

Proof. First check the unary differential. Differentiating the first term of $I_{\kappa}(u)$ gives $\bar{\partial}u \wedge \rho dt + \kappa \partial u \wedge \rho dt$. Differentiating the second gives

$$-(-1)^q (R - 1) \kappa \partial \bar{\partial} u - \kappa \partial u \wedge \rho dt.$$

The last terms cancel, leaving $I_{\kappa}(\bar{\partial}u)$.

Projection to the α summand followed by integration along the normal one-form gives a cochain map P , with $PI_{\kappa} = 1$. To prove that P is a quasi-isomorphism, consider the exact sequence whose subcomplex is the β summand and whose quotient is the α summand. The former has absolute normal forms; the latter has relative normal forms, whose functions vanish at $t = 0$. On the normal complexes the homotopies are

$$\begin{aligned} h_{\text{abs}}(g dt) &= - \int_t^{\infty} g(s) ds, \\ h_{\text{rel}}(g dt) &= \int_0^t g(s) ds - R(t) \int_0^{\infty} g(s) ds. \end{aligned}$$

Both vanish on functions. The absolute homotopy satisfies $dh + hd = 1$. The relative one satisfies $dh + hd = 1 - ip$, where $p(g dt) = \int g$ and $i(1) = \rho dt$ in degree one. Their values are compactly supported, and the relative value vanishes at $t = 0$. Direct differentiation proves all these identities. For a coefficient of tangential form degree r , multiply the normal homotopy by $(-1)^r$; this cancels the tangential cross terms. These formulas also apply to smooth sections depending on z . The β subcomplex is therefore acyclic, and the relative α quotient retracts onto $\mathfrak{j}(U)$ after its displayed shift. The long exact sequence proves that P , hence I_{κ} , is a quasi-isomorphism.

There is also a continuous contraction for the full split complex. Let H be the two normal homotopies just displayed, and put $\partial_{\kappa} = Q_{\kappa} - Q_0$. The correction goes from the α summand to the β summand, so $\partial_{\kappa} H \partial_{\kappa} = 0$ and $P \partial_{\kappa} = 0$. Consequently

$$H_{\kappa} = H - H \partial_{\kappa} H, \quad I_{\kappa} = I_0 - H \partial_{\kappa} I_0, \quad Q_{\kappa} H_{\kappa} + H_{\kappa} Q_{\kappa} = 1 - I_{\kappa} P.$$

To check the last identity, substitute $Q_0 H + H Q_0 = 1 - I_0 P$ and $Q_0 \partial_{\kappa} = -\partial_{\kappa} Q_0$. The linear correction terms cancel except for $H \partial_{\kappa} I_0 P$; the quadratic correction terms vanish. The formula for I_{κ} is the displayed collar map, since the absolute homotopy of ρdt is $R - 1$. This construction uses no division by κ .

It remains to check the quantum contraction, rather than only the unary differential. For antiholomorphic degrees $q + r = 1$, expand the two cross terms in $I_\kappa(u) \wedge I_\kappa(v)$. Moving dt to the right, and integrating by parts on U , gives

$$B(sI_\kappa(u), sI_\kappa(v)) = -2\kappa \left(\int_0^\infty \rho(t)(1 - R(t))dt \right) \int_U^{\text{res}} u \wedge \partial v.$$

The minus sign is the fiber-integration sign fixed above; B itself integrates the two cross terms with positive sign. For the second term use $\int_U \partial u \wedge v = (-1)^{q+1} \int_U u \wedge \partial v$. The normal integral is

$$\int_0^\infty \rho(1 - R)dt = \left[R - \frac{R^2}{2} \right]_0^\infty = \frac{1}{2}.$$

Consequently $B(I_\kappa(u), I_\kappa(v)) = \mu_\kappa(u, v)$, with suspensions understood. This proves compatibility on quadratic monomials. The pair-contraction definitions prove it on every symmetric power.

At the classical level, the preceding continuous linear homotopies extend to r tensor factors by the explicit map

$$\mathcal{H}_r = \sum_{j=1}^r (I_\kappa P)^{\otimes(j-1)} \otimes H_\kappa \otimes 1^{\otimes(r-j)}.$$

Use the Koszul sign when the degree-minus-one map H_κ passes the preceding factors. In $d\mathcal{H}_r + \mathcal{H}_r d$, the terms $1 - I_\kappa P$ telescope, giving $1 - (I_\kappa P)^{\otimes r}$. Averaging over the finite symmetric group gives the result on symmetric powers; characteristic zero permits this averaging. The integrations preserve compact supports in the product variables and extend to the completed bornological tensors. Thus $\text{Sym}(I_\kappa)$ is a classical quasi-isomorphism.

Finally filter both polynomial quantum complexes by the sum of the power of $\hbar$ and the number of fields. The unary differential preserves that filtration degree. A quantum contraction reduces it by one. The associated graded map is the classical quasi-isomorphism tensored with $\mathbb{C}[\hbar]$. The filtration starts in degree zero, and every element has finite filtration degree. Induction on the finite filtration, followed by the exact direct limit, proves the quantum quasi-isomorphism. All formulas use the same ρ and integrate only the normal variable. They therefore commute with extension by zero and with products on disjoint boundary opens. $\square$

The observables also satisfy the gluing property needed for a factorization algebra. The relevant covers $\{U_i\}$ of U are those for which every finite subset of U lies in some U_i . For every r , the sets U_i^r then cover U^r . At a fixed field degree the augmented Čech complex of compactly supported smooth kernels for this cover is exact: a locally finite partition of unity gives its contraction, and only finitely many terms meet a given compact support. Multiplication by the partition functions preserves the linear boundary condition for bulk fields. For fixed r the form degrees are bounded, so the finite filtration by form degree upgrades this Čech exactness to descent for the unary differential. Passing to symmetric coinvariants preserves exactness because averaging over the finite symmetric group is available. The polynomial filtration used above then proves quantum descent. Here factorization algebra means precisely these gluing complexes together with the compatible products for disjoint opens. Thus the theorem is an equivalence of quantum factorization algebras on the boundary. At $\kappa = 0$ its image contains only α fields, so its pair contraction vanishes directly.

1.7 A COMPACT CALCULATION OF THE CURRENT LEVEL

The quantum differential determines how a product changes when its insertions are moved. A compact two-current calculation fixes the sign before any mode presentation is chosen. For a point a in a disk, choose a cutoff χ_a supported in a small disk about a and equal to one near its center. The form

$$u_a = -\frac{\bar{\partial}\chi_a}{z-a}$$

is smooth and compactly supported. Stokes' theorem on an annulus about a gives, for every holomorphic function F on the disk,

$$\int^{\text{res}} u_a \wedge F(z) dz = F(a).$$

Thus u_a is a smooth representative of evaluation at a .

Choose distinct points a, b , disjoint small disks about them, and cutoffs χ_a, χ_b as above. Choose a common radial cutoff $\chi(z) = \psi(|z|^2)$, equal to one on both small disks and supported in a larger disk centered at zero. Put

$$e_a = -\frac{\bar{\partial}\chi}{z-a}, \quad e_b = -\frac{\bar{\partial}\chi}{z-b}, \quad f = \frac{\chi - \chi_a}{z-a}, \quad g = \frac{\chi - \chi_b}{z-b}.$$

The functions f, g are smooth and compactly supported: the apparent poles are removable. Direct differentiation gives $\bar{\partial}f = u_a - e_a$ and $\bar{\partial}g = u_b - e_b$. On the support of u_b , $f(z) = (z-a)^{-1}$. Integration by parts and the evaluation formula give

$$\mu_\kappa(f, u_b) = -\kappa \int^{\text{res}} u_b \wedge \partial f = -\kappa f'(b) = \frac{\kappa}{(a-b)^2}.$$

The other replacement has zero contraction. On the common cutoff annulus, $g = \chi/(z-b)$ and

$$e_a \wedge \partial g = -\frac{\bar{\partial}\chi \wedge \partial\chi}{(z-a)(z-b)} + \frac{\chi \bar{\partial}\chi \wedge dz}{(z-a)(z-b)^2}.$$

Relative to $d\bar{z} \wedge dz$, its coefficient is

$$-\frac{(\psi')^2 |z|^2}{(z-a)(z-b)} + \frac{\psi \psi' z}{(z-a)(z-b)^2}.$$

Both Laurent expansions have strictly negative angular Fourier powers, beginning with z^{-2} . Their angular integrals vanish, so $\mu_\kappa(e_a, g) = 0$.

The function generators have degree -1 and the form generators have degree zero. With the specified positive unary differential, $D = \bar{\partial} + \hbar \Delta_{\mu_\kappa}$, the Leibniz signs and these contractions give the compact chain relation

$$D(f u_b + e_a g) = u_a u_b - e_a e_b + \frac{\hbar \kappa}{(a-b)^2}.$$

Consequently the actual separated product satisfies

$$[u_a u_b] = [e_a e_b] - \frac{\hbar \kappa}{(a-b)^2} \mathbf{1}.$$

The radial term has the regular expansion $e_a = \sum_{n \geq 0} a^n u_n$, where $u_n = -\bar{\partial}\chi/z^{n+1}$, and similarly for e_b . These series converge in every smooth seminorm on the outer annulus. They cannot contribute a diagonal pole. The current determined by the fixed state $q_1 = [u_0]$ therefore has parameter $-\kappa$:

$$J(a)J(b) \sim -\frac{\hbar \kappa}{(a-b)^2}.$$

The remaining problem is to obtain every field and regular product from the supported complex, keeping this normalization.

A direct annular evaluation of the local cocycle has a different role. For holomorphic Laurent functions F, G and a cutoff equal to one at the inner boundary and zero at the outer boundary,

$$\mu_\kappa(F, \bar{\partial}\chi G) = -\kappa \int^{\text{res}} \bar{\partial}\chi \wedge G dF = \kappa \text{Res } G dF.$$

Only one factor needs compact support in this integral. The minus sign in the separated product comes from moving the quantum contraction to the other side of an exact chain relation. Thus the annular scalar evaluation by itself does not identify the extracted mode action. Multiplying all currents by i changes their parameter, but also changes the prescribed state and its jet representatives.

I.8 CAUCHY CONTRACTION ON SMOOTH OBSERVABLES

The quantum differential can be removed by an explicit linear map, provided its effect on products is retained. For $u \in \Omega_c^{\text{ol}}(U)$, extend u by zero to $\mathbb{C}$ and define its Cauchy transform by

$$(\mathcal{C}u)(z) = \frac{1}{2\pi i} \int_{\mathbb{C}} \frac{u(w) \wedge dw}{z - w}.$$

The kernel $1/(z - w)$ is locally integrable. Stokes' theorem on a small circle gives $\partial_{\bar{z}}(1/(\pi z)) = \delta_0$; the normalization $d\bar{w} \wedge dw = 2i dA_w$ then gives

$$\bar{\partial}\mathcal{C}u = u, \quad \mathcal{C}\bar{\partial}f = f \quad (f \in C_c^\infty(\mathbb{C})).$$

For the second identity, the first implies that the difference is holomorphic. Integration by parts shows that all moments of $\bar{\partial}f$ vanish, so expansion of the Cauchy kernel outside a disk containing the support makes $\mathcal{C}\bar{\partial}f$ compactly supported. The difference is therefore a compactly supported holomorphic function and is zero. The output $\mathcal{C}u$ is generally not compactly supported; it is used below only inside a scalar integral.

On two degree-zero observable generators, define

$$P_\kappa(u, v) = -\kappa \int^{\text{res}} (\mathcal{C}u) \partial v.$$

Define P_κ to be zero on all other generator-degree pairs. The derivative of v has compact support, so the displayed integral exists. If $u = a(w)d\bar{w}$ and $v = b(z)d\bar{z}$, it is

$$P_\kappa(u, v) = \frac{\kappa}{\pi^2} \int_{\mathbb{C}^2} \frac{a(w) \partial_z b(z)}{z - w} dA_z dA_w.$$

This formula also defines a pair contraction on an arbitrary compactly supported smooth kernel, by differentiating in the paired z variable and integrating both paired variables. The output is a smooth kernel on the remaining variables, supported in the compact projection of the input support. For each fixed support K and multi-index α in the remaining variables,

$$\|\partial^\alpha \Delta_P F\|_\infty \leq C_{K, \alpha} \|F\|_{C^{|\alpha|+1}}.$$

Indeed, the integral of $|z - w|^{-1}$ on a fixed compact pair of planar variables is finite. Derivatives in the remaining variables pass under that integral. Thus the contraction is continuous in the smooth-kernel convention already specified; no singular observable or extra completion is introduced.

The pairing P_κ is symmetric. After interchanging z, w , its antisymmetric part is proportional to

$$\int_{\mathbb{C}^2} \frac{(\partial_z + \partial_w)(a(w)b(z))}{z - w} dA_z dA_w.$$

In difference and translation coordinates, the derivative in the numerator differentiates only the translation coordinate. Its integral is zero by compact support, without differentiating the singular difference kernel.

Let $K = \Delta_{P_\kappa}$ sum unordered pair contractions, with the shifted Koszul signs, and write $d = \bar{\partial}$. The operator K has degree zero and lowers field number by two. Its commutators are

$$[d, K] = -\Delta_{\mu_\kappa}, \quad [K, \Delta_{\mu_\kappa}] = 0.$$

To verify the first equation, it suffices first to use a function and a form generator:

$$[d, K](fu) = -P_\kappa(\bar{\partial}f, u) = -\mu_\kappa(f, u).$$

The other quadratic generator degrees give zero. Both sides are constant pair-contraction operators, so the identity holds on all monomials. For the second equation, overlapping contractions vanish because the shared factor disappears; disjoint contractions commute because K is even. The same identities hold on smooth kernels by continuity. One may verify density of decomposable kernels on product boxes by Fourier expansion after inserting compact cutoffs: rapid decay of Fourier coefficients gives convergence in every derivative seminorm. A finite partition of unity on the input support gives the general case.

Proposition 1.5 (Finite quantum conjugation). *For every open $U \subseteq \mathbb{C}$ and every $\kappa \in \mathbb{C}$, the map*

$$N_U = e^{-\hbar K} : (\mathcal{B}_\kappa(U), D) \longrightarrow (\text{Sym}(\Omega_c^{\circ, \bullet}(U)[1])[\hbar], d)$$

is a chain isomorphism, with inverse $e^{\hbar K}$. It commutes with extension by zero, translations and rotations.

Proof. The two commutators give

$$e^{\hbar K} d e^{-\hbar K} = d + \hbar[K, d] = d + \hbar\Delta_{\mu_\kappa} = D.$$

On field number r , $K^j = 0$ for $2j > r$, so all exponentials terminate. They preserve the polynomial complexes and their filtration by field number plus $\hbar$ -power. The global Cauchy formula uses the same zero extension for every open set. Its kernel and measure are translation invariant and transform compatibly under rotations. These facts prove the stated naturality. $\square$

The map N_U is not asserted to preserve multiplication at overlapping supports. Its cross contractions carry the quantum products.

1.9 THE VACUUM STATES AND THEIR SMOOTH REPRESENTATIVES

Let $D_R = \{|z| < R\}$ and choose a radial cutoff χ equal to one near zero and zero near the boundary. For $n \geq 0$ put

$$u_n = -\frac{\bar{\partial}\chi}{z^{n+1}}.$$

The support avoids zero. Stokes' theorem gives its divided-jet moments:

$$\int_{D_R}^{\text{res}} u_n \wedge F(z) dz = \frac{F^{(n)}(0)}{n!}, \quad M_j(u_n) = \delta_{jn}, \quad M_j(u) = \int_{D_R}^{\text{res}} u \wedge z^j dz.$$

A compactly supported holomorphic function vanishes. A compact $(0, 1)$ -form with every holomorphic moment zero has a compact primitive: its Cauchy transform vanishes outside a disk containing its support, by expansion in w/z . Conversely, integration by parts kills the moments of $\bar{\partial}f$ for compact f .

The required statement on completed tensors follows from an explicit estimate. If the input is supported in $|z| \leq r_o < R$, choose the cutoff annulus with inner radius $\sigma > r_o$. Then

$$|M_n(u)| \leq C r_o^n, \quad \|u_n\|_{C^d} \leq C_d (n+1)^{d+1} \sigma^{-n-1}.$$

Thus the maps

$$\Pi u = \sum_{n \geq 0} M_n(u) u_n, \quad h_D u = \mathcal{C}(u - \Pi u)$$

are well defined on this fixed support space, with convergence in all derivative seminorms and compact output. On functions set $\Pi = h_D = 0$. The moments and the Cauchy identities prove

$$dh_D + h_D d = 1 - \Pi.$$

For a compact smooth kernel on D_R^r , choose a common inner support radius and a larger cutoff annulus in each variable. The moment bounds multiply, so the resulting multiple series converge in every mixed derivative seminorm. With shifted tensor signs, the map

$$H_r = \sum_{j=1}^r \Pi^{\otimes(j-1)} \otimes h_D \otimes 1^{\otimes(r-j)}$$

satisfies $dH_r + H_r d = 1 - \Pi^{\otimes r}$ by telescoping. Each variable is treated once. No single cutoff operator on all compact supports is required. Averaging over the finite symmetric group proves the same assertion for the completed symmetric powers.

The tensor cohomology is consequently in top antiholomorphic degree r , hence in shifted cohomological degree zero, and is determined by its multivariable moments. This computes the odd generators by their differential; it does not discard them.

Rotations act on the representatives by

$$R_\theta^* u_n = e^{-i(n+1)\theta} u_n.$$

Define the state weight as minus this pullback rotation weight. The radial homotopies commute with rotations, including simultaneous rotation of all variables. At state weight N , a surviving tensor must satisfy

$$\sum_{i=1}^r (n_i + 1) = N.$$

There are finitely many such tuples, and $r \leq N$. Thus the argument applies to a fixed diagonal weight without assuming a finite Fourier expansion in the individual variables.

Rotation invariance also gives $P_\kappa(u_n, u_m) = 0$ for all n, m : the two inputs have strictly positive total state weight, while a scalar has weight zero. The map N_{D_R} therefore fixes every monomial in these radial representatives. Combining its chain isomorphism with the moment calculation gives

$$V_\kappa = \mathbb{C}[q_1, q_2, \dots][\hbar], \quad q_{n+1} \mapsto [u_n], \quad |q_m| = 0, \quad \text{wt}(q_m) = m.$$

More precisely, V_κ is the direct sum of finite-weight cohomology spaces of $\mathcal{B}_\kappa(D_R)$. Each weight space is finite free over $\mathbb{C}[\hbar]$, and there is no odd cohomology or $\hbar$ -torsion. The full disk cohomology also contains infinite moment sequences and is not identified with this direct sum.

Polynomial multiplication specifies the normal state basis $\prod q_{m_i} \mapsto [\prod u_{m_i-1}]$. It does not assert that multiplication at overlapping supports descends to quantum cohomology. Changing the radial cutoff or enlarging the centered disk preserves this basis: applying N reduces the comparison to the equality of its multivariable moments.

1.10 TRANSPORT OF EVERY SEPARATED PRODUCT

For disjoint opens $U_1, \dots, U_s \subseteq U$, the original factorization product is extension by zero followed by multiplication m . Its mixed μ_κ contractions vanish because the supports are disjoint. Under Cauchy conjugation its exact formula is

$$N_U m(N_{U_1}^{-1} \otimes \dots \otimes N_{U_s}^{-1}) = m \exp \left(-\hbar \sum_{i < j} \Delta_{P_\kappa}^{ij} \right),$$

where $\Delta_{P_\kappa}^{ij}$ contracts a pair with one factor in input i and one in input j . To prove the formula, decompose K on a product into contractions internal to each input and contractions between inputs. All these even operators commute. The internal exponentials cancel $N_{U_i}^{-1}$, leaving exactly the displayed cross contractions.

The exponential terminates on every input. Its terms are matchings of the individual field factors, with edges joining different input groups. A matching with j edges appears $j!$ times, one for each ordering of its disjoint contractions; the exponential coefficient $1/j!$ cancels that multiplicity. The proof extends to all compact smooth kernels by the continuity estimate above. It commutes with nesting, either by conjugating the original factorization maps or by observing that each nested expression contains each cross pair exactly once.

For a small radial cutoff at a , write

$$u_{a,n} = -\frac{\bar{\partial} \chi_a}{(z-a)^{n+1}}, \quad n \geq 0.$$

Its moments are divided derivatives at a . Outside the supporting disk,

$$(\mathcal{C} u_{a,n})(z) = \frac{1}{(z-a)^{n+1}}.$$

This follows by applying the moment formula to the holomorphic function $w \mapsto (z-w)^{-1}$. For disjoint supporting disks at a, b , integration by parts therefore gives

$$\begin{aligned} P_\kappa(u_{a,n}, u_{b,m}) &= -\frac{\kappa}{m!} \partial_z^{m+1} (z-a)^{-n-1} \Big|_{z=b} \\ &= \frac{\kappa}{n!m!} \partial_a^n \partial_b^m (a-b)^{-2}. \end{aligned}$$

Put $\lambda = -\hbar\kappa$. The actual cross contraction of divided jets of orders n, m is

$$C_{n,m}(a, b) = \lambda \frac{(-1)^n (n+m+1)!}{n!m!} (a-b)^{-n-m-2}.$$

In particular, $C_{0,0}(a, b) = \lambda(a-b)^{-2}$, in agreement with the compact two-current calculation.

Holomorphic motion of the remaining state is equally explicit. A common outer radial cutoff gives

$$-\frac{\bar{\partial} \chi}{(z-a)^{n+1}} = \sum_{k \geq 0} \binom{n+k}{k} a^k u_{n+k}.$$

The difference from $u_{a,n}$ has zero moments and a compact primitive. For products of co-centered representatives, all internal P terms vanish by rotation about that center. The same comparison therefore holds for each polynomial state. It gives the translation derivation

$$Tq_m = mq_{m+1}, \quad T1 = 0.$$

The moving state corresponding to q_m is $e^{aT}q_m$.

For arbitrary polynomial states inserted at distinct points, label each of their jet factors by α , with jet order r_α and insertion group $i(\alpha)$. The actual product, expressed by N and the moment coordinates, is

$$\sum_M \left(\prod_{\{\alpha, \beta\} \in M} C_{r_\alpha, r_\beta}(a_{i(\alpha)}, a_{i(\beta)}) \right) \left(\prod_{\gamma \text{ unpaired}} e^{a_{i(\gamma)}T} q_{r_\gamma+1} \right).$$

Here M ranges over matchings joining different insertion groups. There are finitely many matchings, including the empty matching. The empty matching retains the regular terms. For fixed numerical insertion points, the unpaired series have smooth representatives on a common outer annulus and converge in all derivative seminorms, by the geometric moment bounds. Thus the formula is an equality in full disk cohomology. Taking weight coefficients and radial Laurent coefficients of this equality gives polynomial states. This is the extraction from the smooth carrier.

I.II COMPACT HOMOTOPIES FOR NORMAL ORDERING

The Wick expansion gives a cohomology class of supported observables. A chain comparison also needs a compact primitive for its difference from the original product. At three insertions, that primitive itself has a quantum correction. A finite Koszul complex inside the boundary observables constructs these primitives in every polynomial degree.

Fix a disk $U = D_R \subset \mathbb{C}$ and finitely many distinct points in U . Choose disjoint centered disks about these points, with radial cutoffs χ_a equal to one near their centers. Choose a radial cutoff χ about zero which equals one on a centered closed disk containing all the smaller closed disks. All cutoffs have compact support in their stated disks. Choose a finite list of distinct pairs (a_i, n_i) , where $n_i \geq 0$ is a divided holomorphic jet order, and index it by $1 \leq i \leq N$. Several pairs may have the same point. Set

$$u_i = -\frac{\bar{\partial}\chi_{a_i}}{(z - a_i)^{n_i+1}}, \quad e_i = -\frac{\bar{\partial}\chi}{(z - a_i)^{n_i+1}}, \quad f_i = \frac{\chi - \chi_{a_i}}{(z - a_i)^{n_i+1}}.$$

The apparent poles are removable. All three expressions are smooth and compactly supported in U , and $\bar{\partial}f_i = u_i - e_i$. In the boundary observable complex, u_i, e_i have degree zero, while f_i has degree -1 .

Keep the bulk parameter κ and the positive differential $D = \bar{\partial} + \hbar\Delta_{\mu_\kappa}$ fixed. Define the symmetric constants

$$p_{ij} = \begin{cases} \frac{\kappa}{n_i!n_j!} \partial_{a_i}^{n_i} \partial_{a_j}^{n_j} (a_i - a_j)^{-2}, & a_i \neq a_j, \\ 0, & a_i = a_j. \end{cases}$$

In particular, $p_{ii} = 0$. The extracted current has level $-\kappa$, so its Wick contraction is $-\hbar p_{ij}$.

Lemma 1.6 (Contractions of compact representatives). *The Cauchy pairing and the quantum cocycle satisfy*

$$\begin{aligned} P_\kappa(u_i, u_j) &= p_{ij}, & P_\kappa(e_i, u_j) &= P_\kappa(e_i, e_j) = 0, \\ \mu_\kappa(f_i, u_j) &= p_{ij}, & \mu_\kappa(f_i, e_j) &= 0. \end{aligned}$$

The pairing P_κ is zero on pairs involving a function generator. The cocycle μ_κ is zero on two function generators or two form generators.

Proof. For distinct centers, the separated Cauchy calculation gives the displayed derivative formula for $P_\kappa(u_i, u_j)$. For a common center, simultaneous rotation multiplies this pairing by weight $-(n_i + n_j + 2)$. The scalar output has weight zero. Rotation invariance therefore makes the pairing zero.

The Cauchy transform of an outer representative is

$$\mathcal{C}e_i = \frac{1 - \chi}{(z - a_i)^{n_i+1}}.$$

Both sides have Dolbeault derivative e_i and tend to zero at infinity. Their difference is an entire function tending to zero, so it vanishes. The displayed expression is zero near the support of every u_j . Hence $P_\kappa(e_i, u_j) = 0$. On the common annulus, each e_i has the convergent expansion

$$e_i = \sum_{r \geq 0} \binom{n_i + r}{r} a_i^r \left(-\frac{\bar{\partial}\chi}{z^{n_i+r+1}} \right).$$

Each term has strictly positive state weight. Pairing two such terms gives zero by rotation invariance. Smooth convergence and continuity of P_κ give $P_\kappa(e_i, e_j) = 0$.

The Cauchy identity on compact functions gives

$$\mu_\kappa(f_i, v) = P_\kappa(\bar{\partial}f_i, v) = P_\kappa(u_i - e_i, v)$$

for each form generator v . This proves both cocycle formulas. The remaining vanishing statements follow from their specified degrees. $\square$

Let x_i, y_i, θ_i denote the observable generators represented by u_i, e_i, f_i , respectively. The symbols f_i continue to denote the actual compact functions. Thus

$$|x_i| = |y_i| = 0, \quad |\theta_i| = -1, \quad Q = \mathbb{C}[x_1, \dots, x_N, y_1, \dots, y_N] \otimes \Lambda(\theta_1, \dots, \theta_N)[\hbar].$$

The odd variables anticommute. Use left derivatives in them, so $\partial_{\theta_i}(\theta_{j_1} \cdots \theta_{j_s})$ has sign $(-1)^{r-1}$ when $i = j_r$ in increasing order. All tensor substitutions use these cohomological degrees. The map $\rho : Q \rightarrow \mathcal{B}_\kappa(U)$ substitutes the compact representatives and multiplies them as observable polynomials.

Proposition 1.7 (A quantum complex of compact currents). *The map ρ embeds Q as a subcomplex of the full smooth boundary observable complex. Its differential is*

$$D_Q = \sum_i (x_i - y_i) \partial_{\theta_i} + \hbar \sum_{i < j} p_{ij} (\partial_{x_j} \partial_{\theta_i} + \partial_{x_i} \partial_{\theta_j}). \quad (1.1)$$

This assertion concerns a complex of vector spaces over $\mathbb{C}[\hbar]$. The ordinary multiplication on Q is not a differential graded multiplication when some $p_{ij} \neq 0$.

Proof. The preceding lemma computes every possible contraction of its generators. Together with $\bar{\partial}f_i = u_i - e_i$, it gives (1.1) on every polynomial.

The u_i are linearly independent: their supports separate the centers, and the angular Fourier powers separate the orders at one center. Their supports are disjoint from the common outer annulus. The e_i are linearly independent there. Indeed, a relation among them would make a rational function $\sum_i c_i (z - a_i)^{-n_i-1}$ vanish on an open subset of the annulus. It would vanish identically, and its partial fractions give $c_i = 0$. Thus all the form generators are independent. A relation among the f_i would give a relation among $u_i - e_i$ after differentiation, so all function generators are independent as well. Symmetric and exterior powers of these finite-dimensional subspaces embed

into their smooth tensor powers. For example, continuous dual linear functionals selecting their basis vectors supply a left inverse on every tensor power, and then on its finite-group coinvariants. This proves injectivity of ρ in the stated smooth-kernel convention.

Finally, $D_Q(\theta_i x_j) - (D_Q \theta_i) x_j = \hbar p_{ij}$. This proves the asserted failure of the Leibniz identity whenever the displayed scalar is nonzero. $\square$

Write $\delta_i = x_i - y_i$, let $A = \mathbb{C}[y_1, \dots, y_N][\hbar]$ with zero differential, and let $\iota : A \rightarrow Q$ be the inclusion. Let $\pi : Q \rightarrow A$ set all δ_i, θ_i to zero. On Q define

$$d_o = \sum_i \delta_i \partial_{\theta_i}, \quad K_Q = \sum_{i < j} p_{ij} \partial_{x_i} \partial_{x_j}, \quad N_Q = e^{-\hbar K_Q}.$$

The derivatives ∂_{x_i} fix y_j and θ_j . For a monomial with total degree q in the variables δ_i, θ_i , define

$$h_o = \begin{cases} \frac{1}{q} \sum_i \theta_i \partial_{\delta_i}, & q > 0, \\ 0, & q = 0. \end{cases}$$

Extend this definition linearly over A . All exponentials below terminate on each polynomial.

Theorem 1.8 (Compact quantum contraction). *The maps*

$$\Pi = \pi N_Q : Q \longrightarrow A, \quad H = N_Q^{-1} h_o N_Q : Q \longrightarrow Q[-1]$$

and the inclusion ι satisfy

$$\Pi \iota = 1, \quad D_Q H + H D_Q = 1 - \iota \Pi, \quad H \iota = \Pi H = H^2 = 0. \quad (1.2)$$

Both ι and Π are chain maps. Consequently Q has cohomology A in degree zero and zero cohomology in every other degree. These identities hold for every κ , including zero.

Proof. The odd derivation $s_o = \sum_i \theta_i \partial_{\delta_i}$ satisfies

$$d_o s_o + s_o d_o = \sum_i \delta_i \partial_{\delta_i} + \sum_i \theta_i \partial_{\theta_i}.$$

One checks the equality on the generators. Both sides are even derivations, which proves it on every polynomial. The right side acts by q on the specified degree- q summand. Thus $d_o h_o + h_o d_o = 1 - \iota \pi$. The operator s_o preserves q and has square zero, because the odd factors anticommute while the even derivatives commute. This also gives $h_o^2 = h_o \iota = \pi h_o = 0$.

Direct commutation gives

$$[K_Q, d_o] = \sum_{i < j} p_{ij} (\partial_{x_j} \partial_{\theta_i} + \partial_{x_i} \partial_{\theta_j}), \quad [K_Q, [K_Q, d_o]] = 0.$$

Therefore $D_Q = N_Q^{-1} d_o N_Q$. Also $N_Q \iota = \iota = N_Q^{-1} \iota$, since K_Q kills polynomials in the y_i alone. Conjugating the classical contraction gives every equation in (1.2). It also proves $D_Q^2 = 0$ and the chain-map assertions. No inverse of κ or $\hbar$ occurs. $\square$

The inclusion into the full boundary complex preserves these explicit homotopies. For a polynomial $F(x)$ in the local current generators,

$$F(u) - \iota_U \left(e^{-\hbar K_Q} F(x) \Big|_{x=y} \right) = D \rho H F(x), \quad \iota_U = \rho \iota. \quad (1.3)$$

Here $F(u) = \rho F(x)$ and $D_Q F(x) = 0$. The equality is in compactly supported observables, before taking cohomology. Distinct input groups have disjoint small disks. Their actual factorization product is the ordinary polynomial on the left. Expanding the exponential on the right gives every matching between groups, with contraction $-\hbar p_{ij}$. Pairings within a group vanish by the common-center convention. Repeated factors are counted by the polynomial derivatives.

For example, with $\delta_i = x_i - y_i$ and three different indices,

$$\begin{aligned} H(x_1 x_2) &= \frac{1}{2} \theta_1(x_2 + y_2) + \frac{1}{2} \theta_2(x_1 + y_1), \\ \Pi(x_1 x_2) &= y_1 y_2 - \hbar p_{12}, \\ H(x_1 x_2 x_3) &= \sum_{i=1}^3 \theta_i \left(y_j y_k + \frac{\delta_j y_k + y_j \delta_k}{2} + \frac{\delta_j \delta_k}{3} - \frac{2\hbar p_{jk}}{3} \right), \\ \Pi(x_1 x_2 x_3) &= y_1 y_2 y_3 - \hbar(p_{12} y_3 + p_{13} y_2 + p_{23} y_1). \end{aligned}$$

In the cubic sum, $\{j, k\} = \{1, 2, 3\} \setminus \{i\}$. Its summand is symmetric in j, k . The classical Euler homotopy contributes the first three terms. Applying N_Q before it contributes $-\hbar p_{jk} \theta_i$. Applying N_Q^{-1} afterwards contributes $\hbar p_{jk} \theta_i / 3$. Their sum gives the printed correction. In particular, omitting that correction leaves the defect

$$\frac{2\hbar}{3} (p_{12} \delta_3 + p_{13} \delta_2 + p_{23} \delta_1)$$

in the cubic chain identity. For nonzero covariance, independence of the variables δ_i makes this defect nonzero. The binary identity alone cannot fix it.

Proposition 1.9 (Split inclusion in smooth boundary observables). *There is an explicit chain map $L_U : \mathcal{B}_\kappa(U) \rightarrow A$ with $L_U \iota_U = 1$. Thus the annular polynomial states inject into the cohomology of the full smooth boundary observable complex. On the subcomplex ρQ , $L_U \rho = \Pi$.*

Proof. Choose holomorphic polynomials $F_i(z)$ such that

$$\frac{F_i^{(n_j)}(a_j)}{n_j!} = \delta_{ij} \quad (1 \leq i, j \leq N).$$

Such polynomials exist by finite interpolation. To verify existence, let $m_a = 1 + \max\{n_j : a_j = a\}$ and $M = \sum_a m_a$. The map from polynomials of degree less than M to their full jets of orders less than m_a is injective: a polynomial in its kernel is divisible by $\prod_a (z - a)^{m_a}$, of degree M . The domain and target have dimension M , so the map is surjective. Prescribe the stated entries and set all other jet entries to zero.

On the classical observable complex, send a form generator v to

$$R(v) = \sum_i \left(\int_U^{\text{res}} v \wedge F_i(z) dz \right) y_i$$

and a function generator to zero. Extend by multiplication. For a compact smooth kernel, integrate in each variable and take the finite sum over its indices i . These integrals are continuous on each fixed compact support and respect the symmetric coinvariants. Integration by parts and holomorphicity give $Rd = 0$. The moments of both u_j and e_j give $R(u_j) = R(e_j) = y_j$. Thus $R\rho = \pi$ on the classical complexes.

The Cauchy contraction restricts to K_Q on ρQ , by Lemma 1.6. Consequently $N_U \rho = \rho N_Q$. Set $L_U = R N_U$, using the chain isomorphism of Proposition 1.5. Then $L_U \rho = \pi N_Q = \Pi$ and $L_U \iota_U = 1$. $\square$

For a fixed set of centers, increasing the finite jet list preserves D_Q, K_Q, h_o, Π, H on the smaller polynomial algebra. Hence the compact contraction also holds on the union of all finite jet lists. Its elements still have finite polynomial degree. The interpolation projection L_U requires a finite list and is not asserted to be compatible with this union. The union has more jet generators, but it does not include arbitrary smooth kernels or all infinite moment sequences.

The maps D_Q, N_Q, H, Π are polynomial in $\hbar$ on each input polynomial. They therefore extend to $\widehat{Q} = \varprojlim_m Q/\hbar^m Q$ and $\widehat{A} = \varprojlim_m A/\hbar^m A$, where each coefficient has finite polynomial degree. A fixed coefficient of an exponential receives finitely many terms. The same identities hold modulo every $\hbar^m$, hence in the inverse limit. This is a contraction of the specified completed subcomplex. It does not compute the cohomology of the completed full boundary complex.

The compact collar chain map sends ρHF to a primitive for the corresponding bulk–boundary product difference. Its chain identity follows by applying that map to (1.3); its separated-product identity identifies the first term with the actual collar product. Thus the normalization $-\hbar\kappa(a-b)^{-2}$ and its compact primitives pass through the same boundary comparison. Construction of operations on collision chains requires further maps compatible with those chains and their boundary strata.

1.12 FIELDS, MODES AND REGULAR PRODUCTS

Applying the separated-product formula to one current and a monomial at zero gives the operator-valued formal distribution

$$J(z) = \sum_{m \geq 1} q_m z^{m-1} + \lambda \sum_{m \geq 1} m \partial_{q_m} z^{-m-1}.$$

The first sum is the creation part $J_-(z)$, acting by multiplication; the second is the annihilation part $J_+(z)$, acting by derivations. A mode $a_{(n)}$ is the coefficient of z^{-n-1} in its field. For this current,

$$J_{-m} = q_m, \quad J_m = -\hbar\kappa m \partial_{q_m} \quad (m > 0), \quad J_0 = 0.$$

These operators were obtained from compact insertions and their quantum differential with the fixed identification $q_{n+1} = [u_n]$.

Write $J^{[r]}(z) = \partial_z^r J(z)/r!$ for its divided derivatives. For a monomial state $a = \prod_{i=1}^r q_{m_i}$, its field is

$$Y(a, z) =: \prod_{i=1}^r J^{[m_i-1]}(z) :.$$

Normal ordering here has an explicit definition. Split each divided current into creation and annihilation parts, expand the finite product, and put every multiplication operator to the left of every derivation. Both types commute within themselves, so the answer is independent of the ordering of the factors of a . Expanding those derivations against any polynomial state gives precisely the preceding matching formula with the second insertion at zero. Thus Y is the extracted state-field operation, including its regular coefficients.

For example, the complete two-jet product is

$$\begin{aligned} Y(q_m, z)q_n &= \sum_{k \geq 0} \binom{m+k-1}{k} q_{m+k} q_n z^k \\ &\quad + \lambda \frac{(-1)^{m-1} (m+n-1)!}{(m-1)!(n-1)!} z^{-m-n}. \end{aligned}$$

For a quadratic state, the regular coefficient contains a contraction with a translated unpaired current:

$$(q_1^2)_{(-1)}q_1 = q_1^3 + 2\lambda q_3 = q_1^3 - 2\hbar\kappa q_3.$$

Indeed, $Y(q_1^2, z)q_1 = J_-(z)^2 q_1 + 2\lambda z^{-2} J_-(z)$. Since $q_{1(-1)}q_1 = q_1^2$ and $q_{1(-1)}q_1^2 = q_1^3$, this also gives the regular-product associator

$$(q_{1(-1)}q_1)_{(-1)}q_1 - q_{1(-1)}(q_{1(-1)}q_1) = 2\lambda q_3.$$

Polynomial multiplication in the normal state basis and the regular vertex product are therefore different operations at nonzero level.

Every field is lower truncated on every polynomial state: only finitely many annihilation modes can differentiate that state. At any fixed remaining nonnegative power of z , its creation factors have only finitely many index choices. All field coefficients are therefore well-defined polynomial endomorphisms.

1.13 ALL OPERATOR PRODUCTS AND THE VERTEX IDENTITIES

Direct differentiation on the polynomial states gives

$$[J_m, J_n] = \lambda m \delta_{m+n,0}, \quad [J_+(z), J_-(w)] = \iota_{z,w} \frac{\lambda}{(z-w)^2},$$

where $\iota_{z,w}$ means expansion in w/z . Commuting annihilation factors past creation factors in the normal fields proves their complete Wick formula. If the factors of a have jet orders r_α and those of b have jet orders s_β , then

$$Y(a, z)Y(b, w) = \sum_M \iota_{z,w} \left(\prod_{\{\alpha, \beta\} \in M} C_{r_\alpha, s_\beta}(z, w) \right) : \prod_{\alpha \text{ unpaired}} J^{[r_\alpha]}(z) \prod_{\beta \text{ unpaired}} J^{[s_\beta]}(w) : .$$

Each matching joins a factor of a to a factor of b ; unpaired fields keep their original positions and divided derivatives. Every commutation removes a paired creation and annihilation factor, so induction on the number of factors proves the formula and its matching multiplicities. It also proves the analogous formula for any number of insertion groups. For instance,

$$Y(q_1^2, z)Y(q_1^2, w) = J(z)^2 J(w)^2 : + \frac{4\lambda}{(z-w)^2} : J(z)J(w) : + \frac{2\lambda^2}{(z-w)^4},$$

with each rational coefficient expanded by $\iota_{z,w}$. Taylor expansion of the unpaired factors gives every regular as well as every singular operator-product coefficient.

Set $\mathbf{1} = \mathbf{1}$. The annihilation operators kill the vacuum, and the creation part of $J^{[m-1]}(z)$ is $e^{zT} q_m$. Hence

$$Y(\mathbf{1}, z) = \text{id}, \quad Y(a, z)\mathbf{1} = e^{zT} a, \quad T\mathbf{1} = 0.$$

The translation identities follow from

$$[T, q_m] = m q_{m+1}, \quad [T, \partial_{q_m}] = -(m-1) \partial_{q_{m-1}},$$

where the second expression is zero at $m = 1$. They give

$$[T, Y(a, z)] = \partial_z Y(a, z) = Y(Ta, z).$$

Locality also follows from the same matching formula. The reverse ordered product has the same normally ordered unpaired factors and the reverse expansion of the same rational contractions. If a, b have r, s factors, respectively, put

$$L = \sum_{\alpha} r_{\alpha} + \sum_{\beta} s_{\beta} + 2 \min(r, s).$$

Every matching denominator divides $(z - w)^L$. After multiplication by this factor the two radial expansions agree, proving $(z - w)^L [Y(a, z), Y(b, w)] = 0$.

The iterate identity requires compatibility of the regular terms as well. Introduce a formal displacement x and put $z = w + x$. Each unpaired jet expands as

$$J^{[r]}(w + x) = \sum_{k \geq 0} \binom{r+k}{k} x^k J^{[r+k]}(w).$$

These are exactly the translated jet states in $Y(a, x)b$. For any third polynomial state c , matchings between the three groups a, b, c , at $z, w, 0$, define one expression

$$F_{a,b,c}(z, w) \in V_{\kappa}[[z, w]][z^{-1}, w^{-1}, (z - w)^{-1}]$$

with finite pole orders. Its three expansions are

$$\iota_{z,w} F = Y(a, z)Y(b, w)c,$$

$$\iota_{w,z} F = Y(b, w)Y(a, z)c,$$

$$\iota_{w,x} F(w + x, w) = Y(Y(a, x)b, w)c.$$

For the last identity, group a matching first by its a - b edges and then by its edges to c . This retains each matching and its multiplicity. At fixed output weight, the expression is a rational function with polynomial numerator in a finite free weight space. This makes all three expansions legitimate coefficient calculations.

For completeness, Jacobi follows from an elementary formal identity. Write $\delta(t) = \sum_{n \in \mathbb{Z}} t^n$ and $\delta(z, w) = z^{-1} \delta(w/z)$. With the corresponding binomial expansion orders,

$$x^{-1} \delta\left(\frac{z-w}{x}\right) - x^{-1} \delta\left(\frac{w-z}{-x}\right) = w^{-1} \delta\left(\frac{z-x}{w}\right).$$

The coefficients of negative powers of x cancel on the left, because they come from nonnegative powers of $z - w$. For $k \geq 0$, the coefficient of x^k on each side is $\partial_w^k \delta(z, w)/k!$; this is the binomial identity $\binom{-l-1}{k} = (-1)^k \binom{l+k}{k}$. Multiply the identity by each rational monomial of a fixed-weight component of $F_{a,b,c}$, using $z = w + x$ in its collision expansion. The same binomial calculation, including negative exponents, gives

$$\begin{aligned} x^{-1} \delta\left(\frac{z-w}{x}\right) Y(a, z)Y(b, w) - x^{-1} \delta\left(\frac{w-z}{-x}\right) Y(b, w)Y(a, z) \\ = w^{-1} \delta\left(\frac{z-x}{w}\right) Y(Y(a, x)b, w). \end{aligned}$$

This is the full vertex Jacobi identity. In modes it reads, for all integers m, n, k ,

$$\sum_{j \geq 0} \binom{m}{j} (a_{(n+j)} b_{(m+k-j)}) = \sum_{j \geq 0} (-1)^j \binom{n}{j} (a_{(m+n-j)} b_{(k+j)}) - (-1)^n b_{(n+k-j)} a_{(m+j)}.$$

The sums are locally finite on polynomial states by lower truncation. All extracted states are even, as the cohomology calculation proved.

Theorem 1.10 (The boundary current at every level). *For every $\kappa \in \mathbb{C}$, the polynomial finite-weight chiral vacuum extracted from $\mathcal{B}_\kappa$ is a vertex algebra over $\mathbb{C}[\hbar]$. Its state space, translation and fields are*

$$V_\kappa = \mathbb{C}[q_1, q_2, \dots][\hbar], \quad Tq_m = mq_{m+1}, \quad Y\left(\prod_i q_{m_i}, z\right) =: \prod_i J^{[m_i-1]}(z) ;,$$

where $J_{-m} = q_m$, $J_m = -\hbar\kappa m \partial_{q_m}$ for $m > 0$, and $J_0 = 0$. The identification $q_{n+1} = [u_n]$ preserves every mode, divided jet, regular product and separated factorization product. Its Heisenberg level is $-\kappa$ with this fixed current.

Proof. The continuous Cauchy conjugation and compact moment homotopies give the stated cohomology and its normal state basis. Transport through that conjugation gives the actual separated-product matching formula. Expanding it with one input at zero yields exactly the fields above, including their regular terms. The preceding vacuum, translation, locality and Jacobi calculations prove all vertex identities. The chain relation and the mode calculation both give level $-\kappa$ without changing the chosen state representatives. $\square$

Theorem 1.11 (The zero-level current). *At $\kappa = 0$, the extracted vertex algebra is the commutative differential algebra $\mathbb{C}[q_1, q_2, \dots][\hbar]$ with $Tq_m = mq_{m+1}$ and $Y(a, z)b = (e^{zT}a)b$. It agrees with the algebraic zero-level Heisenberg vacuum in every mode and product.*

Proof. Both quantum contractions and the annihilation part of J vanish at zero. Only the empty matching remains. Its state-field formula is precisely $(e^{zT}a)b$. $\square$

Corollary 1.12 (Augmentation of the extracted current). *The zero-level polynomial vertex algebra has the augmentation $q_m \mapsto 0$ for all $m \geq 1$. At nonzero level there is no augmentation to $\mathbb{C}[\hbar]$ as a vertex algebra over $\mathbb{C}[\hbar]$.*

Proof. At zero, evaluation at $q_m = 0$ preserves multiplication, translation and the vacuum. At nonzero level $q_{1(1)}q_1 = -\hbar\kappa 1$. A vertex map to the scalar vertex algebra would send its left side to zero and its right side to a nonzero scalar. $\square$

1.14 FORMAL COMPLETION AND THE BULK-BOUNDARY COMPARISON

The preceding constructions are polynomial in $\hbar$ and in field number. The Cauchy exponentials terminate on every observable, and every fixed tuple of polynomial states has finitely many matchings and finite pole orders. Smooth moment expansions converge on a common outer annulus before their finite-weight coefficients are extracted.

Only after this polynomial extraction take the coefficientwise $\hbar$ -completion

$$\widehat{V}_\kappa = \varprojlim_M V_\kappa / \hbar^M V_\kappa = \mathbb{C}[q_1, q_2, \dots][[\hbar]].$$

Every coefficient of $\hbar$ is a finite polynomial, while its field number and weight may increase with that coefficient. The completed state-field operation has target

$$\varprojlim_M (V_\kappa / \hbar^M V_\kappa)((z)).$$

Modulo each $\hbar^M$, the inputs are polynomial states and all the preceding identities hold. The operations commute with reduction in M . Vacuum, translation, Jacobi and locality modulo each $\hbar^M$ therefore pass to this inverse limit.

There need not be a uniform Laurent pole bound for completed inputs. At $\kappa \neq 0$, take $a = \sum_{n \geq 0} \hbar^n q_{n+1}$ and $b = q_1$. Its singular terms are

$$-\kappa \sum_{n \geq 0} (-1)^n (n+1) \hbar^{n+1} z^{-n-2}.$$

The pole orders increase with the $\hbar$ -order. Thus the inverse limit above is the correct field target, rather than $\widehat{V}_\kappa((z))$ with a single Laurent bound. This passage completes the extracted state and field algebra; it makes no claim that cohomology commutes with completion of the entire factorization algebra.

The supported quantum collar map of Theorem 1.4 has its original fields, differential and contraction. It commutes with translations, rotations and every separated product. Hence the current states in the bulk–boundary pushforward are

$$q_m \mapsto [I_\kappa(u_{m-1})], \quad \prod_i q_{m_i} \mapsto \left[\prod_i I_\kappa(u_{m_i-1}) \right].$$

Preservation of separated products passes to their moving-point coefficients, so this comparison preserves every extracted field and mode. The nonzero BF–Chern–Simons field change has bulk parameter κ , whereas the fixed boundary current has level $-\kappa$ with the printed positive unary differential. At zero, this same supported construction realizes the commutative current while retaining the additional bulk gauge and field cohomology.

1.15 SEPARATED PRODUCTS AND TOPOLOGICAL HIGHER OPERATIONS

For distinct holomorphic insertion points, the zero-level Wick product has no contraction terms. Its normal-ordering map sends

$$x_{i,r} \mapsto \partial_{z_i}^r J(z_i), \quad x_{1,0} \star x_{2,0} \star x_{3,0} = x_{1,0} x_{2,0} x_{3,0}.$$

These are equalities on the actual boundary field representation from Theorem 1.11.

An auxiliary topological interval gives the same higher operations on these fields as on their algebraic presentation. To specify the statement, let e_0, e_1 be endpoint cochains and η the oriented edge, with $de_0 = -\eta$, $de_1 = \eta$. Their polynomial-form inclusion and homotopy are

$$i(e_0) = 1 - t, \quad i(e_1) = t, \quad i(\eta) = dt, \quad h(gdt) = \int_0^t g - t \int_0^1 g.$$

Projection takes endpoint values and the integral of a one-form. Direct differentiation gives $dh + hd = 1 - i p$ and $hi = ph = h^2 = 0$. Extend these maps by the identity on boundary field coefficients.

To check all higher identities, let $\mathcal{W}$ be the interval cochains with these field coefficients and let $\mathcal{A}$ be their polynomial-form resolution. The coefficient algebra is the commutative algebra of represented zero-level current fields at the chosen separated points, interpreted coefficientwise in the insertion variables. Write sa for a suspension, with $|sa| = |a| - 1$. Put

$$D_1(sa) = s(da), \quad D_2(sa, sb) = (-1)^{|a|} s(ab).$$

Extend these maps to the direct sum of finite tensor words by inserting them into each consecutive block, with the Koszul sign from the preceding suspended inputs. This is a coderivation for the coproduct

which splits words into initial and final segments. Associativity and the Leibniz rule say $(D_1 + D_2)^2 = 0$. Use the same letters i, p, h for the suspended linear maps and define

$$F_1 = i, \quad R_n = \sum_{a+b=n} D_2(F_a \otimes F_b), \quad F_n = -hR_n, \quad b_n = pR_n \quad (n \geq 2),$$

with $b_1 = sds^{-1}$. An $\mathcal{A}_\infty$ structure here means that the resulting degree-one coderivation b on the tensor words in sW satisfies $b^2 = 0$. The ordinary operations are fixed by

$$b_n(sa_1, \dots, sa_n) = (-1)^{\sum_j (n-j)|a_j|} sm_n(a_1, \dots, a_n).$$

Here is a direct verification of the recursion. The maps F_n extend uniquely to a coalgebra map F . Induct on word length, assuming $DF = Fb$ and $b^2 = 0$ in smaller lengths, where $D = D_1 + D_2$. With the length- n terms temporarily zero, the length- n morphism defect is

$$Q_n = R_n - L_n, \quad L_n = \sum_{\substack{r+q+t=n \\ 2 \leq q \leq n-1}} F_{r+1+t}(1^{\otimes r} \otimes b_q \otimes 1^{\otimes t}).$$

The tensor substitutions carry the insertion signs just specified. Let O_n be the length- n term of b^2 with $b_n = 0$. The identity

$$D(DF - Fb) + (DF - Fb)b = -Fb^2$$

gives $\delta Q_n = -iO_n$, where $\delta Q_n = D_1 Q_n + Q_n b_1$ and b_1 also denotes its extension to words. For $k \geq 2$, $pF_k = hF_k = 0$, by $ph = h^2 = 0$. It follows that $pL_n = hL_n = 0$. Setting $b_n = pQ_n = pR_n$ gives $\delta b_n = -O_n$, which is the missing length- n identity for b^2 . Setting $F_n = -hQ_n = -hR_n$ and using $D_1 h + hD_1 = 1 - ip$ gives

$$\delta F_n = -Q_n + ipQ_n + h\delta Q_n = ib_n - Q_n,$$

because $hi = 0$. This is the missing morphism identity. The induction starts with the cochain maps i, p , so it proves every arity without an infinite sum on any word. The linear map i is a quasi-isomorphism by the explicit contraction.

Every field comparison intertwines each multiplication vertex and each interval homotopy separately. It therefore intertwines these operations in every arity, before summing their binary trees. For example,

$$h(tdt) = \frac{t^2 - t}{2}, \quad \int_0^1 \frac{t - t^2}{2} dt = \frac{1}{12}$$

gives, in ordinary cohomological degrees $|m_n| = 2 - n$,

$$m_3(e_1 J(z_1), \eta J(z_2), \eta J(z_3)) = -\frac{\eta}{12} J(z_1) J(z_2) J(z_3).$$

The degree-one output reflects the interval cochains. It is compatible with the strict degree-zero vertex algebra because this interval resolution has nonzero differential. It is not a cubic interaction in the BF action.

The quantum boundary map above identifies actual supported observables on the boundary with their bulk–boundary pushforward. The current comparison identifies its polynomial chiral vacuum, and the interval argument identifies the transferred separated products. These three maps have different sources. In particular, the last map alone does not describe a bulk action, a collision-chain compactification, or higher operations of an interacting theory. The zero-level BF construction supplies a physical realization of the zero-level current while retaining the additional bulk gauge and field cohomology computed above.

Free fields on an interval

A free action determines Gaussian contractions. It does not determine a multiplication on a chosen complex of topological cochains. The distinction already appears when a field has values at two endpoints and a descendant integrated along the interval between them. Linear interpolation gives a binary product which is associative only up to a specified ternary operation. For a Heisenberg current, that operation retains the double pole of the operator product expansion. Its coefficient is an elementary integral.

The construction below keeps the holomorphic insertion points distinct. It gives all ordered higher operations on the polynomial de Rham model of an interval, with coefficients in the multilocal Wick algebra. An explicit comparison identifies the Wick factors with radially ordered free fields. Topological cochain degree, holomorphic jet order, and powers of the quantum parameter remain separate throughout.

2.1 THE CURRENT AND ITS SEPARATED PRODUCTS

Work on the affine complex curve with coordinate z . Put $K = \mathbb{C}[\kappa][[\hbar]]$, where both parameters have cohomological degree zero. The rank-one Heisenberg vacuum module over K is

$$V = \mathbb{C}[\kappa][q_1, q_2, \dots][[\hbar]].$$

Each coefficient of $\hbar$ is a polynomial in finitely many q_m . All vacuum generators and holomorphic jet symbols have cohomological degree zero. For $m > 0$, define operators

$$J_{-m} = q_m, \quad J_m = \hbar \kappa m \frac{\partial}{\partial q_m}, \quad J_0 = 0.$$

The vacuum is $\mathbf{1} = 1$. Direct differentiation gives

$$[J_m, J_n] = \hbar \kappa m \delta_{m+n,0} \text{ id}.$$

The field $J(z) = \sum_{n \in \mathbb{Z}} J_n z^{-n-1}$ splits as $J_-(z) + J_+(z)$, with creation modes in J_- and annihilation modes in J_+ . In the formal expansion $|z| > |w|$,

$$[J_+(z), J_-(w)] = \hbar \kappa \sum_{m \geq 1} m z^{-m-1} w^{m-1} = \frac{\hbar \kappa}{(z-w)^2}.$$

Here and below, a rational expression denotes its specified formal Laurent expansion. No norm convergence of unbounded operators is asserted.

Normal ordering places every creation mode to the left of every annihilation mode. The field attached to $q_{r_1+1} \cdots q_{r_a+1} \mathbf{I}$ is

$$: \prod_{\nu=1}^a \frac{\partial_z^{r_\nu} J(z)}{r_\nu!} :.$$

The denominators identify mode indices with divided holomorphic jets. All holomorphic derivatives are retained.

For a finite set S of insertion labels, define

$$R_S = \mathbb{C}[z_i \ (i \in S), (z_i - z_j)^{-1} \ (i \neq j)], \quad W_S = (R_S[\kappa][x_{i,r} \mid i \in S, r \geq 0])[[\hbar]].$$

Tensor products of field modules are over K , completed in $\hbar$, with the displayed extension of rational scalars on the output. The symbol $x_{i,r}$ represents the undivided derivative $\partial_{z_i}^r J(z_i)$ inside one globally normal-ordered expression. Every coefficient is a polynomial in finitely many jet symbols. For disjoint sets S, T , extend scalars to $R_{S \cup T}$ and set

$$P_{S,T} = \sum_{i \in S, j \in T} \sum_{r,s \geq 0} C_{ij}^{rs} \frac{\partial}{\partial x_{i,r}} \frac{\partial}{\partial x_{j,s}},$$

$$C_{ij}^{rs} = \hbar \kappa \partial_{z_i}^r \partial_{z_j}^s (z_i - z_j)^{-2}, \quad f \star g = \exp(P_{S,T})(fg).$$

The derivatives in $P_{S,T}$ act only on jet symbols. They do not act on its rational coefficients. On polynomial inputs the exponential is finite. For formal inputs it is defined coefficientwise in $\hbar$.

Fix a total order $i_1, \dots, i_a$ on S , with $|z_{i_1}| > \cdots > |z_{i_a}|$ as a formal expansion convention. Writing $V_o = \mathbb{C}[\kappa][q_1, q_2, \dots]$, define

$$N_S : W_S \longrightarrow \text{Hom}_{\mathbb{C}[\kappa]}(V_o, V_o((z_{i_1})) \cdots ((z_{i_a})))[[\hbar]]$$

by replacing each jet monomial with its globally normal-ordered field and expanding each rational coefficient in that order. A fixed vacuum polynomial involves only finitely many annihilation modes, so the indicated iterated Laurent series are defined coefficientwise.

Proposition 2.1 (Associativity and the field comparison). *For pairwise disjoint S, T, U , the two maps*

$$W_S \otimes W_T \otimes W_U \longrightarrow W_{S \cup T \cup U}, \quad (f \star g) \star h, \quad f \star (g \star h)$$

are equal. In a radial ordering with every point of S outside every point of T , the normal-ordering map N satisfies

$$N(f \star g) = N(f)N(g).$$

Both identities hold for every holomorphic jet and every power of $\hbar$.

Proof. All contraction operators commute: their coefficients are independent of the jet symbols being differentiated. Moreover, $P_{S \cup T, U} = P_{S, U} + P_{T, U}$. Each parenthesization is therefore

$$\exp(P_{S,T} + P_{S,U} + P_{T,U})(fgh).$$

This proves associativity without identifying distinct insertion points.

For the comparison, commute each annihilation mode in f past each creation mode in g . The displayed annihilation-creation commutator gives the contribution of one commutation. Its derivatives give C_{ij}^{rs} . Every commutator is central, so further commutations do not alter it. A collection of a disjoint contractions occurs $a!$ times in $P_{S,T}^a$; division by $a!$ counts it once. Uncontracted modes remain in normal order. This proves the formula on monomials, hence on polynomials. Formal extension is coefficientwise: each input coefficient has finite polynomial degree, and only finitely many powers of $\hbar$ contribute to a fixed coefficient. $\square$

For each fixed radial order, put $W_S^{\text{fld}} = \text{im } \mathcal{N}_S$. The displayed comparison identifies its product with composition of the actual field operators on ordered blocks. If $\mathcal{N}_S(f) = \mathbf{o}$, then $\mathcal{N}_{S \cup T}(f \star g) = \mathbf{o}$ for every following block T . The same holds for a zero factor in the second position. Thus the kernels are preserved by all ordered products. Every interval operation constructed below is a scalar cochain operation multiplied by such a product. It therefore descends to $C_I \otimes W_S^{\text{fld}}$, with the same contraction and proofs. This quotient supplies the operations on the actual free fields, including any relations among their normally ordered expressions.

Write $x_i = x_{i,\mathbf{o}}$ and $C_{ij} = \hbar\kappa/(z_i - z_j)^2$. The first products are

$$\begin{aligned} x_1 \star x_2 &= x_1 x_2 + C_{12}, \\ x_1 \star x_2 \star x_3 &= x_1 x_2 x_3 + C_{12} x_3 + C_{13} x_2 + C_{23} x_1. \end{aligned}$$

Vacuum expectation means taking the scalar term after normal ordering. Two copies of J at the first insertion give

$$\langle : J^2 : (z_1) J(z_2) J(z_3) \rangle_{\text{conn}} = 2C_{12} C_{13}.$$

For three quadratic insertions the connected expectation is $8C_{12} C_{13} C_{23}$. There are two assignments of the first pair of fields to the other two vertices. At each remaining vertex there are two choices of the field used for its first edge. This gives $2 \cdot 2 \cdot 2 = 8$ triangle pairings. Thus a Gaussian theory has nonzero connected composite-field products.

2.2 ENDPOINT VALUES AND AN INTEGRATED DESCENDANT

Give the interval $I = [\mathbf{o}, \mathbf{1}]$ its increasing orientation. Its polynomial forms are the differential graded algebra

$$A_I = \mathbb{C}[t, dt], \quad |t| = \mathbf{o}, \quad |dt| = \mathbf{1}, \quad dt = dt, \quad (dt)^2 = \mathbf{o}.$$

The complex of cellular cochains is

$$C_I^{\mathbf{o}} = \mathbb{C}e_{\mathbf{o}} \oplus \mathbb{C}e_{\mathbf{1}}, \quad C_I^{\mathbf{1}} = \mathbb{C}\eta, \quad de_{\mathbf{o}} = -\eta, \quad de_{\mathbf{1}} = \eta, \quad d\eta = \mathbf{o}.$$

Here $e_{\mathbf{o}}, e_{\mathbf{1}}$ record endpoint values and η records integration along the oriented edge. There are no endpoint vanishing conditions. Define inclusion and projection by

$$\begin{aligned} i(e_{\mathbf{o}}) &= \mathbf{1} - t, & i(e_{\mathbf{1}}) &= t, & i(\eta) &= dt, \\ p(f) &= f(\mathbf{o})e_{\mathbf{o}} + f(\mathbf{1})e_{\mathbf{1}}, \\ p(g \, dt) &= \left(\int_{\mathbf{o}}^{\mathbf{1}} g(t) \, dt \right) \eta. \end{aligned}$$

The degree-minus-one homotopy is

$$h(f) = \mathbf{o}, \quad h(g \, dt) = \int_{\mathbf{o}}^t g(u) \, du - t \int_{\mathbf{o}}^{\mathbf{1}} g(u) \, du.$$

It subtracts the linear function with the same endpoint difference.

Lemma 2.2 (The interval contraction). *The maps above satisfy*

$$pi = \mathbf{1}, \quad dh + hd = \mathbf{1} - ip, \quad hi = ph = h^2 = \mathbf{o}.$$

Proof. The inclusion and projection commute with d by the fundamental theorem of calculus. On $f(t)$, the operator hd gives $f(t) - f(o) - t(f(i) - f(o))$. On $g(t)dt$, the operator dh gives $(g(t) - \int_0^1 g)dt$. These are exactly $1 - i$ and p in their respective degrees. The function $h(gdt)$ vanishes at both endpoints, proving $ph = o$. Also $h(dt) = o$, h vanishes on functions, and its image consists of functions. These observations give $hi = h^2 = o$. $\square$

Use the complexes

$$A(S) = A_I \otimes W_S, \quad C(S) = C_I \otimes W_S,$$

completed coefficientwise in $\hbar$. Their differential acts on the interval factor only. For disjoint supports their product on A is

$$(\alpha f)(\beta g) = (\alpha \wedge \beta)(f \star g).$$

Proposition 2.1 makes this an associative system of differential graded products. The maps i, p, h act on its interval factor and preserve each holomorphic support. These are topological forms with chiral field coefficients. This specified algebraic model does not identify the interval differential with a quantum Batalin–Vilkovisky differential.

The projected binary product $m_2 = p\mu(i \otimes i)$ obeys

$$\begin{aligned} m_2(e_a, e_b) &= \delta_{ab}e_a, & m_2(e_a, \eta) &= m_2(\eta, e_a) = \tfrac{1}{2}\eta, \\ m_2(\eta, \eta) &= o & (a, b \in \{o, i\}). \end{aligned}$$

Every entry is multiplied by the corresponding Wick product of its field coefficients. In particular,

$$m_2(m_2(e_i, e_i), \eta) - m_2(e_i, m_2(e_i, \eta)) = \tfrac{1}{4}\eta.$$

Linear interpolation therefore forces a higher homotopy.

2.3 THE HIGHER OPERATIONS AND THEIR SIGNS

For a cochain a , write sa for its suspension, with $|sa| = |a| - 1$. A sequence of ordered operations is encoded by maps b_n of degree one on suspended inputs. For pairwise disjoint supports these satisfy

$$\sum_{r+q+t=n} (-1)^{|u_1|+\dots+|u_r|} b_{r+t}(u_1, \dots, u_r, b_q(u_{r+1}, \dots, u_{r+q}), u_{r+q+1}, \dots, u_n) = o.$$

All exponents use suspended degrees. The ordinary operations m_n of degree $2 - n$ are defined by

$$b_n(sa_1, \dots, sa_n) = (-1)^{\sum_{j=1}^n (n-j)|a_j|} sm_n(a_1, \dots, a_n).$$

Here the tensor coalgebra has a precise support convention. Fix a finite ambient label set E and extend rational scalars to R_E . Take the direct sum of tensor words whose entries have pairwise disjoint supports $S_1, \dots, S_n \subseteq E$. The coproduct cuts a word into two consecutive pieces. Each operation replaces a consecutive block by one entry supported on its union. Thus all displayed compositions are defined over a common coefficient ring and remain in this coalgebra. The formulas commute with further extensions of E . An A_∞ structure here means a square-zero coderivation of these supported words. The words are finite, so no sum over arities is needed on an element.

On sA , put $D_1(sa) = s(da)$ and $D_2(sa, sb) = (-1)^{|a|}s(ab)$. The differential graded algebra identities say $D^2 = o$, where $D = D_1 + D_2$ is the induced coderivation. Write I, P, H for the suspended maps i, p, h . Starting with $F_1 = I$, define recursively, for $n \geq 2$,

$$R_n = \sum_{a+b=n} D_2(F_a \otimes F_b), \quad F_n = -HR_n, \quad b_n = PR_n, \quad b_1 = sds^{-1}.$$

Each F_n has degree zero. Thus its tensor substitution contributes no extra sign. The only signs come from D_2 and the coderivation insertion rule in the displayed higher identity.

Theorem 2.3 (Ordered transfer with free-field coefficients). *The recursion defining R_n, F_n, b_n gives an uncurved A_∞ structure on the collection $\{C(S)\}$, with operations for every ordered list of pairwise disjoint supports. The maps F_n define an A_∞ quasi-isomorphism to $\{A(S)\}$. All identities preserve the rational holomorphic coefficients and their formal $\hbar$ expansion. The element $e_0 + e_1$ with empty field support is a strict unit.*

Proof. The proof takes place at each finite word length. Suppose the morphism equation $DF = Fb$ and $b^2 = 0$ hold in all smaller lengths. Set the length- n components temporarily to zero. The length- n morphism defect is

$$Q_n = R_n - L_n, \quad L_n = \sum_{\substack{r+q+t=n \\ 2 \leq q \leq n-1}} F_{r+1+t}(1^{\otimes r} \otimes b_q \otimes 1^{\otimes t}),$$

with the tensor signs fixed above. Let O_n be the length- n component of b^2 with $b_n = 0$. The identity

$$D(DF - Fb) + (DF - Fb)b = -Fb^2$$

gives $\delta Q_n = -IO_n$, where $\delta Q_n = D_1 Q_n + Q_n b_1$ and b_1 on the right denotes its extension to the input word. Lower defects vanish by induction. For $k \geq 2$ the recursion gives $PF_k = HF_k = 0$. Consequently $PL_n = HL_n = 0$.

Set $b_n = PQ_n = PR_n$. Applying P to the defect identity gives $\delta b_n = -O_n$, which is exactly the length- n equation $b^2 = 0$. Next set $F_n = -HQ_n = -HR_n$. The contraction identity and $HI = 0$ give

$$\delta F_n = -Q_n + IPQ_n + H\delta Q_n = Ib_n - Q_n.$$

This is the missing length- n morphism equation. Induction proves both assertions. The linear term I is a quasi-isomorphism by Lemma 2.2.

For the unit assertion, consider a binary transfer tree with a unit leaf. At its first vertex the unit removes that multiplication. If the other branch is a leaf, the next homotopy kills it by $HI = 0$. If that branch has an internal vertex, the next map gives $H^2 = 0$ or $PH = 0$. Thus every operation of arity greater than two vanishes on a unit input. The binary and unary unit identities follow from the displayed table and $d(e_0 + e_1) = 0$.

Every tree has finitely many vertices. Associativity makes its field coefficient the same ordered Wick product, independently of the tree. The polynomial contraction acts over K and neither changes a holomorphic denominator nor a power of $\hbar$. The construction therefore extends to the stated coefficients and all holomorphic jets. $\square$

The Wick exponential supplies every field coefficient of the transfer. The explicit interval contraction fixes the higher operations, including choices which a boundary identity alone cannot fix.

2.4 THE TERNARY OPERATION AND THE NEXT IDENTITIES

Write $v = se_1$, $w = s\eta$, and $\mathbf{e} = s(e_0 + e_1)$. Their degrees are $-1, 0, -1$, respectively. The first suspended operations, with field coefficients suppressed, are

$$\begin{aligned} b_1(v) &= w, & b_1(w) &= 0, & b_2(v, v) &= v, & b_2(v, w) &= \frac{1}{2}w, & b_2(w, v) &= -\frac{1}{2}w, \\ b_3(v, w, w) &= \frac{1}{12}w, & b_3(w, v, w) &= -\frac{1}{6}w, & b_3(w, w, v) &= \frac{1}{12}w. \end{aligned}$$

Every other ternary value on v, w is zero. Values involving se_0 follow from $se_0 = \mathbf{e} - v$ and strict unitality.

The integral producing the first coefficient is

$$h(t \, dt) = \frac{t^2 - t}{2}, \quad F_2(v, w) = s \frac{t - t^2}{2}, \quad b_3(v, w, w) = s \left(\int_0^1 \frac{t - t^2}{2} \, dt \right) \eta = \frac{1}{12}w.$$

For the middle input there are two trees, each with the opposite sign. This gives $-w/6$. The last input gives the same value as the first.

Example 2.4 (The Heisenberg ternary operation). For three distinct holomorphic insertions, ordinary unsuspension gives

$$\begin{aligned} m_3(e_1 x_1, \eta x_2, \eta x_3) \\ = -\frac{\eta}{12} \left(x_1 x_2 x_3 + \frac{\hbar \kappa x_3}{(z_1 - z_2)^2} + \frac{\hbar \kappa x_2}{(z_1 - z_3)^2} + \frac{\hbar \kappa x_1}{(z_2 - z_3)^2} \right). \end{aligned}$$

The output has topological degree one. The inputs have total degree two, so m_3 has degree -1 as required. The first term survives at $\hbar = 0$; the other terms retain the quantum central contraction. Applying N identifies the parenthesis with $J(z_1)J(z_2)J(z_3)$ in the chosen radial expansion.

The arity-three identity on (v, v, w) is an exact cancellation:

$$\underbrace{\frac{1}{2}w - \frac{1}{4}w}_{b_2 \text{ insertions}} + \underbrace{\left(-\frac{1}{6}w\right) - \frac{1}{12}w}_{b_1 \text{ insertions into } b_3} = 0.$$

It supplies the homotopy for the binary associator $\eta/4$. For (v, w, v, w) the arity-four identity is

$$-\frac{1}{12}w + \frac{1}{24}w - \frac{1}{24}w + \frac{1}{12}w = 0.$$

The first three summands insert b_2 into b_3 . The last inserts b_3 into b_2 . The remaining summand is zero. With field coefficients, every term multiplies the same fourfold Wick product by Proposition 2.1. These cancellations therefore verify the identities on the actual holomorphic expressions.

The quartic operation for this contraction vanishes identically. The following arity nevertheless requires a new operation. The next integrations give

$$\begin{aligned} F_3(v, w, w) &= s \left(\frac{t^3}{6} - \frac{t^2}{4} + \frac{t}{12} \right), \\ F_4(v, w, w, w) &= -s \frac{t^2(1-t)^2}{24}, \\ b_4(v, w, w, w) &= 0, \quad b_5(v, w, w, w, w) = -\frac{1}{720}w. \end{aligned}$$

Here $\int_0^1 t^2(1-t)^2 dt = 1/30$. On (v, w, w, v, w) , the three b_3 -into- b_3 terms sum to

$$-\frac{1}{72}w - \frac{1}{144}w + \frac{1}{72}w = -\frac{1}{144}w.$$

The two nonzero insertions of b_1 into b_5 give $w/180 + w/720 = w/144$. Their coefficients follow from the five possible positions of the function input, computed below. Their cancellation proves the arity-five identity on an input where setting $b_3 = 0$ would fail.

2.5 EVERY ARITY FROM ONE POLYNOMIAL INTEGRAL

Let the Bernoulli numbers be defined by

$$\frac{u}{e^u - 1} = \sum_{k \geq 0} B_k \frac{u^k}{k!}, \quad B_1 = -\frac{1}{2}.$$

For $k \geq 1$, put $c_k = (-1)^k B_k / k!$.

Proposition 2.5 (Complete interval operations). *For $k \geq 2$ and $0 \leq j \leq k$,*

$$b_{k+1}(w^{\otimes j}, v, w^{\otimes(k-j)}) = (-1)^j \binom{k}{j} c_k w.$$

All other operations of arity at least three on v, w vanish. These formulas, the binary table, and strict unitality determine every operation. For disjoint field supports each value is multiplied by the full ordered Wick product of its coefficients.

Proof. Every nonzero higher transfer tree has only one function input. Indeed, every nonroot internal edge applies h , so its nonzero output is a function vanishing at the endpoints. A vertex cannot multiply two such outputs and then survive another h . Thus the tree must be a comb, with one leaf attached at each successive vertex. Every attached leaf before the root must be dt , since h kills functions. If the last leaf is a function, the root product is a function vanishing at both endpoints, which p kills. An all- dt tree vanishes at its first vertex. This proves the asserted restriction on inputs.

Define $T(q) = h(q dt)$ on polynomials. For the word $(v, w, \dots, w)$ there is one comb, and its root coefficient is

$$c_k = \int_0^1 (-T)^{k-1} t dt.$$

To evaluate all these integrals, set $G(u, t) = \sum_{a \geq 0} u^a (-T)^a t$ and $M(u) = \int_0^1 G(u, t) dt$. The equation $G = t - uTG$ implies

$$\partial_t G = 1 - uG + uM, \quad G(u, 0) = 0, \quad G(u, 1) = 1.$$

Solving this formal differential equation gives

$$G(u, t) = \frac{1 - e^{-ut}}{1 - e^{-u}}, \quad M(u) = \frac{1}{1 - e^{-u}} - \frac{1}{u} = \sum_{k \geq 1} \frac{(-1)^k B_k}{k!} u^{k-1}.$$

The apparent poles at $u = 0$ cancel, so these are formal power series. Finally, placing j copies of w before v gives $\binom{k}{j}$ combs. Each left attachment contributes a minus sign from $D_2(s dt, sf) = -s(f dt)$. Right attachments contribute no sign. Their common integral is c_k , proving the formula. $\square$

The scalar formulas follow from the interval integrals, while the field coefficients enter through the proved Wick comparison. At $\kappa = 0$ or $\hbar = 0$, the contraction terms disappear while the topological higher products remain. Thus this higher arity is not a measurement of quantum loop order.

2.6 THE BETA-GAMMA FIELDS AND THE CURVE OPERATION

For the bosonic beta-gamma system use the base $\mathbb{C}[[\hbar]]$ and replace the Heisenberg symbols by $x_{i,r}^\beta, x_{i,r}^\gamma$. Its elementary contractions are

$$C_{ij}^{\beta\gamma} = \frac{\hbar}{z_i - z_j}, \quad C_{ij}^{\gamma\beta} = -\frac{\hbar}{z_i - z_j}, \quad C_{ij}^{\beta\beta} = C_{ij}^{\gamma\gamma} = 0.$$

Jet contractions are their corresponding holomorphic derivatives. These coefficients are symmetric under simultaneous exchange of the point and field labels, as bosonic locality requires.

For completeness, take modes $\beta(z) = \sum_m \beta_m z^{-m-1}$ and $\gamma(z) = \sum_n \gamma_n z^{-n}$ with $[\beta_m, \gamma_n] = \hbar \delta_{m+n,0}$. The vacuum is killed by β_m for $m \geq 0$ and by γ_n for $n \geq 1$. Explicitly, the vacuum module is $\mathbb{C}[q_m (m \geq 1), p_n (n \geq 0)] [[\hbar]]$, with

$$\beta_{-m} = q_m (m \geq 1), \quad \gamma_{-n} = p_n (n \geq 0), \quad \beta_m = \hbar \partial_{p_m} (m \geq 0), \quad \gamma_n = -\hbar \partial_{q_n} (n \geq 1).$$

Summing the two geometric series gives the displayed contractions. The proof of Proposition 2.1 therefore applies without change to the enlarged symbol set.

In particular, writing β_i, γ_i for its undifferentiated symbols, the ternary operation is

$$m_3(e_1\beta_1, \eta\gamma_2, \eta\beta_3) = -\frac{\eta}{12} \left(\beta_1\gamma_2\beta_3 + \frac{\hbar\beta_3}{z_1 - z_2} - \frac{\hbar\beta_1}{z_2 - z_3} \right).$$

The same scalar higher operations therefore act on both a double-pole current and a simple-pole first-order system, with their different holomorphic contractions intact.

There is a further operation when insertion points collide. In the Heisenberg vertex algebra, the regular coefficient is $a_{(-1)}b = \text{Res}_{z=0} z^{-1}Y(a, z)b dz$. This residue extracts a Laurent coefficient and has contour normalization $1/(2\pi i)$. For three currents, direct normal ordering and Taylor expansion give

$$((J_{(-1)}J)_{(-1)}J) - J_{(-1)}(J_{(-1)}J) = \hbar\kappa \partial^2 J.$$

Indeed, $:JJ:(z)J(w)$ contains $2\hbar\kappa J(z)/(z-w)^2$, whose regular coefficient contributes $\hbar\kappa \partial^2 J$. The reversed iterated regular product has no such term. Thus this regular product cannot be the binary multiplication of a degree-zero ordinary minimal $\mathcal{A}_\infty$ algebra. The nonzero result also exhibits the precise information lost by replacing the separated Wick product with coincident commutative symbols.

2.7 TOPOLOGICAL RESOLUTION AND BOUNDARY OBSERVABLES

The interval construction has a strict comparison to the original multilocal field system. The map

$$\mathcal{W}_S \longrightarrow C(S), \quad f \longmapsto (e_o + e_i)f$$

is a strict $\mathcal{A}_\infty$ quasi-isomorphism. Its image is the constant cochain, the differential kills it, and every higher operation on it vanishes by strict unitality in the interval factor. For $a = o, i$, evaluation $\epsilon_a : C(S) \rightarrow \mathcal{W}_S$ is also a strict quasi-isomorphism: higher outputs have edge degree, and evaluation kills them. Writing $\iota(f) = (e_o + e_i)f$, the composite $\epsilon_a \iota$ is the identity. The other composite has the explicit chain homotopy

$$dk_a + k_a d = 1 - \iota \epsilon_a, \quad k_o(\eta) = e_i, \quad k_i(\eta) = -e_o, \quad k_a(e_b) = o.$$

For example, on e_o, e_i, η , the first identity with $a = o$ reads $-e_i, e_i, \eta$ on both sides. This explains how the same free fields admit a strict model and the explicit nonzero higher operations calculated above.

For a physical realization of the algebraic specialization $\kappa = k \in \mathbb{C}$, use the split holomorphic BF family of Chapter 1 with bulk parameter $\ell = -k$. Its fields on $\mathbb{C} \times \mathbb{R}_{\geq o}$ have separate partial-form and holomorphic-one-form components (α, β) ; write $d' = \bar{\partial} + d_t$ for the partial de Rham differential. They carry a fixed nondegenerate cross pairing and differential

$$Q_\ell(\alpha, \beta) = (d' \alpha, d' \beta + \ell \partial \alpha).$$

The chiral boundary condition is $\iota^* \alpha = o$. The integrated pairing requires at least one compactly supported argument, and the integrated action is defined on compact fields; local densities are defined on unrestricted smooth fields. For $k \neq o$, the field change $\alpha + \ell^{-1} \beta$ identifies this theory with abelian Chern–Simons theory whose symmetric coefficient pairing is $\langle e, e \rangle = \ell = -k$.

The boundary quantum observable complex has the positive unary operator and contraction

$$D = \bar{\partial} + \hbar \Delta_{\mu_\ell}, \quad \mu_\ell(u, v) = -\ell \int_{\mathbb{C}}^{\text{res}} u \wedge \partial v, \quad \int^{\text{res}} = \frac{1}{2\pi i} \int.$$

Choose a smooth radial compact cutoff χ equal to one near the origin. Its current is fixed by $q_{n+1} = [-\bar{\partial} \chi / z^{n+1}]$, $n \geq o$. For separated evaluation forms u_a, u_b and their representatives e_a, e_b on a common outer radial annulus, the compact chain calculation gives

$$[u_a u_b] = [e_a e_b] - \frac{\hbar \ell}{(a-b)^2} \mathbf{1} = [e_a e_b] + \frac{\hbar k}{(a-b)^2} \mathbf{1}.$$

Thus it is the extracted level $-\ell = k$ that matches the algebraic parameter of this chapter. The Cauchy conjugation and moment calculation in Chapter 1 transport every separated product to its Wick formula and recover the complete mode action

$$J_{-m} = q_m, \quad J_m = \hbar k m \partial_{q_m} \quad (m > o), \quad J_o = o, \quad J(z)J(w) \sim \frac{\hbar k}{(z-w)^2}.$$

The compact collar map intertwines the actual quantum differentials and all separated products. Consequently the represented fields have precisely the normalization used by the interval transfer. The extraction first uses polynomial $\hbar$ and the finite-weight vacuum space. Formal completion is applied afterwards to the extracted modes and Wick products, coefficient by coefficient; on arbitrary completed

states the field target is the inverse limit of the Laurent-series spaces modulo $\hbar^M$. This does not identify full disk cohomology with its finite-weight vacuum or interchange cohomology and completion of the full observable complex.

At $k = 0$, the split field pairing remains nondegenerate and the boundary current becomes commutative, with all polynomial jet fields retained. The field change to the nonzero de Rham presentation contains ℓ^{-1} and has no zero specialization. In particular, the zero symmetric pairing on a single Chern–Simons coefficient space supplies no inverse BV pairing or propagator. The zero-level physical realization uses the two separate BF field components.

The compact quantum contraction of Theorem 1.8 supplies a chain map from the finite current part of this interval model to the full boundary observable complex. Fix its finite list of points and divided jets, and set $\kappa = -k$ in that theorem. Its normal polynomial algebra $A = \mathbb{C}[y_1, \dots, y_N][\hbar]$ represents the common-annulus currents. An undivided jet of order n_i is represented by $n_i! y_i$. Insertion groups remain at distinct holomorphic points.

Proposition 2.6 (The supported interval chain map). *Let $\iota_U : A \rightarrow \mathcal{B}_{-k}(U)$ and $L_U : \mathcal{B}_{-k}(U) \rightarrow A$ be the compact inclusion and moment projection of Proposition 1.9. For $a = 0, 1$, define*

$$\Phi_a : C_I \otimes A \longrightarrow \mathcal{B}_{-k}(U), \quad \Phi_a(\alpha \otimes F) = \epsilon_a(\alpha) \iota_U(F), \quad \Psi(b) = (e_0 + e_1) \otimes L_U(b).$$

These are chain maps for the interval differential and the actual quantum differential, respectively. They satisfy

$$\Phi_a \Psi = \iota_U L_U, \quad 1 - \Psi \Phi_a = d(k_a \otimes 1) + (k_a \otimes 1) d \quad \text{on } C_I \otimes A.$$

Their induced map identifies $H^\bullet(C_I \otimes A) = A$ with the annular polynomial subspace of $H^\bullet(\mathcal{B}_{-k}(U))$. For every tuple of polynomial currents at distinct points, the comparison of its supported product with the annular Wick expression has the compact primitive $\rho H F$ in (1.3).

Proof. Endpoint evaluation kills dC_I and $\iota_U(F)$ is a quantum cycle. The map L_U intertwines the quantum differential with zero. These statements prove the chain-map equations. The equality $L_U \iota_U = 1$ and the displayed interval contraction prove both composite identities. The split inclusion proves the cohomology assertion. The product statement is (1.3), with $-\hbar\kappa = \hbar k$. $\square$

One can retain the interval factor during the compact contraction. On $C_I \otimes Q$, use the total differential $d(\alpha \otimes q) = d_I \alpha \otimes q + (-1)^{|\alpha|} \alpha \otimes D_Q q$. Then

$$\mathcal{H}(\alpha \otimes q) = (-1)^{|\alpha|} \alpha \otimes Hq, \quad d\mathcal{H} + \mathcal{H}d = 1 - (1 \otimes \iota_U \Pi).$$

The two terms containing $d_I \alpha$ cancel by their opposite signs. The remaining terms are exactly the compact quantum contraction. Thus this extension keeps the interval differential distinct from the quantum differential while intertwining both.

The map Φ_a is a map of complexes. The product comparison uses compact homotopies because common-annulus representatives overlap. In particular, the second-order quantum differential does not define an ordinary differential graded multiplication on that full target. An ordered higher morphism requires compatible Taylor maps on these same supported complexes. A collision comparison also requires operations on collision chains and their boundary strata. Neither construction follows from the linear chain map alone.

Ordered defect operations and symmetric perturbative brackets likewise require distinct insertion rules: the former retain the order of inputs along the defect, while the latter sum over permutations with

their graded signs. The Wick field comparison preserves ordered products at distinct points. Its supported realization has the compact chain homotopies above. A comparison with renormalized configuration integrals must construct compatible Taylor maps on the same complex and verify their equations at every arity. Neither the Gaussian contractions nor the interval identities alone supply the quantum master equation for an interacting theory.

2.8 SYMMETRIC TAYLOR MAPS AND COINCIDENT SUPPORTS

The quantum differential fails the Leibniz identity when supports overlap. Its defect supplies a binary bracket. The Cauchy conjugation then has higher Taylor coefficients which intertwine that bracket with zero. At three inputs, a quadratic current already has a nonzero coefficient. The same binary coefficient extends the double pole as a distribution on the diagonal of the curve.

Fix an open disk $U \subset \mathbb{C}$, the complex orientation, and $\int^{\text{res}} = (2\pi i)^{-1} \int$. Let B be the graded commutative polynomial algebra of compact smooth kernels in $\Omega_c^{\bullet}(U)[1]$, with coefficients in $\mathbb{C}[\hbar]$. A function generator has degree -1 and a form generator has degree zero. Symmetric powers use graded coinvariants of compact smooth kernels on U^r . Their direct sum has finite field number. Write

$$D = d + \hbar \Delta_\mu, \quad d = \bar{\partial}, \quad \mu(f, v) = -\kappa \int_U^{\text{res}} f \partial v.$$

The cocycle is zero except on one function and one form generator. Its pair contraction has degree one. Define

$$\mathcal{C}v(z) = \frac{1}{2\pi i} \int \frac{v(w) \wedge dw}{z - w}, \quad P(u, v) = -\kappa \int^{\text{res}} (\mathcal{C}u) \partial v.$$

The pairing P is zero unless both generators are forms. Let $K = \Delta_P$ and $N = e^{-\hbar K}$. The Cauchy identities give the chain equation

$$dN = ND.$$

This is Proposition 1.5. All products in the following formulas are the graded commutative products of kernels. They need not have disjoint supports.

For homogeneous $a, b \in B$, set

$$\ell_1(a) = Da, \quad \ell_2(a, b) = D(ab) - (Da)b - (-1)^{|a|} a(Db).$$

Thus ℓ_2 is the actual second-order Leibniz defect, including its factor $\hbar$. It is graded symmetric and has degree one. Use the convention in which an $L_\infty[1]$ structure consists of graded symmetric degree-one operations, encoded by a square-zero coderivation on the coaugmented symmetric coalgebra. Its morphism has graded symmetric Taylor coefficients of degree zero. There is no suspension sign hidden in this convention.

For an ordered list of homogeneous inputs $a_1, \dots, a_n$, let $\mathcal{P}_n$ be its set of partitions into nonempty blocks. Order the entries in each block increasingly, and order the blocks by their smallest entries. Let $\epsilon(\pi; a)$ be the Koszul sign of this rearrangement. Put

$$F_n(a_1, \dots, a_n) = \sum_{\pi \in \mathcal{P}_n} (-1)^{|\pi|-1} (|\pi|-1)! \epsilon(\pi; a) \prod_{C \in \pi} N \left(\prod_{i \in C} a_i \right). \quad (2.1)$$

In particular,

$$F_1 = N, \quad F_2(a, b) = N(ab) - N(a)N(b).$$

Theorem 2.7 (Supported symmetric Taylor comparison). *The operations ℓ_1, ℓ_2 , with $\ell_n = 0$ for $n \geq 3$, define an $L_\infty[1]$ structure on B . The maps (2.1) define an $L_\infty[1]$ isomorphism from this structure to the abelian complex (B, d) . They are continuous on each fixed compact support and fixed polynomial degree and commute with extension by zero. Their first higher identity is*

$$dF_2(a, b) - F_2(Da, b) - (-1)^{|a|} F_2(a, Db) = N\ell_2(a, b). \quad (2.2)$$

Proof. Use the direct sum of finite symmetric words $S^c(B)$, together with its coaugmentation. Each individual word has finite length. Let E be the coalgebra map whose Taylor coefficient of every positive arity is multiplication. Its action on a word is the sum over partitions of that word, with the stated Koszul signs. Its linear part is the identity. It is triangular in word length, so it has a unique coalgebra inverse.

The inverse Taylor coefficient in arity n is $(-1)^{n-1}(n-1)!$ times multiplication. One verifies this by substituting partitions: a set with c components contributes

$$\sum_{\pi \in \mathcal{P}_c} (-1)^{|\pi|-1} (|\pi| - 1)! = \begin{cases} 1, & c = 1, \\ 0, & c > 1. \end{cases}$$

The identity follows, for example, by comparing coefficients in $\log(\exp t) = t$ as formal power series over $\mathbb{C}$. Graded reordering supplies precisely the Koszul sign already fixed.

Let $\widehat{D}$ denote the coderivation applying D to one entry of a word. The conjugate $E^{-1}\widehat{D}E$ has first Taylor coefficient D and second coefficient ℓ_2 . Every contraction in D uses at most two input blocks. Partition inversion cancels a contribution unless it uses every block of its Taylor input. Thus the coefficients of arity at least three vanish. This argument also applies to the derivation part, which uses only one block. Since $D^2 = 0$, the conjugate squares to zero. This proves the asserted bracket identities with their signs.

The same conjugation for the derivation d has only its unary coefficient. The chain isomorphism N induces the coalgebra isomorphism $S^c(N)$. Therefore

$$F = E^{-1}S^c(N)E, \quad \widehat{d}F = F(E^{-1}\widehat{D}E).$$

Partition inversion gives exactly (2.1). It also proves every higher morphism identity, without an infinite arity sum on a word. Its inverse is $E^{-1}S^c(N^{-1})E$. Expanding the equation in arity two gives (2.2) directly.

For fixed polynomial inputs, all contractions terminate. The Cauchy estimate bounds a contracted smooth kernel on its compact support by one additional derivative seminorm. Every iterated contraction uses disjoint individual field variables. Hence it uses tensor products of those continuous contractions, not products of distributions in a shared variable. Finite sums and ordinary kernel products preserve continuity. The global Cauchy formula and zero extension prove naturality. $\square$

For separated radial current representatives, write $x_i = u_{a_i,0}$ and $p_{ij} = \kappa(a_i - a_j)^{-2}$. Then $N(x_i) = x_i$ and

$$F_2(x_1, x_2) = -\hbar p_{12}, \quad F_3(x_1, x_2, x_3) = 0, \quad F_3(x_1^2, x_2, x_3) = 2\hbar^2 p_{12} p_{13}.$$

For the last identity, expand

$$\begin{aligned} N(x_1^2 x_2 x_3) &= x_1^2 x_2 x_3 - \hbar(2p_{12} x_1 x_3 + 2p_{13} x_1 x_2 + p_{23} x_1^2) \\ &\quad + 2\hbar^2 p_{12} p_{13}. \end{aligned}$$

The three two-block terms in F_3 cancel all single contractions. The five uncontracted terms have total coefficient $1 - 3 + 2 = 0$. The remaining term joins all three input groups. In general, partition inversion retains exactly the matchings whose graph on the input groups is connected. Individual field factors still occur in at most one edge. This explains why three linear currents have zero coefficient while a quadratic current changes it.

Proposition 2.8 (The distributional collision homotopy). *The binary map on form generators extends the separated double pole to all compact smooth two-variable kernels by the distribution*

$$T(z, w) = -\partial_z \frac{1}{z - w}.$$

It satisfies

$$\partial_{\bar{z}} T = -\pi \partial_z \delta^{(2)}(z - w)$$

in coefficient distributions with respect to area measure. For a compact function generator f and form generator v , its Taylor identity is the contact cancellation

$$F_2(\bar{\partial}f, v) = -\hbar\mu(f, v), \quad F_2(f, v) = 0, \quad \ell_2(f, v) = \hbar\mu(f, v).$$

Proof. The function $(z - w)^{-1}$ is locally integrable on $\mathbb{C}^2$. Its distributional derivative therefore exists. For $u = a(w)d\bar{w}$ and $v = b(z)d\bar{z}$, the actual Cauchy pairing is

$$P(u, v) = \frac{\kappa}{\pi^2} \int \frac{a(w)\partial_z b(z)}{z - w} dA_z dA_w = \frac{\kappa}{\pi^2} \langle T, a(w)b(z) \rangle.$$

The first formula extends to every compact smooth two-variable kernel by differentiation in z before integration. Away from the diagonal, $T = (z - w)^{-2}$. Differentiating $\partial_{\bar{z}}(z - w)^{-1} = \pi \delta^{(2)}(z - w)$ proves the displayed contact term. Adjoining $d\bar{z}$ gives the corresponding form identity.

On two form generators, $F_2 = -\hbar P$. On a function and a form, both Cauchy contractions vanish, so $F_2(f, v) = 0$. Finally, $P(\bar{\partial}f, v) = \mu(f, v)$ by the compact Cauchy identity. These give the contact cancellation and verify (2.2) on the first inputs where the quantum bracket is nonzero. $\square$

The proposition concerns smooth test kernels, including overlapping supports. A diagonal distribution need not admit evaluation on point-supported distributions. At nonzero κ , the scalar

$$F_2(u_{a,0}, u_{b,0}) = -\hbar\kappa(a - b)^{-2}$$

has no finite limit as $b \rightarrow a$. The cutoffs must shrink to retain disjoint supports, and their smooth seminorms are then unbounded. Thus continuity on fixed smooth support spaces does not give a collision limit of these point-labelled states.

The maps in Theorem 2.7 refine the actual Cauchy chain comparison and retain its quantum differential. They supply the symmetric bracket comparison required by the supported boundary observables. The interval operations earlier in this chapter are ordered operations on a different coalgebra. A morphism from that ordered structure requires ordered Taylor maps and their compatibility with the collision chains. The symmetric construction does not assert such a morphism. Nor does it assign a product to coincident distributional currents.

2.9 THE HIGHER COMPACT CURRENT RETRACT

The compact projection $\Pi = \pi N_Q$ is the linear term of a symmetric higher morphism. Its homotopy $H = N_Q^{-1}h_0N_Q$ also extends to higher arities. A polynomial path which scales the contractible variables constructs all the required coefficients. The points, cutoffs, and compact supports remain fixed along this path.

Use the finite supported current complex Q of Theorem 1.8, with $\delta_i = x_i - y_i$, odd variables θ_i , and $A = \mathbb{C}[\gamma_1, \dots, \gamma_N][\hbar]$. Its differential is $D_Q = N_Q^{-1}d_0N_Q$, where

$$d_0 = \sum_i \delta_i \partial_{\theta_i}, \quad N_Q = e^{-\hbar K_Q}, \quad K_Q = \sum_{i < j} p_{ij} \partial_{\theta_i} \partial_{\theta_j}.$$

The symmetric degree-one brackets on Q are D_Q and its second-order Leibniz defect. All higher brackets vanish. Give A the zero differential and zero brackets. For a degree-zero linear map T between graded commutative algebras, write $C_n(T)$ for its partition formula (2.1), with N replaced by T . Thus $C_1(T) = T$ and $C_2(T)(a, b) = T(ab) - T(a)T(b)$.

Proposition 2.9 (Higher moment projection). *For the finite list of current jets, the maps*

$$\Lambda_n = C_n(L_U) : \mathcal{B}_\kappa(U)^{\otimes n} \longrightarrow A$$

define a graded symmetric $L_\infty[1]$ morphism. Here $L_U = RN_U$ is the full smooth moment chain map of Proposition 1.9. The annular inclusion $\iota_U : A \rightarrow \mathcal{B}_\kappa(U)$ is a strict morphism and $\Lambda \iota_U = 1$ as coalgebra maps. On the embedded subcomplex Q , the Taylor maps are $\Lambda_n|_Q = C_n(\Pi)$. This restriction is an $L_\infty[1]$ quasi-isomorphism.

Proof. The moment map R is a classical algebra map and chain map to A . It therefore induces a strict morphism from the abelian complex (B, d) to the abelian complex A . Compose it with the symmetric Taylor morphism of Theorem 2.7. Since R preserves every product, its composite has coefficients $RC_n(N_U) = C_n(RN_U)$. This proves the asserted morphism on the full smooth carrier.

The differential and its binary defect vanish on annular polynomials. The inclusion is therefore strict. The identities $N_U \iota_U = \iota_U$ and $R \iota_U = 1$ make its unary composite the identity. Partition inversion makes every higher composite zero. Thus the composite is the identity coalgebra map, including all arities. On Q , the equality $L_U \rho = \Pi$ gives the stated restriction. Its linear coefficient is the previously proved quasi-isomorphism $\Pi : Q \rightarrow A$. $\square$

The full boundary complex need not retract by a chain homotopy onto this finite polynomial algebra. The explicit homotopy below is on the actual subcomplex Q , which contains all the chosen compact current representatives and their odd primitives.

Let $\Omega = \mathbb{C}[\tau, d\tau]$ be polynomial forms on $[0, 1]$, with $|\tau| = 0$, $|d\tau| = 1$, and $(d\tau)^2 = 0$. The parameter τ is a homotopy parameter, separate from the physical interval coordinate and the curve coordinate. Place the form factor first. On $\Omega \otimes Q$, use

$$(\alpha \otimes q)(\beta \otimes r) = (-1)^{|q||\beta|} \alpha \beta \otimes qr, \quad D_{\text{tot}}(\alpha \otimes q) = d_\tau \alpha \otimes q + (-1)^{|\alpha|} \alpha \otimes D_Q q.$$

The classical total differential replaces D_Q by d_τ . Extend N_Q to $N_{\text{out}} = 1 \otimes N_Q$. Define the classical algebra map

$$P : Q \longrightarrow \Omega \otimes Q, \quad P(y_i) = y_i, \quad P(\theta_i) = \tau \theta_i, \quad P(\partial_i) = \tau \partial_i + d\tau \theta_i.$$

All displayed expressions have the degree of their input. Set

$$T = N_{\text{out}}^{-1} P N_Q, \quad \mathcal{T}_n = C_n(T).$$

Let $j_A : A \rightarrow \Omega \otimes A$ include the constant forms, and let $\Lambda_\Omega = C(1 \otimes \Pi)$ be the higher moment map on the target complex.

Theorem 2.10 (Polynomial higher homotopy). *The maps $\mathcal{T}_n$ define an $L_\infty[1]$ morphism from Q to $\Omega \otimes Q$, both with the brackets determined by their actual quantum differentials. Endpoint evaluation gives*

$$\text{ev}_0 \mathcal{T} = \iota \Lambda|_Q, \quad \text{ev}_1 \mathcal{T} = 1_Q.$$

The homotopy fixes the inclusion of A and satisfies $\Lambda_\Omega \mathcal{T} = j_A \Lambda|_Q$. Its unary coefficient recovers the compact chain homotopy:

$$\int_0^1 [d\tau] T = H.$$

Here $[d\tau]$ means the coefficient of the leftmost $d\tau$. Every formula is polynomial in $\hbar$ on each polynomial input. After substitution of the compact representatives, all outputs remain supported in the same disk U .

Proof. The classical total differential satisfies

$$d_{\text{tot}} P(\theta_i) = d\tau \theta_i + \tau \delta_i = P(d_\circ \theta_i), \quad d_{\text{tot}} P(\delta_i) = d\tau \delta_i - d\tau \delta_i = 0.$$

It also kills $P(y_i)$. Both classical differentials are derivations, so P is a chain map on the whole polynomial algebra. The Gaussian conjugation commutes with the parameter differential. Consequently T is a chain map for the quantum differentials.

For any such chain map, the coalgebra construction $C(T) = E^{-1} S^c(T) E$ intertwines the descendant coderivations. Apply the partition argument of Theorem 2.7 to source and target. The target differential still has order at most two. It therefore has only the stated unary and binary brackets. This proves every Taylor identity for $\mathcal{T}$.

Endpoint evaluation sets $d\tau$ to zero. It gives $P_0 = \iota\pi$ and $P_1 = 1$. Since $N_Q^{\pm 1} \iota = \iota$, one obtains $T_0 = \iota\Pi$ and $T_1 = 1$. Cumulants commute with composition, as follows by cancellation of the middle EE^{-1} in their coalgebra expressions. Thus $C(T_0) = \iota C(\Pi) = \iota\Lambda|_Q$, while $C(T_1)$ is the identity with no higher coefficients. Also $P\iota = \iota$ and $\pi P = \pi$. These equations prove the assertions about inclusion and projection after Gaussian conjugation and the same coalgebra composition.

For a monomial q of positive total degree r in δ_i, θ_i , substitution gives

$$P(q) = \tau^r q + d\tau \tau^{r-1} \sum_i \theta_i \delta_{\delta_i} q.$$

The left odd factors carry exactly the Koszul sign of the substitution. A degree-zero monomial in these variables has no $d\tau$ coefficient. Integrating τ^{r-1} gives $1/r$, so $\int [d\tau] P = h_\circ$. Conjugation then gives $\int [d\tau] T = N_Q^{-1} h_\circ N_Q = H$. No operation changes the chosen compact representatives or their support. The Gaussian exponentials terminate, proving the final assertions. $\square$

The first higher coefficients display the correction required during the homotopy. The chain rule gives $K_{\text{out}} P = \tau^2 P K_Q$, and therefore

$$T = P \exp(-\hbar(1 - \tau^2) K_Q).$$

Put $a_i = y_i + \tau \delta_i + d\tau \theta_i = T(x_i)$. For two distinct current indices,

$$\mathcal{T}_2(x_i, x_j) = -\hbar(1 - \tau^2) p_{ij}.$$

Its total differential is $2\hbar\tau d\tau p_{ij}$. The target binary bracket on (a_i, a_j) is $-2\hbar\tau d\tau p_{ij}$. They cancel in the arity-two morphism equation, whose source bracket on (x_i, x_j) is zero. Thus keeping only the linear path would violate that equation at nonzero covariance and an interior value of τ .

The first connected three-input coefficient is

$$\mathcal{T}_3(x_1^2, x_2, x_3) = 2\hbar^2(1 - \tau^2)^2 p_{12} p_{13}.$$

It equals the moment-projection coefficient at $\tau = 0$ and vanishes at $\tau = 1$. The full polynomial path supplies its compatibility with every other Taylor coefficient.

The higher moment map extends the same L_U used by Proposition 2.6. Its restriction supplies a symmetric higher inverse to the annular current inclusion, and the polynomial path refines the same compact homotopy used in the supported product comparison. The ordered interval operations and distributional collision domains still require their own compatible maps.

Ordered Taylor maps for supported currents

The quantum differential permits multiplication of observables with disjoint supports. A finite change of current representatives makes their Wick products strict. Interval interpolation then gives ordered Taylor maps into the actual quantum complex. Their linear term has an explicit compact homotopy to the common-annulus current inclusion.

3.1 THE SUPPORT OF AN ORDERED PRODUCT

Fix a centered disk $U = D_R \subset \mathbb{C}$, distinct points $a_1, \dots, a_m \in U$, and disjoint centered disks $D_i \Subset U$ about these points. Choose smooth radial cutoffs $\chi_i \in C_c^\infty(D_i)$ equal to one near a_i . Choose an outer radial cutoff $\chi \in C_c^\infty(U)$ equal to one on a disk containing every $\overline{D_i}$. For each i , fix a finite set $J_i \subset \mathbb{Z}_{\geq 0}$ of jet orders. Let $E = \{(i, n) : 1 \leq i \leq m, n \in J_i\}$. For $v = (i, n) \in E$, put

$$u_v = -\frac{\bar{\partial}\chi_i}{(z - a_i)^{n+1}}, \quad e_v = -\frac{\bar{\partial}\chi}{(z - a_i)^{n+1}}, \quad f_v = \frac{\chi - \chi_i}{(z - a_i)^{n+1}}.$$

These are compact smooth representatives, and $\bar{\partial}f_v = u_v - e_v$. Their observable degrees are $|u_v| = |e_v| = 0$ and $|f_v| = -1$. Use the increasing orientation on $[0, 1]$ and the complex orientation on U , with residue integral $(2\pi i)^{-1} \int$.

Fix the bulk parameter $\ell \in \mathbb{C}$ and work first over $R = \mathbb{C}[\hbar]$. The boundary complex is

$$\mathcal{B}_\ell(V) = \left(\text{Sym}(\Omega_c^{\circ, \bullet}(V)[1])[\hbar], D = \bar{\partial} + \hbar \Delta_{\mu_\ell} \right), \quad \mu_\ell(v, w) = -\ell \int_V^{\text{res}} v \wedge \partial w.$$

The shifted Dolbeault differential has the sign specified in Lemma 1.3. Each symmetric power uses compact smooth kernels on V^r and graded coinvariants. The sum over symmetric powers is algebraic. No field-number completion is used. Define symmetric constants, with a total order on E , by

$$p_{(i,n),(j,r)} = \begin{cases} \frac{\ell}{n!r!} \partial_{a_i}^n \partial_{a_j}^r (a_i - a_j)^{-2}, & i \neq j, \\ 0, & i = j. \end{cases}$$

The extracted current level is $k = -\ell$. Thus its divided-jet Wick contraction is $-\hbar p_{v\omega}$.

Write x_ν, y_ν, θ_ν for the generators represented by u_ν, e_ν, f_ν , respectively. Their finite compact complex is

$$Q = R[x_\nu, y_\nu \mid \nu \in E] \otimes \Lambda(\theta_\nu \mid \nu \in E), \quad |\theta_\nu| = -1,$$

with left odd derivatives and differential

$$D_Q = \sum_\nu (x_\nu - y_\nu) \partial_{\theta_\nu} + \hbar \sum_{\nu < \omega} p_{\nu\omega} (\partial_{x_\omega} \partial_{\theta_\nu} + \partial_{x_\nu} \partial_{\theta_\omega}). \quad (3.1)$$

The substitution $\rho : Q \hookrightarrow \mathcal{B}_\ell(U)$ is the embedding of Proposition 1.7. For a graded R -module M , write $T^c(M) = \bigoplus_{n \geq 1} M^{\otimes n}$, with coproduct given by splitting words between consecutive entries.

Proposition 3.1 (The overlapping product obstruction). *Suppose $\hbar p_{\nu\omega} \neq 0$ in R . On $T^c(sQ)$ there is no uncurved square-zero coderivation whose unary and binary terms are*

$$B_1(sa) = sD_Q a, \quad B_2(sa, sb) = (-1)^{|a|} s(ab), \quad |sa| = |a| - 1.$$

The assertion holds regardless of the terms of arity at least three.

Proof. The component from length two to length one of the square is

$$(B_1 B_2 + B_2 B_1)(sa, sb) = (-1)^{|a|} s(D_Q(ab) - (D_Q a)b - (-1)^{|a|} a D_Q b).$$

Here the second unary insertion has sign $(-1)^{|sa|}$. For $a = \theta_\nu, b = x_\omega$, equation (3.1) makes the right side $-\hbar p_{\nu\omega} 1$. It is nonzero. An operation of arity at least three vanishes on a word of length two. Hence no such operation changes this square. $\square$

For $S \subseteq \{1, \dots, m\}$, define $U_S = \coprod_{i \in S} D_i$ and $E_S = \{(i, n) \in E : i \in S\}$. The binary factorization product

$$\mu_{S,T} : \mathcal{B}_\ell(U_S) \otimes_R \mathcal{B}_\ell(U_T) \longrightarrow \mathcal{B}_\ell(U_{S \cup T})$$

exists for $S \cap T = \emptyset$. It extends each input by zero and multiplies its observable kernels. Every cross contraction is zero: the local pairing evaluates two generators at a common point, and their supports are disjoint. Thus D obeys the graded Leibniz rule for this product. The same conclusion holds on compact smooth kernels by restriction to the relevant diagonals. These products are associative and commute with the indicated extensions by zero.

3.2 A COMPACT LIFT OF EVERY WICK POLYNOMIAL

Let $\mathcal{W}_S = R[y_\nu \mid \nu \in E_S]$, concentrated in degree zero. The variables are normal symbols for divided holomorphic jets. This is the finite-current specialization of the rational Wick symbols of Chapter 2, under

$$z_i \longmapsto a_i, \quad \kappa \longmapsto -\ell, \quad x_{i,n} \longmapsto n! y_{(i,n)}.$$

Here κ is the algebraic current level of that chapter. The map evaluates rational coefficients before any radial Laurent expansion. Distinct a_i make all denominators defined. For disjoint S, T , set

$$K_{S,T} = \sum_{\nu \in E_S, \omega \in E_T} p_{\nu\omega} \partial_{y_\nu} \partial_{y_\omega}, \quad f \star g = e^{-\hbar K_{S,T}} (fg), \quad K_S = \sum_{\substack{\nu < \omega \\ \nu, \omega \in E_S}} p_{\nu\omega} \partial_{y_\nu} \partial_{y_\omega}.$$

All derivatives fix the coefficients. Each exponential terminates on a polynomial. The commuting operators satisfy $K_{S \cup T} = K_S + K_T + K_{S,T}$. It follows that $\star$ is associative on pairwise disjoint labels.

Let $\rho_S^x : \mathcal{W}_S \rightarrow \mathcal{B}_\ell(U_S)$ substitute u_ν for y_ν and multiply the resulting form generators. Define the degree-zero linear map

$$\Gamma_S(f) = \rho_S^x(e^{\hbar K_S} f). \quad (3.2)$$

Theorem 3.2 (Separated Wick lift). *The maps Γ_S are chain maps and satisfy*

$$\mu_{S,T}(\Gamma_S f, \Gamma_T g) = \Gamma_{S \cup T}(f \star g) \quad (S \cap T = \emptyset). \quad (3.3)$$

Their values lie in the actual polynomial current subcomplex $R[u_\nu \mid \nu \in E_S]$ of $\mathcal{B}_\ell(U_S)$. They are linear isomorphisms onto that subcomplex. For $S \subseteq T$, let $j_{S,T}$ extend observables by zero from U_S to U_T , and include W_S into W_T by the same normal variables. Then $j_{S,T}\Gamma_S = \Gamma_T|_{W_S}$.

Proof. The differential kills every polynomial in form generators: their Dolbeault derivatives vanish, and μ_ℓ pairs no two such generators. This proves the chain-map assertion. For the product, compute before substitution:

$$e^{\hbar K_{S \cup T}} e^{-\hbar K_{S,T}}(f g) = e^{\hbar(K_S + K_T)}(f g) = (e^{\hbar K_S} f)(e^{\hbar K_T} g).$$

Substitution turns the last product into the disjoint factorization product. This proves (3.3). The u_ν are independent by their disjoint disks and angular Fourier powers. Their polynomial algebra embeds in smooth kernels, as in Proposition 1.7. The inverse exponential $e^{-\hbar K_S}$ proves the isomorphism assertion. On a polynomial using only E_S , every derivative in $K_T - K_S$ kills that polynomial and every successive derivative of it. Thus $e^{\hbar K_T}|_{W_S} = e^{\hbar K_S}$, which proves the extension-by-zero identity. $\square$

For currents at two and three distinct points, with $u_i = u_{(i,o)}$,

$$\begin{aligned} \Gamma_{\{1,2\}}(y_1 y_2) &= u_1 u_2 + \hbar p_{12}, \\ \Gamma_{\{1,2\}}(y_1 \star y_2) &= u_1 u_2, \\ \Gamma_{\{1,2,3\}}(y_1 \star y_2 \star y_3) &= u_1 u_2 u_3. \end{aligned}$$

The nonzero contractions appear when these supported products are expressed in common-annulus coordinates.

3.3 THE ORDERED BAR AND ITS INTERVAL MAPS

Let $A_I = \mathbb{C}[t, dt]$, with $|t| = 0$, $|dt| = 1$, $d_I t = dt$, $d_I(dt) = 0$, and $(dt)^2 = 0$. Let $C_I = \mathbb{C}e_0 \oplus \mathbb{C}e_1 \oplus \mathbb{C}\eta$ have degrees 0, 0, 1 and differential $de_0 = -\eta$, $de_1 = \eta$, $d\eta = 0$. The contraction is

$$\begin{aligned} i(e_0) &= 1 - t, & i(e_1) &= t, & i(\eta) &= dt, \\ p(f) &= f(0)e_0 + f(1)e_1, & p(gdt) &= \left(\int_0^1 g\right)\eta, & h(gdt) &= \int_0^t g(u)du - t \int_0^1 g(u)du, \end{aligned}$$

with $h(f) = 0$. Direct differentiation gives $dh + hd = 1 - i p$ and $hi = ph = h^2 = 0$. There are no endpoint vanishing conditions. Set

$$C(S) = C_I \otimes W_S, \quad A(S) = A_I \otimes W_S, \quad \mathcal{T}(S) = A_I \otimes \mathcal{B}_\ell(U_S).$$

The source differential acts on the interval factor. On the actual target, use

$$d_{\mathcal{T}}(\alpha \otimes q) = d_I \alpha \otimes q + (-1)^{|\alpha|} \alpha \otimes Dq,$$

and, for disjoint S, T , the product

$$(\alpha \otimes q)(\beta \otimes r) = (-1)^{|q||\beta|}(\alpha \wedge \beta) \otimes \mu_{S,T}(q, r).$$

The total differential is a derivation for these disjoint products. Define $G_S : A(S) \rightarrow \mathcal{T}(S)$ by $G_S(\alpha \otimes f) = \alpha \otimes \Gamma_S f$. Theorem 3.2 makes G a strict morphism of these systems of differential graded products.

For any such collection $V(S)$, its supported tensor coalgebra is

$$T_{\text{disj}}^c(sV) = \bigoplus_{n \geq 1} \bigoplus_{S_1, \dots, S_n \text{ pairwise disjoint}} sV(S_1) \otimes_R \cdots \otimes_R sV(S_n).$$

The coproduct cuts a word between consecutive entries. Empty label sets may occur and carry the unit. Each word has finite length. A Taylor term replaces a consecutive block by its union label. It preserves disjointness from all remaining entries.

For a system with differential d and product μ , define $B_1(sa) = sda$ and $B_2(sa, sb) = (-1)^{|a|} s\mu(a, b)$. Its ordered bar differential on $u_1 | \cdots | u_n$ is

$$B(u_1 | \cdots | u_n) = \sum_{q=1}^2 \sum_{r=0}^{n-q} (-1)^{\sum_{j=1}^r |u_j|} u_1 | \cdots | u_r | B_q(u_{r+1}, \dots, u_{r+q}) | u_{r+q+1} | \cdots | u_n. \quad (3.4)$$

The degree is one. The differential, Leibniz, and associativity identities give $B^2 = 0$, in lengths one, two, and three, respectively. There are no other components of the square. This is the suspended convention for the tensor bar construction described by Keller [1], §3.6, pp. 11–12.

Write I, P, H for the suspended interval maps on sA . Let B^A and B^T be the bar differentials on A and $\mathcal{T}$. On C put $b_1 = sd_C s^{-1}$ and define recursively

$$\begin{aligned} f_1 &= I, & R_n &= \sum_{a+b=n} B_2^A(f_a \otimes f_b), \\ f_n &= -H R_n, & b_n &= P R_n \quad (n \geq 2), & \mathcal{F}_n &= (sG s^{-1}) f_n. \end{aligned}$$

Each f_n and $\mathcal{F}_n$ has suspended degree zero.

Theorem 3.3 (Ordered current Taylor comparison). *The maps b_n define an uncurved A_∞ structure on $C(S)$. The maps $\mathcal{F}_n$ define a morphism of differential graded supported tensor coalgebras*

$$\mathcal{F} : (T_{\text{disj}}^c(sC), b) \longrightarrow (T_{\text{disj}}^c(s\mathcal{T}), B^T), \quad B^T \mathcal{F} = \mathcal{F} b.$$

The linear term is interval interpolation followed by the compact current lift Γ_S . It induces an injection on cohomology into the full target. Its restriction to the polynomial current subcomplex of the target is an A_∞ quasi-isomorphism.

Proof. The interval recursion can be checked at each finite word length. Suppose its morphism equation and $b^2 = 0$ hold below length n . With the length- n terms temporarily zero, write the morphism defect as $Q_n = R_n - L_n$, where

$$L_n = \sum_{\substack{r+q+t=n \\ 2 \leq q < n}} f_{r+1+t} (1^{\otimes r} \otimes b_q \otimes 1^{\otimes t}).$$

Tensor substitution includes the sign of passing the degree-one map b_q through the preceding suspended inputs. Let O_n be the length- n term of b^2 . The identity

$$B^A(B^A f - f b) + (B^A f - f b) b = -f b^2$$

gives $\delta Q_n = -I O_n$, with $\delta Q_n = B_1^A Q_n + Q_n b_1$. For $j \geq 2$, the recursion implies $P f_j = H f_j = 0$. Thus $P L_n = H L_n = 0$. Applying P gives $\delta(P R_n) = -O_n$. Set $b_n = P R_n$ to make b^2 vanish in length n . Since $B_1^A H + H B_1^A = 1 - I P$ and $H I = 0$, setting $f_n = -H R_n$ gives

$$B_1^A f_n - f_n b_1 = -Q_n + I P Q_n + H \delta Q_n = I b_n - Q_n.$$

This completes the length- n morphism equation. Induction proves $B^A f = f b$ and $b^2 = 0$. All operations preserve the support labels, so the induction uses only defined products.

The strict map G induces a coalgebra map commuting with bar differentials. Composing it with f proves the displayed identity for $\mathcal{F}$ at every length. The contraction makes I a quasi-isomorphism. The lift Γ is an isomorphism onto the polynomial current subcomplex, proving the restricted quasi-isomorphism. For the full target, let j_S extend by zero from U_S to U . Let τ_S set normal variables outside E_S to zero. The moment projection L_U of Proposition 1.9 satisfies $L_U \rho = \pi e^{-\hbar K_Q}$, where $K_Q = \sum_{\nu < \omega} p_{\nu\omega} \partial_{x_\nu} \partial_{x_\omega}$ and $\pi(x_\nu) = y_\nu$, $\pi(\theta_\nu) = 0$, $\pi(y_\nu) = y_\nu$. This exponential cancels $e^{\hbar K_S}$ on an input using only E_S . Thus $\tau_S L_U j_S \Gamma_S = 1$ and $p \otimes (\tau_S L_U j_S)$ is a chain left inverse to $i \otimes \Gamma_S$. This proves its cohomology injection. $\square$

3.4 THE FIRST TWO TAYLOR COEFFICIENTS

Put $v = s e_1$, $w = s \eta$, and $\mathbf{e} = s(e_0 + e_1)$, of degrees $-1, 0, -1$, respectively. For $f_j \in \mathcal{W}_{S_j}$ with disjoint labels, write

$$U(f_1, \dots, f_n) = \mu(\Gamma_{S_1} f_1, \dots, \Gamma_{S_n} f_n) = \Gamma_{\cup S_j}(f_1 \star \dots \star f_n).$$

Define $c(t) = (t - t^2)/2$ and $g(t) = t^3/6 - t^2/4 + t/12$. All nonzero binary and ternary Taylor terms on v, w are

$$\begin{aligned} \mathcal{F}_2(v f_1, w f_2) &= s(c(t) U(f_1, f_2)), \\ \mathcal{F}_2(w f_1, v f_2) &= -s(c(t) U(f_1, f_2)), \\ \mathcal{F}_3(v f_1, w f_2, w f_3) &= s(g(t) U(f_1, f_2, f_3)), \\ \mathcal{F}_3(w f_1, v f_2, w f_3) &= -2s(g(t) U(f_1, f_2, f_3)), \\ \mathcal{F}_3(w f_1, w f_2, v f_3) &= s(g(t) U(f_1, f_2, f_3)). \end{aligned} \tag{3.5}$$

Every other such value is zero. Values with an interval unit $\mathbf{e}$ are zero in arities at least two. Linearity and $s e_0 = \mathbf{e} - v$ determine every value on C_I . For example, $B_2^A(st, sdt) = s(tdt)$, while $B_2^A(sdt, st) = -s(tdt)$. Hence $f_2(v, w) = sc$ and $f_2(w, v) = -sc$. The three nonzero cubic expressions R_3 are $s(cdt)$, $-2s(cdt)$, and $s(cdt)$. Since $-h(cdt) = g$, these give (3.5). All remaining expressions either vanish or are functions, which h kills. A unit input reduces a tree to HI , H^2 , or PH , so its higher Taylor terms vanish.

For homogeneous suspended inputs u, v, w , the full first morphism equations, including the unary input terms, are

$$\begin{aligned} B_1^T \mathcal{F}_2(u, v) + B_2^T(\mathcal{F}_1 u, \mathcal{F}_1 v) &= \mathcal{F}_1 b_2(u, v) + \mathcal{F}_2(b_1 u, v) \\ &\quad + (-1)^{|u|} \mathcal{F}_2(u, b_1 v), \end{aligned} \tag{3.6}$$

and

$$\begin{aligned} B_1^T \mathcal{F}_3(u, v, w) + B_2^T(\mathcal{F}_1 u, \mathcal{F}_2(v, w)) + B_2^T(\mathcal{F}_2(u, v), \mathcal{F}_1 w) \\ = \mathcal{F}_1 b_3(u, v, w) + \mathcal{F}_2(b_2(u, v), w) + (-1)^{|u|} \mathcal{F}_2(u, b_2(v, w)) \\ + \mathcal{F}_3(b_1 u, v, w) + (-1)^{|u|} \mathcal{F}_3(u, b_1 v, w) \\ + (-1)^{|u|+|v|} \mathcal{F}_3(u, v, b_1 w). \end{aligned} \tag{3.7}$$

On (v, w, w) the first and third terms on the left are $s(g' dt U)$ and $s(cdt U)$. The right side is $s(dt U/12)$. Thus this equation is exactly $g'(t) + c(t) = 1/12$. On (v, v, w) it is the function identity $tc = c/2 - 3g$. These equations also check the sign contributed by the first suspended input.

The ordinary degree-1 – n Taylor map F_n is specified by

$$\mathcal{F}_n(s a_1, \dots, s a_n) = (-1)^{\sum_{j=1}^n (n-j)|a_j|} s F_n(a_1, \dots, a_n).$$

In particular, on singleton currents, $F_2(e_1 y_1, \eta y_2) = c(t) u_1 u_2$ and $F_3(e_1 y_1, \eta y_2, \eta y_3) = -g(t) u_1 u_2 u_3$. The negative ternary sign comes from the middle input of degree one.

3.5 COMPARISON WITH THE ANNULAR INTERVAL INCLUSION

Let $\delta_\nu = x_\nu - y_\nu$ and $K_Q = \sum_{\nu < \omega} p_{\nu\omega} \partial_{x_\nu} \partial_{x_\omega}$. The compact contraction has

$$\Pi = \pi e^{-\hbar K_Q}, \quad H_Q = e^{\hbar K_Q} h_o e^{-\hbar K_Q}, \quad D_Q H_Q + H_Q D_Q = 1 - \iota \Pi,$$

where π sets δ_ν, θ_ν to zero and $h_o = q^{-1} \sum_\nu \theta_\nu \partial_{\delta_\nu}$ in positive total (δ, θ) -degree q , and is zero in degree zero. These are the maps of Theorem 1.8. Let $j_S : \mathcal{B}_\ell(U_S) \rightarrow \mathcal{B}_\ell(U)$ extend by zero. Define a degree-minus-one map on W_S by

$$T_S f = \rho H_Q \left(e^{\hbar K_S} f \Big|_{y=x} \right).$$

The notation $y = x$ replaces normal symbols by the local variables, while the annular variables inside Q remain fixed.

Proposition 3.4 (The compact linear comparison). *For every $f \in W_S$,*

$$D T_S f = j_S \Gamma_S f - \iota_U f. \quad (3.8)$$

On the interval tensor product, the map $\mathcal{K}_S(\alpha f) = (-1)^{|\alpha|} \alpha \otimes T_S f$ satisfies

$$d_{\mathcal{T}} \mathcal{K}_S + \mathcal{K}_S d_A = (1 \otimes j_S) G_S - (1 \otimes \iota_U).$$

At endpoint $a = 0, 1$, the linear part of the ordered comparison therefore has a chain homotopy to $\Phi_a(\alpha f) = \epsilon_a(\alpha) \iota_U(f)$.

Proof. The polynomial $e^{\hbar K_S} f(x)$ is a D_Q -cycle. Applying Π cancels its exponential and gives $f(y)$. The compact contraction proves (3.8). For the interval identity, the terms containing $d_I \alpha$ have opposite signs and cancel. The other terms equal $\alpha \otimes D T_S f$. Compose with interval inclusion and endpoint evaluation to obtain the final statement. Endpoint evaluation kills $d C_I$. $\square$

For two and three distinct current indices, the primitives are

$$T(y_1 y_2) = \rho \left[\frac{\theta_1(x_2 + y_2) + \theta_2(x_1 + y_1)}{2} \right],$$

$$T(y_1 y_2 y_3) = \rho \left[\sum_{i=1}^3 \theta_i \left(y_j y_k + \frac{\delta_j y_k + y_j \delta_k}{2} + \frac{\delta_j \delta_k}{3} + \frac{\hbar p_{jk}}{3} \right) \right],$$

where $\{j, k\} = \{1, 2, 3\} \setminus \{i\}$. Indeed $H_Q e^{\hbar K_Q} = e^{\hbar K_Q} h_o$ on these inputs. The second derivative of $\delta_j \delta_k / 3$ produces the positive quantum correction $\hbar p_{jk} / 3$. Their differentials are, respectively,

$$u_1 u_2 + \hbar p_{12} - e_1 e_2,$$

$$u_1 u_2 u_3 + \hbar(p_{12} u_3 + p_{13} u_2 + p_{23} u_1) - e_1 e_2 e_3.$$

For the actual product $u_1 u_2 u_3$, its common-annulus projection is

$$e_1 e_2 e_3 - \hbar(p_{12} e_3 + p_{13} e_2 + p_{23} e_1).$$

Thus projecting the coefficient of $F_3(e_1 y_1, \eta y_2, \eta y_3)$ gives $-g(t)$ times the complete three-current Wick expression. Here e_i in the current products denotes the annular form, whereas e_i in the input denotes the interval endpoint cochain.

The constants $p_{\nu\omega}$ are rational functions of the distinct insertion points, with every jet derivative retained. An undivided jet of order n is obtained by multiplying its divided generator by $n!$. With $\ell = -k$, the elementary contraction becomes $\hbar k(a_i - a_j)^{-2}$, as in the free-field interval construction. At $\ell = 0$, all exponential contractions vanish and every displayed map still exists.

For each fixed finite jet list, all formulas extend coefficientwise to $\hbar$ -adic completions with finite polynomial degree in each coefficient. Complete each supported word separately and retain the direct sum over finite word lengths. A fixed coefficient receives finitely many terms. The equations hold modulo every $\hbar^M$, hence in the inverse limit. No commutation of full-boundary cohomology with completion is required.

Proposition 3.5 (The binary equation for annular representatives). *Suppose $\hbar p_{12} \neq 0$. The common-annulus linear map cannot satisfy the binary ordered morphism equation from the Wick source if the target binary operation is imposed as ordinary polynomial multiplication in $\mathcal{B}_\ell(U)$.*

Proof. Use the constant interval cochains $(e_0 + e_1) y_1$ and $(e_0 + e_1) y_2$. Their differentials vanish and their Wick product is $(e_0 + e_1)(y_1 y_2 - \hbar p_{12})$. The binary morphism equation would require a degree-minus-one observable q with

$$Dq = \iota_U(y_1 y_2 - \hbar p_{12}) - \iota_U(y_1) \iota_U(y_2) = -\hbar p_{12} \mathbf{1}.$$

Applying the chain projection L_U gives $0 = -\hbar p_{12} \mathbf{1}$ in the polynomial normal algebra, a contradiction. $\square$

The target of Theorem 3.3 retains the unions U_S of the original disjoint disks. Its operations use their actual factorization products. Extending all entries to a common disk and replacing them by annular representatives changes these support conditions. Proposition 3.4 gives the resulting linear chain homotopy, but it does not turn overlapping multiplication into an ordered bar differential. An ordered comparison allowing arbitrary disk changes requires coherent maps for those changes. Collision chains require their own domains and boundary-stratum equations.

Distributional products of three boundary currents

A double pole has a distributional extension, but its Dolbeault derivative is a contact term. Three quadratic currents also produce a triangle of poles. Its three collision normals are dependent. Partial fractions construct an extension without multiplying those three distributions. Complex scaling then proves that the extension preserves rational identities.

4.1 THE DOUBLE POLE AND ITS RESIDUE

Work on $\mathbb{C}$ with $z = x + iy$, $dA = dx dy$, and $\partial_z = (\partial_x - i\partial_y)/2$. Distributions act complex linearly on compactly supported smooth densities. The residue convention is

$$\int^{\text{res}} \alpha = \frac{1}{2\pi i} \int \alpha, \quad d\bar{z} \wedge dz = 2i dA.$$

Fix $\kappa \in \mathbb{C}$, put $\lambda = -\hbar\kappa$, and give $\hbar$ degree zero. All formulas are polynomial in $\hbar$ on polynomial inputs. For $n \geq 1$, define scalar distributions

$$E_1 = \frac{1}{z} \in L^1_{\text{loc}}(\mathbb{C}), \quad E_n = \frac{(-1)^{n-1}}{(n-1)!} \partial_z^{n-1} E_1, \quad E_0 = 1. \quad (4.1)$$

Define the $(0, 1)$ -currents

$$R_n = \frac{\pi(-1)^{n-1}}{(n-1)!} \partial_z^{n-1} \delta_0 d\bar{z}.$$

Here δ_0 is normalized by $\langle \delta_0, \phi dA \rangle = \phi(0)$.

Lemma 4.1 (Distributional residue). *The restrictions of E_n to $\mathbb{C} \setminus \{0\}$ equal z^{-n} . They satisfy*

$$zE_n = E_{n-1}, \quad \partial_z E_n = -nE_{n+1}, \quad \bar{\partial} E_n = R_n. \quad (4.2)$$

For every $t \in \mathbb{C}^\times$, pullback under $z \mapsto tz$ gives $E_n(tz) = t^{-n} E_n(z)$. In particular, the current contraction is

$$C(z) = \lambda E_2(z), \quad \bar{\partial} C = -\pi \lambda \partial_z \delta_0 d\bar{z}. \quad (4.3)$$

Proof. Local integrability follows from $\int_0^\epsilon r^{-1}r dr < \infty$. For a test function ϕ , apply Stokes' theorem outside $|z| = \epsilon$. The inner circle has the clockwise orientation. It gives

$$\langle \partial_{\bar{z}} E_1, \phi dA \rangle = -\lim_{\epsilon \downarrow 0} \int_{|z| > \epsilon} \frac{\partial_{\bar{z}} \phi}{z} dA = \pi \phi(o).$$

The integral over the omitted disk tends to zero for the differentiated test function. Differentiating this identity proves the residue formula. The derivative identity follows directly from the definition. Also $zE_1 = 1$. Differentiating this identity $n - 1$ times gives $zE_n = E_{n-1}$ for $n > 1$. Change of variables proves complex scaling for E_1 . Distributional differentiation gives it for every E_n . $\square$

The coefficient $\lambda = -\hbar\kappa$ follows from the compact quantum boundary differential. With the conventions above, that differential is

$$\begin{aligned} \mathcal{B}_\kappa(U) &= \text{Sym}(\Omega_c^{\circ, \bullet}(U)[1])[\hbar], \\ D &= \bar{\partial} + \hbar \Delta_{\mu_\kappa}, \quad \mu_\kappa(u, v) = -\kappa \int^{\text{res}} u \wedge \partial v. \end{aligned} \tag{4.4}$$

A function generator has degree -1 , and a $(o, 1)$ -form generator has degree zero. The symmetric powers contain compactly supported smooth kernels on U^r , with all derivative seminorms on each fixed compact support. The algebra is a direct sum over r . A point-supported current is not an element of this smooth observable carrier.

The scalar contraction on smooth form kernels has exactly the extension in (4.1). For $u = a(w)d\bar{w}$ and $v = b(z)d\bar{z}$,

$$P_\kappa(u, v) = \frac{\kappa}{\pi^2} \langle E_2(z - w), a(w)b(z) dA_z dA_w \rangle.$$

Indeed $E_2 = -\partial_z E_1$, so integration by parts gives $\kappa \pi^{-2} \int (z - w)^{-1} a(w) \partial_z b(z) dA_z dA_w$. This is the Cauchy contraction in the smooth observable complex. The Wick contraction has the opposite factor, $-\hbar P_\kappa$. No extension of D to arbitrary distributional observable kernels is used.

4.2 TRANSVERSE PRODUCTS AND THE TRIANGLE

For a scalar distribution T , write $\text{WF}(T)$ for its real wavefront set. The multiplication criterion requires that no nonzero covector of one factor is the negative of a covector of the other. Brouder–Dang–Hélein prove the corresponding continuity statement in [2, Theorem 6.1, p. 21].

For each $n \geq 1$,

$$\text{WF}(E_n) = T_o^* \mathbb{C} \setminus \{o\}. \tag{4.5}$$

The distribution is smooth away from zero and singular at zero. Its singularity follows from its unbounded restriction z^{-n} . Its wavefront set over zero is nonempty and invariant under all rotations, by complex scaling. Rotations act transitively on real covector directions, which proves (4.5).

If u, v are independent complex linear coordinates on $\mathbb{C}^2$, define $E_a(u)E_b(v)$ as the pullback of the tensor product $E_a \otimes E_b$. This is also the wavefront product. Its two conormal spaces meet only at zero. No extension choice at $(u, v) = (o, o)$ is needed for this product. The same construction permits one or both E factors to be replaced by R .

Put $x = z_1 - z_2$, $y = z_2 - z_3$, and $c = z_3$. Thus $z_1 - z_3 = x + y$. The first shared-edge scalar coefficient is

$$\langle : J^2 : (z_1) J(z_2) J(z_3) \rangle = 2\lambda^2 E_2(x) E_2(x + y).$$

The two linear coordinates $x, x + y$ are independent. For three quadratic insertions, the connected coefficient away from the collision divisors is

$$\frac{8\lambda^3}{x^2 y^2 (x + y)^2}.$$

At $x = y = 0$, nonzero covectors from the three conormal spaces sum to zero: choose ξdx , ξdy , and $-\xi d(x + y)$ in real covector notation. Thus the wavefront multiplication criterion fails for the three factors. Failure of this sufficient criterion alone does not prove that every regularization fails. The following formula specifies the extension.

Proposition 4.2 (Triangle extension). *The distribution*

$$\begin{aligned} T_\Delta &= E_2(x)E_4(y) + E_2(x + y)E_4(y) \\ &\quad - 2E_1(x)E_5(y) + 2E_1(x + y)E_5(y) \end{aligned} \quad (4.6)$$

extends $x^{-2}y^{-2}(x + y)^{-2}$. Each product displayed is transverse. It has complex scaling degree -6 : $T_\Delta(tx, ty) = t^{-6}T_\Delta(x, y)$. It is independent of the chosen partial-fraction decomposition, provided that each smooth point of a collision divisor uses its Laurent extension (4.1) and each term has this exact complex scaling.

Proof. Ordinary partial fractions give

$$\frac{1}{x^2 y^2 (x + y)^2} = \frac{1}{x^2 y^4} + \frac{1}{(x + y)^2 y^4} - \frac{2}{x y^5} + \frac{2}{(x + y) y^5}.$$

Every pair of denominators on the right is linearly independent. The preceding tensor construction therefore defines the right side as a distribution. The formulas agree on the complement of the divisors. They also agree near every smooth point of a divisor under the stated Laurent prescription. A difference between two such extensions is supported at the origin. Both satisfy the displayed complex scaling. Lemma 4.3 below makes their difference zero. $\square$

Lemma 4.3 (Complex scaling excludes supported ambiguity). *Let $d \geq 1$ and $m \in \mathbb{Z}$. A distribution S on $\mathbb{C}^d$, supported at zero, cannot satisfy $S(tz) = t^{-m}S(z)$ for every $t \in \mathbb{C}^\times$, unless $S = 0$.*

Proof. A distribution supported at zero is a finite sum $\sum_{\alpha, \beta} s_{\alpha\beta} \partial_z^\alpha \partial_{\bar{z}}^\beta \delta_0$. To obtain this description, use its finite order on a compact neighborhood. Taylor subtraction shows that it vanishes on test functions with sufficiently many zero derivatives at zero. It therefore factors through that finite jet space, whose dual consists of the displayed derivatives.

Write $a = |\alpha|$ and $b = |\beta|$. Under $t = r > 0$, a displayed derivative scales by $r^{-2d-a-b}$. Under $t = e^{i\theta}$, it scales by $e^{-i(a-b)\theta}$. Linear independence of these radial and angular characters requires

$$2d + a + b = m, \quad a - b = m$$

for every nonzero coefficient. Subtraction gives $2d + 2b = 0$, which is impossible. Hence every coefficient vanishes. $\square$

Real scaling alone would permit second derivatives of δ_0 in the triangle example. Their angular weights fail the required weight -6 . The exact complex scaling is therefore a substantive extension condition. The general scaling-degree extension theorem and its supported ambiguity are given by Brunetti–Fredenhagen [3, Theorems 5.2–5.3, pp. 23–24]. The proof above also excludes ambiguities at higher pole orders in this holomorphic setting.

4.3 ALL RATIONAL COEFFICIENTS AT THREE INSERTIONS

Let

$$R = \mathbb{C}[x, y, x^{-1}, y^{-1}, (x + y)^{-1}], \quad R_3 = \mathbb{C}[c] \otimes R.$$

Grade R by simultaneous complex dilation of x, y . Every element is a finite sum of homogeneous rational functions.

Theorem 4.4 (Three-insertion distributional extension). *There is a unique complex linear injection*

$$\mathcal{E}_3 : R_3 \longrightarrow \mathcal{D}'(\mathbb{C}^3)$$

with these properties. It extends each rational function away from its collision divisors. It is polynomial linear in c . Near a smooth point of a divisor it uses the transverse Laurent extension $z^{-n} \mapsto E_n(z)$. On each homogeneous relative-coordinate component of degree $-m$, it has exact complex scaling t^{-m} .

It commutes with holomorphic derivatives, multiplication by holomorphic polynomials, simultaneous translations, and permutations of the insertions. On products with independent collision normals it agrees with the tensor product construction. Formula (4.6) is its value on the triangle coefficient.

Proof. It is enough to construct the extension of a homogeneous summand. For positive integers a, b and $c_0 \geq 0$, the identity

$$\frac{1}{x^a y^b (x + y)^{c_0}} = \frac{1}{x^{a-1} y^b (x + y)^{c_0+1}} + \frac{1}{x^a y^{b-1} (x + y)^{c_0+1}}$$

reduces $a + b$ by one in each term. Repetition expresses the denominator as a finite sum involving at most two independent linear factors. Polynomial numerators can be multiplied into these distributions. This constructs a finite sum of pullbacks of $E_p \otimes E_q$, possibly with smooth polynomial factors. The homogeneity is unchanged.

The construction has the prescribed Laurent extension away from the origin. To check this assertion, use the singular linear form as a local coordinate u . Every other denominator is holomorphic and invertible there. For a holomorphic multiplier h , divide its Taylor expansion as $h(u) = \sum_{j=0}^{n-1} h_j u^j + u^n h_n(u)$. The identities $u^j E_n(u) = E_{n-j}(u)$ for $j < n$, and $u^n E_n(u) = 1$, give exactly its Laurent expansion as a distribution. The coefficients can depend holomorphically on the other coordinate.

Two partial-fraction decompositions therefore differ by a distribution supported at $(x, y) = 0$. They have the same complex homogeneity. Lemma 4.3, with $d = 2$, proves equality. This also proves linearity within each homogeneous component and independence of polynomial numerator presentations. Sum the components and then extend polynomial linearly in c . Restriction to the dense complement of the divisors proves injectivity.

A holomorphic derivative of the constructed extension has the Laurent prescription of the differentiated rational function. Its homogeneity is the differentiated degree. The same uniqueness argument proves commutation with relative holomorphic derivatives. The argument also applies to multiplication by homogeneous polynomials. Linearity gives all polynomial multipliers and all holomorphic derivatives, including c .

For a rational function of differences, every permutation acts by an invertible complex linear map of (x, y) preserving the three divisors. Pullback preserves both the Laurent prescription and complex homogeneity, so uniqueness proves covariance. Polynomial multiplication then gives covariance on R_3 , since a permuted c equals c plus a linear combination of x, y . Simultaneous translation changes only c . The transverse tensor prescription has the same defining properties, which proves the remaining assertion. $\square$

The image $\mathcal{R}_3 = \mathcal{E}_3(R_3)$ carries the specified product

$$\mathcal{E}_3(f) \circ \mathcal{E}_3(g) = \mathcal{E}_3(fg). \quad (4.7)$$

Injectivity makes this a well-defined associative algebra over $\mathbb{C}$. This product is a prescription on $\mathcal{R}_3$. It does not assert multiplication of arbitrary distributions. In particular, adjoining the residue currents requires further product definitions.

If Γ is the union of the conormal bundles of the three pair diagonals and the full conormal bundle of the total diagonal, then

$$\mathrm{WF}(\mathcal{E}_3(f)) \subseteq \Gamma \setminus \{\circ\}.$$

The bound follows termwise from the transverse tensor formula. At a point where only one divisor is present, only its normal covectors occur. At the total diagonal all relative covectors are allowed. No covector in the simultaneous translation direction occurs.

4.4 CONTACT TERMS AND THREE-CURRENT COMPATIBILITY

Pull back R_n with its displayed $d\bar{z}$ factor under each linear coordinate map. Distributional differentiation gives the explicit triangle contact current

$$\begin{aligned} \bar{\partial}T_\Delta &= R_2(x)E_4(y) + E_2(x)R_4(y) \\ &\quad + R_2(x+y)E_4(y) + E_2(x+y)R_4(y) \\ &\quad - 2R_1(x)E_5(y) - 2E_1(x)R_5(y) \\ &\quad + 2R_1(x+y)E_5(y) + 2E_1(x+y)R_5(y). \end{aligned} \tag{4.8}$$

Every term is a transverse tensor product. Thus the equation includes the total collision and does not require multiplication on one divisor. For independent u, v ,

$$\bar{\partial}(R_a(u)E_b(v) + E_a(u)R_b(v)) = -R_a(u) \wedge R_b(v) + R_a(u) \wedge R_b(v) = \circ.$$

Consequently (4.8) is closed, including its contact terms on the total diagonal.

To obtain actual compact observable outputs, choose $0 < r < \rho < R$, put $A = D_r$, $U = D_R$, and choose a smooth radial cutoff χ . Require $\chi = 1$ on D_ρ and $\mathrm{supp} \chi \subseteq D_R$. For $a \in A$, define

$$e_a(z) = -\frac{\bar{\partial}_z \chi(z)}{z - a}, \quad e_{a,n}(z) = \partial_a^n e_a(z).$$

These are compactly supported smooth $(0, 1)$ -forms on a fixed outer annulus. They depend holomorphically on a in every smooth seminorm. No support shrinks to a point as a moves.

Proposition 4.5 (Compactly smeared collision products). *For three elementary insertions, the expression*

$$\mathcal{W}_3(a, b, c) = e_a e_b e_c + \lambda(E_2(a - b)e_c + E_2(a - c)e_b + E_2(b - c)e_a) \tag{4.9}$$

defines a continuous linear map

$$C_c^\infty(\mathcal{A}^3, \mathrm{Dens})[\hbar] \longrightarrow \mathcal{B}_\kappa(U)^\circ, \quad F \longmapsto \langle \mathcal{W}_3, F \rangle.$$

Its values are D -closed compact smooth observable kernels. More generally, the three-insertion Wick formula for any finite monomials in the jets $e_{a,n}$ defines such a map by applying $\mathcal{E}_3$ to each rational contraction coefficient. The field number of the output is finite.

For the elementary product, the external Dolbeault derivative is

$$\bar{\partial}_{a,b,c} \mathcal{W}_3 = \lambda(R_2(a - b)e_c + R_2(a - c)e_b + R_2(b - c)e_a). \tag{4.10}$$

Here, for example, $R_2(a - b) = -\pi \partial_{a-b} \bar{\partial}(a - b)(d\bar{a} - d\bar{b})$. The two parenthesizations of the complete three-insertion Wick product have zero difference as distributions under $\mathcal{E}_3$.

Proof. A product of the e forms is a smooth kernel on a fixed product of compact annuli. For each compact parameter support, all parameter and field derivatives are bounded in every required seminorm. Each coefficient distribution has finite order on this support. Applying it to the parameter variables therefore gives a smooth kernel in the remaining field variables, with the same annular support. For example, a bound of distribution order N gives

$$\|\langle T, Fe \rangle\|_{C^k} \leq C_{T,K,k} \|F\|_{C^N} \sup_K \|e\|_{C^{N+k}}.$$

The same estimate applies to finitely many field factors. It proves continuity on every fixed compact support, hence on the test function direct limit. This construction also handles the holomorphic smooth coefficients multiplying the rational Wick coefficients. There are finitely many matchings, so the sum has finite field number.

Every output factor has antiholomorphic degree one. Its internal Dolbeault derivative is zero, and the quantum cocycle vanishes on pairs of such factors. Thus D kills the output smooth kernels. This uses the fixed differential (4.4). Holomorphic dependence of the e forms and Lemma 4.1 give (4.10).

For any polynomial inputs, label each individual current factor. A Wick matching pairs factors in different insertion groups. Grouping first the first two groups, or first the last two, enumerates the same matchings with the same coefficients. Therefore the complete rational expressions agree before extension. Linearity of $\mathcal{E}_3$ preserves this equality. Multiplication by the same smooth unpaired kernels and smearing preserve it as well. $\square$

For separated centers, choose disjoint inner cutoff disks and set $u_a = -\bar{\partial}\chi_a/(z-a)$. The compact quantum homotopy identifies the product $u_a u_b u_c$ with (4.9) in cohomology. This gives the separated boundary comparison with the fixed sign $\lambda = -\hbar\kappa$. At collisions, Proposition 4.5 extends its smooth annular output as an insertion-parameter distribution. It does not identify a Dirac current with a smooth observable generator. Equation (4.10) specifies the contact obstruction to regarding this extension alone as a Dolbeault chain map.

The regular vertex product has a different three-point identity. With $q_{n+1} = [-\bar{\partial}\chi/z^{n+1}]$, the field on polynomial states is

$$J(z) = \sum_{m \geq 1} q_m z^{m-1} + \lambda \sum_{m \geq 1} m \partial_{q_m} z^{-m-1}.$$

The regular product $a_{(-1)}b$ is the coefficient of z^0 in $Y(a, z)b$. Direct normal ordering gives

$$\begin{aligned} q_{1(-1)}q_1 &= q_1^2, & q_{1(-1)}q_1^2 &= q_1^3, \\ Y(q_1^2, z)q_1 &= J_-(z)^2 q_1 + 2\lambda z^{-2} J_-(z). \end{aligned}$$

The constant coefficient of the last expression is $q_1^3 + 2\lambda q_3$. Consequently its first associativity defect is

$$(q_{1(-1)}q_1)_{(-1)}q_1 - q_{1(-1)}(q_{1(-1)}q_1) = 2\lambda q_3 = -2\hbar\kappa q_3. \quad (4.11)$$

A rational coefficient extension on the insertion space does not identify iterated Laurent constant terms with multiplication at a point. Thus (4.11) is compatible with the zero grouping defect in Proposition 4.5.

4.5 SMOOTH INTERVAL COEFFICIENTS AND THE REMAINING CHAIN CONDITION

Let $I = [0, 1]$ have its increasing orientation. Use the full smooth de Rham complex $\Omega^\bullet(I) = C^\infty(I) \oplus C^\infty(I) dt$, where smooth functions extend to a neighborhood of the closed interval. There are no endpoint vanishing conditions. The finite rational Wick coefficients in three insertion variables form $R_3[x_{i,n}][\hbar]$, with each polynomial involving finitely many jets. The distributional coefficient map is

$$1 \otimes \mathcal{E}_3 : \Omega^\bullet(I) \otimes R_3[x_{i,n}][\hbar] \longrightarrow \Omega^\bullet(I) \otimes \mathcal{R}_3[x_{i,n}][\hbar].$$

For jointly smooth interval dependence, apply $\mathcal{E}_3$ to each fixed rational coefficient and retain its full smooth interval amplitude. The finite-order estimate in Proposition 4.5 permits interval derivatives, endpoint evaluation, and integration to pass through the parameter distribution. The map therefore commutes with d_I . It preserves every three-insertion identity formed from interval-linear maps and the prescribed rational Wick product. This statement includes all smooth interval amplitudes, independently of any polynomial subcomplex.

All three-insertion formulas extend coefficientwise to $\hbar$ -adic series whose individual coefficients have finite polynomial degree and finitely many jets. A fixed output coefficient receives finitely many input coefficients and matching terms. No uniform distribution order or field number across the entire formal series is asserted.

If a current convention uses $+\hbar\kappa_I(z-w)^{-2}$, the boundary comparison requires $\kappa_I = -\kappa$. All jet contractions then become $\lambda\partial_{z_i}^r\partial_{z_j}^sE_2(z_i-z_j)$ with the same undivided-jet convention. Divided jets add the factors $(r!s!)^{-1}$.

On insertion-parameter forms with observable values, the total differential is

$$\mathbb{D}(\omega \otimes b) = \bar{\partial}\omega \otimes b + (-1)^{|\omega|}\omega \otimes Db.$$

It squares to zero. Formula (4.10) is the value of $\mathbb{D}$ on the elementary extended product. The corresponding triangle contact is (4.8). To construct operations on collision chains, one must supply maps for these residue currents and prove compatibility with their boundary strata. The associative product (4.7) does not itself supply products involving these (0, 1)-currents. Nor does the three-insertion construction specify extensions for four or more insertion variables. These are separate mathematical obligations.

Residue traces and three-current chains

The double pole contributes a derivative of the test function on a pair diagonal. That derivative must enter the source differential of a current comparison. For three elementary currents, a pair contraction leaves one current. The three pair diagonals therefore supply a finite complex and an explicit map to compact boundary observables. The regular vertex product has a separate obstruction: its associator survives in boundary cohomology.

5.1 THE DIFFERENTIAL ON TEST CHAINS

Fix $0 < r < \rho < R$, put $A = D_r$, $U = D_R$, and let χ be a smooth radial function supported compactly in U , equal to one on D_ρ . Use the complex orientation, $dA = dx dy$, and $\int^{\text{res}} = (2\pi i)^{-1} \int$. Work over $\mathbb{C}[\hbar]$, with fixed $\kappa \in \mathbb{C}$ and $\lambda = -\hbar\kappa$ of degree zero. The target is the actual smooth boundary complex

$$B = \mathcal{B}_\kappa(U), \quad D = \bar{\partial} + \hbar\Delta_{\mu_\kappa}, \quad \mu_\kappa(f, v) = -\kappa \int_U^{\text{res}} f \partial v.$$

A function generator has degree -1 and a form generator has degree zero. Each symmetric power uses compact smooth kernels and graded coinvariants. The direct sum in field number remains algebraic. Set

$$e_{a,n} = \partial_a^n \left(-\frac{\bar{\partial}\chi}{z-a} \right), \quad e_a = e_{a,0}, \quad S = \text{supp } \bar{\partial}\chi.$$

These forms have support in $S \subseteq U$ and depend holomorphically on $a \in A$ in every smooth seminorm. For three elementary insertions, fix the distribution

$$W = e_{z_1} e_{z_2} e_{z_3} + \lambda \sum_{i < j} E_2(z_i - z_j) e_{z_k}, \quad \{k\} = \{1, 2, 3\} \setminus \{i, j\}.$$

Here $E_2 = -\partial_z(1/z)$ acts on test densities, as in Theorem 4.4. The products of form generators are symmetric smooth kernels in the field variables. The insertion variables remain distinct from those field variables.

For an open $Y \subset \mathbb{C}^d$ with the displayed affine coordinates, define the complex of compact Dolbeault test chains by

$$K_Y^{-j} = C_c^\infty(Y, \text{Dens}_{\mathbb{C}} \otimes \Lambda^j T^{0,1} Y)[\hbar], \quad \partial_Y = - \sum_{a=1}^d \partial_{\bar{z}_a} \iota_{d\bar{z}_a}. \quad (5.1)$$

The index satisfies $0 \leq j \leq d$. Densities are complexified real densities, with coefficient derivatives taken in the coordinate area density. Exterior contraction acts on the multivector factor. The differential has degree one, and $\delta_Y^2 = 0$ because coefficient derivatives commute while exterior contractions anticommute. Use the test-function direct-limit topology on each degree. The distributional pairing satisfies

$$\langle \delta T, \alpha \rangle = \langle T, \delta_Y \alpha \rangle$$

for a distribution-valued antiholomorphic form T and a test chain of the matching exterior degree. This equality fixes every transpose sign.

Put $X = A^3$. For $i < j$, let $Z_{ij} = \{z_i = z_j\} \cap X$ and fix coordinates

$$x = z_i - z_j, \quad s = z_j, \quad w = z_k.$$

The real Jacobian has absolute determinant one. On Z_{ij} use coordinates (s, w) and density $dA_s dA_w$. In these coordinates define a degree-one residue trace

$$r_{ij} : K_X \longrightarrow K_{Z_{ij}}, \quad r_{ij} \alpha = \pi \partial_x \iota_{d\bar{x}} \alpha|_{x=0}. \quad (5.2)$$

The formula removes the normal density factor dA_x after differentiation and restriction. Exterior factors remaining after contraction are tangent to Z_{ij} . In particular r_{ij} vanishes on degree zero.

Lemma 5.1 (Residue trace and Dolbeault differential). *The traces are continuous on fixed compact supports, preserve compactness, and satisfy*

$$\delta_{Z_{ij}} r_{ij} + r_{ij} \delta_X = 0.$$

For $\alpha \in K_X^{-1}$,

$$\langle W, \delta_X \alpha \rangle = \lambda \sum_{i < j} \langle e_w, r_{ij} \alpha \rangle. \quad (5.3)$$

The right side integrates both variables of Z_{ij} .

Proof. Differentiate a compact test coefficient once in x and restrict to $x = 0$. The result has support in the intersection of its compact support with Z_{ij} , and its seminorms require one additional derivative. Write $\iota_x = \iota_{d\bar{x}}$. The normal term of $r_{ij} \delta_X$ vanishes by $\iota_x^2 = 0$. For a tangential coordinate $v = s, w$, one has $\iota_x \iota_{d\bar{v}} = -\iota_{d\bar{v}} \iota_x$. Tangential derivatives commute with ∂_x and restriction. This proves the anticommutation identity.

The normalized residue is $\bar{\delta} E_2(x) = -\pi \partial_x \partial_0(x) d\bar{x}$. Its action on the normal test coefficient is $\pi \partial_x \alpha_x(0, s, w)$. The surviving form e_w is independent of x and holomorphic in w . Consequently differentiation of W gives precisely the right side of (5.3). The uncontracted term is holomorphic in all three insertion variables and contributes zero. $\square$

5.2 A COMPARISON INCLUDING THE PAIR DIAGONALS

On the direct sum of test-chain complexes define

$$K_{\text{res}} = K_X \oplus \bigoplus_{i < j} K_{Z_{ij}}, \quad (5.4)$$

$$d_{\text{res}}(\alpha, (\beta_{ij})) = \left(\partial_X \alpha, (\delta_{Z_{ij}} \beta_{ij} - \lambda r_{ij} \alpha)_{i < j} \right).$$

Each diagonal summand records a contracted pair and its surviving current. It has the same test-chain grading as its geometric dimension prescribes. The quotient by the diagonal summands is the original complex K_X . The inclusion of K_X as a graded summand is not a chain map when λr_{ij} is nonzero.

Define $F : K_{\text{res}} \rightarrow B$ to be zero in negative degrees and to have degree-zero component

$$F(\alpha, (\beta_{ij})) = \langle W, \alpha \rangle + \sum_{i < j} \langle e_w, \beta_{ij} \rangle. \quad (5.5)$$

All test coefficients in this equation are smooth compact densities.

Theorem 5.2 (Three-current residue comparison). *The differential in (5.4) has square zero, and $DF = Fd_{\text{res}}$. The image consists of compact smooth observable kernels supported in S and S^3 , together with their finite linear sums. The maps are continuous on fixed compact test supports. Their test supports may meet every pair diagonal and the total diagonal.*

Let $A_I = \mathbb{C}[t, dt]$ with $d_I t = dt$, $(dt)^2 = 0$, and the increasing orientation on $[0, 1]$. With the form factor first, the map

$$\mathfrak{F} = 1 \otimes F : (A_I \otimes K_{\text{res}}, d_I + d_{\text{res}}) \longrightarrow (A_I \otimes B, d_I + D)$$

is a chain map. In each total differential the second summand acts with the sign $(-1)^{|\gamma|}$ on $\gamma \otimes a$. Endpoint evaluation gives the chain map F on constant interval forms.

Proof. The diagonal component of d_{res}^2 is $-\lambda(\partial_Z r + r\partial_X)\alpha = 0$ by Lemma 5.1. The other terms vanish by $\partial_X^2 = \partial_Z^2 = 0$. Thus the differential includes its entire degree-two compatibility. No additional contraction is possible after pairing two of three elementary currents: the contracted pair is scalar and only one field remains. This is why this complex needs no second contraction term.

Every output in (5.5) has only form generators. Both its internal Dolbeault differential and its quantum contraction vanish. Thus $DF = 0$. In degree -1 , the contact identity gives

$$Fd_{\text{res}}(\alpha, \beta) = \lambda \sum_{i < j} \langle e_w, r_{ij} \alpha \rangle + \sum_{i < j} \langle e_w, \partial_Z \beta_{ij} \rangle - \lambda \sum_{i < j} \langle e_w, r_{ij} \alpha \rangle = 0.$$

The middle sum vanishes by holomorphic dependence and integration by parts. In lower degrees, the target component of F is zero on both sides. In degree zero, the source differential vanishes. This proves the chain equation in every degree.

The estimate $|z - a| \geq \rho - r$ on the field annulus gives uniform smooth bounds for the unpaired forms. Each coefficient distribution has finite order on a compact insertion support. Pairing therefore produces a smooth field kernel with support in the same annulus product, as in Proposition 4.5. The trace terms have the analogous estimate with one normal derivative. No parameter support excludes the total diagonal. Finally tensoring two cochain maps, with the stated sign, proves the interval equation. Endpoint evaluation kills dt and commutes with the total differential. $\square$

The same construction retains every holomorphic jet of each elementary current. For fixed $n_1, n_2, n_3 \geq 0$, replace $\mathcal{W}$ by

$$W_{\mathbf{n}} = \prod_{i=1}^3 e_{z_i, n_i} + \lambda \sum_{i < j} (-1)^{n_i} (n_i + n_j + 1)! E_{n_i + n_j + 2}(z_i - z_j) e_{z_k, n_k}.$$

Replace the trace by

$$r_{ij}^{\mathbf{n}} \alpha = \pi(-1)^{n_i} \partial_x^{n_i + n_j + 1} \iota_{d\tilde{x}} \alpha \Big|_{x=0}.$$

The action of $\bar{\partial} E_m = \pi(-1)^{m-1} \partial_x^{m-1} \bar{\partial}_0 d\tilde{x} / (m-1)!$ cancels both factorials and transpose signs, giving this trace. It anticommutes with $\bar{\partial}$ by the same exterior calculation. Thus the theorem holds for the direct sum over all finite jet choices. For divided jets, use $e_{z_i, n_i} / n_i!$ for each uncontracted input. Divide $r_{ij}^{\mathbf{n}}$ by

$n_i!n_j!$, and use $e_{w,n_k}/n_k!$ in the diagonal output. The complete degree-zero Wick distribution is then $W_{\mathbf{n}}/(n_1!n_2!n_3!)$.

This source specifies elementary three-current contractions and their jets. For three composite insertions, two contractions can survive, and three quadratic insertions contain the triangle coefficient $8\lambda^3 T_{\Delta}$. The derivative of its distributional extension is already defined by (4.8). A source for those composite collision chains also requires maps between its contraction strata. The complex (5.4) does not supply those additional maps.

5.3 THE MAXIMAL TEST SUBCOMPLEX FOR A FIXED WICK SMEARING

The same contact condition has an exact formulation for all finite polynomial insertions, without choosing products of residue currents. Let $V = \mathbb{C}[q_1, q_2, \dots][\hbar]$ be the normal polynomial state space, and put $M = V \otimes_{\mathbb{C}[\hbar]} V \otimes_{\mathbb{C}[\hbar]} V$. All states have degree zero, and every state has finite polynomial degree. Let W_M be the complete three-insertion Wick distribution with these inputs, applying $\mathcal{E}_3$ to each rational coefficient and using the same annular forms for the unpaired fields. Specifically, the divided state q_{n+1} at a uses $e_{a,n}/n!$. This is precisely the map of Proposition 4.5. Put $K = K_X \otimes_{\mathbb{C}[\hbar]} M$, and define

$$\Phi : K^{\circ} \rightarrow B^{\circ}, \quad \Phi(\alpha) = \langle W_M, \alpha \rangle, \quad C : K^{-1} \rightarrow B^{\circ}, \quad C(\alpha) = \Phi(\delta \alpha).$$

Equivalently C pairs δW_M with α . Since $\delta^2 = 0$, one has $C\delta = 0$ on K^{-2} .

Lemma 5.3 (Detection of annular boundaries). *Suppose $b \in B^{\circ}$ is a finite sum of outputs of Φ or C . If $b = Dq$ for some $q \in B^{-1}$, then $b = 0$ as a smooth observable. The same conclusion holds if $1 \otimes b$ is exact in $A_I \otimes B$.*

Proof. Let $u_m = -\delta \chi / z^{m+1}$ for $m \geq 0$. On the annulus,

$$e_a = \sum_{m \geq 0} a^m u_m.$$

Choose r_0 strictly above the radii of a compact parameter support, with $0 < r_0 < r$, and choose σ between r and ρ . For a multi-index $\mathbf{m}$, distributional finite order and the annular denominator give bounds

$$|c_{\mathbf{m}}| \leq C(1 + |\mathbf{m}|)^N r_0^{|\mathbf{m}|}, \quad \|u_{m_1} \cdots u_{m_s}\|_{C^k} \leq C_k(1 + |\mathbf{m}|)^{N_k} \sigma^{-|\mathbf{m}|}.$$

Jets add a fixed power of $|\mathbf{m}|$. The ratio $r_0/\sigma < 1$ proves convergence after every smooth derivative. In field number s , each output therefore has a convergent symmetric multiple series in $u_{m_1} \cdots u_{m_s}$.

The Cauchy pairing of two u_m vanishes by rotation invariance. Continuity then implies $N_U b = b$, where $dN_U = N_U D$ is Proposition 1.5. For $L \geq 0$, define the classical moment map to the polynomial algebra $\mathbb{C}[\gamma_0, \dots, \gamma_L][\hbar]$ by

$$R_L(v) = \sum_{m=0}^L \left(\int_U^{\text{res}} v \wedge z^m dz \right) \gamma_m, \quad R_L(f) = 0.$$

On each smooth symmetric kernel, integrate the indicated moment in every field variable. Holomorphicity and compact integration by parts give $R_L d = 0$. Hence $L_L = R_L N_U$ satisfies $L_L D = 0$. The residue moments obey $R_L(u_m) = \gamma_m$ for $m \leq L$ and vanish otherwise. Thus the family R_L detects every coefficient of the convergent symmetric series. If $b = Dq$, then $R_L b = L_L Dq = 0$ for every L . Every coefficient is zero, so the smooth kernel is zero in each field number. Only finitely many field numbers occur. For the interval assertion, evaluate an exact expression at either endpoint, obtaining $b = D(\text{ev}_a q)$, and apply the first part. $\square$

Define a subcomplex $K_{\ker} \subseteq K$ by

$$K_{\ker}^0 = K^0, \quad K_{\ker}^{-1} = \ker C, \quad K_{\ker}^{-j} = K^{-j} \quad (j = 2, 3).$$

The equation $C\delta = 0$ proves that the displayed spaces form a subcomplex. Equip the kernel with the induced test-chain topology.

Theorem 5.4 (Maximal domain with unchanged test differential). *The map Φ , extended by zero in negative degrees, is a continuous chain map $K_{\ker} \rightarrow B$. If a subcomplex $L \subseteq K$ contains K^0 and a chain map $G : L \rightarrow A_I \otimes B$ satisfies $G|_{K^0} = 1 \otimes \Phi$, then $L \subseteq K_{\ker}$. The same conclusion holds if this degree-zero equality is changed by an exact observable for each input. Thus $K_{\ker}$ is the largest subcomplex with this degree-zero smearing and the unchanged test differential, even when negative-degree comparison maps are allowed.*

Proof. Every Φ output is a D -cycle. On degree -1 , the chain equation is precisely $\Phi\delta = C = 0$. Lower degrees contribute zero, proving the first assertion. Continuity follows from the fixed-support finite order estimates for $\mathcal{E}_3$ and the annular kernels. For $\alpha \in L^{-1}$, the chain equation for G implies that $1 \otimes C\alpha$ is exact. The same implication holds if the degree-zero map is changed by an exact observable. Lemma 5.3 gives $C\alpha = 0$. Thus $L^{-1} \subseteq \ker C$. All other degree inclusions follow from the definition of $K_{\ker}$. $\square$

The restriction is strict at nonzero level. Choose a small coordinate box around $x = 0, s = s_0, w = w_0$, with $s_0 \neq w_0$. Require its closure to lie in X and avoid both other pair diagonals. Choose compact smooth functions $\theta(x)$ and $\psi(s, w)$ there, with $\theta = 1$ near zero and $\int \psi dA_s dA_w = 1$. For three elementary currents take

$$\alpha = x\theta(x)\psi(s, w) dA_x dA_s dA_w \otimes \partial_{\bar{x}}.$$

The only residue is $r_{12}\alpha = \pi\psi dA_s dA_w$. Consequently $C\alpha = \pi\lambda \int \psi(s, w)e_w dA_s dA_w$. Its first constant holomorphic moment is $\pi\lambda \neq 0$ when $\kappa \neq 0$. Thus neither a negative-degree boundary observable nor an interval primitive can extend the smearing to this test chain. Tests supported off the pair diagonals have zero contact and belong to $K_{\ker}$ in every degree.

5.4 THE REGULAR-PRODUCT ASSOCIATOR

The preceding source differential carries a contact to a surviving current. An ordinary ternary homotopy with three closed state inputs has a different equation. Let $\iota : V \rightarrow B$ send q_{m+1} to u_m . The regular vertex product is the constant Laurent coefficient of $Y(a, z)b$. Its elementary calculation gives

$$(q_1(-1)q_1)(-1)q_1 - q_1(-1)(q_1(-1)q_1) = 2\lambda q_3.$$

Indeed $J_-(z) = q_1 + q_2z + q_3z^2 + \dots$, and $:J(z)^2: q_1 = J_-(z)^2 q_1 + 2\lambda z^{-2} J_-(z)$. The constant coefficient is $q_1^3 + 2\lambda q_3$. The other parenthesization gives q_1^3 .

Proposition 5.5 (Nonexistence of a ternary boundary primitive). *If $\kappa \neq 0$, there is no $h \in (A_I \otimes B)^{-1}$ satisfying*

$$(d_I + D)h = 1 \otimes 2\lambda\iota(q_3). \quad (5.6)$$

In particular, an uncurved A_∞ algebra on $C_I \otimes V$ with unary operation $d_I \otimes 1$ cannot have the regular vertex product as its binary operation on constant interval cochains at nonzero level. The same obstruction excludes an A_∞ structure on a boundary resolution whose induced binary product on the injected states is this regular product.

Proof. Evaluate (5.6) at $t = 0$ and apply $L_2 = R_2 N_U$ from Lemma 5.3, where the subscript here denotes the moment bound. The left side becomes zero. The right side becomes $2\lambda y_2$, which is nonzero over $\mathbb{C}[h]$ when $\kappa \neq 0$. Equivalently the z^2 moment of $\iota(q_3) = u_2$ equals one.

For three closed degree-zero inputs, the arity-three A_∞ identity states that the binary associator is the differential of a degree-minus-one ternary value. The unary input terms vanish. Take the three constant interval inputs $(e_0 + e_1)q_1$. Their regular-product associator is $(e_0 + e_1)2\lambda q_3$. It is nonzero in $H^0(C_I \otimes V) = V$. Thus the identity cannot hold. On any boundary resolution with the specified injected cohomology classes, the same identity would make its cohomological product associative, contradicting the detected associator. This is the arity-three consequence of the suspended bar equations in Keller [1, §3.6, pp. 11–12]. $\square$

For fixed disjoint disk unions, the ordered Taylor maps of Theorem 3.3 retain the target $A_I \otimes \mathcal{B}_\kappa(U_S)$ and the complete Wick product. Their nonzero higher coefficients do not supply a primitive for (5.6). Endpoint evaluation kills their coefficients of arity greater than one. The residue comparison instead changes the source differential by its explicit diagonal terms. Extending it to composite collision strata and relating those strata to iterated regular Laurent coefficients remains a separate construction. The nonzero class $2\lambda q_3$ must be retained in that comparison.

Bibliography

- [1] B. Keller, *Introduction to A -infinity algebras and modules*, Homology, Homotopy and Applications **3** (2001), 1–35. <https://arxiv.org/abs/math/9910179>.
- [2] C. Brouder, N. V. Dang, and F. Hélein, *Continuity of the fundamental operations on distributions having a specified wave front set*, arXiv:1409.7662, 2016 revision, Theorem 6.1, p. 21. <https://arxiv.org/abs/1409.7662v2>.
- [3] R. Brunetti and K. Fredenhagen, *Microlocal analysis and interacting quantum field theories: Renormalization on physical backgrounds*, arXiv:math-ph/9903028, 1999, Theorems 5.2–5.3, pp. 23–24. <https://arxiv.org/abs/math-ph/9903028v1>.