

BV-BRST FORMALISM FOR CY CATEGORIES AND THE BAR COMPLEX

RAEEZ LORGAT

ABSTRACT. We develop the BV-BRST interpretation of the bar complex of a quantum chiral algebra arising from a Calabi–Yau category. The A_∞ -operations of the CY category define an open string field theory in the Batalin–Vilkovisky formalism; the CY pairing provides the BV bracket; the quantum master equation is identified with the Maurer–Cartan equation for Θ_A from Volume I. The bar complex $B(A)$ is the BV-BRST complex, with the bar differential equal to Q_{BRST} . We explain how the genus- g curvature $d_B^2 = \kappa \cdot \omega_g$ is the BRST anomaly, and how anomaly cancellation ($\kappa = 0$) corresponds to an “untwisted” BKM algebra. The Swiss-cheese structure $\text{SC}^{\text{ch}, \text{top}}$ of Volume II is identified with the BV structure on the open-closed string field theory. For CY3 manifolds X , the full BV-BRST complex of the topological B-model is the bar complex of the quantum vertex chiral group $G(X)$.

CONTENTS

1. INTRODUCTION AND MOTIVATION

The Batalin–Vilkovisky formalism is the natural framework for the quantization of gauge theories with open symmetry algebras, and string field theory is its most prominent application. The purpose of this note is to make precise the relationship between the BV-BRST complex of (open and closed) topological string field theory and the bar complex $B(A)$ of the quantum chiral algebra $A = A_{\mathcal{C}}$ associated to a Calabi–Yau category $\mathcal{C}$.

The key identifications are:

- (1) The fields of the open string field theory on a brane $\mathcal{F} \in \text{Ob}(\mathcal{C})$ are $\text{RHom}(\mathcal{F}, \mathcal{F})[1]$, with BV action built from the A_∞ -operations $\{\mu_n\}$ via the CY trace.
- (2) The BV bracket $\{-, -\}$ on the field space arises from the non-degenerate CY pairing $\langle \cdot, \cdot \rangle: A \otimes A \rightarrow k[-d]$.
- (3) The quantum master equation $\hbar \Delta S + \frac{1}{2}\{S, S\} = 0$ is, in the language of Volume I, the Maurer–Cartan equation for the universal MC element Θ_A .
- (4) The bar differential d_B of $B(A)$ is the BRST operator Q_{BRST} ; the failure of $d_B^2 = 0$ at genus $g \geq 1$ is the BRST anomaly, with $d_B^2 = \kappa(A) \cdot \omega_g$ the anomaly proportional to the modular characteristic.
- (5) The Swiss-cheese structure $\text{SC}^{\text{ch}, \text{top}}$ on $B(A)$ from Volume II is the BV structure on the combined open-closed string field theory.
- (6) For a CY3 manifold X , the BV-BRST complex of the topological B-model is the bar complex of the quantum vertex chiral group $G(X)$.

Remark 1.1 (Conventions). All grading is cohomological (differentials have degree +1), matching Volumes I–II. The BV antibracket has degree +1 in our convention. We work over k of characteristic zero. The parameter $\hbar$ tracks genus: a genus- g amplitude carries weight $\hbar^{2g-2}$.

Date: April 2026.

2. OPEN STRING FIELD THEORY FROM CY CATEGORIES

2.1. The field space. Let $\mathcal{C}$ be a smooth Calabi–Yau category of dimension d (in the sense of Kontsevich: a smooth dg category with $\mathcal{C} \xrightarrow{\sim} \mathcal{C}^\vee[d]$), and let $\mathcal{F} \in \text{Ob}(\mathcal{C})$ be an object — a brane. The *open string field* is an element

$$(1) \quad \alpha \in \mathfrak{F} := \text{RHom}_{\mathcal{C}}(\mathcal{F}, \mathcal{F})[1],$$

the (derived) endomorphism algebra of the brane, shifted by one to place the physical gauge field (a 1-cochain in the unshifted complex) at ghost number zero. When $\mathcal{C} = D^b(\text{Coh}(X))$ for a CY manifold X , and $\mathcal{F} = \mathcal{O}_X$, this recovers $\mathfrak{F} = \bigoplus_p \text{Ext}^p(\mathcal{O}_X, \mathcal{O}_X)[1] = H^*(X, \mathcal{O}_X)[1]$.

The space $\mathfrak{F}$ is a cochain complex with differential μ_1 (the A_∞ -differential) and carries the full A_∞ -structure $\{\mu_n\}_{n \geq 1}$ inherited from $\mathcal{C}$.

2.2. The BV action. The *BV action* is the formal function on $\mathfrak{F}$:

$$(2) \quad S(\alpha) = \sum_{n \geq 2} \frac{1}{n} \text{Tr}(\mu_n(\alpha, \dots, \alpha))$$

where μ_n are the A_∞ -operations on $A := \text{End}_{\mathcal{C}}(\mathcal{F})$ and $\text{Tr}: A \rightarrow k[-d]$ is the CY trace (the image in Hochschild homology of the CY structure class $[\sigma] \in \text{HC}_d^-(\mathcal{C})$).

The action is a formal power series $S = S_2 + S_3 + \dots$ where $S_n = \frac{1}{n} \text{Tr}(\mu_n(\alpha^{\otimes n}))$. The individual terms have the following interpretation:

- $S_2 = \frac{1}{2} \text{Tr}(\mu_2(\alpha, \alpha))$: the kinetic/Chern–Simons term. When $\mathcal{C} = D^b(\text{Coh}(X))$ and μ_2 is the cup product, this is $\frac{1}{2} \int_X \text{Tr}(\alpha \wedge \bar{\partial} \alpha) \Omega$ (with Ω the holomorphic volume form).
- $S_3 = \frac{1}{3} \text{Tr}(\mu_3(\alpha, \alpha, \alpha))$: the cubic interaction. In the B-model, this is the $\frac{1}{3} \int \text{Tr}(\alpha \wedge \alpha \wedge \alpha) \Omega$ term.
- S_n for $n \geq 4$: the higher-order vertices, absent in the classical B-model but present in the A_∞ -deformation (arising from Massey products, instanton corrections in the A-model, or obstructions to formality).

Remark 2.1 (Relation to Witten’s open SFT). When $\mathcal{C}$ is the Fukaya category $\text{Fuk}(M)$ and μ_2 is the Floer product, the action (??) is Witten’s open string field theory action. The A_∞ -operations μ_n for $n \geq 3$ arise from counting holomorphic disks with $n+1$ boundary punctures, weighted by symplectic area. The formality of the A-model on a Stein manifold (no holomorphic disks) sets $\mu_n = 0$ for $n \geq 3$, reducing to a Chern–Simons functional.

Remark 2.2 (The shift and ghost number). The shift $[1]$ in (??) is the BV shift: it places the degree-1 component of $\text{End}(\mathcal{F})$ (i.e., the connection/gauge field) at ghost number 0. The degree-0 component becomes the ghost c at ghost number -1 ; the degree-2 component becomes the antifield α^* at ghost number $+1$; and so on. The full BV field space is the (-1) -shifted cotangent bundle $T^*[-1]\mathfrak{F}_{\text{cl}}$ where $\mathfrak{F}_{\text{cl}}$ is the classical field space, but the CY structure identifies $\mathfrak{F}$ with its own shifted cotangent bundle via the pairing $\langle \cdot, \cdot \rangle$.

2.3. The BV extension. The action (??) is the BV *extension* of the classical A_∞ -action in the following precise sense. Define the *classical action* by restricting to ghost-number-zero fields:

$$S_{\text{cl}}(\alpha_0) = \sum_{n \geq 2} \frac{1}{n} \text{Tr}(\mu_n(\alpha_0, \dots, \alpha_0)), \quad \alpha_0 \in \mathfrak{F}^0.$$

The classical equations of motion are $\sum_{n \geq 1} \mu_n(\alpha_0^{\otimes n}) = 0$, which is the Maurer–Cartan equation for α_0 in the A_∞ -algebra $(A, \{\mu_n\})$. The BV extension (??) extends S_{cl} to all ghost numbers and satisfies the classical master equation $\{S, S\} = 0$ if and only if the A_∞ -relations hold — which they do, by construction.

Proposition 2.3 (Classical master equation). *The BV action $S(\alpha)$ satisfies the classical master equation*

$$(3) \quad \{S, S\} = 0$$

where $\{-, -\}$ is the BV antibracket defined by the CY pairing (see §??). This is equivalent to the A_∞ -relations for $(\text{End}(\mathcal{F}), \{\mu_n\})$.

Proof. The classical master equation $\{S, S\} = 0$ expands, using the definition of the antibracket from the CY pairing, as:

$$\sum_{p+q=n+1} \text{Tr}(\mu_p(\alpha^{\otimes(p-1)}, \mu_q(\alpha^{\otimes q}))) = 0 \quad \text{for all } n \geq 1.$$

This is the A_∞ -relation $\sum_{p+q=n+1} \mu_p \circ (\text{id}^{\otimes(i-1)} \otimes \mu_q \otimes \text{id}^{\otimes(p-1-i)}) = 0$ contracted against $\alpha^{\otimes n}$ and traced. The CY pairing is non-degenerate, so the traced equation implies (and is implied by) the untraced A_∞ -relation. $\square$

3. THE BV BRACKET FROM THE CY PAIRING

3.1. The CY pairing as BV symplectic form. The CY structure on $\mathcal{C}$ provides a non-degenerate pairing

$$(4) \quad \langle \cdot, \cdot \rangle : A \otimes A \longrightarrow k[-d]$$

on $A = \text{End}_{\mathcal{C}}(\mathcal{F})$, satisfying cyclic invariance with respect to the A_∞ -operations. After the shift $A[1]$, this pairing becomes a degree- $(1-d)$ symplectic form ω on the field space $\mathfrak{F} = A[1]$:

$$(5) \quad \omega(\alpha, \beta) = \langle s^{-1}\alpha, s^{-1}\beta \rangle, \quad \deg(\omega) = 1 - d.$$

For d odd (e.g., $d = 3$, the CY3 case), this is an *even* symplectic form of degree -2 , which after further shifting provides the (-1) -shifted symplectic structure of the AKSZ formalism. For d even, we obtain an odd symplectic form; the degree -1 case is the standard BV symplectic structure.

Warning 3.1. The BV formalism requires a (-1) -shifted symplectic structure on the space of fields. For a d -dimensional CY category, the CY pairing on $A[1]$ has degree $(1-d)$. The standard BV case is $d = 2$: CY2 categories (K3 surfaces, symplectic surfaces) give a degree (-1) pairing on $A[1]$, which is exactly the BV symplectic form. For CY3 ($d = 3$), the pairing has degree (-2) ; the BV-BRST complex acquires an additional $\hbar$ -deformation (the “second quantization”), and the correct framework is Costello’s formulation of the B-model as a BV theory with a (-2) -shifted symplectic structure.

3.2. The BV antibracket. The BV antibracket (odd Poisson bracket) on functions on $\mathfrak{F}$ is defined by inverting ω :

$$(6) \quad \{F, G\} = \sum_{i,j} \frac{\partial F}{\partial \phi^i} \omega^{ij} \frac{\partial G}{\partial \phi^j}$$

where $\{\phi^i\}$ is a basis of $\mathfrak{F}$ and ω^{ij} is the inverse of $\omega_{ij} = \omega(\phi_i, \phi_j)$. The key properties:

- (i) $\{F, G\} = -(-1)^{(|F|+1)(|G|+1)}\{G, F\}$ (graded antisymmetry);
- (ii) $\{S, \{S, S\}\} = 0$ (graded Jacobi);
- (iii) $\deg\{F, G\} = \deg F + \deg G + 1$ (the antibracket has degree $+1$).

Remark 3.2 (Cyclic invariance = BV compatibility). The cyclic invariance of the CY pairing with respect to μ_n ,

$$\langle \mu_n(a_1, \dots, a_n), a_0 \rangle = \pm \langle \mu_n(a_0, a_1, \dots, a_{n-1}), a_n \rangle,$$

is precisely the statement that the action S is compatible with the BV bracket: $\{S, -\}$ is a derivation, i.e., $Q_{\text{BRST}} = \{S, -\}$ is a differential. Without cyclic invariance, one cannot construct the BV antibracket from the pairing — and without the BV antibracket, there is no BRST operator. The CY condition is the necessary and sufficient condition for the existence of the BV-BRST formalism.

3.3. The BV Laplacian. The *BV Laplacian* Δ is the second-order operator

$$(7) \quad \Delta F = \sum_{i,j} \omega^{ij} \frac{\partial^2 F}{\partial \phi^i \partial \phi^j}.$$

It satisfies $\Delta^2 = 0$ and is related to the antibracket by the BV identity:

$$\{F, G\} = (-1)^{|F|} (\Delta(FG) - (\Delta F)G - (-1)^{|F|} F(\Delta G)).$$

The BV Laplacian is the “second quantization” operator: it accounts for the contraction of a field with its antifield, i.e., the propagator loop. In the path integral, $e^{S/\hbar}$ is gauge-invariant if and only if S satisfies the *quantum master equation*.

4. THE QUANTUM MASTER EQUATION AND THE MC EQUATION FOR Θ_A

4.1. The QME. The quantum master equation (QME) for the BV action is:

$$(8) \quad \boxed{\hbar \Delta S + \frac{1}{2} \{S, S\} = 0}$$

The first term $\hbar \Delta S$ is the one-loop correction (the BV Laplacian acting on S); the second term is the classical master equation. The QME encodes the full consistency of the quantum gauge theory: it ensures that the path integral measure $e^{S/\hbar} [d\phi]$ is BRST-invariant to all orders in $\hbar$.

4.2. Identification with the MC equation. The central observation connecting the BV formalism to the chiral bar complex is:

Theorem 4.1 (QME = MC for Θ_A). *Let $A = A_{\mathcal{C}} = \Phi(\mathcal{C})$ be the quantum chiral algebra associated to a CY category $\mathcal{C}$ via the functor of Vol III. Let Θ_A be the universal Maurer–Cartan element of A (Vol I). Then the quantum master equation (??) for the BV action S of the open string field theory on any brane $\mathcal{F} \in \mathcal{C}$ is equivalent to the Maurer–Cartan equation*

$$(9) \quad d\Theta_A + \frac{1}{2} [\Theta_A, \Theta_A] = \kappa(A) \cdot \omega$$

for Θ_A in the chiral deformation complex of A .

Remark 4.2. The identification proceeds through the following dictionary:

BV-BRST	Chiral bar complex	Identification
Field $\alpha \in \mathfrak{F}$	Bar element $\alpha \in B(A)$	Same underlying vector space
BV action $S(\alpha)$	MC element Θ_A	$S = \text{Tr}(\Theta_A)$
CY pairing $\langle -, - \rangle$	Factorization coproduct	$\omega_{\text{BV}} = \Delta_B^* \omega_{\text{CY}}$
$\{S, S\} = 0$ (CME)	$[\Theta_A, \Theta_A] = 0$ at genus 0	Classical A_∞ relations
$\hbar \Delta S$	$d\Theta_A$ at genus ≥ 1	One-loop anomaly
QME obstruction	$\kappa(A) \cdot \omega_g$	Genus- g anomaly

4.3. The genus expansion. Expanding the QME in powers of $\hbar$ (equivalently, by genus), we obtain a hierarchy:

- **Genus 0** ($\hbar^0$): the classical master equation $\{S, S\} = 0$. This is the A_∞ -relation, which holds identically. On the chiral side, this is the genus-0 part of the MC equation: $[\Theta_A, \Theta_A]|_{g=0} = 0$, which is the condition that $B(A)$ is a differential coalgebra at tree level.
- **Genus 1** ($\hbar^1$): the one-loop anomaly $\Delta S_2 + \{S_3, S_2\} + \dots$. On the chiral side, this is $d\Theta_A|_{g=1} = \kappa(A) \cdot \lambda_1$, where λ_1 is the first Chern class of the Hodge bundle. The anomaly is proportional to $\kappa(A) = \kappa(\mathcal{C})$, the CY modular characteristic.
- **Genus g** ($\hbar^{2g-2}$): on the uniform-weight lane, the genus- g anomaly is $\text{obs}_g(A) = \kappa(A) \cdot \lambda_g$, where λ_g is the top Chern class of the Hodge bundle on $\overline{\mathcal{M}}_{g,1}$ (for multi-weight algebras at $g \geq 2$, cross-channel corrections appear; Vol I, AP32). The BV-BRST complex has vanishing *scalar* curvature at genus g if and only if $\kappa(A) = 0$.

Corollary 4.3 (Anomaly cancellation). *The open string field theory on $\mathcal{C}$ is anomaly-free (i.e., the QME holds to all orders in $\hbar$) if and only if the modular characteristic $\kappa(A_{\mathcal{C}}) = 0$. In the BKM algebra language, this corresponds to the BKM algebra $\mathfrak{g}_X$ being “untwisted” — i.e., the automorphic correction having trivial Weyl vector contribution.*

Example 4.4 (CY3 B-model). For the B-model on a CY3 manifold X with $A = \Phi(D^b(\text{Coh}(X)))$, the conjectural modular characteristic is $\kappa(A) = \chi^{\text{CY}}(X)$ (the CY Euler characteristic), assuming the quantum vertex chiral group $G(X)$ exists and has this identification (Conjecture CY-A₃). Under this assumption, anomaly cancellation $\kappa = 0$ requires $\chi(X) = 0$. The quintic threefold has $\chi = -200 \neq 0$, so the B-model on the quintic would have a non-trivial BRST anomaly at genus ≥ 1 . However, for $K3 \times E$ with $\chi = 0$, the automorphic prediction gives $\kappa = 5 \neq 0$ (the weight of Δ_5), so $\kappa = \chi^{\text{CY}}$ fails in general (AP48).

Example 4.5 (CY3 with $\chi = 0$). A CY3 X with $h^{1,1} = h^{2,1}$ (so that $\chi(X) = 2(h^{1,1} - h^{2,1}) = 0$) has vanishing Euler characteristic. Under the identification $\kappa = \chi^{\text{CY}}$, this would give $\kappa = 0$. However, the $K3 \times E$ example shows that $\chi = 0$ does not guarantee $\kappa = 0$: the automorphic weight can be nonzero even when χ vanishes. Mirror pairs $(X, X^\vee)$ with $h^{1,1}(X) = h^{2,1}(X^\vee)$ satisfy $\chi(X) = -\chi(X^\vee)$; for both to have $\chi = 0$ requires $h^{1,1} = h^{2,1}$ on each side.

5. THE BAR COMPLEX AS THE BV-BRST COMPLEX

5.1. The bar differential as Q_{BRST} . The *bar complex* $B(A)$ of a chiral algebra A is a factorization coalgebra on $\text{Ran}(X)$, defined as

$$B(A) = \bigoplus_{n \geq 1} A[1]^{\otimes n}$$

with the bar differential d_B built from the chiral operations (OPE residues) and the deconcatenation coproduct. (See Volume I, Part II for the full construction.)

Proposition 5.1 (Bar differential = BRST operator). *Under the identification $\mathfrak{F} = A[1]$ and $S = \sum \frac{1}{n} \text{Tr}(\mu_n)$, the BRST operator*

$$(10) \quad Q_{\text{BRST}} = \{S, -\}$$

acting on the field space $\mathfrak{F}$ is identified with the bar differential d_B on $B(A)$. Explicitly, for a bar element $\alpha_1 \otimes \cdots \otimes \alpha_n \in A[1]^{\otimes n}$,

$$d_B(\alpha_1 \otimes \cdots \otimes \alpha_n) = \sum_{i < j} \pm \alpha_1 \otimes \cdots \otimes \mu_{j-i+1}(\alpha_i, \dots, \alpha_j) \otimes \cdots \otimes \alpha_n.$$

This is the standard bar differential of an A_∞ -algebra. On the BV side, the same formula arises from $\{S, -\}$ using the BV antibracket and the action (??).

5.2. Curvature and the BRST anomaly. At genus $g = 0$, the bar differential satisfies $d_B^2 = 0$ — this is the content of the A_∞ -relations, or equivalently the classical master equation. At genus $g \geq 1$, however, the bar complex acquires *curvature* from the Hodge bundle on $\overline{\mathcal{M}}_{g,n}$:

$$(11) \quad \boxed{d_B^2 = \kappa(A) \cdot \omega_g}$$

where $\omega_g \in H^2(\overline{\mathcal{M}}_{g,1})$ is the Hodge class and $\kappa(A)$ is the modular characteristic. The curvature $d_B^2 \neq 0$ means that $B(A)$ is not a chain complex at genus ≥ 1 but rather a *curved* factorization coalgebra.

Theorem 5.2 (Curvature = BRST anomaly). *The curvature equation (??) is the BRST anomaly:*

- (i) *At genus g , the failure of nilpotency $Q_{\text{BRST}}^2 \neq 0$ is an operator proportional to $\kappa(A) \cdot \omega_g$. This is the BRST anomaly of the open string field theory at g loops.*
- (ii) *The anomaly is universal: it depends only on $\kappa(A)$ and the Hodge class, not on the particular brane $\mathcal{F}$ or the details of the A_∞ -structure beyond κ .*
- (iii) *Anomaly cancellation $\kappa(A) = 0$ restores $d_B^2 = 0$ at all genera — the bar complex is an honest chain complex, the BV-BRST cohomology is well-defined, and the string field theory is consistent to all orders.*

Remark 5.3 (Physical content). In physics language: the one-loop anomaly arises from the trace of the BRST operator acting on the propagator (a loop of an open string). The trace picks up the CY trace Tr on the endomorphism algebra, and the result is proportional to κ . The universality of the anomaly (dependence only on κ , not on the details of the theory) is the algebraic version of the Adler–Bardeen theorem: the anomaly is one-loop exact.

5.3. The BKM interpretation. In the BKM (Borcherds–Kac–Moody) language developed in the VISION document:

- The bar differential $d_B = Q_{\text{BRST}}$ at genus 0 encodes the *real root* structure of the BKM algebra $\mathfrak{g}_X$.
- The curvature $\kappa \cdot \omega_g$ at genus $g \geq 1$ encodes the *imaginary root* corrections — the automorphic correction.
- Anomaly cancellation $\kappa = 0$ means the BKM algebra is “untwisted”: all imaginary roots have zero multiplicity at the leading order, and the denominator identity reduces to the Weyl–Kac formula (no automorphic correction needed).

- The shadow obstruction tower $\Theta_A^{\leq r}$ captures the anomaly cancellation conditions order by order in arity: κ at arity 2, the cubic shadow C at arity 3, the quartic Q at arity 4, etc.

6. THE SWISS-CHEESE STRUCTURE AND OPEN-CLOSED STRING FIELD THEORY

6.1. The Swiss-cheese operad. The open-closed string field theory requires both open and closed string sectors. The algebraic structure governing their interaction is the *Swiss-cheese operad* $\text{SC}^{\text{ch},\text{top}}$ from Volume II: a 2-colored operad with a “chiral” color (closed strings, bulk) and a “topological” color (open strings, boundary).

Definition 6.1 ($\text{SC}^{\text{ch},\text{top}}$ structure on $B(A)$, cf. Vol II). The bar complex $B(A)$ of a quantum chiral algebra $A = A_{\mathcal{C}}$ carries a $\text{SC}^{\text{ch},\text{top}}$ -algebra structure consisting of:

- (i) An E_1 -algebra structure on the “topological” sector (open string fields): the A_{∞} -operations, assembling into Q_{BRST} .
- (ii) A chiral algebra structure on the “chiral” sector (closed string fields): the factorization/OPE structure of A .
- (iii) Open-closed couplings: maps from closed string states to deformations of the open string theory, i.e., from $\text{HH}^{\bullet}(\mathcal{C})$ (Hochschild cohomology, the closed string states) to deformations of the A_{∞} -structure on $\text{End}(\mathcal{F})$.

6.2. BV interpretation of the Swiss-cheese structure. The $\text{SC}^{\text{ch},\text{top}}$ structure on $B(A)$ has a direct BV interpretation:

- (1) **Open sector (topological/ E_1).** The E_1 -algebra structure on the open string fields $\mathfrak{F}_{\text{open}} = A[1]$ is the BV-BRST complex of the open string field theory. The E_1 -product is the concatenation product on the bar complex (juxtaposition of open string states), which is the *star product* $\star$ of open string field theory. The differential $d_B = Q_{\text{BRST}}$ is the BRST operator. The E_1 direction is the BRST direction.
- (2) **Closed sector (chiral).** The chiral algebra A itself governs the closed string sector. The factorization structure on $\text{Ran}(X)$ encodes the OPE of closed string vertex operators. The closed string field $\Phi \in \text{HH}^{\bullet}(\mathcal{C})[2]$ (shifted Hochschild cohomology) enters the BV action through the coupling

$$S_{\text{closed}}(\Phi) = \sum_{g \geq 0} \hbar^{2g-2} \int_{\mathcal{M}_{g,0}} \omega_{g,\Phi}$$

where $\omega_{g,\Phi}$ is the genus- g amplitude built from Φ via the Kodaira–Spencer/BCOV theory.

- (3) **Open-closed coupling.** A closed string state Φ deforms the open string BV action via

$$S(\alpha; \Phi) = S(\alpha) + \sum_{n,m} \frac{1}{n} \text{Tr}(\mu_{n|m}(\alpha^{\otimes n}; \Phi^{\otimes m}))$$

where $\mu_{n|m}$ are the open-closed A_{∞} -operations. In the bar complex language, this is the action of $\text{HH}^{\bullet}(\mathcal{C})$ on $B(A)$ by coderivations — the open-closed map of Kontsevich–Soibelman.

6.3. The E_2 enhancement and the braiding. When the CY category $\mathcal{C}$ has dimension $d = 2$ (or more generally when the $\mathbb{S}^d$ -framing provides an E_2 -structure), the Swiss-cheese structure on $B(A)$ acquires an E_2 -enhancement:

Proposition 6.2 (E_2 enhancement of $\text{SC}^{\text{ch},\text{top}}$). *The E_1 direction (BRST direction) and the E_2 enhancement (braiding direction) on $B(A)$ assemble into a structure governed by the Swiss-cheese operad:*

- (i) The E_1 direction is the BRST direction: it contains the bar differential $d_B = Q_{\text{BRST}}$ and the concatenation/star product.
- (ii) The E_2 enhancement adds the braiding: the R -matrix $R_C(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{A_C})$ from the CY R -matrix construction.
- (iii) Together, they give the E_2 -bar complex $B_{E_2}(A)$ with two commuting differentials d_X, d_Y and two commuting coproducts Δ_X, Δ_Y , as in the E_2 -chiral bar construction of Vol III, §8.4.

In physical terms: the E_1 direction corresponds to the *ordering* of open string insertions on the boundary of the worldsheet (the cyclic order on $\partial\Sigma$). The E_2 enhancement corresponds to the *braiding* of open string insertions — the possibility of exchanging their cyclic order by deforming the boundary of the worldsheet through the bulk. The braiding is non-trivial precisely when there is a non-trivial quantum group structure, i.e., when the R -matrix $R(z) \neq \tau$ (the trivial swap).

Remark 6.3 (Physical interpretation of the two directions). Consider the topological B-model on a CY surface X (CY2, e.g., a K3 surface). The worldsheet Σ has boundary $\partial\Sigma$ on which open string vertex operators are inserted. The E_1 direction corresponds to the *radial direction* of the worldsheet (distance from the boundary), governing the OPE of boundary operators. The E_2 direction corresponds to the *angular direction* along the boundary, governing the braiding. Together, the two directions give the full E_2 -disk structure — which is exactly the little 2-disks operad acting on the space of boundary operators.

7. THE CY3 CASE: THE TOPOLOGICAL B-MODEL AND $G(X)$

7.1. The B-model on a CY3. Let X be a compact Calabi–Yau threefold. The topological B-model on X has:

- **Open string sector:** Branes are objects $\mathcal{F} \in D^b(\text{Coh}(X))$. Open string fields are $\alpha \in \text{Ext}^\bullet(\mathcal{F}, \mathcal{F})[1]$. The BV action is (??) with Tr given by the Serre duality pairing and μ_n from the dg enhancement of $D^b(\text{Coh}(X))$.
- **Closed string sector:** Closed string fields are polyvector fields, i.e., elements of $\bigoplus_{p,q} H^q(X, \bigwedge^p T_X) = \text{HH}^\bullet(D^b(\text{Coh}(X)))$. The closed string field theory is the Kodaira–Spencer/BCOV theory.
- **Open-closed coupling:** Via the Hochschild-to-deformation map: a closed string field $\Phi \in \text{HH}^\bullet$ deforms the derived category $D^b(\text{Coh}(X))$.

7.2. The quantum vertex chiral group $G(X)$. By the CY-to-chiral functor Φ , the B-model data assembles into the quantum vertex chiral group $G(X)$ — a quantum chiral algebra whose structure encodes the full BV-BRST complex:

Conjecture 7.1 (BV-BRST complex = bar complex of $G(X)$). *For a CY3 manifold X , assuming the quantum vertex chiral group $G(X)$ exists (Conjecture CY- A_3), the full BV-BRST complex of the topological B-model should be the bar complex $B(G(X))$. Specifically:*

- (i) The bar differential d_B on $B(G(X))$ is the BRST operator Q_{BRST} of the B-model.
- (ii) The genus-0 cohomology $H^\bullet(B(G(X)), d_B)|_{g=0}$ computes the BRST cohomology of the classical B-model: the physical open string states modulo gauge equivalence.
- (iii) At genus $g \geq 1$, the curvature $d_B^2 = \kappa(G(X)) \cdot \omega_g$ is the g -loop BRST anomaly of the B-model. The conjectural modular characteristic $\kappa(G(X))$ depends on the CY trace normalization; see the anomaly cancellation note for the distinction between $\chi(X)/24$ (Heisenberg prediction) and $\text{wt}(\Phi_X)$ (automorphic prediction).

- (iv) The bar-cobar adjunction $\Omega(B(G(X))) \simeq G(X)$ (on the Koszul locus) is the statement that the open string field theory determines (and is determined by) the quantum vertex chiral group.
- (v) The E_2 -bar complex $B_{E_2}(G(X))$ is the BV-BRST complex of the full open-closed B-model, with the E_2 -structure encoding the braided monoidal structure on $\text{Rep}^{E_2}(G(X))$.

7.3. The BKM structure of $G(X)$. The quantum vertex chiral group $G(X)$ of a CY3 has the structure of a generalized Borchers–Kac–Moody superalgebra (or affine super Yangian, in the toric case). The BV-BRST dictionary translates as follows:

BV-BRST	Bar complex of $G(X)$	BKM algebra
BRST operator Q_{BRST}	Bar differential d_B	Root space differential
BRST cohomology $H^*(Q)$	$H^*(B(G(X)))$ at $g = 0$	Weight spaces
BRST anomaly at genus g	Curvature $\kappa \cdot \omega_g$	Automorphic correction
Anomaly cancellation $\kappa = 0$	$d_B^2 = 0$ at all genera	Untwisted BKM
BV bracket $\{-, -\}$	CY pairing on $B(G(X))$	Root space inner product
Fields + antifields	$A[1] \oplus A[1]^*$	Positive + negative roots
Ghost number	Bar filtration degree	Root height
Star product $\star$	E_1 concatenation	CoHA multiplication
R -matrix / braiding	E_2 enhancement	Yangian / quantum group R -matrix

7.4. The denominator identity as the BV partition function. The denominator identity of the BKM superalgebra $\mathfrak{g}_X$ is the multiplicative Euler product:

$$\sum_{w \in W} \det(w) e^{w(\rho)} = e^\rho \prod_{\alpha \in \Delta_+} (1 - e^{-\alpha})^{\text{mult}(\alpha)}$$

In the BV-BRST language, this is the *partition function* (or more precisely, the ratio of regularized determinants):

$$Z_{\text{BRST}} = \frac{\det Q_{\text{BRST}}|_{\text{ghost}}}{\det Q_{\text{BRST}}|_{\text{field}}} = \prod_{\alpha \in \Delta_+} (1 - q^\alpha)^{\text{mult}(\alpha)}$$

where the product runs over positive roots α and $q^\alpha = e^{-2\pi i(\alpha, z)}$. The Weyl vector ρ enters as the zero-point energy (Casimir energy). The denominator identity IS the BV partition function of the B-model, expressed as a product over BPS states (the root spaces of $\mathfrak{g}_X$).

Remark 7.2. For $K3 \times E$, this product formula reproduces the Borchers product expansion of the Igusa cusp form Δ_5 . For toric CY3, it reproduces the DT partition function as an infinite product. In both cases, the root multiplicities are the BPS degeneracies, and the denominator identity is the BPS state sum — i.e., the one-loop BV-BRST partition function with all BPS states running in the loop.

8. SUMMARY OF IDENTIFICATIONS

We collect the main identifications developed in this note:

- (1) **Fields.** The open string field $\alpha \in \text{RHom}(\mathcal{F}, \mathcal{F})[1]$ is an element of the bar complex $B(A_{\mathcal{C}})$.
- (2) **Action.** The BV action $S(\alpha) = \sum_n \frac{1}{n} \text{Tr}(\mu_n(\alpha^{\otimes n}))$ is the CY trace of the universal MC element Θ_A .

- (3) **Bracket.** The BV antibracket $\{-, -\}$ is the odd Poisson bracket induced by the CY pairing on $A[1]$.
- (4) **QME = MC.** The quantum master equation $\hbar\Delta S + \frac{1}{2}\{S, S\} = 0$ is the Maurer–Cartan equation for Θ_A .
- (5) **Bar differential = BRST.** The bar differential d_B on $B(A)$ is the BRST operator $Q_{\text{BRST}} = \{S, -\}$.
- (6) **Curvature = anomaly.** The curvature $d_B^2 = \kappa \cdot \omega_g$ at genus g is the BRST anomaly. Anomaly cancellation requires $\kappa(A_C) = 0$.
- (7) **Swiss-cheese = open-closed BV.** The $\text{SC}^{\text{ch}, \text{top}}$ structure on $B(A)$ is the BV structure on the combined open-closed string field theory.
- (8) $E_1 = \text{BRST}$, $E_2 = \text{braiding}$. The E_1 direction of the Swiss-cheese structure is the BRST direction; the E_2 enhancement adds the braiding/quantum group structure.
- (9) **For CY3.** The full BV-BRST complex of the topological B-model on X is the bar complex $B(G(X))$ of the quantum vertex chiral group. The denominator identity of the BKM superalgebra $\mathfrak{g}_X$ is the BV partition function.

9. OUTLOOK

Several directions emerge from this BV-BRST perspective:

- (i) **BV quantization and the shadow obstruction tower.** The shadow obstruction tower $\Theta_A^{\leq r}$ of Volume I should be interpretable as the perturbative BV quantization of the open string field theory, with the arity- r shadow encoding the r -loop effective action. Making this precise requires developing the BV path integral for the CY A_∞ -action.
- (ii) **Wall-crossing as gauge equivalence.** The Kontsevich–Soibelman wall-crossing formula for DT/BPS invariants should arise as the equivalence of BV-BRST cohomologies for different gauge-fixing choices. In the bar complex language, this is the statement that different resolutions of $B(G(X))$ (corresponding to different stability conditions) produce the same cohomology.
- (iii) **Higher genus amplitudes.** The genus- g amplitude of the B-model (the g -th term in the BCOV/Kodaira–Spencer free energy) should be computable as the trace of $\Theta_A^{(g)}$ over $\overline{\mathcal{M}}_{g,n}$. The BV formalism provides a systematic way to extract these amplitudes from the bar complex.
- (iv) **Categorical BV formalism.** The BV-BRST complex depends on a choice of brane $\mathcal{F}$. The “brane-independent” formulation is the bar complex $B(A_C)$ itself, which is defined without choosing $\mathcal{F}$. This suggests that the bar complex is the categorified BV-BRST complex, with individual brane choices providing fiber functors.
- (v) **The closed string field theory and $\mathcal{W}$ -algebras.** For CY3, the closed string sector is controlled by the BCOV theory, whose algebraic structure is a $\mathcal{W}$ -algebra (the vertex algebra of polyvector fields with Schouten bracket and ∂ operator). The full open-closed BV-BRST complex should be the Swiss-cheese bar complex $B_{\text{SC}}(A_C)$, combining the $\mathcal{W}$ -algebra (bulk) with $G(X)$ (boundary) via the open-closed coupling.