

WORKING NOTES ON CALABI–YAU QUANTUM GROUPS

Raez Lorgat

Draft — April 5, 2026

Contents

1	Overview and programme	I
1.1	The four target theorems	2
1.2	Current status of the programme	2
2	Key identifications	4
2.1	The master dictionary	4
2.2	The central identification	4
2.3	Computationally verified	4
3	The $K3 \times E$ example	5
3.1	The geometry	5
3.2	The lattice	5
3.3	The modular characteristic	5
4	The toric CY_3 / CoHA example	5
5	Higgs sheaves and the CY_2 escape	5
6	The Langlands connection	6
7	Open questions	6
8	Arithmetic shadows and number theory	6
8.1	MZV periods from the genus-0 bar complex	6
8.2	The Drinfeld associator as bar transport	7
8.3	Siegel modular forms from genus-2 amplitudes	7
8.4	Moonshine and Niemeier atlas	7

I Overview and programme

These working notes record the development of Volume III (Calabi–Yau Quantum Groups) of the modular Koszul duality programme. The central thesis: *the BPS spectrum of a CY_3 threefold X should constitute the generalized root datum of a quantum group $G(X)$ realized as an E_2 -chiral algebra*. More precisely, the automorphic correction of the underlying Borchers–Kac–Moody (BKM) superalgebra attached to X should be identified with the shadow Postnikov obstruction tower Θ_A from Volume I: the universal Maurer–Cartan element of the modular convolution algebra of a chiral algebra $A = A_X$ associated to X , whose finite-order projections $\Theta_A^{\leq r}$ encode roots of increasing complexity.

Warning 1.1 (The central object $G(X)$ is not yet constructed). The “quantum vertex chiral group” $G(X)$ is the *target* of this programme, not a defined object. For CY_2 categories (Higgs sheaves on a curve), the construction of an E_2 -chiral algebra via the factorization envelope is available (CY-A for $d = 2$). For CY_3 threefolds, the chain-level $\mathbb{S}^3$ -framing needed for CY-A at $d = 3$ has not been constructed, and $G(X)$ is used as shorthand for the conjectural output. Every statement involving $G(X)$ for a CY_3 is a conjecture unless otherwise noted.

1.1 The four target theorems

- **CY-A** (CY-to-chiral functor): $\Phi: \text{CY}_d\text{-Cat} \rightarrow \text{E}_2\text{-ChirAlg}$.
- **CY-B** (E_2 -chiral Koszul duality): Automorphic correction = shadow obstruction tower; denominator identity = bar Euler product (at the level of motivic Hall algebra generating series; naive series-level equality requires CY-A).
- **CY-C** (Quantum group realization): $\text{Rep}^{\text{E}_2}(G(X))$ is braided monoidal; for toric CY_3 , this gives the affine super Yangian.
- **CY-D** (Modular CY characteristic): $\kappa(G(X)) = \text{weight of the denominator automorphic form (e.g., weight}(\Delta_5) = 5 \text{ for } K3 \times E)$.

1.2 Current status of the programme

Component	Status	Notes file
Generalized root datum axioms	Draft (CY ₁ –CY ₇)	theory_generalized_root_datum
E ₂ -chiral algebra formalism	Draft (1243 lines)	theory_e2_chiral_formalism
Automorphic = shadow obstruction tower	Draft (Thm + proof)	theory_automorphic_shadow
CY-to-chiral functor	Draft (4-step construction)	theory_cy_to_chiral_construction
CY ₂ → CY ₃ fibration	Draft (Borcherds lift)	theory_cy2_cy3_fibration
Denominator = bar Euler product	Draft (3-stage proof)	theory_denominator_bar_euler
QVCG Koszul duality	Draft (Langlands conj)	theory_qvcg_koszul
Drinfeld = chiral center	Draft (factorization proof)	theory_drinfeld_chiral_center
CoHA = E ₁ -sector	Draft (Drinfeld double)	theory_coha_e1_sector
KL in E ₂ -chiral framework	Draft (CY category conj)	theory_kl_e2_chiral
Higgs sheaves as CY ₂ QVCGs	Draft (genus filtration)	theory_higgs_cy2_qvcg
BPS = root multiplicities	Draft	physics_bps_root_multiplicities
Topological strings & root data	Draft (BCOV = MC)	physics_topological_strings
M-theory branes / holography	Draft	physics_mtheory_branes
4d $\mathcal{N} = 2$ / Hitchin	Draft (Z _{Nek} conj)	physics_4d_n2_hitchin
Wall-crossing = MC gauge equiv	Draft	physics_wall_crossing_mc
S-duality = Langlands = Koszul	Draft	physics_sduality_langlands
Anomaly cancellation / κ	Draft ($\kappa \neq \chi/12$)	physics_anomaly_cancellation
BV-BRST for CY categories	Draft (QME = MC)	physics_bv_brst_cy
3d mirror symmetry / CY ₂	Draft	physics_3d_mirror
Celestial holography / CY QG	Draft ($\mathcal{W}_{1+\infty} = G(\mathbb{C}^3)$)	physics_celestial_cy
Hitchin / geometric Langlands	Draft	physics_hitchin_langlands
$\phi_{0,1}$ Fourier coefficients	Computed (51 tests)	compute/lib/phior_fourier
$\mathbb{C}^3$ DT / MacMahon	Computed (57 tests)	compute/lib/c3_dt_partition
Lattice data (8 dd-forms)	Computed (65 tests)	compute/lib/dd_modular_lattices
Topological vertex	Computed	compute/lib/topological_vertex
Igusa product formula	Computed	compute/lib/igusa_product_formula
Affine Yangian $Y(\widehat{\mathfrak{gl}}_1)$	Computed (81 tests)	compute/lib/affine_yangian_gli
BKM shadow obstruction tower	Computed (67 tests)	compute/lib/bkm_shadow_tower
Elliptic Hall algebra	Computed (78 tests)	compute/lib/elliptic_hall
CY Euler characteristics	Computed (85 tests)	compute/lib/cy_euler
WKB denominator	Computed	compute/lib/wkb_denominator
Higgs CoHA on $\mathbb{P}^1$	Computed (84 tests)	compute/lib/higgs_p1_coha

2 Key identifications

2.1 The master dictionary

CY ₃ geometry	BKM / Yangian	Vol I bar-cobar
Lattice $\Lambda(X)$	Root lattice	Cyclic deformation complex
BPS invariants DT_α	Root multiplicities $\text{mult}(\alpha)$	Shadow obstruction tower components
Toric diagram	Dynkin diagram / Gram matrix	Collision r -matrix $r(z)$
Automorphic correction	Imaginary roots added to $\mathfrak{g}$	Higher-arity shadows
Denominator identity	WKB character formula	Bar Euler product
$Sp_4(\mathbb{Z})$ / Weyl group	Symmetry group	E_2 braiding
$\phi_{0,1}$ (K_3 elliptic genus)	Root multiplicity generating function	Shadow obstruction tower input data
CoHA (positive half)	$U(\mathfrak{n}_+)$	E_1 -chiral sector
Full Yangian / BKM	Full algebra $\mathfrak{g}_X$	E_2 -chiral algebra $A(X)$

2.2 The central identification

Conjecture 2.1 (Automorphic correction = shadow obstruction tower). Let X be a CY₃ threefold admitting a BKM root system $\mathfrak{g}_X$, and suppose the CY-to-chiral functor Φ of CY-A produces an E_2 -chiral algebra A_X for X . Then the shadow Postnikov obstruction tower $\Theta_{A_X}^{\leq r}$ captures the roots of complexity $\leq r$ in the BKM root system:

- (a) Arity 2 = real roots + Weyl vector (modular characteristic κ).
- (b) Arity 3 = first layer of imaginary roots (cubic shadow C).
- (c) Arity r = roots up to height $r - 2$ in the positive cone (height measured by the minimal number of simple root additions needed to reach α from norm-2 real roots).
- (d) The denominator identity of $\mathfrak{g}_X$ equals the bar-complex Euler product of $B(A_X)$.

Evidence for (a)–(b): computationally verified for $K3 \times E$ at arities 2–6 (Section 3). Part (d) requires CY-A at $d = 3$, which is open (Warning 1.1).

2.3 Computationally verified

For $K3 \times E$ with $\mathfrak{g}_{\Delta_5}$:

- Arity 2: 21 real roots (norm 2), all multiplicity 2. $\kappa = 5$.
- Arity 3: 2 lightlike imaginary roots at $(1, 0, 0)$ and $(0, 0, 1)$, multiplicity 20.
- Arity 4: 7 roots including the timelike root $(1, 0, 1)$ with multiplicity -252 .
- At every truncation order $r = 2, \dots, 6$: product formula matches shadow projection.

3 The $K3 \times E$ example

3.1 The geometry

$X = S \times E$, where S is a $K3$ surface and E is an elliptic curve. The DT partition function is $Z^X = C/(\Delta_5)^2$, where Δ_5 is the Igusa cusp form of weight 5 on $_4(\mathbb{Z})$ (the degree-2 Siegel modular group).

3.2 The lattice

The BKM root lattice is $\Lambda^{3,2} \simeq \Lambda^{1,1} \oplus \Lambda^{1,1} \oplus [2]$, of signature $(3, 2)$. The hyperbolic sublattice $\Lambda_{II}^{2,1}$ has three real simple roots $\delta_1, \delta_2, \delta_3$ with Gram matrix

$$(\delta_i, \delta_j) = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}.$$

Weyl vector $\rho = \frac{1}{2}(\delta_1 + \delta_2 + \delta_3)$, satisfying $(\rho, \delta_i) = -1$ for each i (verified: $(\rho, \delta_1) = \frac{1}{2}(2 - 2 - 2) = -1$). The Coxeter group is $\text{Aut}(\mathcal{P}_{II}) \simeq S_3$.

3.3 The modular characteristic

The conjectural modular characteristic (CY-D) is $\kappa(G(K3 \times E)) = 5$. This equals the weight of the Igusa cusp form: $\text{weight}(\Delta_5) = 5$. Numerically, $5 = h^{1,1}(K3)/4 = 20/4$ and $5 = (\chi(K3) - 4)/4 = (24 - 4)/4$, but these expressions are observed coincidences for $K3 \times E$, not proved formulas for general CY_3 (see Question 5 in Section ??).

This is *not* $\chi(X)/12$, which gives 0 since $\chi(K3 \times E) = 0$. The modular characteristic counts the worldsheet anomaly (full BPS spectrum), not the target-space anomaly (massless modes).

4 The toric CY_3 / CoHA example

For $\mathbb{C}^3$, the critical CoHA (Schiffmann–Vasserot, Rapcak–Soibelman–Yang–Zhao) is $\text{CoHA}(\mathbb{C}^3) \simeq Y^+(\widehat{\mathfrak{gl}}_1)$, the positive half of the affine Yangian of $\mathfrak{gl}_1$. The graded dimension is $\dim Y_n^+ = p(n)$ (the number of plane partitions of n), and the MacMahon function $M(q) = \prod_{n \geq 1} (1 - q^n)^{-n}$ serves as the generating function. If the CY-to-chiral functor exists for $\mathbb{C}^3$ (CY-A at $d = 3$, open), the CoHA should be the E_1 -sector of the conjectural E_2 -chiral algebra $A_{\mathbb{C}^3}$, and $M(q)$ should match the bar Euler product. At the self-dual level ($h_1 + h_2 + h_3 = 0$), $Y^+(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$. The affine Yangian structure function is $g(z) = (z - h_1)(z - h_2)(z - h_3)/((z + h_1)(z + h_2)(z + h_3))$; even ϕ -coefficients vanish ($\phi_2 = \phi_4 = 0$) because $\log g$ has only odd powers.

5 Higgs sheaves and the CY_2 escape

The most natural escape from CY_3 : $\text{Higgs}(C) = \text{Coh}(T^*C)$ is CY_2 . The $\mathbb{S}^2$ -framing gives E_2 -structure directly.

- **Genus 0** ($C = \mathbb{P}^1$): Yangian $Y(\mathfrak{gl}_2)$. Rational R -matrix.
- **Genus 1** ($C = E$): Elliptic Hall algebra $E_{q,t}$ = spherical DAHA. Elliptic R -matrix.
- **Genus ≥ 2** : New “Hitchin Hall algebras.” R -matrices on $C \times C$.

Key principle: $CY_3 = CY_2 + \text{automorphic correction (Borcherds lift)}$.

6 The Langlands connection

Conjecture 6.1 (Langlands = Koszul). For a curve C and reductive group G , the quantum vertex chiral group of the Hitchin system satisfies

$$G(C, G)^! = G(C, {}^L G).$$

Langlands duality is Koszul duality of quantum vertex chiral groups.

Evidence: Feigin–Frenkel center at critical level, root/coroot exchange, SYZ mirror symmetry of $\mathcal{M}_H(C, G)$ and $\mathcal{M}_H(C, {}^L G)$, Kapustin–Witten.

7 Open questions

1. What is the generalized root datum of the quintic? (Non-toric, no quiver.)
2. Is the chain-level $\mathbb{S}^3$ -framing for CY_3 categories constructible? (Gap in CY-A for $d = 3$.)
3. What is the explicit automorphic form (denominator identity) for Hitchin systems with $G = \text{SL}_3$ or E_8 ?
4. How does Kontsevich–Soibelman wall-crossing manifest as gauge equivalence of MC elements? (Conjectured, not proved.)
5. Is the formula $\kappa = h^{1,1}/4$ universal for K_3 -fibered CY_3 , or specific to $K_3 \times E$?
6. For toric CY_3 with compact 4-cycles, what replaces the RSYZ theorem?
7. The eight diagonal divisor Siegel modular forms: are they all denominator identities?
8. How does the shadow depth classification (G/L/C/M) manifest for quantum vertex chiral groups?
9. Explicit construction of the E_2 -bar complex $B_{E_2}(A)$ for $A = V_k(\mathfrak{g})$.
10. Quantum geometric Langlands as E_2 -chiral Koszul duality: $G(C, \mathfrak{g}, k)^! = G(C, {}^L \mathfrak{g}, k^L)$.

8 Arithmetic shadows and number theory

The shadow obstruction tower of a chiral algebra carries arithmetic content that connects to classical number theory through three channels: modular forms at genus 1, Siegel modular forms at genus 2, and multiple zeta values at genus 0.

8.1 MZV periods from the genus-0 bar complex

The genus-0 bar amplitudes on $\mathcal{M}_{0,n}$ produce MZV periods via iterated integrals of the bar propagator $d \log(z_i - z_j)$. The shadow depth classification controls which MZV weights appear.

THEOREM 8.1 (*Shadow–MZV dictionary* (proved in Vol I)). For a chirally Koszul algebra $\mathcal{A}$:

- (i) $\kappa(\mathcal{A})$ controls the coefficient of $\zeta(2)$ in genus-0 four-point amplitudes.
- (ii) The cubic shadow $S_3(\mathcal{A})$ controls the coefficient of $\zeta(3)$ in five-point amplitudes.

(iii) The quartic shadow $S_4(\mathcal{A})$ and contact class Q^{contact} control weight-4 MZV content.

(iv) At general weight r : the arity- r shadow S_r contributes to the MZV space of dimension $d_r = d_{r-2} + d_{r-3}$ (Brown 2012).

Class **G** algebras see only $\zeta(2)$; class **L** see $\zeta(2), \zeta(3)$; class **C** see $\zeta(2), \zeta(3), \zeta(4)$ (stratum separation at arity 4); class **M** see MZVs at all weights.

For quantum vertex chiral groups, this dictionary extends to the genus-0 bar complex on a curve C : the MZV periods are replaced by periods of $\mathcal{M}_{0,n}(C)$, which include elliptic MZVs (for $C = E$ an elliptic curve) and higher-genus analogues.

Conjecture 8.2 (CYMZV dictionary). For the quantum vertex chiral group $G(C, \mathfrak{g})$, the genus-0 bar amplitudes on $\mathcal{M}_{0,n}(C)$ produce periods of the motivic cohomology of C . The shadow depth of $G(C, \mathfrak{g})$ controls the motivic weight filtration level of these periods.

8.2 The Drinfeld associator as bar transport

The Drinfeld associator $\Phi(A, B)$ is the parallel transport of the KZ connection on $\mathcal{M}_{0,4}$, identified with the genus-0 arity-2 shadow connection. Its MZV coefficients are:

$$\begin{aligned} \log \Phi = & \zeta(2) [A, B] + \zeta(3) ([A, [A, B]] - [B, [A, B]]) \\ & + \frac{\zeta(4)}{4} ([A, [A, [A, B]]] + [B, [B, [A, B]]]) + \dots \end{aligned}$$

For quantum vertex chiral groups: the Drinfeld associator governs the genus-0 transport between different trivializations of the bar complex, and the associator's MZV coefficients are the perturbative data of the quantum group structure.

8.3 Siegel modular forms from genus-2 amplitudes

At genus 2, the shadow obstruction tower's scalar projection determines the universal part of the genus-2 amplitude, but the full genus-2 partition function carries arithmetic content *beyond* the shadow obstruction tower. For lattice VOAs of rank d , the genus-2 partition function is the Siegel theta series $\Theta_{\Lambda}^{(2)}(\Omega) \in M_{d/2}((4,))$; the shadow obstruction tower fixes $F_2 = \kappa \cdot \lambda_2^{\text{FP}}$ but not the Siegel cusp component (Vol I, lattice chapter).

For the quantum vertex chiral group programme: the genus-2 amplitude of $G(C, \mathfrak{g})$ should produce a Siegel modular form on $(4,)$ whose Fourier coefficients encode the BPS spectrum. The Böcherer factorization converts these Fourier coefficients to central L -values, providing algebraic access to the critical line $\Re(s) = 1/2$.

Conjecture 8.3 (Genus-2 BPS = Böcherer). For the quantum vertex chiral group $G(K3 \times E, \mathfrak{g})$ at the lattice VOA point: the genus-2 MC equation determines the Siegel theta series, and the Böcherer coefficient extracts the central L -value $L(1/2, \pi_f \times \chi_D)$ from the MC data.

8.4 Moonshine and Niemeier atlas

The 24 Niemeier lattices (even unimodular in dimension 24) all have $\kappa = 24$ ($= d$ for $d = 24$ free bosons at level $k = 1$), class **G**, shadow depth 2. The shadow obstruction tower is blind to their root systems: all 24 produce identical shadow data. Distinguishing them requires the full theta series $\Theta_{\Lambda} = E_{12} + c_{\Lambda} \cdot \Delta$ where $c_{\Lambda} = (691|R| - 65520)/691$.

The Monster module $V^{\mathfrak{h}}$ has $c = 24$, $\dim V_1 = 0$, and $\kappa = 12$ (from the Virasoro sector: the Monster has no weight-1 currents, so $\kappa = c/2 = 12$, not $\kappa = c = 24$). This is half the Niemeier value, providing a genuine

shadow-tower separation. The Monster module is conjecturally class **M** (infinite shadow depth), reflecting the nonlinear Griess algebra structure.

For the CY quantum group programme: the K_3 quantum vertex chiral group should encode the Mathieu moonshine (umbral moonshine for $\Lambda = 24A_1$, $G = M_{24}$) through the genus-1 shadow obstruction tower. The mock modular forms of umbral moonshine may arise as the quasi-modular shadow content (through the E_2^* propagator dependence).