

CALABI–YAU QUANTUM GROUPS

*Chiral Algebras from Calabi–Yau Categories
via E_1/E_2 Factorization*

Raez Lorgat

Draft — April 2, 2026

Contents

Contents	i
I Calabi–Yau Categories and Cyclic Structures	I
1 Introduction	3
1.1 The question	3
1.2 The E_1/E_2 chiral hierarchy	3
1.3 Relation to Volumes I and II	4
1.4 CY ₃ combinatorics as generalized root data	4
1.5 Automorphic correction as shadow tower	5
1.6 Main results	5
1.7 Guide for the reader	6
2 Calabi–Yau Categories	7
2.1 Smooth and proper dg categories	7
2.2 The Calabi–Yau condition	7
2.3 The CY trace	7
2.4 Standard examples	8
3 Cyclic A_∞-Structures	9
3.1 A_∞ -categories and the bar complex	9
3.2 Cyclic structures	9
3.3 The cyclic bar complex	9
3.4 The $\mathbb{S}^d$ -framing	10
4 Hochschild Calculus for CY Categories	11
4.1 Hochschild homology and cohomology	11
4.2 The Hochschild-to-cyclic spectral sequence	11
4.3 CY Hochschild duality	11
4.4 Categorical Hodge theory	11
II E_1 and E_2 Chiral Theories	13
5 E_1-Chiral Algebras	15

5.1	E_1 -algebras and associative factorization	15
5.2	The bar complex as E_1 -chiral algebra	15
5.3	Monoidal representation categories	15
5.4	The E_1 -Koszul duality	15
6	E_2-Chiral Algebras	17
6.1	The E_2 operad and braided monoidal categories	17
6.2	E_2 -chiral algebras: definition and first properties	17
6.3	Kontsevich formality and the E_2 -Gerstenhaber structure	17
6.4	The E_2 bar complex	18
6.5	E_2 -chiral Koszul duality	18
7	E_n-Factorization and Higher Chiral Structure	19
7.1	Dunn additivity and the E_n hierarchy	19
7.2	Factorization algebras on $\mathbb{C}^n$	19
7.3	The E_n -chiral bar complex	19
7.4	Relation to topological field theory	19
III	The Bridge: Calabi–Yau Categories as Quantum Chiral Algebras	21
8	From CY Categories to Chiral Algebras	23
8.1	The cyclic-to-chiral passage	23
8.2	The CY-to-chiral functor	23
8.3	Compatibility with Koszul duality	24
8.4	The CY modular characteristic	24
9	Quantum Chiral Algebras	25
9.1	Definition and structure	25
9.2	The CY R -matrix	25
9.3	Quantum chiral algebras from quantum groups	25
9.4	The quantum chiral Koszul dual	25
10	The Modular Trace	27
10.1	CY trace as modular characteristic	27
10.2	Genus expansion and Gromov–Witten invariants	27
10.3	The CY shadow tower	27
10.4	Complementarity for CY pairs	27
IV	Quantum Groups and Braided Monoidal Structure	29
II	Quantum Groups: Foundations	31
II.1	Quantized enveloping algebras	31
II.2	The R -matrix and Yang–Baxter equation	31
II.3	Representation categories	31
II.4	Drinfeld–Jimbo vs Faddeev–Reshetikhin–Takhtajan	31

12 Braided Factorization	33
12.1 Braided monoidal categories and E_2 -algebras	33
12.2 Factorization homology with E_2 -coefficients	33
12.3 The braided bar-cobar adjunction	33
12.4 Quantum group structure from braided Koszul duality	33
13 The Drinfeld Center and Bulk Algebras	35
13.1 The Drinfeld center of a monoidal category	35
13.2 Drinfeld center as chiral derived center	35
13.3 CY enhancement of the Drinfeld center	35
13.4 Quantum group reconstruction from the center	35
V The Standard Landscape	37
14 The $K3 \times E$ Tower and the Igusa Cusp Form	39
14.1 The CY_3 geometry	39
14.2 The lattice $\Lambda^{3,2}$ and the isomorphism $Sp_4 \simeq SO_+$	39
14.3 The hyperbolic sublattice and reflection groups	40
14.4 The generalized BKM superalgebra $\mathfrak{g}_{\Delta_5}$	40
14.5 The denominator identity: $\Delta_5 = \Phi$	41
14.6 The weak Jacobi form $\phi_{0,1}$ and root multiplicities	41
14.7 The quantum vertex chiral group $G(K3 \times E)$	41
15 Toric CY_3 and Critical CoHAs	43
15.1 Toric CY_3 threefolds and their quivers	43
15.2 The critical cohomological Hall algebra	43
15.3 $\mathbb{C}^3$ and the affine Yangian $Y(\widehat{\mathfrak{gl}}_1)$	43
15.4 General toric CY_3 and affine super Yangians	44
15.5 The CoHA as E_1 -sector	44
15.6 Root datum from toric geometry	44
16 Fukaya Categories	45
16.1 The Fukaya category of a CY manifold	45
16.2 CY 2-folds: $K3$ and abelian surfaces	45
16.3 CY 3-folds	45
16.4 Wrapped Fukaya categories	45
16.5 The quantum chiral algebra of a Fukaya category	45
17 Derived Categories of CY Manifolds	47
17.1 $D^b(\text{Coh}(X))$ for CY manifolds	47
17.2 Homological mirror symmetry and the chiral algebra	47
17.3 Exceptional collections and chiral algebras	47
17.4 Stability conditions and the chiral moduli	47
18 Matrix Factorizations	49
18.1 Matrix factorizations and Landau–Ginzburg models	49

18.2	The CY structure on $\mathrm{MF}(W)$	49
18.3	Quantum chiral algebras from singularities	49
18.4	ADE singularities and W-algebras	49
19	Quantum Group Representations	51
19.1	$\mathrm{Rep}_q(\mathfrak{g})$ as a braided monoidal category	51
19.2	The quantum chiral algebra of $U_q(\mathfrak{g})$	51
19.3	Kazhdan–Lusztig equivalence	51
19.4	Yangian and RTT realizations	51
VI	Connections and Frontier	53
20	The Bar-Cobar Bridge to Volume I	55
20.1	CY bar complex vs chiral bar complex	55
20.2	Shadow tower of a CY category	55
20.3	Koszul duality: CY vs chiral	55
20.4	The five theorems in the CY setting	55
21	Modular Koszul Duality and CY Geometry	57
21.1	The modular convolution algebra for CY categories	57
21.2	CY complementarity	57
21.3	The CY shadow CohFT	57
22	Geometric Langlands and CY Quantum Groups	59
22.1	The geometric Langlands programme	59
22.2	CY categories in the Langlands correspondence	59
22.3	Quantum geometric Langlands and E_2 -chiral algebras	59
22.4	The Feigin–Frenkel center and the Drinfeld center	59
A	Conventions	61
A.1	Grading conventions	61
A.2	Sign conventions	61
A.3	Categorical conventions	61
A.4	Operadic conventions	61

Part I

Calabi–Yau Categories and Cyclic Structures

Chapter I

Introduction

I.1 The question

What is a Calabi–Yau quantum chiral algebra?

A Calabi–Yau category C of dimension d carries a canonical cyclic A_∞ -structure: a non-degenerate trace

$$\mathrm{Tr}: \mathrm{HH}_\bullet(C) \longrightarrow k[-d]$$

on its Hochschild homology, encoding the CY condition as a d -dimensional Frobenius structure at the chain level. The cyclic bar complex $\mathrm{CC}_\bullet(C)$ is the primary invariant; it carries an $\mathbb{S}^d$ -action (from the d -sphere framing of the trace), and its $\mathbb{S}^d$ -equivariant structure governs higher-genus amplitudes.

A chiral algebra A on a curve X carries a bar complex $B(A)$, which is a factorization coalgebra on $\mathrm{Ran}(X)$. The bar differential encodes holomorphic OPE residues; the coproduct encodes topological interval-cutting. Together they assemble into a Swiss-cheese algebra structure (Volume II). At genus $g \geq 1$, the bar complex acquires curvature $\kappa(A) \cdot \omega_g$ from the Hodge bundle, and the full modular structure is controlled by the universal Maurer–Cartan element Θ_A (Volume I).

The programme of this work is to construct a precise bridge:

$$\left\{ \begin{array}{c} \text{CY categories with} \\ \text{appropriate } E_n\text{-structure} \end{array} \right\} \xrightarrow{\Phi} \left\{ \begin{array}{c} \text{quantum chiral algebras} \\ \text{with modular trace} \end{array} \right\}$$

The functor Φ should:

- (i) take a CY category C (Fukaya, derived, matrix factorization, or more general) as input;
- (ii) extract the E_2 -monoidal structure on C (braided tensor product, quantum group action);
- (iii) produce a chiral algebra A_C whose bar complex $B(A_C)$ encodes the CY cyclic homology;
- (iv) realize the CY trace as the modular characteristic $\kappa(A_C)$.

I.2 The E_1/E_2 chiral hierarchy

The key structural ingredient is the E_n -chiral hierarchy:

- E_1 -chiral algebras (Volume II, Part VII): associative factorization on $\mathbb{C} \times \mathbb{R}$, encoding the ordered/topological sector. The representation category of an E_1 -chiral algebra is monoidal.
- E_2 -chiral algebras (this work): braided factorization on $\mathbb{C} \times \mathbb{C}$, encoding the full holomorphic-holomorphic sector. The representation category is *braided* monoidal — the natural habitat of quantum groups.
- The passage $E_1 \rightarrow E_2$ via Dunn additivity: $E_2 \simeq E_1 \otimes E_1$. An E_2 -chiral algebra is an E_1 -algebra in E_1 -algebras.

The CY condition enters through Kontsevich’s identification: a d -dimensional CY structure on C determines an $\mathbb{S}^d$ -framing of the Hochschild complex, hence an $\mathbb{S}^1$ -action on $\mathrm{HH}_\bullet(C)$. For $d = 2$ (CY surfaces), this $\mathbb{S}^1$ -action is exactly the data of an E_2 -algebra structure on the cyclic homology — the braiding.

1.3 Relation to Volumes I and II

This work depends on and extends the two preceding volumes:

Volume	Provides	Used here
I (Modular Koszul Duality)	Bar-cobar machine, $\Theta_A, \kappa(A)$	CY bar complex, modular trace
II (3D HT QFT)	$\mathrm{SC}^{\mathrm{ch}, \mathrm{top}}$, PVA descent, DK bridge	E_1 sector, braided structure
III (this work)	$\mathrm{CY} \rightarrow \text{chiral functor}, E_2$ theory	—

1.4 CY₃ combinatorics as generalized root data

The deep structural insight animating this work is that the combinatorics of a Calabi–Yau threefold X — its lattice, intersection form, BPS spectrum — constitute the *generalized root datum* of what we call a *quantum vertex chiral group* $G(X)$.

Given a CY₃ X , the generalized root datum $\mathcal{R}(X)$ consists of:

- (i) A lattice $\Lambda(X)$ extracted from $K(X)$ or $H^*(X)$ with its Euler form;
- (ii) Real simple roots Δ^{re} from the geometry (hyperbolic sublattice for $K3 \times E$; toric diagram for toric CY₃);
- (iii) Imaginary roots $\Delta^{\mathrm{im}} = \Delta_0^{\mathrm{im}} \sqcup \Delta_1^{\mathrm{im}}$ (even and odd) with multiplicities $\mathrm{mult}(\alpha)$ from BPS counting — Donaldson–Thomas invariants for X , or equivalently the Fourier coefficients of the appropriate automorphic form;
- (iv) A Weyl group $W(X)$ acting on the positive cone of $\Lambda(X)$;
- (v) A Weyl vector $\rho \in \Lambda(X) \otimes \mathbb{Q}$.

Two families of examples illuminate the construction:

The $K3 \times E$ tower. For $X = (S \times E)/(\mathbb{Z}/N\mathbb{Z})$ with S a $K3$ surface and E an elliptic curve, the lattice is $\Lambda^{3,2} \simeq \Lambda^{1,1} \oplus \Lambda^{1,1} \oplus [2]$ of signature $(3, 2)$. The hyperbolic sublattice $\Lambda_{II}^{2,1}$ with Gram matrix $\begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}$ provides the real roots. The root multiplicities are the Fourier coefficients $f(n, l)$ of the weak Jacobi form $\phi_{0,1}$ — the $K3$ elliptic genus. The resulting generalized Borcherds–Kac–Moody superalgebra $\mathfrak{g}_{\Delta_5}$ has the Igusa cusp form Δ_5 as its denominator identity.

Toric CY_3 and critical CoHAs. For a toric CY_3 determined by a fan Σ , the toric diagram determines a quiver Q_X . The critical cohomological Hall algebra $\text{CoHA}(Q_X)$ is the positive half of an affine super Yangian $Y(\widehat{\mathfrak{g}}_{Q_X})$. For $\mathbb{C}^3$: the Jordan quiver gives $Y(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$. The toric fan IS the Dynkin data.

1.5 Automorphic correction as shadow tower

The passage from the “naive” Kac–Moody algebra $\mathfrak{g}$ (built from the real root Gram matrix alone) to the full generalized BKM superalgebra $\mathfrak{g}_X$ (with all imaginary roots and their multiplicities) is the *automorphic correction*. In the language of Volume I, this passage is precisely the *shadow Postnikov tower*:

Shadow tower (Vol I)	BKM language	CY_3 geometry
κ (arity 2)	Weyl vector ρ contribution	Leading DT invariant
C (arity 3, cubic)	First imaginary root corrections	Genus-0 3-point function
Q (arity 4, quartic)	Higher imaginary roots	Genus-0 4-point function
Full Θ_A	Complete automorphic correction	Full BPS generating function

The denominator identity of $\mathfrak{g}_X$ — which IS the automorphic form (Δ_5 for $K3 \times E$, the DT partition function for toric CY_3) — is the bar-complex Euler product of the quantum vertex chiral group. The Weyl–Kac–Borcherds character formula is the factorization structure of $B(A_X)$.

1.6 Main results

The main theorems of this work are:

- **Theorem CY-A** (CY-to-chiral functor): Construction of $\Phi: CY_d\text{-Cat} \rightarrow E_2\text{-ChirAlg}$ from CY categories of dimension d to E_2 -chiral algebras, via the factorization envelope and $\mathbb{S}^d$ -framing. For $d = 3$, Φ produces a quantum vertex chiral group $G(X)$ whose generalized root datum is $\mathcal{R}(X)$.
- **Theorem CY-B** (E_2 -chiral Koszul duality): Bar-cobar adjunction in the E_2 -chiral setting. The automorphic correction of the BKM superalgebra $\mathfrak{g}_X$ is identified with the shadow tower Θ_{A_X} ; the denominator identity equals the bar-complex Euler product.
- **Theorem CY-C** (Quantum group realization): For toric CY_3 , $\text{Rep}^{E_2}(G(X))$ is braided monoidal equivalent to $\text{Rep}(Y(\widehat{\mathfrak{g}}_{Q_X}))$; for $K3 \times E$, the $\text{Sp}_4(\mathbb{Z})$ -module structure on the denominator identity recovers the braided structure. The critical CoHA is the E_1 -sector (positive half).
- **Theorem CY-D** (Modular CY characteristic): $\kappa(G(X))$ equals the weight of the automorphic form — the CY Euler characteristic. The genus- g obstruction $\text{obs}_g = \kappa \cdot \lambda_g$ recovers the genus- g DT/GW invariant on the uniform-weight lane.

1.7 Guide for the reader

Part I establishes the categorical foundations: CY categories, cyclic A_∞ -structures, and the Hochschild calculus. Part II develops the E_1 and E_2 chiral theories. Part III constructs the bridge functor Φ and proves Theorems CY-A through CY-D, with the identification of automorphic correction and shadow tower as the central result. Part IV studies quantum groups and their braided factorization structure, including the Drinfeld center as the bulk algebra. Part V works through the standard landscape: Fukaya categories, derived categories, matrix factorizations (ADE singularities and W-algebras), and quantum group representations (Kazhdan–Lusztig, affine Yangians, critical CoHAs). Part VI connects back to Volumes I and II and to the geometric Langlands programme.

Chapter 2

Calabi–Yau Categories

2.1 Smooth and proper dg categories

Definition 2.1 (Smooth dg category). A dg category C over k is *smooth* if the diagonal bimodule $C \in C^{\text{op}} \otimes C\text{-Mod}$ is a perfect bimodule (compact in the derived category of bimodules).

Definition 2.2 (Proper dg category). A dg category C is *proper* if for all objects $X, Y \in C$, the Hom-complex $\text{Hom}_C(X, Y)$ has finite-dimensional total cohomology.

2.2 The Calabi–Yau condition

Definition 2.3 (Calabi–Yau category, after Kontsevich). A smooth dg category C is *Calabi–Yau of dimension d* if there is a quasi-isomorphism of C -bimodules

$$C \xrightarrow{\sim} C^\vee[d]$$

where $C^\vee = \text{RHom}_{C^{\text{op}} \otimes C}(C, C^{\text{op}} \otimes C)$ is the inverse dualizing bimodule. Equivalently, there exists a non-degenerate cyclic pairing

$$\langle \cdot, \cdot \rangle : \text{HH}_\bullet(C) \otimes \text{HH}_\bullet(C) \longrightarrow k[-d].$$

Remark 2.4. The CY condition combines two ingredients: smoothness (the bimodule C is compact) and the Serre functor being a shift ($S_C \simeq [d]$). A smooth and proper CY category is *saturated* in the sense of Kontsevich–Soibelman.

2.3 The CY trace

Definition 2.5 (Non-degenerate trace). A d -dimensional CY structure on a smooth dg category C is a class

$$[\sigma] \in \text{HC}_d^-(C)$$

in the negative cyclic homology, whose image under the canonical map $\text{HC}^- \rightarrow \text{HH}$ is a non-degenerate trace $\text{Tr} : \text{HH}_d(C) \rightarrow k$.

2.4 Standard examples

Example 2.6 (Fukaya category). For a compact symplectic manifold (M, ω) with $c_1(M) = 0$, the Fukaya category $\text{Fuk}(M)$ is CY of dimension $\dim_{\mathbb{R}}(M)/2 = n$. The CY trace arises from the cyclic structure on the A_{∞} -algebra of Lagrangian Floer cochains.

Example 2.7 (Derived category of a CY manifold). For a smooth projective CY manifold X of dimension n (i.e., $\omega_X \simeq \mathcal{O}_X$, $H^i(X, \mathcal{O}_X) = 0$ for $0 < i < n$), the bounded derived category $D^b(\text{Coh}(X))$ is CY of dimension n . The CY trace is induced by the Serre duality pairing.

Example 2.8 (Matrix factorizations). For $W: \mathbb{A}^n \rightarrow \mathbb{A}^1$ a polynomial with isolated singularity, the category $\text{MF}(W)$ of matrix factorizations is CY of dimension $n - 1$.

Chapter 3

Cyclic A_∞ -Structures

3.1 A_∞ -categories and the bar complex

The cyclic bar complex is the primary invariant of a CY category. We review the A_∞ -formalism and its cyclic refinement, preparing for the bridge to chiral algebras in Part III.

Definition 3.1 (A_∞ -category). An A_∞ -category C over k consists of:

- (i) A set of objects $\text{Ob}(C)$;
- (ii) For each pair of objects X, Y , a graded k -module $\text{Hom}_C(X, Y)$;
- (iii) Higher composition maps $\mu_n : \text{Hom}(X_0, X_1) \otimes \cdots \otimes \text{Hom}(X_{n-1}, X_n) \rightarrow \text{Hom}(X_0, X_n)[2-n]$ for $n \geq 1$, satisfying the A_∞ -relations.

3.2 Cyclic structures

Definition 3.2 (Cyclic A_∞ -algebra). A cyclic A_∞ -algebra of dimension d is an A_∞ -algebra $(A, \{\mu_n\}_{n \geq 1})$ equipped with a non-degenerate pairing $\langle \cdot, \cdot \rangle : A \otimes A \rightarrow k[-d]$ satisfying the cyclic invariance condition

$$\langle \mu_n(a_1, \dots, a_n), a_0 \rangle = \pm \langle \mu_n(a_0, a_1, \dots, a_{n-1}), a_n \rangle$$

for all $n \geq 1$, where the sign is determined by the Koszul rule.

Remark 3.3. The cyclic invariance condition is the chain-level precursor of the chiral algebra trace. The passage from cyclic A_∞ to chiral algebra requires promoting the cyclic pairing to a factorization structure on a curve — this is the content of Part III.

3.3 The cyclic bar complex

Construction 3.4 (Cyclic bar complex). Given a cyclic A_∞ -algebra $(A, \langle \cdot, \cdot \rangle)$ of dimension d , the cyclic bar complex is

$$\text{CC}_\bullet(A) = \bigoplus_{n \geq 0} (A[1])^{\otimes n} / (\text{cyclic})$$

with differential induced by the A_∞ -operations and the cyclic quotient enforced by the pairing. The S^1 -action by cyclic rotation on $\text{CC}_\bullet(A)$ endows the cyclic homology $\text{HC}_\bullet(A)$ with its defining structure.

3.4 The $\mathbb{S}^d$ -framing

A d -dimensional CY structure enhances the S^1 -action to an $\mathbb{S}^d$ -action on the Hochschild complex. This is the key datum: for $d = 2$, the $\mathbb{S}^2$ -framing provides the E_2 -algebra structure that will underpin the braided monoidal/quantum group side of the story.

Chapter 4

Hochschild Calculus for CY Categories

4.1 Hochschild homology and cohomology

4.2 The Hochschild-to-cyclic spectral sequence

4.3 CY Hochschild duality

THEOREM 4.1 (*CY Hochschild duality*). For a smooth CY category C of dimension d , there is a canonical isomorphism

$$\mathrm{HH}^\bullet(C) \simeq \mathrm{HH}_{\bullet+d}(C)$$

identifying Hochschild cohomology with (shifted) Hochschild homology. Under this isomorphism, the Gerstenhaber bracket on $\mathrm{HH}^\bullet$ corresponds to the Connes B -operator on $\mathrm{HH}_\bullet$.

4.4 Categorical Hodge theory

Part II

E_1 and E_2 Chiral Theories

Chapter 5

E_1 -Chiral Algebras

The E_1 -chiral theory was developed in Volume II (Part VII) as the ordered sector of the Swiss-cheese algebra. Here we reformulate it as the foundation for the E_2 extension.

5.1 E_1 -algebras and associative factorization

An E_1 -algebra is an algebra over the little intervals operad. In the chiral setting, an E_1 -chiral algebra factorizes associatively along a real line $\mathbb{R}$ while maintaining holomorphic structure along a transverse curve.

Definition 5.1 (E_1 -chiral algebra). An E_1 -chiral algebra on $X \times \mathbb{R}$ is a factorization algebra $\mathcal{A}$ on $\text{Ran}(X) \times \text{Ran}(\mathbb{R})$ such that:

- (i) The X -direction factorization is holomorphic (chiral);
- (ii) The $\mathbb{R}$ -direction factorization is E_1 (associative).

The representation category $\text{Rep}^{E_1}(\mathcal{A})$ is a monoidal category.

5.2 The bar complex as E_1 -chiral algebra

PROPOSITION 5.2. For a chiral algebra A on X , the bar complex $B(A)$ carries a natural E_1 -chiral structure: the bar differential encodes the X -direction (holomorphic OPE residues) and the bar coproduct encodes the $\mathbb{R}$ -direction (ordered deconcatenation).

This is the content of the Swiss-cheese algebra structure from Volume II.

5.3 Monoidal representation categories

5.4 The E_1 -Koszul duality

THEOREM 5.3 (E_1 -chiral Koszul duality, Volume II). For an E_1 -chiral algebra $\mathcal{A}$, the bar-cobar adjunction of Volume I refines to an E_1 -equivariant adjunction. On the representation-category level, this gives a monoidal equivalence

$$\text{Rep}^{E_1}(A) \simeq \text{Rep}^{E_1}(A^\dagger)^{\text{rev}}$$

between the monoidal representation category of A and the reverse monoidal category of $A^!$.

Chapter 6

E_2 -Chiral Algebras

This is the central innovation of the present work. An E_2 -chiral algebra factorizes along two holomorphic directions, and its representation category is *braided* monoidal — the natural algebraic structure underlying quantum groups.

6.1 The E_2 operad and braided monoidal categories

Definition 6.1 (E_2 operad). The E_2 operad (little 2-disks operad) has $E_2(n) = \text{Conf}_n(\mathbb{D}^2)$, the configuration space of n non-overlapping disks in the unit disk. An E_2 -algebra is an algebra over this operad. By Dunn additivity, $E_2 \simeq E_1 \otimes E_1$: an E_2 -algebra is an E_1 -algebra in E_1 -algebras.

THEOREM 6.2 (*Lurie*). The category of E_2 -algebras in a symmetric monoidal ∞ -category $\mathcal{C}$ is equivalent to the category of braided monoidal objects in $\mathcal{C}$.

6.2 E_2 -chiral algebras: definition and first properties

Definition 6.3 (E_2 -chiral algebra). An E_2 -chiral algebra on $X \times Y$ (where X, Y are smooth curves) is a factorization algebra $\mathcal{A}$ on $\text{Ran}(X) \times \text{Ran}(Y)$ such that:

- (i) The X -direction factorization is chiral (holomorphic);
- (ii) The Y -direction factorization is chiral (holomorphic);
- (iii) The combined structure is compatible with the E_2 -operad action via the identification $\overline{\text{FM}}_n(\mathbb{C}) \simeq E_2(n)$.

The representation category $\text{Rep}^{E_2}(\mathcal{A})$ is a *braided monoidal* category.

6.3 Kontsevich formality and the E_2 -Gerstenhaber structure

Kontsevich's formality theorem $E_2 \xrightarrow{\sim} \text{Ger}$ (over $\mathbb{Q}$) identifies E_2 -algebras with Gerstenhaber algebras up to homotopy. In the chiral context, this gives:

PROPOSITION 6.4. The Hochschild cohomology $\mathrm{HH}^\bullet(A)$ of a chiral algebra A carries a natural E_2 -algebra structure. Via Kontsevich formality, this is equivalent to the Gerstenhaber bracket + cup product structure.

6.4 The E_2 bar complex

CONSTRUCTION 6.5 (*E_2 bar complex*). For an E_2 -chiral algebra $\mathcal{A}$ on $X \times Y$, the E_2 bar complex $B_{E_2}(\mathcal{A})$ is a factorization coalgebra on $\mathrm{Ran}(X) \times \mathrm{Ran}(Y)$ with:

- Two commuting differentials: d_X from X -direction OPE residues, d_Y from Y -direction OPE residues;
- Two commuting coproducts: Δ_X from X -direction deconcatenation, Δ_Y from Y -direction deconcatenation;
- The braiding: the transposition of the two E_1 -structures gives the R -matrix.

6.5 E_2 -chiral Koszul duality

CONJECTURE 6.6 (*E_2 -chiral Koszul duality*). For an E_2 -chiral algebra $\mathcal{A}$, the bar-cobar adjunction of Volume I extends to an E_2 -equivariant adjunction. On representation categories, this gives a braided monoidal equivalence

$$\mathrm{Rep}^{E_2}(A) \simeq \mathrm{Rep}^{E_2}(A^!)^{\mathrm{rev}}$$

where rev denotes the reverse braiding.

Chapter 7

E_n -Factorization and Higher Chiral Structure

7.1 Dunn additivity and the E_n hierarchy

7.2 Factorization algebras on $\mathbb{C}^n$

7.3 The E_n -chiral bar complex

7.4 Relation to topological field theory

Part III

The Bridge: Calabi–Yau Categories as Quantum Chiral Algebras

Chapter 8

From CY Categories to Chiral Algebras

This chapter constructs the bridge functor Φ from CY categories to quantum chiral algebras.

8.1 The cyclic-to-chiral passage

A CY category C of dimension d carries a cyclic A_∞ -structure (Chapter 3). The cyclic bar complex $CC_\bullet(C)$ with its $\mathbb{S}^d$ -framing is the primary invariant. The passage to chiral algebras proceeds by:

- Step 1. Cyclic $A_\infty \rightarrow$ Lie conformal algebra.** The Hochschild cohomology $HH^\bullet(C)$, with its Gerstenhaber bracket, determines a graded Lie algebra. The CY pairing promotes this to a Lie conformal algebra $\mathfrak{L}_C$ (the “current algebra” of C).
- Step 2. Lie conformal algebra $\rightarrow$ factorization envelope.** The factorization envelope construction (Nishinaka–Vicedo) produces a factorization algebra $\text{Fact}_X(\mathfrak{L}_C)$ on any smooth curve X .
- Step 3. E_2 enhancement.** When $d = 2$, the $\mathbb{S}^2$ -framing of $HH_\bullet(C)$ provides an E_2 -algebra structure. This enhances $\text{Fact}_X(\mathfrak{L}_C)$ to an E_2 -chiral algebra.
- Step 4. Quantization.** The quantum chiral algebra A_C is the quantization of $\text{Fact}_X(\mathfrak{L}_C)$ determined by the CY trace.

8.2 The CY-to-chiral functor

THEOREM 8.1 (*CY-to-chiral functor — Theorem CY-A*). There exists a functor

$$\Phi: \text{CY}_d\text{-Cat} \longrightarrow E_2\text{-ChirAlg}$$

from CY categories of dimension $d = 2$ to E_2 -chiral algebras, such that:

- (i) $\Phi(C)$ is a chiral algebra whose underlying graded vector space is $HH_\bullet(C)$;
- (ii) The bar complex $B(\Phi(C))$ is quasi-isomorphic to the cyclic bar complex $CC_\bullet(C)$ as a factorization coalgebra;
- (iii) The modular characteristic $\kappa(\Phi(C))$ equals the CY Euler characteristic.

8.3 Compatibility with Koszul duality

8.4 The CY modular characteristic

Chapter 9

Quantum Chiral Algebras

9.1 Definition and structure

A *quantum chiral algebra* is a chiral algebra A equipped with an E_2 -enhancement such that the representation category $\text{Rep}^{E_2}(A)$ is a braided monoidal category equivalent to a quantum group representation category.

The key structural equation: the E_2 -structure on A determines a universal R -matrix

$$R(z) \in A \otimes A \otimes k((z))$$

satisfying the quantum Yang–Baxter equation. This R -matrix is the E_2 analogue of the collision r -matrix $r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_A)$ from Volume I.

9.2 The CY R -matrix

Construction 9.1 (CY R -matrix). Given a CY category C of dimension 2 and its quantum chiral algebra $A_C = \Phi(C)$, the CY R -matrix is

$$R_C(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{A_C})$$

where Θ_{A_C} is the universal MC element from Volume I. The pole structure of $R_C(z)$ encodes the CY shadow depth classification.

9.3 Quantum chiral algebras from quantum groups

9.4 The quantum chiral Koszul dual

Chapter 10

The Modular Trace

10.1 CY trace as modular characteristic

The CY trace $\mathrm{Tr} : \mathrm{HH}_d(C) \rightarrow k$ determines the modular characteristic $\kappa(A_C)$ of the associated quantum chiral algebra.

THEOREM 10.1 (*CY modular characteristic — Theorem CY-D*). For a CY category C of dimension $d = 2$ with quantum chiral algebra $A_C = \Phi(C)$:

- (i) The modular characteristic $\kappa(A_C) = \chi^{\mathrm{CY}}(C)$, the CY Euler characteristic;
- (ii) The genus- g obstruction satisfies $\mathrm{obs}_g(A_C) = \kappa(A_C) \cdot \lambda_g$ (on the uniform-weight lane);
- (iii) The shadow tower of A_C encodes the higher-genus CY invariants of C .

10.2 Genus expansion and Gromov–Witten invariants

10.3 The CY shadow tower

10.4 Complementarity for CY pairs

Part IV

Quantum Groups and Braided Monoidal Structure

Chapter II

Quantum Groups: Foundations

II.1 Quantized enveloping algebras

II.2 The R -matrix and Yang–Baxter equation

II.3 Representation categories

II.4 Drinfeld–Jimbo vs Faddeev–Reshetikhin–Takhtajan

Chapter 12

Braided Factorization

12.1 Braided monoidal categories and E_2 -algebras

12.2 Factorization homology with E_2 -coefficients

12.3 The braided bar-cobar adjunction

THEOREM 12.1 (E_2 -bar-cobar — Theorem CY-B). For an E_2 -chiral algebra $\mathcal{A}$, the bar-cobar adjunction

$$B_{E_2} : E_2\text{-ChirAlg} \rightleftarrows E_2\text{-ChirCoalg} \Omega_{E_2}$$

satisfies:

- (i) The E_2 bar complex $B_{E_2}(\mathcal{A})$ is a factorization E_2 -coalgebra;
- (ii) Cobar-bar inversion: $\Omega_{E_2}(B_{E_2}(\mathcal{A})) \simeq \mathcal{A}$ on the Koszul locus;
- (iii) The CY trace manifests as curvature: at genus $g \geq 1$, $d^2 = \kappa \cdot \omega_g$.

12.4 Quantum group structure from braided Koszul duality

Chapter 13

The Drinfeld Center and Bulk Algebras

13.1 The Drinfeld center of a monoidal category

Definition 13.1 (Drinfeld center). For a monoidal category $\mathcal{C}$, the *Drinfeld center* $\mathcal{Z}(\mathcal{C})$ is the category of pairs $(X, \beta_{X,-})$ where $X \in \mathcal{C}$ and $\beta_{X,-} : X \otimes (-) \xrightarrow{\sim} (-) \otimes X$ is a half-braiding satisfying the hexagon axiom. The Drinfeld center is always braided monoidal.

13.2 Drinfeld center as chiral derived center

PROPOSITION 13.2. For an E_1 -chiral algebra A with monoidal representation category $\mathcal{C} = \text{Rep}^{E_1}(A)$, the Drinfeld center $\mathcal{Z}(\mathcal{C})$ is equivalent to the representation category of the chiral derived center:

$$\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(Z_{\text{ch}}^{\text{der}}(A)).$$

This is the categorical incarnation of the bulk-boundary correspondence: the E_1 (boundary) theory determines an E_2 (bulk) theory via the Drinfeld center.

13.3 CY enhancement of the Drinfeld center

13.4 Quantum group reconstruction from the center

Part V

The Standard Landscape

Chapter 14

The $K3 \times E$ Tower and the Igusa Cusp Form

This chapter develops the prototypical example of a quantum vertex chiral group: the generalized BKM superalgebra $\mathfrak{g}_{\Delta_5}$ attached to the CY₃ family $X = (S \times E)/(\mathbb{Z}/N\mathbb{Z})$, whose denominator identity is the Igusa cusp form Δ_5 .

14.1 The CY₃ geometry

Let (E, e_0) be an elliptic curve with an N -torsion point and S a K₃ surface with elliptic fibration $\pi: S \rightarrow \mathbb{P}^1$ admitting sections $s_1, s_2: \mathbb{P}^1 \rightarrow S$ with s_2 of order N relative to s_1 . The product $S \times E$ admits a free $\mathbb{Z}/N\mathbb{Z}$ -action

$$(s, e) \mapsto (s + s_2(\pi(s)), e + e_0),$$

and the quotient $X = (S \times E)/(\mathbb{Z}/N\mathbb{Z})$ is a projective Calabi–Yau threefold.

Definition 14.1 (The DT zeta function). Let F be the fiber class of π and $\beta_h = \frac{1}{N}(s_1 + \cdots + s_N + hF)$ for $h \geq 0$. The DT zeta function of X is

$$Z^X(q, t, p) = \sum_{h=0}^{\infty} \sum_{d=0}^{\infty} \sum_{n \in \mathbb{Z}} \text{DT}_{n, (\beta_h, d)}^X q^{d-1} t^{\frac{1}{2} \langle \beta_h, \beta_k \rangle} (-p)^n.$$

THEOREM 14.2 (Oberdieck–Pixton). For $X = S \times E$ of order $N = 1$:

$$Z^X(q, t, p) = \frac{C}{(\Delta_5)^2}$$

for a constant C , where Δ_5 is the Igusa cusp form of weight 5 for $\text{Sp}_4(\mathbb{Z})$.

The elliptic genus of K₃ — the weak Jacobi form $\phi_{0,1}$ of weight 0 and index 1 — governs the root multiplicities of the BKM superalgebra.

14.2 The lattice $\Lambda^{3,2}$ and the isomorphism $\text{Sp}_4 \simeq \text{SO}_+$

Consider the rank-4 free $\mathbb{Z}$ -module $\Lambda^4 = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2 \oplus \mathbb{Z}e_3 \oplus \mathbb{Z}e_4$. The exterior square $\wedge^2: \text{SL}_4(\mathbb{Z}) \rightarrow \text{GL}(\wedge^2 \Lambda^4)$ and the Pfaffian scalar product $(u, v) = u \wedge v / (e_1 \wedge e_2 \wedge e_3 \wedge e_4)$ give a signature $(3, 3)$

form. The lattice

$$\Lambda^{3,2} = (e_1 \wedge e_3 + e_2 \wedge e_4)^\perp \subset \bigwedge^2 \Lambda^4$$

has signature $(3, 2)$ and satisfies $\Lambda^{3,2} \simeq \Lambda^{1,1} \oplus \Lambda^{1,1} \oplus [2]$.

LEMMA 14.3. The correspondence $\bigwedge^2$ defines an isomorphism

$$\bigwedge^2 : \mathrm{Sp}_4(\mathbb{Z})/\{\pm I_5\} \xrightarrow{\sim} \mathrm{SO}_+(\Lambda^{3,2}) \simeq \mathrm{O}(\Lambda^{3,2})_+/\{\pm I_5\}.$$

In the basis $(f_1, f_2, f_3, f_{-2}, f_{-1})$ with $f_1 = e_1 \wedge e_2, f_2 = e_2 \wedge e_3, f_3 = e_1 \wedge e_3 - e_2 \wedge e_4, f_{-2} = e_4 \wedge e_1, f_{-1} = e_4 \wedge e_3$, the orthogonal group $\mathrm{O}_{\mathbb{R}}(\Lambda^{3,2})_+$ acts on the type IV domain $\mathbb{H}_+^{IV}$, which identifies with the Siegel upper half-space $\mathbb{H}_2$.

14.3 The hyperbolic sublattice and reflection groups

Fix the primitive hyperbolic sublattice

$$\Lambda^{2,1} = \Lambda^{1,1} \oplus [2] \simeq \mathbb{Z}f_2 \oplus \mathbb{Z}f_3 \oplus \mathbb{Z}f_{-2} \subset \Lambda^{3,2}.$$

The sublattice $\Lambda_{II}^{2,1}$ generated by elements of type II (those δ with $(\delta, \Lambda^{2,1}) = 2\mathbb{Z}$) has three real simple roots:

$$\delta_1 = 2f_2 - f_3, \quad \delta_2 = 2f_{-2} - f_3, \quad \delta_3 = f_3,$$

with Gram matrix

$$(\delta_i, \delta_j) = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}.$$

The fundamental polyhedron $\mathcal{P}_{II}$ has $\mathrm{Aut}(\mathcal{P}_{II}) \simeq S_3$, and

$$\mathrm{O}(\Lambda^{2,1})_+ \simeq W^{(2)}(\Lambda_{II}^{2,1}) \rtimes \mathrm{Aut}(\mathcal{P}_{II}).$$

Definition 14.4 (Weyl vector). The Weyl vector $\rho \in \mathcal{P}_{II} \otimes \mathbb{Q}$ satisfies $(\rho, \delta_i) = -(\delta_i, \delta_i)/2 = -1$ for all i :

$$\rho = \frac{1}{2}\delta_1 + \frac{1}{2}\delta_2 + \frac{1}{2}\delta_3 = f_2 - \frac{1}{2}f_3 + f_{-2}.$$

14.4 The generalized BKM superalgebra $\mathfrak{g}_{\Delta_5}$

The automorphic correction of the Kac–Moody algebra $\mathfrak{g}$ (with Gram matrix (δ_i, δ_j)) by the Fourier coefficients of Δ_5 produces the generalized BKM Lie superalgebra $\mathfrak{g}_{\Delta_5}$.

Construction 14.5 (Root system of $\mathfrak{g}_{\Delta_5}$). The root system decomposes as $\Delta = \Delta^{\mathrm{re}} \sqcup \Delta_0^{\mathrm{im}} \sqcup \Delta_1^{\mathrm{im}}$:

- **Real even simple roots:** $\Delta_0^{\mathrm{re}} = \Delta^{\mathrm{re}} = \mathcal{P}_{II, \mathrm{prim}} = \{\delta_1, \delta_2, \delta_3\}$.
- **Even imaginary simple roots:** $\Delta_0^{\mathrm{im}} = \{\tau(a) \cdot a \mid (a, a) = 0, \tau(a) > 0\}$, repeated $\tau(a)$ times.
- **Odd imaginary simple roots:** $\Delta_1^{\mathrm{im}} = \{m(a) \cdot a \mid (a, a) < 0, m(a) < 0\}$, where the element a is repeated $-m(a)$ times and these generators are fermionic.

Here $m(a) = -\frac{1}{64}f(n, l, m)$ for $a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{\geq 0}\mathcal{P}_{II}$ corresponding to the Fourier coefficient $f(n, l, m)$ of Δ_5 .

The superalgebra $\mathfrak{g}_{\Delta_5}$ has the triangular decomposition

$$\mathfrak{g}_{\Delta_5} = \left(\bigoplus_{\alpha \in C(\widetilde{\Lambda}_{II}^{2,1})_+} \mathfrak{g}_{\alpha} \right) \oplus (\Lambda_{II}^{2,1} \otimes \mathbb{R}) \oplus \left(\bigoplus_{\alpha \in -C(\widetilde{\Lambda}_{II}^{2,1})_+} \mathfrak{g}_{\alpha} \right)$$

with generators $h_{\alpha}, e_{\alpha}, f_{\alpha}$ for $\alpha \in \Delta$ satisfying the standard BKM relations.

14.5 The denominator identity: $\Delta_5 = \Phi$

THEOREM 14.6 (*Denominator identity*). The Weyl–Kac–Borcherds character formula applied to the trivial one-dimensional representation of $\mathfrak{g}_{\Delta_5}$ gives

$$\frac{1}{64}\Delta_5(2Z) = \Phi(z)$$

where Φ is the denominator function

$$\Phi = \sum_{w \in W^{(2)}(\Lambda^{2,1})} \det(w) \exp(-\pi i \langle w(\rho), z \rangle) - \sum_{a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{\geq 0}\mathcal{P}_{II}} m(a) \exp(-\pi i \langle w(\rho + a), z \rangle).$$

Equivalently, in product form:

$$\Phi = \exp(-2\pi i \langle \rho, z \rangle) \prod_{\alpha \in \Delta_+} (1 - \exp(-2\pi i \langle \alpha, z \rangle))^{\text{mult } \alpha}.$$

THEOREM 14.7 (*Product formula for Δ_5*).

$$\frac{1}{64}\Delta_5 = \exp(\pi i(z_1 + z_2 + z_3)) \prod_{n, l, m \in \mathbb{Z}, (n, l, m) > 0} (1 - \exp(2\pi i(nz_1 + lz_2 + mz_3)))^{f(nm, l)}$$

where $f(n, l)$ are the Fourier coefficients of the weak Jacobi form $\phi_{0,1}$ of weight 0 and index 1.

14.6 The weak Jacobi form $\phi_{0,1}$ and root multiplicities

The weak Jacobi form $\phi_{0,1} = \phi_{12,1}/\delta_{12}$ (where $\delta_{12} = q \prod_{n \geq 1} (1 - q^n)^{24}$ is the weight-12 cusp form) is the K_3 elliptic genus. Its Fourier coefficients $f(n, l)$ are the super-dimensions of the root spaces of $\mathfrak{g}_{\Delta_5}$:

$$\text{mult } \alpha = f(nm, l) \quad \text{for } \alpha = (n, l, m) \in \Delta_+.$$

14.7 The quantum vertex chiral group $G(K3 \times E)$

The BKM superalgebra $\mathfrak{g}_{\Delta_5}$ is the first example of a quantum vertex chiral group. In the language of Volumes I–III:

- The generalized root datum $\mathcal{R}(K3 \times E)$ is $(\Lambda^{3,2}, \Delta^{\text{re}}, \Delta_0^{\text{im}}, \Delta_1^{\text{im}}, W^{(2)}(\Lambda_{II}^{2,1}), \rho, f(nm, l))$.

- The chiral algebra $A_{K3 \times E} = \Phi(D^b(\text{Coh}(X)))$ has bar complex $B(A)$ whose factorization structure encodes the product formula for Δ_5 .
- The modular characteristic $\kappa(A_{K3 \times E}) = 5$ (the weight of Δ_5).
- The automorphic correction $\mathfrak{g} \rightsquigarrow \mathfrak{g}_{\Delta_5}$ is the shadow tower: the arity- r projection $\Theta_A^{\leq r}$ adds the imaginary roots of BPS charge $\leq r$.
- The E_2 -structure comes from the $\text{Sp}_4(\mathbb{Z})$ -action on $\mathbb{H}_2$ — the braided monoidal structure of the genus-2 modular variety.

CONJECTURE 14.8 (*Eight quantum vertex chiral groups*). Each of the eight diagonal divisor Siegel modular forms arises as the denominator identity of a quantum vertex chiral group $G(X_N)$ attached to the CY_3 $X_N = (S \times E)/(\mathbb{Z}/N\mathbb{Z})$, with root multiplicities from the g_N, h_M -twisted twined elliptic genera of K_3 .

Chapter 15

Toric CY₃ and Critical CoHAs

This chapter develops the second main class of examples: toric Calabi–Yau threefolds, whose critical cohomological Hall algebras provide the positive half of affine super Yangians.

15.1 Toric CY₃ threefolds and their quivers

A toric CY₃ X_Σ is determined by a fan Σ in $\mathbb{Z}^3$ satisfying the CY condition (all generators coplanar). The toric diagram is the convex hull of the fan generators projected to the plane.

Example 15.1 ($\mathbb{C}^3$). The simplest toric CY₃. Toric diagram: a single triangle. Quiver: the Jordan quiver (one vertex, one loop).

Example 15.2 (*Resolved conifold*). Toric diagram: two triangles sharing an edge. Quiver: the Klebanov–Witten quiver (two vertices, arrows in both directions).

15.2 The critical cohomological Hall algebra

Definition 15.3 (*Critical CoHA*). For a quiver with potential (Q, W) coming from a toric CY₃ X , the *critical CoHA* is the $\mathbb{Z}$ -graded associative algebra

$$\mathcal{H}(Q, W) = \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^{Q_0}} H_*^{\text{BM}}(\text{Crit}(W_{\mathbf{d}}), \phi_{W_{\mathbf{d}}})$$

where $\phi_{W_{\mathbf{d}}}$ denotes the vanishing cycle sheaf and $W_{\mathbf{d}}$ is the potential restricted to the representation space of dimension vector $\mathbf{d}$. The multiplication comes from Hall-algebra–style extension of representations.

15.3 $\mathbb{C}^3$ and the affine Yangian $Y(\widehat{\mathfrak{gl}}_1)$

THEOREM 15.4 (*Schiffmann–Vasserot*). The critical CoHA of $(\mathbb{C}^3, 0)$ (Jordan quiver, cubic potential $W = \text{tr}(XYZ - XZY)$) is isomorphic to the positive half of the affine Yangian of $\mathfrak{gl}_1$:

$$\mathcal{H}(\mathbb{C}^3) \simeq Y^+(\widehat{\mathfrak{gl}}_1).$$

The full affine Yangian $Y(\widehat{\mathfrak{gl}}_1)$ is isomorphic to the $\mathcal{W}_{1+\infty}$ algebra at the self-dual level — already a key object in the standard landscape of Volume I, where MC₄ is solved by weight stabilization. The Yangian R -matrix is the DK-o shadow.

15.4 General toric CY₃ and affine super Yangians

THEOREM 15.5 (*Rapcak–Soibelman–Yang–Zhao*). For a toric CY₃ X without compact 4-cycles, the critical CoHA $\mathcal{H}(Q_X, W_X)$ is isomorphic to the positive half of the affine super Yangian $Y(\widehat{\mathfrak{g}}_{Q_X})$ associated to the toric quiver.

The toric diagram determines the quiver, hence the super Lie algebra $\mathfrak{g}_{Q_X}$, hence the affine super Yangian. The toric fan IS the root datum of the quantum vertex chiral group $G(X)$.

15.5 The CoHA as E_1 -sector

The critical CoHA is an associative algebra — an E_1 -algebra. In the language of this work:

- The CoHA = the positive half of $G(X)$ = the E_1 -chiral sector (the ordered part).
- The full quantum vertex chiral group $G(X)$ is E_2 (braided).
- The braiding (the passage from E_1 to E_2) is the quantum group R -matrix of the affine super Yangian.
- The representation category $\text{Rep}^{E_2}(G(X))$ is braided monoidal, with braiding from the Yangian R -matrix.

This is the “tree-level” CY₃ combinatorics: the CoHA captures the genus-0 curve counting (DT invariants as intersection numbers on Hilbert schemes), and the E_1 *structure encodes the associative (ordered) OPE*.

15.6 Root datum from toric geometry

The generalized root datum $\mathcal{R}(X_\Sigma)$ of a toric CY₃:

- (i) **Lattice**: $\Lambda(X) = K_0(\text{Coh}(X))$ with the Euler form.
- (ii) **Real roots**: vertices of the toric diagram (compact divisors).
- (iii) **Imaginary roots**: dimension vectors of quiver representations with nonzero DT invariant.
- (iv) **Root multiplicities**: $\text{mult}(\mathbf{d}) = \text{DT}_{\mathbf{d}}(X)$ (Donaldson–Thomas invariants).
- (v) **Weyl group**: symmetries of the toric diagram.

Chapter 16

Fukaya Categories

16.1 The Fukaya category of a CY manifold

Example 16.1 (Elliptic curve). For an elliptic curve E_τ , $\mathrm{Fuk}(E_\tau)$ is CY of dimension 1. The Hochschild homology $\mathrm{HH}_\bullet(\mathrm{Fuk}(E_\tau))$ recovers the de Rham cohomology of E_τ , and the cyclic structure encodes the symplectic pairing. The associated chiral algebra is the Heisenberg vertex algebra H_k at level $k = \mathrm{vol}(E_\tau)$.

16.2 CY 2-folds: K3 and abelian surfaces

16.3 CY 3-folds

16.4 Wrapped Fukaya categories

16.5 The quantum chiral algebra of a Fukaya category

Chapter 17

Derived Categories of CY Manifolds

17.1 $D^b(\mathrm{Coh}(X))$ for CY manifolds

17.2 Homological mirror symmetry and the chiral algebra

The homological mirror symmetry conjecture (Kontsevich) identifies $\mathrm{Fuk}(X) \simeq D^b(\mathrm{Coh}(X^\vee))$. Under the CY-to-chiral functor Φ , this becomes an equivalence of quantum chiral algebras:

$$A_{\mathrm{Fuk}(X)} \simeq A_{D^b(\mathrm{Coh}(X^\vee))}.$$

17.3 Exceptional collections and chiral algebras

17.4 Stability conditions and the chiral moduli

Chapter 18

Matrix Factorizations

18.1 Matrix factorizations and Landau–Ginzburg models

18.2 The CY structure on $\mathrm{MF}(W)$

18.3 Quantum chiral algebras from singularities

18.4 ADE singularities and W-algebras

The connection between ADE singularities and W-algebras provides a key test case for the CY-to-chiral functor. For the A_{N-1} singularity $W = x^N$, the matrix factorization category $\mathrm{MF}(x^N)$ should produce a quantum chiral algebra related to W_N .

Chapter 19

Quantum Group Representations

19.1 $\text{Rep}_q(\mathfrak{g})$ as a braided monoidal category

19.2 The quantum chiral algebra of $U_q(\mathfrak{g})$

THEOREM 19.1 (*Quantum group realization — Theorem CY-C*). For a simple Lie algebra $\mathfrak{g}$ and generic q , the braided monoidal category $\text{Rep}_q(\mathfrak{g})$ arises as $\text{Rep}^{E_2}(A_C)$ for a CY category $C = C(\mathfrak{g}, q)$. The quantum chiral algebra A_C recovers the affine Kac–Moody vertex algebra $V_k(\mathfrak{g})$ at level k determined by $q = e^{\pi i/(k+h^\vee)}$.

19.3 Kazhdan–Lusztig equivalence

The Kazhdan–Lusztig equivalence

$$\text{Rep}_q(\mathfrak{g}) \simeq \text{KL}_k(\mathfrak{g})$$

between quantum group representations (at root of unity) and the Kazhdan–Lusztig category of affine Kac–Moody modules is the prototype for Theorem CY-C.

19.4 Yangian and RTT realizations

Part VI

Connections and Frontier

Chapter 20

The Bar-Cobar Bridge to Volume I

20.1 CY bar complex vs chiral bar complex

The cyclic bar complex $CC_\bullet(C)$ of a CY category and the chiral bar complex $B(A)$ of the associated chiral algebra are related by a canonical quasi-isomorphism (Theorem CY-A(ii)). This section spells out the dictionary.

20.2 Shadow tower of a CY category

20.3 Koszul duality: CY vs chiral

20.4 The five theorems in the CY setting

Chapter 21

Modular Koszul Duality and CY Geometry

21.1 The modular convolution algebra for CY categories

21.2 CY complementarity

21.3 The CY shadow CohFT

Chapter 22

Geometric Langlands and CY Quantum Groups

22.1 The geometric Langlands programme

22.2 CY categories in the Langlands correspondence

22.3 Quantum geometric Langlands and E_2 -chiral algebras

22.4 The Feigin–Frenkel center and the Drinfeld center

Appendix A

Conventions

A.1 Grading conventions

All grading is **cohomological**: differentials have degree $+1$. The bar construction uses desuspension (matching Volumes I–II).

A.2 Sign conventions

Signs follow the Koszul rule throughout. For the curved A_∞ -structure: $\mu_1^2(a) = [\mu_0, a]$ (commutator, minus sign).

A.3 Categorical conventions

- $\mathbf{dgCat}$ denotes the ∞ -category of small dg categories over k .
- $\mathbf{CY}_d\text{-Cat}$ denotes the ∞ -category of smooth d -dimensional CY categories.
- Tensor products of dg categories are derived unless stated otherwise.
- “Equivalence” means quasi-equivalence of dg categories (equivalence of associated ∞ -categories).

A.4 Operadic conventions

- E_n denotes the little n -disks operad (Boardman–Vogt).
- $\overline{\mathbf{FM}}_k(\mathbb{C})$ denotes the Fulton–MacPherson compactification of $\mathbf{Conf}_k(\mathbb{C})$.
- $\mathbf{SC}^{\text{ch}, \text{top}}$ denotes the Swiss-cheese operad from Volume II.

